

RESEARCH ARTICLE

Ant–fungus species combinations engineer physiological activity of fungus gardens

J. N. Seal^{1,2,*}, M. Schiøtt³ and U. G. Mueller²**ABSTRACT**

Fungus-gardening insects are among the most complex organisms because of their extensive co-evolutionary histories with obligate fungal symbionts and other microbes. Some fungus-gardening insect lineages share fungal symbionts with other members of their lineage and thus exhibit diffuse co-evolutionary relationships, while others exhibit little or no symbiont sharing, resulting in host–fungus fidelity. The mechanisms that maintain this symbiont fidelity are currently unknown. Prior work suggested that derived leaf-cutting ants in the genus *Atta* interact synergistically with leaf-cutter fungi (*Attamyces*) by exhibiting higher fungal growth rates and enzymatic activities than when growing a fungus from the sister-clade to *Attamyces* (so-called ‘Trachymyces’), grown primarily by the non-leaf cutting *Trachymyrmex* ants that form, correspondingly, the sister-clade to leaf-cutting ants. To elucidate the enzymatic bases of host–fungus specialization in leaf-cutting ants, we conducted a reciprocal fungus-switch experiment between the ant *Atta texana* and the ant *Trachymyrmex arizonensis* and report measured enzymatic activities of switched and sham-switched fungus gardens to digest starch, pectin, xylan, cellulose and casein. Gardens exhibited higher amylase and pectinase activities when *A. texana* ants cultivated *Attamyces* compared with *Trachymyces* fungi, consistent with enzymatic specialization. In contrast, gardens showed comparable amylase and pectinase activities when *T. arizonensis* cultivated either fungal species. Although gardens of leaf-cutting ants are not known to be significant metabolizers of cellulose, *T. arizonensis* were able to maintain gardens with significant cellulase activity when growing either fungal species. In contrast to carbohydrate metabolism, protease activity was significantly higher in *Attamyces* than in *Trachymyces*, regardless of the ant host. Activity of some enzymes employed by this symbiosis therefore arises from complex interactions between the ant host and the fungal symbiont.

KEY WORDS: Attini, Cellulose, Co-evolution, Enzyme activity, Pectin, Starch, Symbiosis, Xylan

INTRODUCTION

Co-operation among unrelated organisms (mutualistic symbioses) has led to many key evolutionary innovations with far-reaching ecological consequences. One of the larger mysteries is understanding how such complex organisms maintain homeostasis and are able to function as distinct units (Douglas, 2010; Kültz et al., 2013; McFall-Ngai et al., 2013). Of the numerous examples of

such complexity, the fungus-gardening insects have evolved obligate macro-symbioses with specific clades of fungi, and use fungal symbionts essentially as an external digestive organ that allows the insect to thrive on otherwise non-digestible substrates, such as structural carbohydrates of plants (e.g. cellulose) (Aanen et al., 2002; Aylward et al., 2012a; Aylward et al., 2012b; Bacci et al., 1995; Farrell et al., 2001; De Fine Licht and Biedermann, 2012; Martin, 1987a; Mueller et al., 2005). One of the most striking attributes of these symbioses is the degree of physiological integration: the insect host functions as a distributor of fungal enzymes, which digest plant fibers external to the insect’s body (Aanen and Eggleton, 2005; Aylward et al., 2012b; De Fine Licht and Biedermann, 2012; De Fine Licht et al., 2013; Martin, 1987b; Schiøtt et al., 2010).

The fungus-gardening (attine) ants exhibit several macroevolutionary trends with regard to their farmed symbionts. Although vertical transmission of symbionts and generalized fidelity between clades of ants and clades of fungi is the norm, horizontal transmission is quite extensive, especially among the phylogenetically early branching lineages (Green et al., 2002; Kellner et al., 2013; Mehdiabadi et al., 2012; Mikheyev et al., 2010; Mueller et al., 2010; Mueller et al., 1998; Schultz and Brady, 2008). One of the more profound evolutionary and ecological transitions occurred in the derived lineages (the ‘higher Attini’), the most complex of which are found in the clade containing leaf-cutting ants (genera *Atta* and *Acromyrmex*). These ants exhibit large queens, pronounced variation in worker size and colony sizes of immense proportions (>1 million workers); as a result, fungus-growing ants may exert enormous ecological impacts (Hölldobler and Wilson, 2011; Meyer et al., 2011; Seal, 2009; Wilson, 1980; Wirth et al., 2003). Leaf-cutter ants generally cultivate a single species of fungus, called *Attamyces bromatificus* (hereafter, *Attamyces*) in its anamorph (vegetative, clonal) growth form and *Leucocoprinus gongylophorus* in its teleomorph (sexual) form (Mueller et al., 2010; Mueller et al., 2011b; Mueller et al., 1998). All leaf-cutting ants are thought to have descended from a common ancestor shared with several *Trachymyrmex* species in the *Trachymyrmex septentrionalis* group, of which many inhabit North America (Rabeling et al., 2007; Schultz and Brady, 2008). Ants in the genus *Trachymyrmex* are characterized by relatively small colonies (hundreds of workers), modest or no worker polymorphism, and occasional population sizes that may make them ecologically relevant (Beshers and Traniello, 1994; Leal and Oliveira, 1998; Seal and Tschinkel, 2006; Seal and Tschinkel, 2010; Torres et al., 1999). *Trachymyrmex* ants typically cultivate a more diverse assemblage of closely related *Leucocoprinus* lineages (Mikheyev et al., 2010; Mikheyev et al., 2008; Mueller et al., 1998; Schultz and Brady, 2008). None of these have been taxonomically resolved; hence we refer to them provisionally as ‘Trachymyces’.

The leaf-cutting ant fungi are thought to be specialized toward the rapid metabolism of starches, hemicelluloses and proteins found in

¹Department of Biology, University of Texas at Tyler, 3900 University Blvd, Tyler, TX 75799, USA. ²Integrative Biology, University of Texas at Austin, 1 University Station C0930, Austin, TX 78712, USA. ³Centre for Social Evolution, Department of Biology, University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen, Denmark.

*Author for correspondence (trachymyrmex@gmail.com)

Received 15 October 2013; Accepted 17 April 2014

leaves, whereas the non-leaf-cutting fungi have lower metabolic activities toward these substrates and may have a greater ability to digest cellulose (Bacci et al., 2013; D'Ettorre et al., 2002; Erthal et al., 2004; Erthal et al., 2009; De Fine Licht et al., 2013; De Fine Licht et al., 2010; Richard et al., 2005; Schiøtt et al., 2008; Schiøtt et al., 2010). These physiological differences seem to be linked with dietary preferences associated with the ants; leaf-cutter ants provide their garden with fresh leaves, whereas the non-leaf-cutters provide their fungus with various dried plant debris, flower parts and caterpillar excrement (Hölldobler and Wilson, 2011; Leal and Oliveira, 2000; Seal and Tschinkel, 2007b).

Although the two higher-attine ant lineages generally exhibit fidelity to their fungal lineages (i.e. leaf-cutter ants tend to cultivate *Attamyces*; *Trachymyrmex* ants tend to cultivate *Trachymyces*), several observations are inconsistent with a strict 1:1 co-evolution. Colonies can be experimentally switched to non-native fungi, and surveys indicate that switches can occur naturally on ecological and evolutionary scales (Mehdiabadi et al., 2012; Mikheyev et al., 2010; Mueller et al., 2011b; Mueller et al., 1998; Seal and Tschinkel, 2007a; Stradling and Powell, 1986; Weber, 1956) (supplementary material Fig. S1). The only replicated experiments that addressed whether higher-attine fungal lineages were adapted to their ant hosts suggested that diseases constrain symbiont switching in two species of *Trachymyrmex* ants (Seal and Mueller, 2014; Seal and Tschinkel, 2007a). One preliminary study suggested that some *Trachymyces* cultivars can lower the performance of leaf-cutting ant colonies, but this study suffered from a low sample size (Sánchez-Peña, 2005). These findings prompted our study of the enzymatic properties of gardens in fungus-switched colonies of *Atta* and *Trachymyrmex* ants. We investigated the potential for constraints in symbiont switching by testing three hypotheses, described below.

The 'host-independent hypothesis' postulates that ant species' identity is of lesser importance or not important in influencing fungal enzyme activity in gardens, and consequently predicts that *Attamyces* gardens will exhibit higher enzymatic activities regardless of the ant species growing them (and conversely, *Trachymyces* should always show lower enzyme activities regardless of the ant host growing them). Evidence for this hypothesis derives from an *in vitro* study (fungus grown on artificial media) (Stradling and Powell, 1986) and *in vivo* quantification of enzymatic activity of native ant–fungus associations (De Fine Licht et al., 2010) showing that *Attamyces* has indeed higher growth rates and enzymatic activities than non-leaf-cutting fungi. Additional support derives from non-replicated experiments where the addition of fresh *Attamyces* garden to a garden-deprived *Trachymyrmex urichi* colony resulted in increased ant colony growth (Stradling and Powell, 1986), as well as reduced colony and garden growth when *Atta* colonies were forced to cultivate *Trachymyces* gardens (Sánchez-Peña, 2005). A problem with this hypothesis is that by itself it does not explain why *Trachymyrmex* ants would persist in growing a physiologically inferior fungal symbiont (rather than switch to *Attamyces* strains cultivated by sympatric leaf-cutter ants); it only explains why *Atta* ants may avoid the apparently 'inferior' *Trachymyces*. Because *Attamyces* appears to have swept by lateral transfer through leaf-cutter ant populations (Mikheyev et al., 2010), a generally superior *Attamyces* could have spread likewise through *Trachymyrmex* populations.

The 'adaptive combination hypothesis' postulates that only the specific ant–fungus combinations observed in nature confer high fitness (i.e. *Atta* ants cultivating *Attamyces* fungus; *Trachymyrmex* ants cultivating *Trachymyces* fungi). Under this hypothesis, these

natural combinations exhibit higher fungal enzymatic activities than novel combinations (*Trachymyces* × *Atta*, or *Attamyces* × *Trachymyrmex*). While enzymatic activity may be typically higher in *Attamyces* than in *Trachymyces*, artificial experimentally induced combinations would result in lower activities. Evidence supporting this hypothesis derives from fungus-switch experiments where *T. septentrionalis* colonies growing *Attamyces* did not increase their reproductive output compared with colonies growing *Trachymyces* (Seal and Tschinkel, 2007a), which was possibly due to infections by pathogens (Seal and Mueller, 2014). These experiments did not examine the reverse switch (*Atta* colonies growing *Trachymyces*) (but see Sánchez-Peña, 2005) or investigate the effects of switches in other *Trachymyrmex* species.

A third possibility, termed the 'complex synergism hypothesis', is that enzymatic responses to ant–fungus combinations may depend on complex interactions between ants and symbiotic fungi. According to this hypothesis, the combinations that exhibit higher enzymatic activities do not reflect phylogeny, so higher activities may result when *Atta* ants are growing *Trachymyces* or *Trachymyrmex* ants are growing *Attamyces*.

We tested these three hypotheses by performing a 2×2 reciprocal symbiont-switch experiment with colonies of the leaf-cutting ant *Atta texana* Buckley 1860 and the non-leaf-cutting ant *Trachymyrmex arizonensis* (Wheeler 1907) that were growing either *Attamyces* or *Trachymyces*. The weight of the evidence supports the complex synergism hypothesis because enzymatic activity is influenced by ant–fungus interactions.

RESULTS

Enzymatic activity

The two species of fungi exhibited different catabolic properties toward carbohydrates and proteins when cultivated by the two species of ants. Amylase (starch hydrolysis) activity was significantly dependent on fungal species – *Attamyces* had much higher amylase activity than *Trachymyces* – but was not dependent on the species of ant host cultivating either fungus. There appeared to be a synergistic effect of fungal and ant species (significant interaction term), which was caused by the lower amylase activity of *Trachymyces* gardens when cultivated by *A. texana* compared with the amylase activity of the three other types of garden (*Attamyces* cultivated by *A. texana*, or either *Trachymyces* or *Attamyces* cultivated by *T. arizonensis* (Fig. 1, Table 1). Thus, amylase activity was significantly reduced when *Trachymyces* was grown by *A. texana*. *Trachymyrmex arizonensis* had similar amylase activity regardless of the type of fungus this ant species was growing.

Pectinase activity was significantly influenced by ant and fungal species and also exhibited significant synergistic effects (Table 1). Pectinase activity was highest in the *A. texana* × *Attamyces* combination where it was ~30% higher than in all other groups (Fig. 2, Table 1). *Trachymyrmex arizonensis* colonies had comparable pectinase activity regardless of fungal species grown, not unlike the pattern of amylase activity (Fig. 1). In contrast to the patterns for amylase and pectinase, cellulase activity of the fungus garden appeared to be influenced by the ant species growing it. Cellulase activity was significantly higher in *T. arizonensis* colonies regardless of fungal species than in *A. texana* growing *Attamyces* (Fig. 3, Table 1). Xylanase activity did not appear to be influenced by ant or fungal species (Table 1; supplementary material Fig. S2). Furthermore, all xylanase effects except fungal symbiont species were significantly heteroscedastic. Thus, the xylanase *P*-values reported here may be inflated.




Fig. 1. Mean amylase activity in gardens for each experimental combination of ant (*Atta texana* or *Trachymyrmex arizonensis*) and fungal species (*Attamyces* or *Trachymyces*). Amylase activity is given as µg reducing sugar per mg ml⁻¹ of fungus garden extract. Significant differences are denoted by different letters ($P<0.05$, Scheffé's test). Data were analyzed on log₁₀ transformed data. Error bars are ± 1 s.d.

Attamyces gardens exhibited protease activity that was approximately twice that of *Trachymyces* gardens and this appeared to be independent of ant species [*Attamyces* 1545(\pm 536) $\times 10^3$ U, *Trachymyces* 834(\pm 384) $\times 10^3$ U; Table 1; supplementary material Fig. S3]. There might have been a weak interaction between ant and fungal species in protease activity (protease activity being higher in *T. arizonensis* colonies growing *Attamyces* than in conspecific colonies growing *Trachymyces*, $F_{1,19}=5.95$, $P=0.03$); however, because variances were heteroscedastic, attempts to make firm conclusions here introduce the risk of a Type I Error. Consequently, the interaction was removed from the model.

Fungus garden growth rates

Despite differences among enzymatic activities, fungus garden growth rates did not differ among the colonies of either ant species.

Trachymyrmex arizonensis

Over the course of the year, the growth rate of gardens of colonies collected in 2011 was not dependent on fungal species when cultivated by *T. arizonensis* (fungus: $F_{1,6}=0.964$, $P=0.361$; Fig. 4A). Because gardens grew in size steadily during the 1 year experiment, time was a significant variable ($F_{10,60}=72.25$, $P=0.0001$, Greenhouse–Geisser $E=0.145_{145,871}$, $P<0.0001$, Huynh–Feldt $E=0.211_{2,113,12,68}$, $P<0.0001$), but time did not interact significantly




Fig. 2. Mean pectinase activity in gardens for each experimental combination of ant (*Atta texana* or *Trachymyrmex arizonensis*) and fungal species (*Attamyces* or *Trachymyces*). Pectinase activity is given as µg reducing sugar per mg ml⁻¹ of fungus garden extract. Significant differences are denoted by different letters ($P<0.05$, Scheffé's test). Data were analyzed on log₁₀ transformed data. Error bars are ± 1 s.d.

with fungal species. Colonies therefore exhibited similar growth patterns regardless of fungal species ($F_{10,60}=3.06$, $P=0.003$, Greenhouse–Geisser $E=0.145_{145,871}$, $P=0.11$, Huynh–Feldt $E=0.211_{2,113,12,68}$, $P=0.08$).

Atta texana

Colonies growing *Trachymyces* grew slightly larger gardens over the first 12 months of their lives than those growing *Attamyces*, but this was statistically insignificant ($F_{1,10}=4.02$, $P=0.07$; Fig. 4B). Years were pooled because of low sample size. Time was significant ($F_{8,80}=43.96$, $P=0.0001$, Greenhouse–Geisser $E=0.388_{3,1,31,03}$, $P<0.0001$, Huynh–Feldt $E=0.639_{5,11,51,08}$, $P<0.0001$) and there was a significant interaction between garden volume and time ($F_{8,80}=3.22$, $P<0.01$, Greenhouse–Geisser $E=0.388_{3,1,31,03}$, $P<0.05$, Huynh–Feldt $E=0.639_{5,11,51,08}$, $P<0.05$). However, closer examination indicated that the significant interaction was largely due to differences between the first 2 months and all successive measurements. Thus, time was the most significant variable explaining the variation in the increase of garden volume.

DISCUSSION

The activity of particular enzymes in gardens of *A. texana* and *T. arizonensis* appears to be an emergent feature of ant–fungal interaction. Amylase and pectinase activities were higher when *A.*

Table 1. Summary of factorial ANOVA tests of enzyme activities of gardens of *Atta texana* and *Trachymyrmex arizonensis* ants cultivating either *Attamyces* or *Trachymyces* fungus

Substrate	Ant species	Fungal species	Ant \times fungus interaction
Starch	$F_{1,19}=3.23$, $P=0.09$	$F_{1,19}=11.71$, $P=0.003$	$F_{1,19}=6.91$, $P=0.02$
Pectin	$F_{1,19}=9.42$, $P=0.006$	$F_{1,19}=14.95$, $P<0.001$	$F_{1,19}=8.09$, $P=0.01$
Cellulose	$F_{1,19}=23.64$, $P=0.0001$	$F_{1,19}=0.0003$, $P=0.99$	$F_{1,19}=5.50$, $P=0.03$
Xylan	$F_{1,19}=1.36$, $P=0.27$	$F_{1,20}=0.006$, $P=0.94$	$F_{1,20}=1.23$, $P=0.31$
Protein	$F_{1,20}=0.245$, $P=0.63$	$F_{1,20}=7.89$, $P=0.01$	–

Interactions with $P>0.25$ were pooled and only main effects are listed here. Significant tests ($\alpha\leq 0.05$) are highlighted in bold. To account for the unbalanced design (only five colonies of *T. arizonensis* were growing *Attamyces*, whereas all other sample sizes were $N=6$) and to make up for the ‘missing’ colony, a dummy variable was instead used (see Materials and methods); as a result, one degree of freedom had to be removed from the corresponding F -tests. This was done only in cases where the factorial test was significant to avoid inflating the significance of the interaction.




Fig. 3. Mean cellulase activity in gardens for each experimental combination of ant (*Atta texana* or *Trachymyrmex arizonensis*) and fungal species (*Attamyces* or *Trachymyces*). Cellulase activity is given as μg reducing sugar per mg ml^{-1} of fungus garden extract. Significant differences are denoted by different letters ($P < 0.05$, Scheffé's test). Data were analyzed on square-root transformed data. Error bars are ± 1 s.d.

texana grew *Attamyces* than when growing *Trachymyces*, consistent with both the host-independent hypothesis (*Attamyces* has higher enzymatic activity regardless of the ant species growing it) and the adaptive combination hypothesis (activities are highest in natural combinations), but the corresponding pattern was not observed for *T. arizonensis*. *Trachymyrmex arizonensis* gardens exhibited similar amylase and pectinase activities regardless of the fungal species being cultivated. The weight of the observations therefore supports the complex synergism hypothesis, because the activity of each enzyme appears to follow idiosyncratic patterns as a result of ant–fungus combinations. In other words, neither fungal species by itself nor ant species by itself determines enzymatic activities of the symbiosis; rather, activities result from an interaction between ant and fungal species.

The weight of the accumulated evidence therefore does not support Stradling and Powell's (Stradling and Powell, 1986) hypothesis (i.e. the host-independent hypothesis) that *Attamyces* have intrinsically higher enzymatic activities and growth rates than *Trachymyces*, and that this superiority of *Attamyces* played a key role in the evolution of leaf cutting (Figs 1–4). *Trachymyrmex arizonensis* exhibited similar fungus garden growth rates regardless of fungal species cultivated. The equivalence of growth of the two fungal species when grown by *T. arizonensis* may explain why, while *Attamyces* may be more productive *in vitro* (Stradling and Powell, 1986), it has not swept through the *T. arizonensis* population. This may also explain why some *T. arizonensis* colonies in the field grow *Attamyces* (U.G.M., unpublished data; supplementary material Fig. S1). However, the host-independent hypothesis was supported by observations in the *A. texana* colonies. The reduced enzymatic performance of *A. texana* growing *Trachymyces* might explain why *A. texana* is known to grow only *Attamyces*, even though our observations show that *A. texana* does not seem to have difficulty in growing *Trachymyces* in the laboratory [unlike an earlier report on *Attamyces mexicana* colonies switched in the lab to a *Trachymyces* fungus (Sánchez-Peña, 2005)].

What then constrains cultivar switching? One possibility is that interactions among the microbiota associated with the ants and/or




Fig. 4. Fungus garden growth rate of each ant species growing either fungal species. (A) Growth of fungus gardens of *Trachymyrmex arizonensis* colonies over the course of one year (August 2011 to July 2012). Sample sizes: *Trachymyces* $N=3$, *Attamyces* $N=5$. Data were from the cohort of founding queens collected in 2011. Fungus garden volume increase was not significantly different for the two fungal species (fungus: $F_{1,6}=0.964$, $P=0.361$). Volumes from April were excluded from analysis because the variances were significantly heteroscedastic. (B) Growth of fungus gardens of *Atta texana* colonies over the course of their first year of life. Sample sizes: *Trachymyces* $N=6$ ($N=3$ from 2010 cohort, $N=3$ from 2011 cohort), *Attamyces* $N=5$ ($N=1$ from 2010 cohort, $N=4$ from 2011 cohort). Fungus garden volume increase was not significantly different for the two fungal species (fungus: $F_{1,10}=4.02$, $P=0.07$). Volumes from August, December and February were excluded from the analysis because the variances were significantly heteroscedastic (see Materials and methods). In both cases, error bars correspond to $\pm 95\%$ confidence intervals. Data depicted are untransformed.

fungi constrain ant and fungal combinations. This possibility was raised when *T. septentrionalis* and *Trachymyrmex turrifex* experienced fitness reductions after adopting *Attamyces* fungus (Seal and Mueller, 2014), which apparently was the result of invasion by weedy fungi destroying gardens, though the exact mechanism of garden decline was not known with certainty in that study. *Trachymyrmex arizonensis* likely associate with different microbial communities that are distinct from other species of *Trachymyrmex*, as do other attine species (Ishak et al., 2011; Rodrigues et al., 2011; Sen et al., 2009), which may or may not be able to withstand invasions by competitor fungi (Haeder et al., 2009; Mueller, 2012; Rodrigues et al., 2009). Another non-exclusive possibility is that ants may not apply the appropriate triggers to novel fungal

symbionts that would stimulate the synthesis of the diverse enzymatic machinery required for the metabolism of carbohydrates, lignocelluloses and proteins that *Attamyces* is known to possess (Aylward et al., 2013). For example, pectinase activity was lower in *T. arizonensis* colonies growing *Attamyces* than in *A. texana* colonies growing *Attamyces* (Fig. 2). Thus ant–fungus switches may result in mismatches at the physiological and genomic level (i.e. negative inter-genomic epistasis (Heath, 2009; Linksvayer, 2007; Wade, 2007).

One of the more surprising findings of this study sustains an ongoing debate on whether *Attamyces* can metabolize cellulose. A number of studies report significant cellulase activity for *Attacultivated* fungi grown on a variety of natural and artificial diets (Bacci et al., 1995; D’Ettorre et al., 2002; Martin and Weber, 1969; Schiøtt et al., 2008; Silva et al., 2003), whereas other studies failed to find significant cellulase activity (Abril and Bucher, 2002; Bucher et al., 2004). The results presented here are intriguing because they suggest that *T. arizonensis* colonies growing *Attamyces* on natural substrates (catkins) can in fact metabolize cellulose because *Attamyces* grown by *T. arizonensis* exhibited higher cellulase activity than *Attamyces* grown by *A. texana*. *Attamyces* is known to possess at least three genes for cellulose degradation (Aylward et al., 2013). Therefore, one possible explanation of our observations is that when *Attamyces* is grown by *Trachymyrmex* ants, cellulase genes are activated or are more actively expressed than when grown by *Atta* ants. Some authors have raised a second possibility that leaf-cutting ants never allow their *Attamyces* gardens to digest cellulose because the ants remove older sections of gardens and place them in refuse depots once simpler compounds (starches, pectins, etc.) are exhausted (De Fine Licht et al., 2010; Moller et al., 2011). Evaluation of this hypothesis requires knowledge of whether *Atta* and *Trachymyrmex* ants differ in how much time they give their fungus to get to a point in the garden cycle when cellulose is extracted from substrates. A third possibility is that switching gardens between host species may cause a reshuffling of auxiliary microbes in garden biofilms such that cellulose-metabolizing microbes [e.g. yeasts (Mendes et al., 2012)] are either more abundant or more active in gardens grown by *T. arizonensis*. Considering recent interest in the role of fungus gardens in cellulose metabolism (Aylward et al., 2012a; Suen et al., 2010), the fungal symbioses of *T. arizonensis* ants might be important sources of cellulases, meriting further study.

MATERIALS AND METHODS

This study employed reciprocal symbiont switch experiments conducted on the leaf-cutter ant *A. texana* and the higher-attine ant *T. arizonensis*. Colonies of both species were obtained by rearing newly mated queens collected immediately after mating flights (Seal, 2009; Seal and Tschinkel, 2007c) in 2010 and 2011. Colonies were maintained in plaster-lined boxes and fed *ad libitum* in a manner similar to previously published methods (Seal and Tschinkel, 2007a; Seal and Tschinkel, 2007b). Queens were fed and nests cleaned daily until the first workers appeared, after which colonies were fed at least twice weekly with oak catkins (staminate flowers of *Quercus shumardii*, a naturally occurring fungal substrate that many fungus-gardening ants will accept and feed their fungus; J.N.S., unpublished observation).

Trachymyrmex arizonensis is a common species along mid-elevations (1000–2000 m) in mixed oak–juniper–pinyon pine forests in the Sonoran Desert and western portions of the Chihuahuan Desert (Rabeling et al., 2007). *Atta texana* is found throughout central, east and south Texas, western Louisiana and northeastern Mexico (Mueller et al., 2011a; Mueller et al., 2011b; Sanchez-Peña, 2010). *Atta texana* cultivates a single fungal species typical of the vast majority of leaf-cutting ants (*Leucocoprinus gongylophorus*) (Fisher et al., 1994; Mikheyev et al., 2006; Mueller et al.,

2011a; Mueller, 2002; Pagnocca et al., 2001); however, this species rarely reproduces sexually – we refer to it by its anamorph (asexual) form, *Attamyces bromatificus* Kreisel (Mueller et al., 2010; Mueller et al., 2011b; Seal et al., 2012; Seal and Mueller, 2014). In contrast, *T. arizonensis* cultivates a *Trachymyces* fungus typical for most (but not all) *Trachymyrmex* species that is placed in a taxonomically unresolved *Leucocoprinus* clade that is the sister clade to the *Attamyces* clade. Ongoing surveys, however, indicate that some populations of *T. arizonensis* cultivate *Attamyces* instead of *Trachymyces* (supplementary material Fig. S1; U.G.M., in preparation). *Trachymyrmex arizonensis* likely exchanges symbionts with the sympatric leaf-cutter ant *Acromyrmex versicolor* (supplementary material Fig. S1; U.G.M., in preparation). Although the two ant species used in this study are not sympatric, these ants and their cultivated fungi are representative of the two major clades of the higher Attini (Schultz and Brady, 2008). Despite population differentiation between *Attamyces* from Texas and Arizona (Mueller et al., 2011b), this genetic differentiation is substantially less than the differentiation between *Attamyces* and *Trachymyces* (Mikheyev et al., 2006; Mikheyev et al., 2008; Mueller et al., 2010; Mueller et al., 2011b) (supplementary material Fig. S1).

Queens were randomly assigned to either the *Attamyces* or *Trachymyces* condition and provided with ~50 g of garden (*T. arizonensis* queens) or 200 g of garden (*A. texana* queens; owing to the much larger size of *Atta* queens) (Seal, 2009). *Atta texana* queens were supplied with either *Attamyces* collected from an *A. texana* nest near Austin, TX, USA, in May 2007 (30°12.622'N, 97°38.469'W, 125 m elevation) or *Trachymyces* fungus collected from a *T. septentrionalis* nest collected in April 2010 at the University of Texas Stengl ‘Lost Pines’ Biological Station (30°05.218'N, 97°10.425'W, 141 m elevation).

The experimental switches on *T. arizonensis* used in this study were conducted on eight colonies started with queens collected on 27–28 July 2011. Five of these were reared on *Attamyces* and three on *Trachymyces*. Because of unequal mortality among these queens, we also include data on 10 colonies reared similarly from queens collected on 25–27 July 2010 (all of which were cultivating *Trachymyces*); three of these colonies were randomly selected for enzymatic activity analysis. All *T. arizonensis* colonies were collected at the Southwest Research Station, near Portal, AZ, USA (31°53.025'N, 109°12.374'W, 1646 m elevation), 1 day after the first heavy rain of the summer monsoon season. Colonies growing *Attamyces* were supplied with fungus collected from an *Acromyrmex versicolor* colony reared from a newly mated queen collected near Tucson, AZ, USA, in 2009 (32°18.971'N, 110°53.562'W, 858 m elevation). Colonies growing *Trachymyces* were supplied with fungus from a *T. arizonensis* colony collected in July 2010 at the Southwest Research Station. Upon collection, all *T. arizonensis* queens were supplied with one melanized pupa obtained from a mature *T. arizonensis* colony in the laboratory, placed with ca. 8 cm³ of fungus into a 4 cm Petri dish (garden chamber) that was inserted inside a 9 cm Petri dish. All of the space between these two Petri dishes was filled with cotton saturated with sterile water. *Trachymyrmex arizonensis* queens were fed and cleaned daily until the first workers appeared (Seal and Tschinkel, 2007c). Colonies were kept in these nesting containers until the first callow workers appeared (ca. 6 weeks later), then colonies were transferred to 7×7 cm plastic boxes with a 5 mm-thick bottom of moistened dental plaster (Marjoy Enterprises, San Antonio, TX, USA). After approximately 1 year of age, colonies were moved to larger nesting containers used in previous studies on *Trachymyrmex* ants (plaster-lined, cylindrically shaped, 196 cm³ depressions in a square plastic box, 11×11×3 cm) (Seal and Mueller, 2014; Seal and Tschinkel, 2007a; Seal and Tschinkel, 2007b). All *T. arizonensis* colonies were fed oak catkins (*Quercus shumardii*) *ad libitum* throughout the duration of the entire experiment. Colonies of the two species were fed the same type of catkins to avoid confounding the results with diet (Kooij et al., 2011).

The 12 *Atta texana* colonies used in this study were similarly reared from newly mated queens collected after mating flights near Hornsby Bend, TX, USA (30°12'37.3"N, 97°38'28.07"W). Of the colonies growing *Trachymyces*, three were collected on 13 May 2011 and three were collected on 16–18 May 2010, whereas the six colonies growing *Attamyces* came from cohorts collected on 18 May 2010 (*N*=1), 13 May 2011 (*N*=4) and April 2007 (*N*=1). Freshly collected queens were placed in square plastic

boxes (7×7×3 cm) lined with 5 mm of dental plaster, which was moistened twice weekly. Because *Atta* queens do not forage during the founding phase (Fernández-Marin et al., 2004; Seal, 2009), they were given substrate only after the first workers emerged. After worker emergence, colonies were connected to a foraging arena via a plastic tube and fed and cleaned at least twice a week. *Atta texana* colonies were fed substrates similar to those fed to the *T. arizonensis* colonies, except they were also periodically supplied with pear leaves (Bradford pear, *Pyrus calleryana*).

Fungus garden growth measurements

Colonies were monitored monthly over the course of the first year of their lives. Volumes of gardens were measured monthly in both species. Fungus garden volumes were estimated by measuring the maximum dimensions (length, width and height) in each nest box at monthly intervals. The top of the plaster nest chamber was completely covered with a piece of Plexiglas. The width and length of each fungus garden were measured with a ruler placed on top of each nest cover (a piece of Plexiglas that contained five pre-drilled 1 mm holes, 2 cm apart, from the center of the cover) (Seal and Mueller, 2014; Seal and Tschinkel, 2007a; Seal and Tschinkel, 2007b).

Fungal enzymatic activity assays

Enzymatic activity assays were conducted in 2012 when all colonies were at least 1 year old. Because fungus gardens are thought to exhibit spatial variation in enzymatic activity (Moller et al., 2011; Suen et al., 2010), fungus garden material was selected from the uppermost 1.5 cm of each fungus garden, which approximated a sample from the top one-third of a garden. Most fungally derived digestion occurs in the upper-most part of the garden, whereas relative activity of enzymes derived from non-cultivar microbes (e.g. bacterial biofilms) seems to be greater in the lower portions (Moller et al., 2011; Suen et al., 2010). Furthermore, because enzymatic activities in gardens depend on the substrates used by the ants (Kooij et al., 2011), all ant colonies of both species were fed a strict diet of oak catkins for 4 weeks prior to enzymatic assays. Finally, gardens were also sampled at least 48 h after the last feeding, so that there were no freshly deposited substrates in the gardens. Enzymes were extracted from the fungus gardens by grinding ca. 120 mg of fungus garden material in an Eppendorf tube containing 500 µl of 20 mmol l⁻¹ phosphate buffer (pH 6.9) after the removal of visible eggs, larvae and pupae. Extracts were centrifuged at 4°C for 15 min at 14,000 rpm. The supernatant was then transferred to a fresh tube, which was used for both the carbohydrase and protease activity assays. Each colony was sampled four times over the course of 2 weeks, and the average enzymatic activity was taken from these four values to provide an estimate for each colony.

Carbohydrase activity was measured using the dinitrosalicylic acid method, which assays reducing sugar concentration (Miller, 1959; Silva et al., 2003). Accordingly, 10 µl of fungus garden extract was added to 40 µl of water and 50 µl of 1% (w/v) (=500 µg of substrate) carbohydrate solution, and incubated at room temperature for 60 min. Four carbohydrate solutions were used: pectin (from apple), starch, carboxymethyl cellulose and xylan (from beech wood) (all purchased from Sigma-Aldrich, St Louis, MO, USA). Prior work has indicated these substrates to be among the most common types of plant carbohydrate digested by the ant fungus (Moller et al., 2011; Schiøtt et al., 2008; Schiøtt et al., 2010). The reactions were terminated by adding 50 µl of 96 mmol l⁻¹ dinitrosalicylic acid (DNS) solution and incubating at 99°C for 15 min. At high temperature, the DNS dye changes color depending on the concentration of reducing sugars (the darker the color, the higher the concentration of reducing sugars). Control samples were treated by adding the DNS and the enzyme extract before immediate incubation at 99°C. After incubation, 50 µl of each sample was added to 150 µl of water and then read in a spectrophotometer at 540 nm. Amounts hydrolyzed were interpolated using a standard curve for glucose.

Protease activity was measured using the Azocasein method (Charney and Tomarelli, 1947) used previously in studies on attine protease activity (Semenova et al., 2011). This method uses azocasein (azo-labeled casein; Sigma-Aldrich). Azocasein is a non-specific protease substrate, which upon hydrolysis releases the azo dye, the concentration of which can be inferred colorimetrically. A 10 µl sample of fungal extract was added to 15 µl of 2%

azocasein solution and incubated for 60 min at room temperature. The reaction was terminated by adding 120 µl of 10% trichloroacetic acid (TCA). The resulting suspension was centrifuged (15,000 g) for 5 min and added to 140 µl of 1 mol l⁻¹ NaOH. Absorbance was measured at 440 nm at 25°C. Control samples were prepared identically except enzyme extract was added immediately before the application of TCA. Relative protease activity was calculated from the difference between treatment and control absorbances.

The methods used to estimate enzyme activity do not differentiate between enzymes secreted by the cultivated fungus or associated microbial biofilms. The metabolic activities of fungus gardens are likely quite diverse (Aylward et al., 2012a; Aylward et al., 2012b). Yeasts inhabiting the garden may be important contributors of cellulase activity and perform crucial functions in extracting energy from pectins (Mendes et al., 2012). Although it is unclear whether the distinction between compounds produced by *Leucocoprinus* spp. versus other microbes inhabiting the fungus gardens is important to the ants, because the ants are potentially consuming whatever they find in the fungus garden, the data presented here correspond to an 'extended phenotype' of enzymatic activity in the symbiosis *sensu lato*. As a result, when the statistical tests report a significant 'fungal' effect (or 'ant' effect), it should be interpreted as an effect influenced by the fungus garden and possibly its associated biofilms.

Statistical analyses

The problem of unequal sample sizes resulting from only five switched *T. arizonensis* colonies (whereas all other ant × fungal combinations had *N*=6) was solved using Underwood's technique of creating a dummy variable calculated from a mean value of each group to which the missing data belonged (Seal and Tschinkel, 2007a; Underwood, 1997). This method alters neither the mean nor the variance (Underwood, 1997), but to avoid rejecting the null hypothesis falsely (Type I Error), the test is made more conservative by subtracting one degree of freedom from the corresponding *F*-test for each of the missing values generated.

Garden volumes were analyzed with a repeated measures ANOVA, with time as the repeated measure and with fungal symbiont species as main effect. Volumes were repeatedly measured on all colonies in this experiment at approximately 1 month intervals. In addition to tests for standard parametric assumptions, we tested for sphericity, which tests for equality of variances among the repeated measures (Keselman et al., 2001; Sokal and Rohlf, 1995). In instances where sphericity tests were rejected, we report Greenhouse–Geisser- and Huynh–Feldt-corrected *F*-statistics and *P*-values. Data were either log₁₀ or square-root transformed to meet parametric assumptions.

Fungus identification

Small samples (hyphae or spores) were collected with sterile forceps from the emerging cultures and placed in Chelex resin (Sigma-Aldrich) and heated to near boiling for 90 min. Pure fungal cultures were identified by PCR amplification of the *ITS* gene using primers ITS4 and ITS5 (Mueller et al., 1998; Sen et al., 2009; White et al., 1990), then sequencing at the University of Texas at Austin University DNA Sequencing Facilities. Sequences were identified using BLAST at NCBI GenBank.

Acknowledgements

Constructive comments by reviewers and editors greatly improved the manuscript. We thank Kevin Anderson at the Hornsby Bend Environmental Research Center, and the staff at the American Museum's Southwest Research Station, especially Barbara Roth. Rebecca Clark, Bob Johnson and Diana Wheeler shared collections and their knowledge of the attines of Arizona. Cecil Harkey at the UT Austin Core Facility supplied the plate reader.

Competing interests

The authors declare no competing financial interests.

Author contributions

J.N.S. and U.G.M. conceived and designed the experiments. M.S. developed and modified the enzymatic assays. J.N.S. executed the experiments and enzymatic assays. J.N.S. and U.G.M. wrote and revised the manuscript, with significant input by M.S.

Funding

Financial support was provided by the National Science Foundation [IOS-0920138] to J.N.S. and U.G.M., and by the Danish National Research Foundation [DNRF57] to M.S.

Supplementary material

Supplementary material available online at
http://jeb.biologists.org/lookup/suppl/doi:10.1242/jeb.098483/-/DC1

References

- Aanen, D. K. and Eggleton, P. (2005). Fungus-growing termites originated in African rain forest. *Curr. Biol.* **15**, 851–855.
- Aanen, D. K., Eggleton, P., Rouland-Lefevre, C., Guldberg-Froslev, T., Rosendahl, S. and Boomsma, J. J. (2002). The evolution of fungus-growing termites and their mutualistic fungal symbionts. *Proc. Natl. Acad. Sci. USA* **99**, 14887–14892.
- Abril, A. B. and Bucher, E. H. (2002). Evidence that the fungus cultured by leaf-cutting ants does not metabolize cellulose. *Ecol. Lett.* **5**, 325–328.
- Aylward, F. O., Burnum, K. E., Scott, J. J., Suen, G., Tringe, S. G., Adams, S. M., Barry, K. W., Nicora, C. D., Piehowski, P. D., Purvine, S. O. et al. (2012a). Metagenomic and metaproteomic insights into bacterial communities in leaf-cutter ant fungus gardens. *ISME J.* **6**, 1688–1701.
- Aylward, F., Currie, C. and Suen, G. (2012b). The evolutionary innovation of nutritional symbioses in leaf-cutter ants. *Insects* **3**, 41–61.
- Aylward, F., Burnum-Johnson, K., Tringe, S., Teiling, C., Tremmel, D., Moeller, J., Scott, J. A., Barry, K. W., Piehowski, P., Nicora, C. et al. (2013). *Leucoagaricus gongylophorus* produces diverse enzymes for the degradation of recalcitrant plant polymers in leaf-cutter ant fungus gardens. *Appl. Environ. Microbiol.* **79**, 3770–3778.
- Bacci, M., Jr, Anversa, M. M. and Pagnocca, F. C. (1995). Cellulose degradation by *Leucocoprinus gongylophorus*, the fungus cultured by the leaf-cutting ant *Atta sexdens rubropilosa*. *Antonie Leeuwenhoek* **67**, 385–386.
- Bacci, M., Jr, Bueno, O. C., Rodrigues, A., Pagnocca, F. C., Somera, A. F. and Silva, A. (2013). A metabolic pathway assembled by enzyme selection may support herbivory of leaf-cutter ants on plant starch. *J. Insect Physiol.* **59**, 525–531.
- Beshers, S. N. and Traniello, J. F. A. (1994). The adaptiveness of worker demography in the attine ant, *Trachymyrmex septentrionalis*. *Ecology* **75**, 763–775.
- Bucher, E. H., Marchesini, V. and Abril, A. (2004). Herbivory by leaf-cutting ants: nutrient balance between harvested and refuse material. *Biotropica* **36**, 327–332.
- Charney, J. and Tomarelli, R. M. (1947). A colorimetric method for the determination of the proteolytic activity of duodenal juice. *J. Biol. Chem.* **171**, 501–505.
- D'Ettorre, P., Mora, P., Dibangou, V., Rouland, C. and Errard, C. (2002). The role of the symbiotic fungus in the digestive metabolism of two species of fungus-growing ants. *J. Comp. Physiol. B* **172**, 169–176.
- De Fine Licht, H. H. and Biedermann, P. H. (2012). Patterns of functional enzyme activity in fungus farming ambrosia beetles. *Front. Zool.* **9**, 13.
- De Fine Licht, H. H., Schiøtt, M., Mueller, U. G. and Boomsma, J. J. (2010). Evolutionary transitions in enzyme activity of ant fungus gardens. *Evolution* **64**, 2055–2069.
- De Fine Licht, H., Schiøtt, M., Rogowska-Wrzęsinska, A., Nygaard, S., Roepstorff, P. and Boomsma, J. (2013). Laccase detoxification mediates the nutritional alliance between leaf-cutting ants and fungus-garden symbionts. *Proc. Natl. Acad. Sci. USA* **110**, 583–587.
- Douglas, A. E. (2010). *The Symbiotic Habit*. Princeton, NJ: Princeton University Press.
- Erthal, M., Jr, Peres Silva, C. and Samuels, R. I. (2004). Digestive enzymes of leaf-cutting ants, *Acromyrmex subterraneus* (Hymenoptera: Formicidae: Attini): distribution in the gut of adult workers and partial characterization. *J. Insect Physiol.* **50**, 881–891.
- Erthal, M., Jr, Silva, C. P., Cooper, R. M. and Samuels, R. I. (2009). Hydrolytic enzymes of leaf-cutting ant fungi. *Comp. Biochem. Physiol.* **152B**, 54–59.
- Farrell, B. D., Sequeira, A. S., O'Meara, B. C., Normark, B. B., Chung, J. H. and Jordal, B. H. (2001). The evolution of agriculture in beetles (Curculionidae: Scolytinae and Platypodinae). *Evolution* **55**, 2011–2027.
- Fernández-Marin, H., Zimmerman, J. K. and Wcislo, W. T. (2004). Ecological traits and evolutionary sequence of nest establishment in fungus-growing ants (Hymenoptera, Formicidae, Attini). *Biol. J. Linn. Soc. Lond.* **81**, 39–48.
- Fisher, P. J., Stradling, D. J. and Pegler, D. N. (1994). *Leucoagaricus basidiomata* from a live nest of the leaf-cutting ant *Atta cephalotes*. *Mycol. Res.* **98**, 884–888.
- Green, A. M., Mueller, U. G. and Adams, R. M. M. (2002). Extensive exchange of fungal cultivars between sympatric species of fungus-growing ants. *Mol. Ecol.* **11**, 191–195.
- Haeder, S., Wirth, R., Herz, H. and Spitter, D. (2009). Candidicin-producing *Streptomyces* support leaf-cutting ants to protect their fungus garden against the pathogenic fungus *Escovopsis*. *Proc. Natl. Acad. Sci. USA* **106**, 4742–4746.
- Heath, K. D. (2009). Intergenic epistasis and coevolutionary constraint in plants and rhizobia. *Evolution* **64**, 1446–1458.
- Hölldobler, B. and Wilson, E. O. (2011). *The Leafcutter Ants*. New York, NY: W.W. Norton.
- Ishak, H. D., Miller, J. L., Sen, R., Dowd, S. E., Meyer, E. and Mueller, U. G. (2011). Microbiomes of ant castes implicate new microbial roles in the fungus-growing ant *Trachymyrmex septentrionalis*. *Scientific Reports* **1**, 204.
- Kellner, K., Fernández-Marin, H., Ishak, H. D., Sen, R., Linksvayer, T. A. and Mueller, U. G. (2013). Co-evolutionary patterns and diversification of ant–fungus associations in the asexual fungus-farming ant *Mycocepurus smithii* in Panama. *J. Evol. Biol.* **26**, 1353–1362.
- Keselman, H. J., Algina, J. and Kowalchuk, R. K. (2001). The analysis of repeated measures designs: a review. *Br. J. Math. Stat. Psychol.* **54**, 1–20.
- Kooij, P., Schiøtt, M., Boomsma, J. J. and de Fine Licht, H. (2011). Rapid shifts in *Atta cephalotes* fungus-garden enzyme activity after a change in fungal substrate (Attini, Formicidae). *Insectes Soc.* **58**, 145–151.
- Kültz, D., Clayton, D. F., Robinson, G. E., Albertson, C., Carey, H. V., Cummings, M. E., Dewar, K., Edwards, S. V., Hofmann, H. A., Gross, L. J. et al. (2013). New frontiers for organismal biology. *BioScience* **63**, 464–471.
- Leal, I. R. and Oliveira, P. S. (1998). Interactions between fungus-growing ants (Attini), fruits, and seeds in Cerrado vegetation in southeast Brazil. *Biotropica* **30**, 170–178.
- Leal, I. R. and Oliveira, P. S. (2000). Foraging ecology of attine ants in a Neotropical savanna: seasonal use of fungal substrate in the cerrado vegetation of Brazil. *Insectes Soc.* **47**, 376–382.
- Linksvayer, T. A. (2007). Ant species differences determined by epistasis between brood and worker genomes. *PLoS ONE* **2**, e994.
- Martin, M. M. (1987a). *Invertebrate-Microbial Interactions: Ingested Fungal Enzymes in Arthropod Biology*. Ithaca, New York, NY: Cornell University Press.
- Martin, M. M. (1987b). The symbioses between the attine ants and the fungi they culture in their nests. In *Invertebrate-microbial Interactions: Ingested Fungal Enzymes in Arthropod Biology* (ed. M. M. Martin), pp. 91–126. Ithaca, NY: Cornell University Press.
- Martin, M. M. and Weber, N. A. (1969). The cellulose-utilizing capability of the fungus cultured by the Attine ant *Atta colombica tonsipes*. *Ann. Entomol. Soc. Am.* **62**, 1386–1387.
- McFall-Ngai, M., Hadfield, M. G., Bosch, T. C., Carey, H. V., Domazet-Lošo, T., Douglas, A. E., Dubilier, N., Eberl, G., Fukami, T., Gilbert, S. F. et al. (2013). Animals in a bacterial world, a new imperative for the life sciences. *Proc. Natl. Acad. Sci. USA* **110**, 3229–3236.
- Mehdabadi, N. J., Mueller, U. G., Brady, S. G., Himler, A. G. and Schultz, T. R. (2012). Symbiont fidelity and the origin of species in fungus-growing ants. *Nat. Commun.* **3**, 840.
- Mendes, T. D., Rodrigues, A., Dayo-Owoyemi, I., Marson, F. A. L. and Pagnocca, F. C. (2012). Generation of nutrients and detoxification: possible roles of yeasts in leaf-cutting ant nests. *Insects* **3**, 228–245.
- Meyer, S., Leal, I., Tabarelli, M. and Wirth, R. (2011). Ecosystem engineering by leaf-cutting ants: nests of *Atta cephalotes* drastically alter forest structure and microclimate. *Ecol. Entomol.* **36**, 14–24.
- Mikheyev, A. S., Mueller, U. G. and Abbot, P. (2006). Cryptic sex and many-to-one coevolution in the fungus-growing ant symbiosis. *Proc. Natl. Acad. Sci. USA* **103**, 10702–10706.
- Mikheyev, A. S., Vo, T. and Mueller, U. G. (2008). Phylogeography of post-Pleistocene population expansion in a fungus-gardening ant and its microbial mutualists. *Mol. Ecol.* **17**, 4480–4488.
- Mikheyev, A. S., Mueller, U. G. and Abbot, P. (2010). Comparative dating of attine ant and lepidoteous cultivar phylogenies reveals coevolutionary synchrony and discord. *Am. Nat.* **175**, E126–E133.
- Miller, G. L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.* **31**, 426–428.
- Moller, I. E., De Fine Licht, H. H., Harholt, J., Willats, W. G. and Boomsma, J. J. (2011). The dynamics of plant cell-wall polysaccharide decomposition in leaf-cutting ant fungus gardens. *PLoS ONE* **6**, e17506.
- Mueller, U. G. (2002). Ant versus fungus versus mutualism: ant–cultivar conflict and the deconstruction of the attine ant–fungus symbiosis. *Am. Nat.* **160** Suppl. 4, S67–S98.
- Mueller, U. G. (2012). Symbiont recruitment versus ant–symbiont co-evolution in the attine ant–microbe symbiosis. *Curr. Opin. Microbiol.* **15**, 269–277.
- Mueller, U. G., Rehner, S. A. and Schultz, T. R. (1998). The evolution of agriculture in ants. *Science* **281**, 2034–2038.
- Mueller, U. G., Gerardo, N. M., Aanen, D. K., Six, D. L. and Schultz, T. R. (2005). The evolution of agriculture in insects. *Annu. Rev. Ecol. Syst.* **36**, 563–595.
- Mueller, U. G., Scott, J. J., Ishak, H. D., Cooper, M. and Rodrigues, A. (2010). Monoculture of leafcutter ant gardens. *PLoS ONE* **5**, e12668.
- Mueller, U. G., Mikheyev, A. S., Hong, E., Sen, R., Warren, D. L., Solomon, S. E., Ishak, H. D., Cooper, M., Miller, J. L., Shaffer, K. A. et al. (2011a). Evolution of cold-tolerant fungal symbionts permits winter fungiculture by leafcutter ants at the northern frontier of a tropical ant–fungus symbiosis. *Proc. Natl. Acad. Sci. USA* **108**, 4053–4056.
- Mueller, U. G., Mikheyev, A. S., Solomon, S. E. and Cooper, M. (2011b). Frontier mutualism: coevolutionary patterns at the northern range limit of the leaf-cutter ant–fungus symbiosis. *Proc. R. Soc. B* **278**, 3050–3059.
- Pagnocca, F. C., Bacci, M., Jr, Fungaro, M. H., Bueno, O. C., Hebling, M. J., Sant'anna, A. and Capelari, M. (2001). RAPD analysis of the sexual state and sterile mycelium of the fungus cultivated by the leaf-cutting ant *Acromyrmex hispidus fallax*. *Mycol. Res.* **105**, 173–176.
- Rabeling, C., Cover, S. P., Johnson, R. A. and Mueller, U. G. (2007). A review of the North American species of the fungus-gardening ant genus *Trachymyrmex* (Hymenoptera: Formicidae). *Zootaxa* **1664**, 1–53.
- Richard, F.-J., Mora, P., Errard, C. and Rouland, C. (2005). Digestive capacities of leaf-cutting ants and the contribution of their fungal cultivar to the degradation of plant material. *J. Comp. Physiol. B* **175**, 297–303.
- Rodrigues, A., Cable, R. N., Mueller, U. G., Bacci, M., Jr and Pagnocca, F. C. (2009). Antagonistic interactions between garden yeasts and microfungi garden pathogens of leaf-cutting ants. *Antonie Leeuwenhoek* **96**, 331–342.

- Rodrigues, A., Mueller, U. G., Ishak, H. D., Bacci, M., Jr and Pagnocca, F. C. (2011). Ecology of microfungal communities in gardens of fungus-growing ants (Hymenoptera: Formicidae): a year-long survey of three species of attine ants in Central Texas. *FEMS Microbiol. Ecol.* **78**, 244-255.
- Sánchez-Peña, S. R. (2005). *Essays on Organismal Aspects of the Fungus-Growing Ant Symbiosis: Ecology, Experimental Symbiont Switches and Fitness of Atta, and a New Theory on the Origin of Ant Fungiculture*, pp. 156. Austin, TX: University of Texas – Austin.
- Sanchez-Peña, S. (2010). Some fungus-growing ants (Hymenoptera: Formicidae) from northeastern Mexico. *Fla. Entomol.* **93**, 501-504.
- Schiøtt, M., De Fine Licht, H. H., Lange, L. and Boomsma, J. J. (2008). Towards a molecular understanding of symbiont function: identification of a fungal gene for the degradation of xylan in the fungus gardens of leaf-cutting ants. *BMC Microbiol.* **8**, 40.
- Schiøtt, M., Rogowska-Wrzesinska, A., Roepstorff, P. and Boomsma, J. J. (2010). Leaf-cutting ant fungi produce cell wall degrading pectinase complexes reminiscent of phytopathogenic fungi. *BMC Biol.* **8**, 156.
- Schultz, T. R. and Brady, S. G. (2008). Major evolutionary transitions in ant agriculture. *Proc. Natl. Acad. Sci. USA* **105**, 5435-5440.
- Seal, J. N. (2009). Scaling of body weight and fat content in fungus-gardening ant queens: does this explain why leaf-cutting ants found claustrally? *Insectes Soc.* **56**, 135-141.
- Seal, J. N. and Mueller, U. G. (2014). Instability of novel ant–fungal associations suggest that microbial interactions constrain horizontal transfer in higher fungus-gardening ants. *Evol. Ecol.* **28**, 157-176.
- Seal, J. N. and Tschinkel, W. R. (2006). Colony productivity of the fungus-gardening ant, *Trachymyrmex septentrionalis* McCook, in a Florida pine forest (Hymenoptera: Formicidae). *Ann. Entomol. Soc. Am.* **99**, 673-682.
- Seal, J. N. and Tschinkel, W. R. (2007a). Co-evolution and the superorganism: switching cultivars does not alter the performance of fungus-gardening ant colonies. *Funct. Ecol.* **21**, 988-997.
- Seal, J. N. and Tschinkel, W. R. (2007b). Complexity in an obligate mutualism: do fungus-gardening ants know what makes their garden grow? *Behav. Ecol. Sociobiol.* **61**, 1151-1160.
- Seal, J. N. and Tschinkel, W. R. (2007c). Energetics of newly mated queens and colony founding in the fungus-gardening ants *Cyphomyrmex rimosus* and *Trachymyrmex septentrionalis* (Hymenoptera: Formicidae). *Physiol. Entomol.* **32**, 8-15.
- Seal, J. N. and Tschinkel, W. R. (2010). Distribution of the fungus-gardening ant, *Trachymyrmex septentrionalis* during and after a record drought. *Insect Conserv. Divers.* **3**, 134-142.
- Seal, J. N., Gus, J. and Mueller, U. G. (2012). Fungus-gardening ants prefer native fungal species: do ants control their crops? *Behav. Ecol.* **23**, 1250-1256.
- Semenova, T., Hughes, D., Boomsma, J. J. and Schiøtt, M. (2011). Evolutionary patterns of proteinase activity in attine ant fungus gardens. *BMC Microbiol.* **11**, 15.
- Sen, R., Ishak, H. D., Estrada, D., Dowd, S. E., Hong, E. and Mueller, U. G. (2009). Generalized antifungal activity and 454-screening of *Pseudonocardia* and *Amycolatopsis* bacteria in nests of fungus-growing ants. *Proc. Natl. Acad. Sci. USA* **106**, 17805-17810.
- Silva, A., Bacci, M., Jr, Gomes de Siqueira, C., Correa Bueno, O., Pagnocca, F. C. and Aparecida Hebling, M. J. (2003). Survival of *Atta sexdens* workers on different food sources. *J. Insect Physiol.* **49**, 307-313.
- Sokal, R. R. and Rohlf, F. J. (1995). *Biometry*. New York, NY: W. H. Freeman and Co.
- Stradling, D. J. and Powell, R. J. (1986). The cloning of more highly productive fungal strains: a factor in the speciation of fungus growing ants. *Experientia* **42**, 962-964.
- Suen, G., Scott, J. J., Aylward, F. O., Adams, S. M., Tringe, S. G., Pinto-Tomás, A. A., Foster, C. E., Pauly, M., Weimer, P. J., Barry, K. W. et al. (2010). An insect herbivore microbiome with high plant biomass-degrading capacity. *PLoS Genet.* **6**, e1001129.
- Torres, J. A., Santiago, M. and Salgado, M. (1999). The effects of the fungus-growing ant, *Trachymyrmex jamaicensis*, on soil fertility and seed germination in a subtropical dry forest. *Trop. Ecol.* **40**, 237-245.
- Underwood, A. J. (1997). *Experiments in Ecology*. Cambridge, UK: Cambridge University Press.
- Wade, M. (2007). The co-evolutionary genetics of ecological communities. *Nat. Rev. Genet.* **8**, 185-195.
- Weber, N. A. (1956). Fungus-growing ants and their fungi: *Trachymyrmex septentrionalis*. *Ecology* **37**, 150-161.
- White, T., Bruns, T., Lee, S. and Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR Protocols: A Guide to Methods and Applications* (ed. M. Innis, D. Gelfand, J. Sninsky and T. White), pp. 315-322. San Diego, CA: Academic Press.
- Wilson, E. O. (1980). Caste and division of labor in leaf-cutter ants (Hymenoptera: Formicidae: Atta). I. The overall pattern in *Atta sexdens*. *Behav. Ecol. Sociobiol.* **7**, 143-156.
- Wirth, R., Herz, H., Ryel, R. J., Beyschlag, W. and Hölldobler, B. (2003). *Herbivory of Leaf-Cutting Ants: A Case Study on Atta colombica in the Tropical Rainforest of Panama*. Berlin, Germany: Springer Verlag.




Figure S1. Preliminary phylogenetic relationships among the fungal lineages used in the experimental switches in our study. Phylogenetic relationships were inferred in a Bayesian analysis from a 475-nucleotide sequence of EF-1 α . Arrows indicate, respectively, the close phylogenetic proximity between *Attamyces* from Arizona and Texas, and between the *Trachymyces* from Arizona and Texas. In addition to other species in the USA (not shown), *Attamyces* is cultivated by *Acromyrmex versicolor*, *Atta texana* and by some *Trachymyrmex arizonensis* nests, whereas *Trachymyces* is cultivated by *T. arizonensis* and *T. septentrionalis*.

Phylogentic reconstruction from Ulrich G. Mueller, Heather D. Ishak, Sofia M. Brushi, Alexander S. Mikheyev, Scott E. Solomon, Jarrod J. Scott, Michael Cooper, Henrik H. DeFine Licht,

Stephen A. Rehner, Ted R. Schultz, Mauricio Bacci Jr. In prep. Biogeography of the leafcutter ant-fungus mutualism. Unpublished manuscript.




Figure S2. Mean xylanase activity (µg reducing sugar per mg/ml of fungus garden extract) in gardens for each experimental combination of ant (*Atta texana* or *Trachymyrmex arizonensis*) and fungal species (*Attamyces* or *Trachymyces*). No effects (ant, fungus, or interactions) were statistically significant (Table 1). Data were analyzed on log₁₀ transformed data. Error bars are +/- 1 standard deviation. All effects except fungal symbiont species were significantly heteroscedastic.




Figure S3. Mean protease activity (relative absorbance per mg/ml of fungus garden extract) in gardens for each experimental combination of ant (*Atta texana* or *Trachymyrmex arizonensis*) and fungal species (*Attamyces* or *Trachymyces*). *Attamyces* gardens showed significant higher protease activity than *Trachymyces* gardens. Ant-host species had no statistically significant effect on protease activity. Although there was a significant interaction between ant-host and fungus ($F_{1,19} = 5.95$, $p=0.025$; gardens of *T. arizonensis* growing *Attamyces* exhibiting higher activities than when growing *Trachymyces* (Scheffé's test)), variances were heteroscedastic and were thus pooled (Table 1). Data were analyzed on \log_{10} transformed data. Error bars are ± 1 standard deviation. All effects except fungal symbiont species were significantly heteroscedastic.