

RESEARCH ARTICLE

Casein kinase 1 α decreases β -catenin levels at adherens junctions to facilitate wound closure in *Drosophila* larvae

Chang-Ru Tsai¹ and Michael J. Galko^{1,2,3,*}**ABSTRACT**

Skin wound repair is essential to restore barrier function and prevent infection after tissue damage. Wound-edge epidermal cells migrate as a sheet to close the wound. However, it is still unclear how cell-cell junctions are regulated during wound closure (WC). To study this, we examined adherens junctions during WC in *Drosophila* larvae. β -Catenin is reduced at the lateral cell-cell junctions of wound-edge epidermal cells in the early healing stages. Destruction complex components, including Ck1 α , GSK3 β and β -TrCP, suppress β -catenin levels in the larval epidermis. Tissue-specific RNAi targeting these genes also caused severe WC defects. The Ck1 α^{RNAi} -induced WC defect is related to adherens junctions because loss of either β -catenin or E-cadherin significantly rescued this WC defect. In contrast, TCF^{RNAi} does not rescue the Ck1 α^{RNAi} -induced WC defect, suggesting that Wnt signaling is not related to this defect. Direct overexpression of β -catenin recapitulates most of the features of Ck1 α reduction during wounding. Finally, loss of Ck1 α also blocked junctional E-cadherin reduction around the wound. Our results suggest that Ck1 α and the destruction complex locally regulate cell adhesion to facilitate efficient wound repair.

KEY WORDS: Wound repair, Adherens junctions, Epithelium, Casein kinase 1 α , β -Catenin, *Drosophila*

INTRODUCTION

Drosophila studies in wound repair and regeneration have provided many insights at the molecular and cellular levels (Nakamura et al., 2018; Tsai et al., 2018; Zulueta-Coarasa and Fernandez-Gonzalez, 2017). Different signaling pathways and actin regulators play crucial roles during this process (Tsai et al., 2018). To close a wound gap, the epidermal cells become motile but at the same time maintain their adhesive contacts with their neighbors. How cells balance motility with adhesion is not yet clear.

Adherens junctions (AJs) are calcium-dependent adhesion complexes that are important for holding cells together within diverse epithelial tissues (Pinheiro and Bellaïche, 2018). Analysis of AJ function *in vivo* during vertebrate wound healing is complicated by the fact that loss of cadherins is generally lethal

(Larue et al., 1994; Riethmacher et al., 1995). During fly embryonic wound healing, AJ components, including E-cadherin, β -catenin and α -catenin, are redistributed around the wound margin (Abreu-Blanco et al., 2013; Hunter et al., 2015; Matsubayashi et al., 2015; Wood et al., 2002; Zulueta-Coarasa et al., 2014). Specifically, AJ components are decreased at the interfaces between wound-edge cells and increased at the wound-edge cellular junctions where these cells are joined (Hunter et al., 2015; Matsubayashi et al., 2015). E-cadherin levels around embryonic wounds are negatively regulated by endocytosis (Hunter et al., 2015), which is in turn required for normal healing (Hunter et al., 2015; Matsubayashi et al., 2015). E-cadherin overexpression delays WC and reduces actin protrusions (Hunter et al., 2015). E-cadherin is also regulated transcriptionally by Toll/NF κ B signaling (Carvalho et al., 2014). Whether regulation of AJ levels and function are important in post-embryonic healing, which employs directed cell migration over contraction of an actin cable (Tsai et al., 2018), is not yet clear.

In addition to its essential role at the adherens junction, β -catenin is also the downstream transcriptional co-activator of Wnt/Wingless signaling. Wnt signaling regulates regenerative repair in *Drosophila* imaginal discs (Hariharan and Serras, 2017; Schubiger et al., 2010; Smith-Bolton et al., 2009). In the absence of Wnt ligand, cytoplasmic β -catenin is phosphorylated and ubiquitinated by a protein complex termed the β -catenin destruction complex. This complex consists of casein kinase 1 alpha (Ck1 α), glycogen synthase kinase 3 β (GSK3 β), Axin, adenomatous polyposis coli (APC), protein phosphatase 2A (PP2A) and the E3-ubiquitin ligase β -TrCP. Within the β -catenin destruction complex, Ck1 α phosphorylates β -catenin and promotes its degradation (Yanagawa et al., 2002; Zhang et al., 2006). Ubiquitinated β -catenin is subsequently degraded by the proteasome (Aberle et al., 1997). Upon Wnt ligand-receptor binding, β -catenin is released from the destruction complex and translocates to the nucleus where it binds to the transcription factor, T cell factor (TCF or Pangolin in *Drosophila*), to activate downstream gene expression (Hecht and Kemler, 2000). The role of Wnt signaling and the destruction complex in larval WC have not been tested to date.


We investigated the role of β -catenin during larval epidermal WC and found that junctional β -catenin is locally reduced after wounding on membranes radial to the wound site. We further showed that the β -catenin destruction complex is crucial to downregulate junctional β -catenin and facilitate WC. The Ck1 α^{RNAi} -induced WC defects are related to its functions at the adherens junctions but not to roles in Wnt signaling.

RESULTS**Reduction of junctional β -catenin in wound-edge epidermal cells**

To examine AJs during wound healing, larvae were poke-wounded (see Materials and Methods), a procedure that produces small

¹Program in Developmental Biology, Baylor College of Medicine, Houston, TX 77030, USA. ²Department of Genetics, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA. ³Genetics & Epigenetics Graduate Program, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

*Author for correspondence (mjgalko@mdanderson.org)

 C.-R.T., 0000-0001-5672-1235; M.J.G., 0000-0002-3759-2017

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

and largely symmetrical wounds. Larval whole mounts were immunostained using anti- β -catenin antibodies (Riggleman et al., 1990) at various times after wounding. In unwounded larvae, β -catenin was apparent at cell-cell junctions (compare membrane-GFP and β -catenin staining in Fig. 1A,A' and Fig. S1A,A'). This signal is specific because expression of an RNAi transgene targeting β -catenin significantly reduced it without reducing an independent membrane label (Fig. S1B',D). Epidermal expression of an *E-cadherin* RNAi transgene also decreased β -catenin levels at the junctions (Fig. S1C',D), and vice versa (Fig. S1E,F), indicating that junctional β -catenin depends upon E-cadherin expression as previously reported (Pai et al., 1996). Ten minutes after wounding, β -catenin was still apparent at the interface between wound-edge epidermal cells (Fig. 1B', arrows). However, β -catenin at most interfaces radial to and immediately proximal to the wound was reduced 1 h (Fig. 1C', arrowheads) and 2 h (Fig. 1D', arrowheads) after wounding. After WC was complete at 5 h (Fig. 1E-F), β -catenin staining became diffuse as the epidermal cells presumably remodeled their junctions. Comparison of the ratio of lateral β -catenin levels in first row versus second row cells (Fig. 1G) revealed a clear decrease over the first hour after wounding (Fig. 1H). These results suggest that regulation of β -catenin levels near the wound may be important during epidermal WC.

The destruction complex regulates junctional β -catenin and is required for wound closure

Because β -catenin levels are decreased near the wound (Fig. 1C',D',H), we tested whether the destruction complex regulates β -catenin levels in the larval epidermis. We used immunofluorescence to examine β -catenin levels in the larval epidermis expressing an RNAi transgene targeting *Ckl1 α* , the key destruction complex kinase that promotes β -catenin degradation. β -Catenin was significantly increased when the larval epidermis expressed *Ckl1 α ^{RNAi}* (Fig. 2B,E) compared with a control RNAi transgene (Fig. 2A,E). Intriguingly, β -catenin upregulation was particularly apparent at AJs. The junctional β -catenin (Fig. S2D,E) and E-cadherin (Fig. S2G,H) were also increased in larval epidermis expressing the *Ckl1 α ^{RNAi}* transgene via another epidermal driver (*e22c-Gal4*) compared with the control (Fig. S2C,E,F,H).

We next measured whether the increase in β -catenin impacted WC. Expression of *Ckl1 α ^{RNAi}* also caused a strong WC defect (Fig. 2G,J). The WC defect was also observed in larvae expressing an RNAi transgene targeting a non-overlapping region of the *Ckl1 α* sequence (Fig. S2A,B), arguing strongly against RNAi off-target effects. These results suggest that *Ckl1 α* negatively regulates β -catenin to facilitate WC. To test whether GSK3 β and β -TrCP, other destruction complex components, also regulate β -catenin, we




Fig. 1. Junctional β -catenin in wound-edge epidermal cells is reduced after wounding. (A-E') Dissected epidermal whole mounts of unwounded (A,A') or poke-wounded (B-E') third instar larvae expressing *UAS-DsRed2nuc* (nuclei, magenta) and *UAS-src-GFP* (cell membranes, green) via the *A58-Gal4* driver 10 min (B,B'), 1 h (C,C'), 2 h (D,D') and 5 h (E,E') after wounding. (A-E) The nuclei and cell membrane. (A'-E') The adherens junctions of the same samples immunostained using anti- β -catenin antibodies (white). Scale bar: 50 μ m. Dotted yellow lines indicate wound borders. Arrows in B,B' highlight examples of clear junctional β -catenin signal (B') and membrane-GFP signal (B). Arrowheads in C-D' highlight examples of reduced junctional β -catenin (C',D') where membrane-GFP is still present (C,D). (F) Quantitation of open poke wounds in control larvae. The epidermal reporter used was *e22c-Gal4, UAS-LifeAct-Cherry, UAS-luciferase^{RNAi}*. $n \geq 20$ for each time point. (G) Schematic of quantitation strategy for measuring β -catenin levels on lateral segments near the wound. (H) Quantitation of junctional β -catenin ratio in first- and second-row wound-edge epidermal cells. Each dot represents one larva. Data are mean \pm s.e.m. **** $P < 0.0001$ (unpaired *t*-test).




Fig. 2. The destruction complex regulates junctional β -catenin levels and wound closure. (A–D) Dissected epidermal whole mounts of unwounded third instar larvae expressing *UAS-DsRed2nuc* (nuclei, magenta), *UAS-src-GFP* (cell membranes, green, not shown) and the indicated transgenes via either the *e22c-Gal4* (A,C,D) or *A58-Gal4* drivers (B). Anti- β -catenin antibody staining is in white. (A) *Control^{RNAi}*. (B) *Ck1α^{RNAi#1}* (*A58-Gal4* used because this line grows slowly with *e22c-Gal4*). (C) *GSK3β^{RNAi}*. (D) *β-TrCP^{RNAi}*. (E) Quantitation of junctional β -catenin intensity in larvae expressing the different transgenes. Each dot represents an average of the β -catenin signal from five junctions of one larva. Data are mean \pm s.e.m. * P <0.05 (one-way ANOVA). (F–I) Dissected epidermal whole mounts of pinched-wounded third instar larvae expressing *UAS-DsRed2nuc* (nuclei, magenta) via the *e22c-Gal4* driver and the indicated RNAi transgenes. Cell boundaries were immunostained using anti-Fasciclin III antibodies (green). Scale bars: 50 μ m in A for A–D; 100 μ m in F for F–I. (F) *Control^{RNAi}*. (G) *Ck1α^{RNAi}*. (H) *GSK3β^{RNAi}*. (I) *β-TrCP^{RNAi}*. (J) Quantitation of the percentage of open wounds in larvae expressing the indicated transgenes via the *e22c-Gal4* driver. Each dot represents one set of $n \geq 8$ larvae for each genotype. Data are mean \pm s.e.m. * P <0.05, ** P <0.01 (one-way ANOVA).

expressed RNAi transgenes targeting them in the larval epidermis. Larvae expressing RNAi transgenes targeting *GSK3β* or β -TrCP also exhibited a significant increased in β -catenin (Fig. 2C–E) and WC defects (Fig. 2H–J). These results indicate that multiple components of the destruction complex regulate β -catenin levels and epidermal WC.

The wound closure defect observed with *Ck1α^{RNAi}* is β -catenin dependent

If β -catenin is a crucial downstream target of *Ck1α* during wound healing, then silencing β -catenin should reduce *Ck1α^{RNAi}*-induced WC defects. Continuous epidermal expression of the RNAi line targeting *Ck1α* that can be genetically combined with other transgenes is lethal (RNAi#3 in Fig. S2A). Therefore, we used a

temperature-sensitive allele of *Gal80* (McGuire et al., 2004) to temporally control transgene expression (see experimental schematic in Fig. 3A). We also verified that the *β-catenin^{RNAi}* transgene indeed knocked down β -catenin in this inducible paradigm (Fig. S3D–F). *Ck1α^{RNAi}* also increased β -catenin levels in the cytoplasm (compare Fig. S3C with S3B). Junctional E-cadherin was similarly upregulated in the epidermis expressing the *Ck1α^{RNAi}* transgene (Fig. S3H,K) compared with the control (Fig. S3G,K), although the trend of increased junctional β -catenin did not reach significance (Fig. S3B,C,F). Importantly, inducible expression of *Ck1α^{RNAi}* also caused a strong WC defect that was unaffected by expression of a control RNAi transgene (Fig. 3C,F); this defect was rescued by co-expressing *UAS-Ck1α-HA* (Fig. 3F), further suggesting the WC defect is due to loss of *Ck1α* rather than




Fig. 3. Silencing β -catenin partially rescues the $Ck1\alpha^{RNAi}$ -induced wound closure defect. (A) Schematic of the experimental design/temperature shift regimen for using Gal80^{TS} to inducibly express UAS-dependent transgenes in the larval epidermis. (B–E) Dissected epidermal whole mounts of pinch-wounded third instar larvae expressing Gal80^{TS} transgene driven by a tubulin promoter, *UAS-DsRed2Nuc* (nuclei, magenta) via the *e22c-Gal4* driver, and the indicated transgenes. Cell boundaries were immunostained using anti-Fasciillin III antibodies (green). (B) Control^{RNAi}, (C) *UAS-Ck1α^{RNAi#3}* and control^{RNAi}, (D) *UAS-β-catenin^{RNAi}*, (E) *Ck1α^{RNAi#3}* and *UAS-β-catenin^{RNAi}*. Scale bar: 100 μm. (F) Quantitation of the percentage of open wounds in larvae expressing indicated transgenes via the *e22c-Gal4* driver. Each dot represents one set of $n \geq 8$ for each genotype. Data are mean \pm s.e.m. * $P < 0.05$, ** $P < 0.01$ (one-way ANOVA).

to off-target effects. Silencing β -catenin by itself did not block WC (Fig. 3D,F) despite a strong reduction in β -catenin (Fig. S3D). However, knocking down β -catenin significantly ameliorated the WC defect observed upon expression of *Ck1α^{RNAi}* (Fig. 3E,F). These results suggest that the WC defect observed upon expression of *Ck1α^{RNAi}* is dependent on β -catenin levels in the larval epidermis.

***Ck1α^{RNAi}*-induced wound closure defects are related to its functions at the adherens junction rather than to Wnt signaling**

As β -catenin is an important downstream transcriptional co-activator of Wnt signaling, we also tested whether Wnt signaling is activated upon wounding. Larvae bearing either *fz3-LacZ* or *6TH-LacZ*, two Wnt signaling reporters (Chang et al., 2008; Sato et al., 1999), were wounded and examined 4 h after wounding. The *fz3-LacZ* reporter was functional as it was activated in the developing wing imaginal discs (Fig. S4A'). However, *fz3-lacZ* was not activated after wounding (compare wounded epidermis in Fig. S4D with unwounded epidermis in Fig. S4A). Similarly, *6TH-LacZ* was also expressed in the wing discs (Fig. S4B') but not activated in the epidermis after wounding (Fig. S4E versus control in S4B). The JNK reporter, *msn-LacZ*, a positive control, was activated around the wound margin after wounding (Fig. S4C,F) as previously reported (Galko and Krasnow, 2004; Lesch et al., 2010). To test whether Wnt signaling is activated in the epidermis expressing a *Ck1α^{RNAi}* transgene, we also examined these two Wnt reporters. The *fz3-LacZ* reporter did not change upon loss of *Ck1α* (Fig. S4I). However, *6TH-LacZ* was activated in the epidermis expressing *Ck1α^{RNAi}* transgene (Fig. S4J), suggesting that the Wnt signaling is activated in this condition.

β -Catenin has roles in both adhesion (Ozawa et al., 1989) and as a transducer of Wnt signaling (Wieschaus et al., 1984). Therefore, we tested which of these two functions impact WC. First, to test genetically whether the *Ck1α^{RNAi}*-induced WC defect is related to a role at AJs (as opposed to some other functions of β -catenin), we co-expressed an RNAi transgene targeting E-cadherin, the core component of AJs, to determine whether it could rescue the WC defect observed upon *Ck1α^{RNAi}* expression. We again used Gal80^{TS} to temporally control the Gal4 driver activity (see experimental schematic in Fig. 4A). Control larvae closed their wounds 24 h after pinch wounding (Fig. 4B,H), while epidermal expression of *Ck1α^{RNAi}* caused a strong WC defect (Fig. 4C,H). As with *β-catenin^{RNAi}* (Fig. 3D,F), epidermal expression of an *E-cadherin^{RNAi}* transgene by itself did not block WC (Fig. 4D,H). This *E-cadherin^{RNAi}* transgene is functional as constitutive expression dramatically reduced junctional E-cadherin (Fig. S1G,H) and β -catenin staining (Fig. S1C',D). The *E-cadherin^{RNAi}* transgene also worked in this inducible paradigm (Fig. S5D,F for β -catenin and Fig. S5I,K for E-cadherin). Importantly, knocking down E-cadherin significantly reduced *Ck1α^{RNAi}*-induced WC defects (Fig. 4E,H), indicating that the WC defect induced by *Ck1α^{RNAi}* is related to its regulation of AJs.

To test genetically whether the *Ck1α^{RNAi}*-induced WC defect might also be related to a role in Wnt signaling, we co-expressed RNAi transgenes targeting T cell factor (TCF). Epidermal expression of the *TCF^{RNAi}* transgene by itself did not block WC (Fig. 4F,H). This *TCF^{RNAi}* transgene is functional as its expression in the wing pouch via *nubbin-Gal4* disrupted normal wing development (compare Fig. S5L with S5M), consistent with interference with Wnt signaling. Epidermal expression of a




Fig. 4. The epidermal *Ck1α*^{RNAi}-induced wound-closure defect is E-cadherin dependent and Wnt signaling independent. (A) Schematic of the experimental design/temperature shift regimen for using Gal80^{ts} to inducibly express UAS-dependent transgenes in the larval epidermis. (B–G) Dissected epidermal whole mounts of pinch-wounded larvae expressing the Gal80^{ts} transgene driven by a tubulin promoter, *UAS-DsRed2Nuc* (nuclei, magenta) via the *e22c-Gal4* driver, and the indicated transgenes 24 h after wounding. Cell boundaries were immunostained using anti-Fasciillin III antibodies (green). (B) Control^{RNAi}, (C) *Ck1α*^{RNAi#3} and control^{RNAi}, (D) *E-cad*^{RNAi}, (E) *Ck1α*^{RNAi#3} and *E-cad*^{RNAi}, (F) *UAS-TCF*^{RNAi}, (G) *Ck1α*^{RNAi} and *UAS-TCF*^{RNAi}. Scale bar: 100 μm. (H) Quantitation of the percentage of open wounds in third instar larvae expressing the indicated transgenes via the *e22c-Gal4* driver. Each dot represents one set of $n \geq 8$ larvae for each genotype. Data are mean \pm s.e.m. ** $P < 0.01$; ns, not significant (one-way ANOVA).

TCF^{RNAi} transgene did not ameliorate the WC defect induced by *Ck1α*^{RNAi} expression (Fig. 4G,H), suggesting that Wnt signaling is not required for the *Ck1α*^{RNAi}-induced WC defect.

β-Catenin overexpression also caused wound closure defects related to its functions at the adherens junction

An alternative way to test the role of β-catenin during WC is to directly overexpress it in the larval epidermis (see schematic Fig. 5A). Similarly to RNAi transgenes targeting β-catenin-negative regulators (Fig. 2G–J), β-catenin overexpression also blocked WC (Fig. 5C,H) relative to controls (Fig. 5B,H). Because β-catenin is a key component of Wnt signaling, we tested whether β-catenin overexpression activates Wnt signaling in the larval epidermis. The Wnt reporter, *fz3-LacZ* (Fig. S4K), was not activated upon β-catenin overexpression. However, another Wnt reporter, *6TH-LacZ*, was activated (Fig. S4L). If β-catenin-induced Wnt activation is responsible for its WC defect, silencing TCF should rescue the observed WC defect. However, epidermal expression of the *TCF*^{RNAi} transgene did not block the β-catenin-induced WC defect (Fig. 5E,H).

To test whether the WC defect caused by β-catenin overexpression is related to its function at AJs, we wounded larvae expressing both *E-cadherin*^{RNAi} and *β-catenin* transgenes in the larval epidermis.

Interestingly, silencing E-cadherin significantly reduced the β-catenin-induced WC defects (Fig. 5G,H). To test the impact of β-catenin overexpression on E-cadherin, we performed immunostaining using an E-cadherin antibody. Surprisingly, overexpression of β-catenin did not increase E-cadherin (Fig. S6H,K) as with the *Ck1α*^{RNAi} transgene (Figs S3H,K and S5H,K). We also examined the sub-cellular locations of β-catenin in these animals and found that although β-catenin is highly expressed in the nuclear and cytoplasmic compartments (Fig. S6C), its levels on the membrane were modest.

Actin distribution in *Ck1α*^{RNAi} and β-catenin overexpression

Both *Ck1α*^{RNAi} and β-catenin overexpression cause WC defects that are dependent on the levels of E-cadherin. However, these genetic manipulations differ in the levels and cellular distributions of both β-catenin and E-cadherin. Because β-catenin is a key protein that bridges E-cadherin at AJs with the actin cytoskeleton, we checked the distribution of F-actin in these two genotypes. In the control epidermis, the F-actin label LifeAct-mCherry is evenly distributed within epidermal cells and membrane levels are low (Fig. 6A). Strikingly, β-catenin overexpression significantly increased cortical actin (Fig. 6B) while expression of the *Ck1α*^{RNAi} transgene did not (Fig. 6C). These results suggest that although both β-catenin overexpression and *Ck1α*^{RNAi} caused E-cadherin-dependent WC




Fig. 5. Epidermal β -catenin overexpression-induced wound closure defect is E-cadherin dependent and Wnt signaling independent. (A) Schematic of the experimental design/temperature shift regimen for using Gal80^{ts} to inducibly express UAS-dependent transgenes in the larval epidermis. (B-G) Dissected epidermal whole mounts of pinch-wounded larvae expressing the Gal80^{ts} transgene driven by a tubulin promoter, *UAS-DsRed2Nuc* (nuclei, magenta) via the *e22c-Gal4* driver, and the indicated transgenes 14 h after wounding. Cell boundaries were immunostained using anti-Fasciclin III antibodies (green). (B) *Control^{RNAi}*, (C) *UAS-β-catenin* and *control^{RNAi}*, (D) *E-cad^{RNAi}*, (E) *UAS-β-cat* and *E-cad^{RNAi}* (F), *UAS-TCF^{RNAi}*, (G) *UAS-β-cat* and *UAS-TCF^{RNAi}*. Scale bar: 100 μ m. (H) Quantitation of the percentage of open wounds in third instar larvae expressing the indicated transgenes via the *e22c-Gal4* driver. Each dot represents one set of $n \geq 8$ larvae for each genotype. Data are mean \pm s.e.m. ****P* < 0.001; ns, not significant (one-way ANOVA).

defects, the precise mechanism of their wound closure defects may differ at the level of actin dynamics.

Ck1 α is required for reduction of lateral E-cadherin around the wound

Finally, we tested whether Ck1 α is important to reduce the lateral AJs in the wound-edge cells (Fig. 1), which was our initial observation. We examined E-cadherin-GFP levels 1 h after poking. Similar to β -catenin, lateral E-cadherin around the wounds was reduced 1 h after poke wounding (Fig. 6D,D',F). However, this reduction was ameliorated in the epidermis expressing a *Ck1 α ^{RNAi}* transgene (Fig. 6E-F), indicating that *Ck1 α* is involved in local downregulation of lateral junctions near the wound.

DISCUSSION

In summary, four lines of experimental evidence support the hypothesis that regulation of junctional β -catenin is crucial for epidermal WC in *Drosophila* larvae. First, junctional β -catenin is reduced in wound-edge epidermal cells during WC. Second, RNAi transgenes targeting the β -catenin destruction complex caused upregulation of junctional β -catenin and strong WC defects. Third, the *Ck1 α ^{RNAi}*-induced WC defect was reduced by silencing E-cadherin, indicating that this defect is related to its function at AJs. Finally, *Ck1 α ^{RNAi}* impedes removal of the lateral E-cadherin from

lateral junctions radial to the wound. β -Catenin overexpression also caused AJ-dependent WC defects although the mechanism may be distinct given differences in E-cadherin levels, and β -catenin and actin distributions. In summary, regulation of cell-cell interactions through AJs may be a crucial mechanism for coordinating wound-induced cell sheet migration.

In the fly embryo, loss of E-cadherin delays WC (Abreu-Blanco et al., 2013). However, silencing E-cadherin did not block WC in third instar larvae. Similarly, knocking down β -catenin did not cause a WC defect, indicating that the requirement of adherens junctions varies at different developmental stages. Given that the embryonic epidermis has not yet secreted cuticle and that the larval epidermis is tightly adherent to the apical cuticle it secretes (Galko and Krasnow, 2004), some differences in adhesion requirements might be expected. The normal morphology of the E-cadherin and β -catenin-deficient larval epidermis is surprising given how important these proteins are for tissue morphology in development. E-cadherin loss in the mouse causes early embryonic lethality as epithelial tissues fail to form properly (Larue et al., 1994; Riethmacher et al., 1995). Likewise, strong E-cadherin alleles in the fly disrupt early morphogenesis (Oda et al., 1994; Tepass and Hartenstein, 1994). E-cadherin loss in skin so profoundly affects morphology (Tinkle et al., 2008) that wound healing studies are essentially precluded, while β -catenin loss using the same *keratin-14 Cre* driver causes defects in the hair follicle cycle




Fig. 6. Actin localization and Ck1 α are required for the reduction of lateral E-cadherin on wound-edge epidermal cell membranes. (A–C) Dissected larval epidermal whole mounts of third instar larvae expressing *UAS-Lifeact-mCherry* (magenta) and the indicated transgenes via the *e22c-Gal4* driver. (D–E') Live image of larval epidermis expressing *E-cadherin-GFP* and *UAS-LifeAct-mCherry* (magenta) via *e22c-Gal4* driver (control, D,D') and (*Ck1 α ^{RNAi}*, E,E') 1 h after wounding. (D,E) *E-cadherin-GFP*, green. (D'–E') *F-actin* to visualize the wound margins. Arrows and arrowheads in D,E indicate examples of first-row and second-row junctions, respectively. Scale bars: 50 μ m. Dotted white lines indicate wound borders. (F) Quantitation of junctional *E-cadherin-GFP* ratio in first- and second-row wound-edge epidermal cells. Each dot represents one animal. Data are mean \pm s.e.m. ** P <0.01; ns, not significant (unpaired t -test).

(Huelsen et al., 2001). Conditional knockout of mouse *Ck1 α* causes hyperplasia and pigmentation defects in skin (Chang et al., 2017), and these mice have not been analyzed for wound healing phenotypes.

In the laser-wounded fly embryo, *E-cadherin* is reduced at the wound perimeter and the lateral interfaces between neighboring wound-edge epidermal cells (Hunter et al., 2015). *E-cadherin* level is simultaneously increased at the vertices between neighboring wound-edge cells along the wound margin. In both embryos (Hunter et al., 2015) and larvae (this study), ubiquitous epidermal upregulation of AJ components interferes with effective WC. Although *Ck1 α* is required for the local reduction of *E-cadherin* near wounds, the specific function of wound-edge adherens junction downregulation in embryos and larvae remains technically challenging to test. During collective cell migration in culture, lateral *E-cadherin* increases tension and slows migration (Suffoletto et al., 2018). A reduction of *E-cadherin* at the migratory leading edge of mouse wounds, similar to that seen here, has been observed (Hudson et al., 2009). At the functional level, loosening of junctions and reduction of *E-cadherin* at the leading edge through ephrin signaling appears important for mouse keratinocytes to efficiently move into the wound gap (Nunan et al., 2015). Thus, at least in principle, reduction of junctional components at the lateral interface may reduce tension between wound-edge cells to facilitate efficient WC.

What regulates the redistribution of adherens junctions? *E-cadherin* genetically interacts with dynamin, an essential regulator of endocytosis, during WC in the fly embryo (Matsubayashi et al., 2015). Notably, endocytosis is also essential for wound healing in the fly embryo (Hunter et al., 2015; Matsubayashi et al., 2015) and *E-cadherin* reduction around the wound is partially endocytosis dependent (Hunter et al., 2015). Endocytosis may also regulate β -

catenin levels in wounded larvae but there is at least one other mechanism, identified here, that seems to do this. That is the activity of the destruction complex – several members of which (*Ck1 α* , *GSK3 β* and β -TrCP), regulate epidermal β -catenin levels and are also required for WC. It is curious that the destruction complex is required for WC but Wnt signaling is not involved. Here, neither knockdown of *TCF* nor β -catenin caused a WC defect, indicating that the transcriptional output of Wnt signaling is not required for WC. The requirement for the destruction complex during WC suggests the possibility that some other signaling pathway required for WC (Baek et al., 2012; Brock et al., 2012; Galko and Krasnow, 2004; Kakanj et al., 2016; Stevens and Page-McCaw, 2012; Tsai et al., 2017; Wu et al., 2009) may regulate β -catenin levels through this complex.

AJs maintain tissue integrity and coordinate cell sheet migration, as these cells are mechanically linked (Pinheiro and Bellaïche, 2018). In embryos, loss of *E-cadherin* impairs actomyosin ring formation and delays WC (Abreu-Blanco et al., 2013). Interestingly, we did not observe WC defects in larvae lacking epidermal *E-cadherin* or β -catenin. What accounts for this stage-dependent difference? Occluding (septate) junctions are also required for efficient WC in fly embryos (Carvalho et al., 2018) as are some Integrin complex components in fly larvae (Park et al., 2018). A specific role for occluding junctions in larval WC has not yet been tested. It is also possible that the apical cuticular attachment of larval epidermal cells mechanically stabilizes them in such a way that makes them less sensitive to loss of AJ functions.

During wound healing in fly embryos, gain of *E-cadherin* reduces wound-edge actin intensity and cell protrusions (Hunter et al., 2015). This suggests an inverse relationship between AJ levels and

wound-edge actin polymerization. In embryos and in single-cell wounds in the larval epidermis the formation of an actomyosin ring is important for WC (Abreu-Blanco et al., 2013; Kakanj et al., 2016). Pinch-wounded larvae polymerize actin at the wound edge in a discontinuous fashion (Brock et al., 2012; Tsai et al., 2017) and larval epidermal cells appear to rely more on protrusive migration (Wu et al., 2009) during WC. Here, we observed that larval epidermal cells overexpressing β -catenin have a higher basal level of cortical actin, which may account, in part, for their inability to efficiently close wounds. It will be interesting to test whether gain of β -catenin also impacts wound-edge actin polymerization and cell protrusion activity during epidermal WC in larvae.

MATERIALS AND METHODS

Fly genetics

Flies were raised on regular corn meal media. All crosses were raised at 25°C unless indicated. The GAL4/UAS system was used to drive tissue-specific gene expression of transgenes under UAS control (Brand and Perrimon, 1993). For the embryonic and larval epidermis, *e22c-Gal4* was used (Lawrence et al., 1995); for the larval epidermis, *A58-Gal4* was used (Galko and Krasnow, 2004). For fly wing imaginal discs, *nubbin-Gal4* was used (Calleja et al., 1996). For the wound closure assay, we used *e22c-Gal4*, *UAS-src-GFP*, *UAS-DsRed2-Nuc* or *A58-Gal4*, *UAS-src-GFP*, *UAS-DsRed2Nuc* (Lesch et al., 2010). *UAS-RNAi* lines from the TRiP Bloomington collection were: *JF01792 (Ck1 α ^{RNAi#1})*, *GL00021 (Ck1 α ^{RNAi#2})*, *HMS02276 (Ck1 α ^{RNAi#4})*, *JF02306 (TCF^{RNAi})*, *JF01252 (β -catenin^{RNAi})*, *HMS00693 (E-cadherin^{RNAi})*, *JF01504 (β -TrCP^{RNAi})* and *JF01355 (Luciferase^{RNAi})*. *UAS-RNAi* lines from the Vienna Drosophila Research Center were: *GD4256 (Ck1 α ^{RNAi#3})* and *KK108994 (GSK3 β ^{RNAi})*. To enhance *GSK3 β ^{RNAi}* and *β -TrCP^{RNAi}* knockdown efficiency, larvae were raised at 29°C. Other transgenic lines from Bloomington Stock Center were: #8369, *P{UAS-arm.Exel}2 (UAS- β -catenin)*; #60584, *TI{TI}shg^{GFP}*; and #8529, *P{UAS-lacZ.Exel}2*. *LacZ* reporters used were: *msn-LacZ* (Spradling et al., 1999); *fz3-LacZ* (Sato et al., 1999); *6TH-LacZ* (Chang et al., 2008). *UAS-LifeAct-mCherry* (Riedl et al., 2008) was used to label actin.

In cases where early expression of UAS transgenes was unhealthy (combination of *UAS- β -catenin* and *UAS-E-cadherin^{RNAi}*) or lethal (*UAS-Ck1 α ^{RNAi#3}*), larvae bearing *tubP-gal80^S* (McGuire et al., 2003), *e22c-Gal4*, *UAS-src-GFP*, *UAS-DsRed2Nuc* and toxic UAS transgene were raised for 6 days at 18°C to begin development, shifted to 32°C for 2 days to reach mid-third-instar, and then allowed to recover at 25°C following pinch wounding.

A list of the genotypes of the flies used in each figure is provided in the supplementary Materials and Methods.

Wounding techniques

Pinch wounding of the mid-third instar larvae was carried out according to our detailed protocol (Burra et al., 2013). Pinch wounds were scored as 'open' if the initial wound gap remained after 24 h (or at the indicated timepoint) and as 'closed' if a continuous epidermal sheet was observed at the wound site. To calculate the percentage of larvae with open wounds, three sets of $n \geq 8$ per genotype were pinched and scored for open wounds under a stereo microscope (Leica MZ16FA).

Poke wounding was carried out using a homemade mechanical probe similar to Von Frey filaments, which generate a constant force when bent. These probes were developed by Roger Lopez-Bellido, Patrick Huang and Thomas Wang in the Galko laboratory and are described elsewhere (Lopez-Bellido et al., 2019). To make a poke wound, the mechanical probe (0.004 inch in diameter/2346 kpa pressure applied) was applied to the dorsal side of the mid-third instar larva, abdominal segment 2-4 (A2-4), until the filament bends for ~1 s. Poked larvae were dissected and stained at various time points after wounding.

Whole-mount immunofluorescence and LacZ staining

The third instar larval epidermis was dissected, fixed and immunostained as detailed previously (Burra et al., 2013). To highlight epidermal cell

boundaries (septate junctions), a mouse monoclonal antibody against Fasciclin III was used (1:50; Developmental Studies Hybridoma Bank) (Patel et al., 1987). Rabbit anti-DsRed (Clontech) was used at 1:1000. Mouse-anti- β -catenin was used at 1:75 (Developmental Studies Hybridoma Bank, N2 7A1) (Riggleman et al., 1990). X-Gal staining was performed as described previously (Galko and Krasnow, 2004). X-Gal development at 37°C was 30 min for *msn-LacZ*, overnight for *fz3-LacZ* and 20 min for *6TH-LacZ*.

Live imaging

Third instar larvae expressing E-cadherin-GFP and *UAS-LifeAct-mCherry* and different UAS transgenes via *e22c-Gal4* were etherized and poke wounded before mounting. 15 min and 1 h after wounding, larvae were etherized, immobilized and immersed in 1:4 (v/v) diethyl ether to halocarbon oil (Sigma, H88098) and covered with glass coverslips. An Olympus FV1000 Confocal microscope and Fluoview software were used to acquire images (Das et al., 2017).

Imaging analysis

An Olympus FV1000 Confocal microscope and Fluoview software were used to acquire images of the dissected epidermal whole mounts. Leica MZ16FA stereomicroscope with Planapo 1.6 \times objective and appropriate filters was used for live imaging of epidermal wounds. ImageJ software was used for image processing.

Quantitation and statistical analysis

Junctional β -catenin levels of unwounded third instar larvae of different genotypes were measured as follows: in ImageJ, a single line was randomly drawn along the anterior-posterior axis across six epidermal cells (cell tri-junctions were avoided). The peak signals at each junction were measured and averaged for each larva. The intensity of different genotypes was normalized to the *UAS-Luciferase^{RNAi}* control. Junctional β -catenin signal was greatly reduced in larvae expressing *UAS- β -catenin^{RNAi}* or *UAS-E-cadherin^{RNAi}* transgenes. Thus, to measure the junctional β -catenin intensities of those larvae, membrane GFP (*UAS-src-GFP*) signal was used to define the cell-cell junctions. The measured junctional β -catenin intensities were further subtracted by the mean intensity of cytoplasmic β -catenin of the same larva. Junctional β -catenin levels in poke-wounded third instar larvae at different time points were measured as follows: in ImageJ, junctional β -catenin intensities of four randomly selected lateral interfaces within the first row or the second row of wound-edge epidermal cells were measured and averaged. The ratios of averaged first row to second row wound-edge cells were compared between different time points. Unpaired two-tailed *t*-test (two groups) or one-way ANOVA (more than two groups, multiple comparisons) were used to test the significance of experiments: ns, not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$.

Acknowledgements

We thank Drs Anna Sokac, Swathi Arur, George Eisenhoffer and Pierre McCrea, and members of M.J.G.'s lab for comments and suggestions. Dr Adriana Paulucci-Holthausen at the Basic Science Research Building (BSRB) Microscopy Facility at MD Anderson Cancer Center provided training and support, and the NIH shared instrumentation grant (1S10OD024976-01) supported the confocal microscope. We also thank Hugo Bellen, Kenji Matsuno and Michael Buszczak for fly stocks; the Bloomington Drosophila Stock Center, the Vienna Drosophila RNAi Center and the Kyoto stock center for fly lines; and Developmental Studies Hybridoma Bank for antibodies.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: C.-R.T., M.J.G.; Validation: C.-R.T.; Investigation: C.-R.T.; Writing - original draft: C.-R.T.; Writing - review & editing: C.-R.T., M.J.G.; Visualization: C.-R.T.; Supervision: M.J.G.; Funding acquisition: C.-R.T., M.J.G.

Funding

C.-R.T was supported by an American Heart Association predoctoral fellowship (16PRE30880004). M.J.G. and this work is supported by the National Institute of General Medical Sciences (NIH R35GM126929). Deposited in PMC for immediate release.

Supplementary information

Supplementary information available online at
<http://dev.biologists.org/lookup/doi/10.1242/dev.175133.supplemental>

References

- Aberle, H., Bauer, A., Stappert, J., Kispert, A. and Kemler, R. (1997). beta-catenin is a target for the ubiquitin-proteasome pathway. *EMBO J.* **16**, 3797-3804. doi:10.1093/emboj/16.13.3797
- Abreu-Blanco, M. T., Verboon, J. M., Liu, R., Watts, J. J. and Parkhurst, S. M. (2013). Drosophila embryos close epithelial wounds using a combination of cellular protrusions and an actomyosin purse string. *J. Cell Sci.* **125**, 5984-5997. doi:10.1242/jcs.109066
- Baek, S. H., Cho, H. W., Kwon, Y.-C., Lee, J. H., Kim, M. J., Lee, H. and Choe, K.-M. (2012). Requirement for Pak3 in Rac1-induced organization of actin and myosin during Drosophila larval wound healing. *FEBS Lett.* **586**, 772-777. doi:10.1016/j.febslet.2012.01.061
- Brand, A. H. and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401-415.
- Brock, A. R., Wang, Y., Berger, S., Renkawitz-Pohl, R., Han, V. C., Wu, Y. and Galko, M. J. (2012). Transcriptional regulation of Profilin during wound closure in Drosophila larvae. *J. Cell Sci.* **125**, 5667-5676. doi:10.1242/jcs.107490
- Burra, S., Wang, Y., Brock, A. R. and Galko, M. J. (2013). Using Drosophila larvae to study epidermal wound closure and inflammation. *Methods Mol. Biol.* **1037**, 449-461. doi:10.1007/978-1-62703-505-7_26
- Calleja, M., Moreno, E., Pelaz, S. and Morata, G. (1996). Visualization of gene expression in living adult Drosophila. *Science* **274**, 252-255. doi:10.1126/science.274.5285.252
- Carvalho, L., Jacinto, A. and Matova, N. (2014). The Toll/NF- κ B signaling pathway is required for epidermal wound repair in Drosophila. *Proc. Natl. Acad. Sci. USA* **111**, E5373-E5382. doi:10.1073/pnas.1408224111
- Carvalho, L., Patricio, P., Ponte, S., Heisenberg, C.-P., Almeida, L., Nunes, A. S., Araújo, N. A. M. and Jacinto, A. (2018). Occluding junctions as novel regulators of tissue mechanics during wound repair. *J. Cell Biol.* **217**, 4267. doi:10.1083/jcb.201804048
- Chang, M. V., Chang, J. L., Gangopadhyay, A., Shearer, A. and Cadigan, K. M. (2008). Activation of wingless targets requires bipartite recognition of DNA by TCF. *Curr. Biol.* **18**, 1877-1881. doi:10.1016/j.cub.2008.10.047
- Chang, C.-H., Kuo, C.-J., Ito, T., Su, Y.-Y., Jiang, S.-T., Chiu, M.-H., Lin, Y.-H., Nist, A., Mernberger, M., Stiewe, T. et al. (2017). CK1 α ablation in keratinocytes induces p53-dependent, sunburn-protective skin hyperpigmentation. *Proc. Natl. Acad. Sci. USA* **114**, E8035-E8044. doi:10.1073/pnas.1702763114
- Das, R., Bhattacharjee, S., Patel, A. A., Harris, J. M., Bhattacharya, S., Letcher, J. M., Clark, S. G., Nanda, S., Iyer, E. P. R., Ascoli, G. A., et al. (2017). Dendritic cytoskeletal architecture is modulated by combinatorial transcriptional regulation in Drosophila melanogaster. *Genetics* **207**, 1401-1421. doi:10.1534/genetics.117.300393
- Galko, M. J. and Krasnow, M. A. (2004). Cellular and genetic analysis of wound healing in Drosophila larvae. *PLoS Biol.* **2**, e239. doi:10.1371/journal.pbio.0020239
- Hariharan, I. K. and Serras, F. (2017). Imaginal disc regeneration takes flight. *Curr. Opin. Cell Biol.* **48**, 10-16. doi:10.1016/j.cob.2017.03.005
- Hecht, A. and Kemler, R. (2000). Curbing the nuclear activities of beta-catenin control over Wnt target gene expression. *EMBO Rep.* **1**, 24-28. doi:10.1093/embo-reports/kvd012
- Hudson, L. G., Newkirk, K. M., Chandler, H. L., Choi, C., Fossey, S. L., Parent, A. E. and Kusewitt, D. F. (2009). Cutaneous wound reepithelialization is compromised in mice lacking functional Slug (Snai2). *J. Dermatol. Sci.* **56**, 19-26. doi:10.1016/j.jdermsci.2009.06.009
- Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G. and Birchmeier, W. (2001). β -Catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. *Cell* **105**, 533-545. doi:10.1016/S0092-8674(01)00336-1
- Hunter, M. V., Lee, D. M., Harris, T. J. C. and Fernandez-Gonzalez, R. (2015). Polarized E-cadherin endocytosis directs actomyosin remodeling during embryonic wound repair. *J. Cell Biol.* **210**, 801-816. doi:10.1083/jcb.201501076
- Kakanj, P., Moussian, B., Grönke, S., Bustos, V., Eming, S. A., Partridge, L. and Leptin, M. (2016). Insulin and TOR signal in parallel through FOXO and S6K to promote epithelial wound healing. *Nat. Commun.* **7**, 12972. doi:10.1038/ncomms12972
- Larue, L., Ohsugi, M., Hirchenhain, J. and Kemler, R. (1994). E-cadherin null mutant embryos fail to form a trophoblast epithelium. *Proc. Natl. Acad. Sci. USA* **91**, 8263-8267. doi:10.1073/pnas.91.17.8263
- Lawrence, P. A., Bodmer, R. and Vincent, J. P. (1995). Segmental patterning of heart precursors in Drosophila. *Development* **121**, 4303-4308.
- Lesch, C., Jo, J., Wu, Y., Fish, G. S. and Galko, M. J. (2010). A targeted UAS-RNAi screen in Drosophila larvae identifies wound closure genes regulating distinct cellular processes. *Genetics* **186**, 943-957. doi:10.1534/genetics.110.121822
- Lopez-Bellido, R., Puig, S., Huang, P. J., Tsai, C.-R., Turner, H. N., Galko, M. J. and Gutstein, H. B. (2019). Growth factor signaling regulates mechanical nociception in flies and vertebrates. *J. Neurosci.* **39**, 6012-6030. doi:10.1523/JNEUROSCI.2950-18.2019
- Matsubayashi, Y., Coulson-Gilmer, C. and Millard, T. H. (2015). Endocytosis-dependent coordination of multiple actin regulators is required for wound healing. *J. Cell Biol.* **210**, 677-679. doi:10.1083/jcb.20141103707282015c
- McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K. and Davis, R. L. (2003). Spatiotemporal rescue of memory dysfunction in Drosophila. *Science* **302**, 1765-1768. doi:10.1126/science.1089035
- McGuire, S. E., Roman, G. and Davis, R. L. (2004). Gene expression systems in Drosophila: a synthesis of time and space. *Trends Genet.* **20**, 384-391. doi:10.1016/j.tig.2004.06.012
- Nakamura, M., Dominguez, A. N. M., Decker, J. R., Hull, A. J., Verboon, J. M. and Parkhurst, S. M. (2018). Into the breach: how cells cope with wounds. *Open Biol.* **8**, 180135. doi:10.1098/rsob.180135
- Nunan, R., Campbell, J., Mori, R., Pitulescu, M. E., Jiang, W. G., Harding, K. G., Adams, R. H., Nobes, C. D. and Martin, P. (2015). Ephrin-Bs drive junctional downregulation and actin stress fiber disassembly to enable wound re-epithelialization. *Cell Rep.* **13**, 1380-1395. doi:10.1016/j.celrep.2015.09.085
- Oda, H., Uemura, T., Harada, Y., Iwai, Y. and Takeichi, M. (1994). A Drosophila homolog of cadherin associated with Armadillo and essential for embryonic cell-cell adhesion. *Dev. Biol.* **165**, 716-726. doi:10.1006/dbio.1994.1287
- Ozawa, M., Baribault, H. and Kemler, R. (1989). The cytoplasmic domain of the cell adhesion molecule uvomorulin associates with three independent proteins structurally related in different species. *EMBO J.* **8**, 1711-1717. doi:10.1002/j.1460-2075.1989.tb03563.x
- Pai, L.-M., Kirkpatrick, C., Blanton, J., Oda, H., Takeichi, M. and Peifer, M. (1996). Drosophila alpha-catenin and E-cadherin bind to distinct regions of Drosophila Armadillo. *J. Biol. Chem.* **271**, 32411-32420. doi:10.1074/jbc.271.50.32411
- Park, S.-H., Lee, C.-W., Lee, J.-H., Park, J. Y., Roshandell, M., Brennan, C. A. and Choe, K.-M. (2018). Requirement for and polarized localization of integrin proteins during Drosophila wound closure. *Mol. Biol. Cell* **29**, 2137-2147. doi:10.1091/mbc.E17-11-0635
- Patel, N. H., Snow, P. M. and Goodman, C. S. (1987). Characterization and cloning of fasciilin III: a glycoprotein expressed on a subset of neurons and axon pathways in Drosophila. *Cell* **48**, 975-988. doi:10.1016/0092-8674(87)90706-9
- Pinheiro, D. and Bellaïche, Y. (2018). Mechanical force-driven adherens junction remodeling and epithelial dynamics. *Dev. Cell* **47**, 3-19. doi:10.1016/j.devcel.2018.09.014
- Riedl, J., Crevenna, A. H., Kessenbrock, K., Yu, J. H., Neukirchen, D., Bista, M., Bradke, F., Jenne, D., Holak, T. A., Werb, Z., et al. (2008). Lifeact: a versatile marker to visualize F-actin. *Nat. Methods* **5**, 605-607. doi:10.1038/nmeth.1220
- Riethmacher, D., Brinkmann, V. and Birchmeier, C. (1995). A targeted mutation in the mouse E-cadherin gene results in defective preimplantation development. *Proc. Natl. Acad. Sci. USA* **92**, 855-859. doi:10.1073/pnas.92.3.855
- Riggelman, B., Schedl, P. and Wieschaus, E. (1990). Spatial expression of the Drosophila segment polarity gene armadillo is posttranscriptionally regulated by wingless. *Cell* **63**, 549-560. doi:10.1016/0092-8674(90)90451-J
- Sato, A., Kojima, T., Ui-Tei, K., Miyata, Y. and Saigo, K. (1999). Dfrizzled-3, a new Drosophila Wnt receptor, acting as an attenuator of Wingless signaling in wingless/hypomorphic mutants. *Development* **126**, 4421-4430.
- Schubiger, M., Sustar, A. and Schubiger, G. (2010). Regeneration and transdetermination: the role of wingless and its regulation. *Dev. Biol.* **347**, 315-324. doi:10.1016/j.ydbio.2010.08.034
- Smith-Bolton, R. K., Worley, M. I., Kanda, H. and Hariharan, I. K. (2009). Regenerative growth in Drosophila imaginal discs is regulated by wingless and *Myc*. *Dev. Cell* **16**, 797-809. doi:10.1016/j.devcel.2009.04.015
- Spradling, A., Stern, A. D., Rhem, E., Laverty, T., Mozden, N., Misra, S. and Rubin, G. (1999). The Berkeley Drosophila Genome Project gene disruption project: Single P-element insertions mutating 25% of vital Drosophila genes. *Genetics* **153**, 135-177.
- Stevens, L. J. and Page-McCaw, A. (2012). A secreted MMP is required for reepithelialization during wound healing. *Mol. Biol. Cell* **23**, 1068-1079. doi:10.1091/mbc.e11-09-0745
- Suffoletto, K., Jetta, D. and Hua, S. Z. (2018). E-cadherin mediated lateral interactions between neighbor cells necessary for collective migration. *J. Biomech.* **71**, 159-166. doi:10.1016/j.jbiomech.2018.02.002
- Tepass, U. and Hartenstein, V. (1994). Epithelium formation in the Drosophila midgut depends on the interaction of endoderm and mesoderm. *Development* **120**, 579-590.
- Tinkle, C. L., Pasolli, H. A., Stokes, N. and Fuchs, E. (2008). New insights into cadherin function in epidermal sheet formation and maintenance of tissue integrity. *Proc. Natl. Acad. Sci. USA* **105**, 15405-15410. doi:10.1073/pnas.0807374105
- Tsai, C.-R., Anderson, A. E., Burra, S., Jo, J. and Galko, M. J. (2017). Yorkie regulates epidermal wound healing in Drosophila larvae independently of cell proliferation and apoptosis. *Dev. Biol.* **427**, 61-71. doi:10.1016/j.ydbio.2017.05.006

- Tsai, C.-R., Wang, Y. and Galko, M. J.** (2018). Crawling wounded: molecular genetic insights into wound healing from *Drosophila* larvae. *Int. J. Dev. Biol.* **62**, 479-489. doi:10.1387/ijdb.180085mg
- Wieschaus, E., Nüsslein-Volhard, C. and Jürgens, G.** (1984). Mutations affecting the pattern of the larval cuticle in *Drosophila melanogaster*. *Wilhelm Roux's Arch. Dev. Biol.* **193**, 296-307. doi:10.1007/BF00848158
- Wood, W., Jacinto, A., Grose, R., Woolner, S., Gale, J., Wilson, C. and Martin, P.** (2002). Wound healing recapitulates morphogenesis in *Drosophila* embryos. *Nat. Cell Biol.* **4**, 907-912. doi:10.1038/ncb875
- Wu, Y., Brock, A. R., Wang, Y., Fujitani, K., Ueda, R. and Galko, M. J.** (2009). A blood-borne PDGF/VEGF-like ligand initiates wound-induced epidermal cell migration in *Drosophila* larvae. *Curr. Biol.* **19**, 1473-1477. doi:10.1016/j.cub.2009.07.019
- Yanagawa, S.-I., Matsuda, Y., Lee, J.-S., Matsubayashi, H., Sese, S., Kadowaki, T. and Ishimoto, A.** (2002). Casein kinase I phosphorylates the Armadillo protein and induces its degradation in *Drosophila*. *EMBO J.* **21**, 1733-1742. doi:10.1093/emboj/21.7.1733
- Zhang, L., Jia, J., Wang, B., Amanai, K., Wharton, K. A. and Jiang, J.** (2006). Regulation of wingless signaling by the CKI family in *Drosophila* limb development. *Dev. Biol.* **299**, 221-237. doi:10.1016/j.ydbio.2006.07.025
- Zulueta-Coarasa, T. and Fernandez-Gonzalez, R.** (2017). Tension (re)builds: Biophysical mechanisms of embryonic wound repair. *Mech. Dev.* **144**, 43-52. doi:10.1016/j.mod.2016.11.004
- Zulueta-Coarasa, T., Tamada, M., Lee, E. J. and Fernandez-Gonzalez, R.** (2014). Automated multidimensional image analysis reveals a role for Abl in embryonic wound repair. *Development* **141**, 2901-2911. doi:10.1242/dev.106898




Fig. S1. Validations of β -catenin antibody, β -catenin^{RNAi} and E-cadherin^{RNAi}.

(A-C') Dissected larval epidermal whole mounts of third instar larvae expressing *UAS-DsRed2nuc* (nuclei, magenta), *UAS-src-GFP* (green) and the indicated transgenes via the *A58-Gal4* driver. (A,A') *Control*^{RNAi}, (B,B') β -catenin^{RNAi}, (C,C') *E-cadherin*^{RNAi}. (A'-C') Anti- β -catenin, white. Scale bar in (A) is 50 μ m for (A-C'). (D) Quantitation of junctional β -catenin intensity in larvae expressing RNAi transgenes targeting components of the adherens junction. Each dot represents the average of the β -catenin signal intensities of five junctions measured within one larva. One-way ANOVA. ****, $p < 0.0001$. (E-F) Dissected larval epidermal whole mounts of third instar larvae expressing *E-cadherin-GFP* and a control *UAS-Luciferase*^{RNAi} transgene (E) or *UAS- β -catenin*^{RNAi} (F) via *e22c-Gal4*. Scale bar in (E) is 50 μ m for (E,F). (G-H) Dissected larval epidermal whole mounts of third instar larvae expressing *UAS-DsRed2nuc* (nuclei, magenta) and a control *UAS-LacZ* transgene (G) or *UAS-E-cadherin*^{RNAi} (H) via *A58-Gal4*. Scale bar in (G) is 50 μ m for (G,H).









Fig. S2. Different *Ck1α*^{RNAi} transgenes and their impacts on wound closure and adherens junctions.

(A) Gene regions targeted by *UAS-Ck1α*^{RNAi} transgenes. Continuous epidermal expression of *Ck1α*^{RNAi} #3 is lethal via *e22c-Gal4* driver. (B) Quantitation of the percent of open wounds in larvae expressing the indicated transgenes via the *e22c-Gal4* driver 16 hours after wounding. Each dot represents one set of $n \geq 8$ larvae for each genotype. One-way ANOVA; ****, $p < 0.0001$; ns = not significant. Error bars, standard error of mean. (C-D) Dissected epidermal whole mounts of unwounded third instar larvae expressing *UAS-DsRed2nuc* (nuclei, not shown), *UAS-src-GFP* (cell membranes, not shown) and the indicated transgenes via *e22c-Gal4*. White, anti- β -catenin. Scale bar in (C) is 50 μ m for (C,D,F,G). (E) Quantitation of junctional β -catenin intensity in larvae expressing *Ck1α*^{RNAi} versus control. Each dot represents an average of the β -catenin signal from five junctions of one larva. Unpaired t-test; ****, $p < 0.0001$. (F-G) Dissected epidermal whole mounts of unwounded third instar larvae expressing *UAS-DsRed2nuc* (nuclei, magenta), *UAS-src-GFP* (cell membranes, green, not shown) and the indicated transgenes via *e22c-Gal4*. White, anti-E-cadherin. (H) Quantitation of junctional E-cadherin intensity in larvae expressing *Ck1α*^{RNAi} versus control. Each dot represents an average of the E-cadherin signal from five junctions of one larva. Unpaired t-test; **, $p < 0.01$.







Fig. S3. Epidermal expression of $Ckl\alpha^{RNAi}$ transgene increases E-cadherin.

(A) Schematic of the experimental design/temperature shift regimen for using Gal80^{ts} to inducibly express UAS-dependent transgenes in the larval epidermis. (B-E,G-J) Dissected larval epidermal whole mounts of third instar larvae expressing *Gal80^{ts}* transgene driven by tubulin promoter, *UAS-DsRed2nuc* (nuclei, not shown), *UAS-src-GFP* (cell membrane, not shown), and the indicated transgenes via the *e22c-Gal4* driver. (B,G) *Luciferase^{RNAi}*, (C,H) *Ckl\alpha^{RNAi#3}*; *Luciferase^{RNAi}*, (D,I) *\beta-catenin^{RNAi}*, (E,J) *Ckl\alpha^{RNAi#3}*; *\beta-catenin^{RNAi}*. (B-E) Anti- β -catenin, white. Scale bar in (B) is 50 μ m for (B-E,G-J). (G-J) Anti-E-cadherin, white. Quantitation of junctional β -catenin intensity (F) and E-cadherin intensity (K) in larvae expressing the indicated transgenes. Each dot represents the average of the junctional signal intensities of five junctions measured within one larva. One-way ANOVA. *, $p < 0.05$; ns = not significant.




Fig. S4. Wnt signaling is not activated after wounding but is activated in the larval epidermis lacking Ck1 α or expressing β -catenin.

(A-F) X-gal staining (blue) of dissected epidermal whole mounts of unwounded (A-C,G-L) or pinch wounded (D-F) third instar larvae expressing the Wnt reporters *fz3-LacZ* (A,A',D), *6TH-lacZ* (B,B',E), or a JNK reporter *msn-lacZ* (C, F). All wounds were examined four hour post-wounding. The wing discs of unwounded larvae expressing the Wnt reporter *fz3-lacZ* (A') or *6TH-LacZ* (B') indicate control staining. Arrows in (B) indicate examples of unrelated signal. Blue dots, LacZ positive nuclei; red dashed line, wound edges. Arrowheads, *msn-lacZ* positive wound-edge cells. (G-L) X-gal staining (blue) of dissected epidermal whole mounts of unwounded third instar larvae expressing the Wnt reporters *fz3-LacZ* (G, I, K) or *6TH-lacZ* (H, J, L) and the indicated transgenes (left margin) via *e22c-Gal4*.








Fig. S5. Validation of conditional *UAS-E-cadherin*^{RNAi} and constitutive *UAS-TCF*^{RNAi} efficacy.

(A) Schematic of the experimental design/temperature shift regimen for using Gal80^{ts} to inducibly express UAS-dependent transgenes in the larval epidermis. (B-E,G-J) Dissected larval epidermal whole mounts of third instar larvae expressing a *Gal80^{ts}* transgene driven by tubulin promoter, *UAS-DsRed2nuc* (nuclei, not shown), *UAS-src-GFP* (not shown), and the indicated transgenes via the *e22c-Gal4* driver. (B,G) *Luciferase*^{RNAi}, (C,H) *Cklα*^{RNAi#3}; *Luciferase*^{RNAi}, (D,I) *E-cadherin*^{RNAi}, (E,J) *Cklα*^{RNAi#3}; *E-cadherin*^{RNAi}. (B-E) Anti-β-catenin, white. (G-J) Anti-E-cadherin, white. Scale bar in (B) is 50 μm for (B-E,G-J). Quantitation of junctional β-catenin intensity (F) and E-cadherin intensity (K) in larvae expressing indicated transgenes. Each dot represents the average of the junctional intensities of five junctions measured within one larva. One-way ANOVA. **, p<0.01; ***, p<0.001; ****, p<0.0001. (L,M) Dissected wings of female adult flies expressing the indicated transgenes via a wing pouch driver, *nub-Gal4*, during wing development. The deformed shape of the wing in the *UAS-TCF*^{RNAi} group indicates that this transgene is inhibiting Wnt signaling.







Fig. S6. β -catenin overexpression and *E-cadherin*^{RNAi} effects on β -catenin and E-cadherin expression

(A) Schematic of the experimental design/temperature shift regimen for using Gal80^{ts} to inducibly express UAS-dependent transgenes in the larval epidermis. (B-E,G-J) Dissected larval epidermal whole mounts of third instar larvae expressing a Gal80^{ts} transgene driven by tubulin promoter, *UAS-DsRed2nuc* (nuclei, not shown), *UAS-src-GFP* (cell membrane, not shown), and the indicated transgenes via the *e22c-Gal4* driver. (B,G) *Luciferase*^{RNAi}, (C,H) *UAS- β -catenin*; *Luciferase*^{RNAi}, (D,I) *E-cadherin*^{RNAi}, (E,J) *UAS- β -catenin*; *E-cadherin*^{RNAi}. (B-E) Anti- β -catenin, white. Scale bar in (B) is 50 μ m for (B-E,G-J). For ease of visualization, panels C and E are imaged at lower laser intensity so as not to be oversaturated. (G-J) Anti-E-cadherin, white. Quantitation of junctional β -catenin intensity (F) and E-cadherin intensity (K) in larvae expressing indicated transgenes. Each dot represents the average of the junctional intensities of five junctions measured within one larva. Unpaired t-test; *, p<0.05; ****, p<0.0001.

Supplementary Materials and methods

Flies used in this study

Please note the genotype of sex chromosome is simplified. The actual genotypes for the sex chromosome could be mixed, depending on the source RNAi collection, UAS transgenes, and the sex of individual larvae as male and female progeny were pooled together in test populations.

Fig. panels – genotypes tested:

Fig. 1A-E'. $w^{1118}; A58-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. 1F. $w^{1118}; e22c-Gal4, UAS-LifeAct-mCherry/+; UAS-Luciferase^{RNAi}$ (JF01355)/+

Fig. 1H. $w^{1118}; A58-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. 2A. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Luciferase^{RNAi}$ (JF01355)/+

Fig. 2B. $w^{1118}; A58-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Ck1\alpha^{RNAi\#1}$ (JF01792)

Fig. 2C. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-GSK3\beta^{RNAi}$ (KK108994)

Fig. 2D. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-\beta-TrCP^{RNAi}$ (JF01504)/+

Fig. 2E. Same as Fig. 2A-D.

Fig. 2F. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Luciferase^{RNAi}$ (JF01355)/+

Fig. 2G. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Ck1\alpha^{RNAi\#1}$ (JF01792)/+

Fig. 2H. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-GSK3\beta^{RNAi}$ (KK108994)

Fig. 2I. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-\beta-TrCP^{RNAi}$ (JF01504)/+

Fig. 2J. Same as Fig. 2F-I.

Fig. 3B. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; tubP-gal80^{\Delta S}/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. 3C. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-Ck1 α ^{RNAi#3}* (GD4256); *tubP-gal80^{ts}/UAS-Luciferase^{RNAi}* (JF01355)

Fig. 3D. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS- β -catenin^{RNAi}* (JF01252)

Fig. 3E. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-Ck1 α ^{RNAi#3}* (GD4256); *tubP-gal80^{ts}/UAS- β -catenin^{RNAi}* (JF01252)

Fig. 3F. Same as Fig. 3B-E.

Fig. 4B. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-Luciferase^{RNAi}* (JF01355)

Fig. 4C. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-Ck1 α ^{RNAi#3}* (GD4256); *tubP-gal80^{ts}/UAS-Luciferase^{RNAi}* (JF01355)

Fig. 4D. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. 4E. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-Ck1 α ^{RNAi#3}* (GD4256); *tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. 4F. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-TCF^{RNAi}* (JF02306)

Fig. 4G. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-Ck1 α ^{RNAi#3}* (GD4256); *tubP-gal80^{ts}/UAS-TCF^{RNAi}* (JF02306)

Fig. 4H. Same as Fig. 4B-G.

Fig. 5B. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-Luciferase^{RNAi}* (JF01355)

Fig. 5C. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/P{UAS-arm.Exel}2* (*UAS- β -catenin*, BL#8369); *tubP-gal80^{ts}/UAS-Luciferase^{RNAi}* (JF01355)

Fig. 5D. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-TCF^{RNAi}* (JF02306)

Fig. 5E. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/P{UAS-arm.Exel}2* (*UAS- β -catenin*, BL#8369); *tubP-gal80^{ts}/UAS-TCF^{RNAi}* (JF02306)

Fig. 5F. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. 5G. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/P{UAS-arm.Exel}2* (*UAS- β -catenin*, BL#8369); *tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. 5H. Same as Fig. 5B-G.

Fig. 6A. w^{1118} ; *e22c-Gal4*, *UAS-LifeAct-mCherry*, *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)/*TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584); *UAS-Luciferase^{RNAi}* (JF01355)/+

Fig. 6B. w^{1118} ; *e22c-Gal4*, *UAS-LifeAct-mCherry*, *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)/*P{UAS-arm.Exel}2* (*UAS- β -catenin*, BL#8369), *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)

Fig. 6C. w^{1118} ; *e22c-Gal4*, *UAS-LifeAct-mCherry*, *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)/*TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584); *UAS-Ck1 α ^{RNAi#1}* (JF01792)/+

Fig. 6D,D'. w^{1118} ; *e22c-Gal4*, *UAS-LifeAct-mCherry*, *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)/*TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584); *UAS-Luciferase^{RNAi}* (JF01355)/+

Fig. 6E,E'. w^{1118} ; *e22c-Gal4*, *UAS-LifeAct-mCherry*, *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)/*TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584); *UAS-Ck1 α ^{RNAi#1}* (JF01792)/+

Fig. 6F. Same as Fig. 6D,E.

Fig. S1A,A'. w^{1118} ; *A58-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-Luciferase^{RNAi}* (JF01355)

Fig. S1B,B'. w^{1118} ; *A58-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS- β -catenin^{RNAi}* (JF01252)

Fig. S1C,C'. w^{1118} ; *A58-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. S1D. Same as Fig. S1A-C'

Fig. S1E. w^{1118} ; *e22c-Gal4*, *TI{TI}shg^{GFP}* (*E-cadherin-GFP*, BL#60584)/+; *UAS-Luciferase^{RNAi}* (JF01355)/+

Fig. S1F. $w^{1118}; e22c-Gal4, TI\{TI\}shg^{GFP} (E-cadherin-GFP, BL\#60584)/+; UAS-\beta-catenin^{RNAi} (JF01252)/+$

Fig. S1G. $w^{1118}; P\{UAS-lacZ.Exel\}2 (UAS-LacZ, BL\#8529)/+; A58-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+$

Fig. S1H. $w^{1118}; A58-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/ UAS-E-cadherin^{RNAi} (HMS00693)$

Fig. S2B

- $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Luciferase^{RNAi} (JF01355)/+$
- $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Ck1\alpha^{RNAi\#2} (GL00021)/+$
- $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Ck1\alpha^{RNAi\#4} (HMS02276)/+$

Fig. S2C,F. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Luciferase^{RNAi} (JF01355)/+$

Fig. S2D,G. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; UAS-Ck1\alpha^{RNAi\#1} (JF01792)/+$

Fig. S2E. Same as Fig. S2C,D

Fig. S2H. Same as Fig. S2F,G

Fig. S3B,G. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; tubP-gal80^{ts}/UAS-Luciferase^{RNAi} (JF01355)$

Fig. S3C,H. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Ck1\alpha^{RNAi\#3} (GD4256); tubP-gal80^{ts}/UAS-Luciferase^{RNAi} (JF01355)$

Fig. S3D,I. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; tubP-gal80^{ts}/UAS-\beta-catenin^{RNAi} (JF01252)$

Fig. S3E,J. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Ck1\alpha^{RNAi\#3} (GD4256); tubP-gal80^{ts}/UAS-\beta-catenin^{RNAi} (JF01252)$

Fig. S3F. Same as Fig. S3B-E

Fig. S3K. Same as Fig. S3G-J

Fig. S4A,D. $w^{1118}; fz3-LacZ/+$

Fig. S4B,E. $w^{1118}; 6TH-LacZ/+$

Fig. S4C,F. $w^{1118}; A58-Gal4, msn-LacZ/+$

Fig. S4G. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; fz3-LacZ/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. S4H. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; 6TH-LacZ/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. S4I. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; fz3-LacZ/UAS-Ck1\alpha^{RNAi\#1}$ (JF01792)

Fig. S4J. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; 6TH-LacZ/ UAS-Ck1\alpha^{RNAi\#1}$ (JF01792)

Fig. S4K. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/ P\{UAS-arm.Exel\}2$ (*UAS- β -catenin*, BL#8369); $fz3-LacZ/+$

Fig. S4L. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/P\{UAS-arm.Exel\}2$ (*UAS- β -catenin*, BL#8369); $6TH-LacZ/+$

Fig. S5B,G. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; tubP-gal80^{ts}/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. S5C,H. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Ck1\alpha^{RNAi\#3}$ (GD4256); $tubP-gal80^{ts}/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. S5D,I. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}$ (HMS00693)

Fig. S5E,J. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/UAS-Ck1\alpha^{RNAi\#3}$ (GD4256); $tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}$ (HMS00693)

Fig. S5F. Same as Fig. S5B-E

Fig. S5K. Same as Fig. S5G-J

Fig. S5L. $w^{1118}; nubbin-Gal4/+; UAS-Luciferase^{RNAi}$ (JF01355)/+

Fig. S5M. $w^{1118}; nubbin-Gal4/+; UAS-TCF^{RNAi}$ (JF02306)/+

Fig. S6B,G. $w^{1118}; e22c-Gal4, UAS-DsRed2Nuc, UAS-src-GFP/+; tubP-gal80^{ts}/UAS-Luciferase^{RNAi}$ (JF01355)

Fig. S6C,H. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/P{UAS-arm.Exel}2*
(*UAS-β-catenin*, BL#8369); *tubP-gal80^{ts}/UAS-Luciferase^{RNAi}* (JF01355)

Fig. S6D,I. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/+*; *tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. S6E,J. w^{1118} ; *e22c-Gal4*, *UAS-DsRed2Nuc*, *UAS-src-GFP/P{UAS-arm.Exel}2* (*UAS-β-catenin*, BL#8369); *tubP-gal80^{ts}/UAS-E-cadherin^{RNAi}* (HMS00693)

Fig. S6F. Same as Fig. S6B,D.

Fig. S6K. Same as Fig. S6G,H