

Mitochondrial dynamics underlying thermal plasticity of cuttlefish (*Sepia officinalis*) hearts

Key words: Temperature sensitivity, cephalopod, evolutionary adaptation, thermal acclimation, proton leak, lactate and octopine dehydrogenase, respiration, systemic and branchial hearts, cardiac fibres

Michael Oellermann, Hans O. Pörtner, Felix C. Mark*

Alfred Wegener Institute for Polar and Marine Research, 27570 Bremerhaven, Germany

*Author for correspondence (e-mail: Felix.Christopher.Mark@awi.de)

1 **ABSTRACT**

2 In the eurythermal cuttlefish *Sepia officinalis*, performance depends on hearts that ensure
3 systemic oxygen supply over a broad range of temperatures. We therefore aimed to identify
4 adjustments in energetic cardiac capacity and underlying mitochondrial function supporting
5 thermal acclimation and adaptation that could be critical for the cuttlefish's competitive
6 success in variable environments. Two genetically distinct cuttlefish populations were
7 acclimated to 11°C, 16°C and 21°C, respectively. Subsequently, skinned and permeabilised
8 heart fibres were used to assess mitochondrial functioning by means of high-resolution
9 respirometry and a substrate-inhibitor protocol, followed by measurements of cardiac citrate
10 synthase and cytosolic enzyme activities. Temperate English Channel cuttlefish had lower
11 mitochondrial capacities but larger hearts than subtropical Adriatic cuttlefish. Warm
12 acclimation to 21°C decreased mitochondrial complex I activity in Adriatic cuttlefish and
13 increased complex IV activity in English Channel cuttlefish. However, compensation of
14 mitochondrial capacities did not occur during cold acclimation to 11°C. In systemic hearts,
15 thermal sensitivity of mitochondrial substrate oxidation was high for proline and pyruvate but
16 low for succinate. Oxygen efficiency of catabolism rose from 11°C to 21°C via shifts to
17 oxygen-conserving oxidation of proline and pyruvate and via reduced relative proton leak.
18 The changes observed for substrate oxidation, mitochondrial complexes, relative proton leak
19 or heart weights improve energetic efficiency and essentially seem to extend tolerance to high
20 temperatures and reduce associated tissue hypoxia. We conclude that cuttlefish sustain cardiac
21 performance and thus, systemic oxygen delivery over short and long-term changes of
22 temperature and environmental conditions by multiple adjustments in cellular and
23 mitochondrial energetics.

24 INTRODUCTION

25 The evolution of modern cephalopods was shaped by the rising competition with marine
26 vertebrates and led to innovations that advance high levels of performance. Concomitant
27 increases of energy demands required extant cephalopods to optimise supply and use of
28 oxygen (O'Dor and Webber, 1986; O'Dor and Webber, 1991). As a result, cephalopods
29 evolved high concentrations of blood pigment optimised for oxygen transport and a closed
30 circulatory system driven by two branchial and one powerful systemic heart (Schipp, 1987;
31 Wells and Smith, 1987; Wells, 1992; Pörtner and Zielinski, 1998). However, design
32 constraints were involved, leading to locomotion by jet propulsion and comparatively low
33 blood oxygen carrying capacities. Optimisation of performance led at least some cephalopod
34 species to operate at their functional limits (O'Dor and Webber, 1986; Pörtner, 2002b).
35 Environmental stressors such as fluctuations in ambient temperature or ambient hypoxia may
36 be particularly challenging for the highly oxygen dependent cephalopods that face high
37 competitive pressure (Rosa and Seibel, 2008).

38 Body functions of animals operate only within a certain thermal range and are set by
39 ambient temperature for ectotherms. According to recent evidence, oxygen supply becomes
40 limiting at high temperatures when oxygen demand increases due to limited functional
41 capacities of the circulation and ventilation system to deliver oxygen (Pörtner and Knust,
42 2007; Pörtner and Farrell, 2008). Adaptive adjustments of e.g. ventilatory musculature and
43 heart rate or stroke volume are suitable to compensate for temperature induced impairments of
44 oxygen supply (Wells, 1992; Frederich and Pörtner, 2000). These adjustments, however,
45 involve changes in the capacity of mitochondria to provide sufficient aerobic energy to vital
46 tissues like the heart (Pörtner, 2002a). Low temperatures also cause capacity limitations,
47 which may involve insufficient release of oxygen to tissues by the blood pigment
48 haemocyanin (Melzner et al., 2007b) and limited mitochondrial energy provision as required
49 to power circulation or ventilation (Pörtner, 2002a).

50 Ectothermic hearts play a major role in defining limits of aerobic performance and thermal
51 tolerance, as demonstrated by impaired cardiac function close to extreme temperatures or
52 exercise levels (Farrell, 2002; Somero, 2010). In cephalopods, systemic hearts cover
53 increased metabolic demands during either exercise or rising temperatures by a 2-3 fold
54 increase of stroke volume or heartbeat frequency (Wells, 1992), but show limited
55 performance at high critical temperatures, as indicated in *Sepia officinalis*, where blood
56 perfusion fails to increase further beyond 23°C (Melzner et al., 2007a; Melzner et al., 2007b).

57 Yet, unlike in fishes, cephalopod systemic hearts receive well oxygenated blood and gain
58 support by two accessory hearts, contractile blood vessels and mantle pressure oscillations
59 (Schipp, 1987; Melzner et al., 2007a). Adversely, high blood viscosity, 2-3 fold lower blood
60 oxygen carrying capacities and higher resting metabolic rates compared to haemoglobin
61 bearing fish with a similar lifestyle (Wells, 1992; Pörtner, 1994) demand higher performance
62 of the systemic heart. While cephalopod hearts are able to meet this workload within limits
63 during acute rises of metabolic demand, it is unknown whether they adjust to seasonal (i.e.
64 acclimation) or long term environmental changes over multiple generations (i.e. evolutionary
65 adaptation). Such temperature acclimation or adaptation has been shown for fish hearts. Here,
66 changes comprise increases in heart size (Goolish, 1987; Kent et al., 1988), increased
67 mitochondrial content (Kleckner and Sidell, 1985; Johnston and Harrison, 1987; Kolok,
68 1991), shifts from carbohydrate to fatty acid oxidation (Sephton and Driedzic, 1991; Sidell et
69 al., 1995) or enhanced activities of enzymes essential in aerobic metabolism (Crockett and
70 Sidell, 1990; Podrabsky et al., 2000), following cold-exposure. Yet, evidence for such
71 temperature related adjustments is still lacking for cephalopod hearts.

72 The eurythermal common cuttlefish *Sepia officinalis* (Linnaeus 1758) lives on the
73 continental shelf from the cold-temperate eastern North Atlantic to warm-subtropical
74 Mediterranean and Atlantic waters off the Senegalese coast and follows a bottom dwelling,
75 migratory life style on rocky to sandy grounds down to 200 m depth. Cuttlefish are rather
76 sluggish, grow fast (up to two kg) and spawn after one or two years in shallow waters and die
77 thereafter (Jereb and Roper, 2005). In this study, we compared a cuttlefish population from
78 the English Channel living between 9°C and 17.5°C with a genetically distinct Mediterranean
79 population (Wolfram et al., 2006) facing a range from 10°C to 25°C, respectively, between
80 winter and summer (Boucaud-Camou and Boismery, 1991; Artegiani et al., 1997; Wang et al.,
81 2003).

82 In this study, we aimed to understand how temperature changes affect the heart of the
83 common cuttlefish *Sepia officinalis* by (1) exploring cardiac adjustments following long term
84 genetic isolation in a temperate and subtropical habitat as well as (2) following thermal
85 acclimation and (3) by investigating the thermal sensitivity of heart mitochondria.

86 MATERIAL AND METHODS

87 Experimental animals

88 Wild laid eggs of the European cuttlefish (*Sepia officinalis*) were collected from the
89 temperate Oosterschelde lagoon, Netherlands (3°56'E, 51°35'N) and the subtropical Venetian
90 lagoon close to Chioggia, Adria, Italy (12°18'E, 45°13'N). Populations from these localities
91 form geographically distanced and genetically distinct clades without evidence of on-going
92 genetic exchange (Wolfram et al., 2006; Perez-Losada et al., 2007). Thus, inter-population
93 differences will most likely reflect evolutionary adaptation or genetic drift. About 300 eggs
94 from five to six different egg masses (50 to 60 eggs per mass) were collected from a
95 frequently visited spawning ground (>100 individuals per day) in the Oosterschelde lagoon
96 and 200 eggs from more than ten egg masses were collected by scuba diving and snorkelling
97 in the Venetian lagoon in 2008 and 2009. After transport to our institute individuals from each
98 population were hatched and raised in separate tanks, connected to a re-circulating
99 aquaculture system at constant temperature (mean \pm s.d., 15.9°C \pm 0.1) at suitable stocking
100 densities (Hanley et al., 1998) under constant 12 h dark:12 h light cycle. Hatchlings were fed
101 daily with live shrimps (*Neomysis integer*, *Palaemonetes varians* and *Crangon crangon*) and
102 after reaching a bigger size, exclusively with frozen brown shrimp (*Crangon crangon*). Water
103 quality parameters were monitored weekly and kept at levels appropriate for cuttlefish culture
104 (Hanley et al., 1998) (mean \pm s.d., O₂ = 10.07 \pm 0.43 mg l⁻¹, pH = 8.05 \pm 0.02, salinity = 32.6
105 psu \pm 0.5, NH₄⁺ < 0.2 mg l⁻¹, NO₂⁻ < 0.2 mg l⁻¹, NO₃⁻ < 80 mg l⁻¹) by means of water
106 treatment (protein skimmers, mechanical and biological filters, UV sterilisation) or water
107 replacement.

108 After animal size had reached 30 - 40 g, each population was divided into three groups
109 that were acclimated to 11°C, 16°C and 21°C from 8 to 22 weeks. These acclimation
110 temperatures were selected to expose European cuttlefish to their seasonal and depth
111 dependent temperature range, 10°C to 17.5°C for the English Channel population and 10°C to
112 25°C for the Adriatic Sea population. Each group was kept in separate tanks connected to
113 independent re-circulating systems for each temperature. Only animals showing normal
114 behaviour without signs of illness (skin infections mostly due to jetting against tank walls)
115 were selected for experiments. Although Adriatic cuttlefish weighed significantly more in the
116 21°C group (Supplementary table S2) at the end of the acclimation period (possibly due to

117 enhanced growth rates), correlation analysis did not show clear evidence that weight
118 differences obscured our data.

119 **Mitochondrial respiration**

120 *Dissection and muscle fibre preparation*

121 Animals were anesthetized in 3% ethanol until non-responsive before culling.
122 Subsequently, total weight, mantle length, total length and sex were recorded. After opening
123 the mantle cavity, samples of blood, gills and ink were taken and frozen for further
124 experiments. The three hearts were excised starting with the branchial hearts and placed
125 immediately into 1 ml ice-cold biopsy buffer (modified after (Kuznetsov et al., 2008)) to
126 preserve mitochondrial function. The biopsy buffer contained (in mmol l⁻¹, 2.77 CaK₂EGTA,
127 7.23 EGTA, 14.46 KOH, 5.77 Na₂ATP, 6.56 MgCl₂, 20 taurine, 20 imidazole, 0.5
128 dithiothreitol (DTT), 50 MES, 588 sucrose, 252 glycine, pH 7.4 at 26°C, 1000 mosmol l⁻¹).

129 After removal of non-cardiac tissues and weighing of all three hearts, approximately 50
130 mg of systemic heart tissue were placed into a drop of biopsy buffer on a Petri dish on ice and
131 then coarsely torn apart and dissected into small fibre bundles using two tweezers. The
132 remainder of heart tissue and the branchial hearts were snap-frozen in liquid nitrogen and
133 stored at -80°C for subsequent enzyme assays. Fibre bundles were immediately transferred to
134 1 ml biopsy buffer in a 12 well multiwell culture plate and permeabilised with 50 µg ml⁻¹
135 saponin (Note: preliminary testing confirmed appropriate saponin concentration) by gentle
136 mixing (115 rpm) on ice for 30 min. The fibres were then removed and washed three times for
137 10 min in 2 ml ice-cold mitochondrial respiration medium (in mmol l⁻¹, 50 HEPES, 25
138 KH₂PO₄, 0.5 EGTA, 50 KCl, 50 NaCl, 10 MgCl₂, 20 taurine, 50 lactobionate, 350 sucrose,
139 150 glycine, 1 g l⁻¹ freshly added fatty acid free BSA, pH 7.4 at 22°C, 1000 mosmol l⁻¹,
140 modified after (Mommensen and Hochachka, 1981; Agnisola et al., 1991; Kuznetsov et al.,
141 2008)) and stored in ice-cold respiration medium until use (protocol modified after (Saks et
142 al., 1998)).

143 *Measurements of mitochondrial respiration*

144 Before each assay, fibres were blotted dry on chilled Whatman® paper, divided into two
145 2-6 mg bundles and transferred to 2 ml duplicate chambers of an Oxygraph-2k respirometer
146 (Oroboros Instruments, Innsbruck, Austria) containing air saturated respiration medium at the
147 experimental temperature. Mitochondrial respiration was then measured online as background
148 corrected weight specific oxygen consumption rate (pmol O₂ s⁻¹ mg⁻¹, i.e. negative time

149 derivative of oxygen concentration) using DatLab analysis software (Oroboros Instruments,
150 Innsbruck, Austria). Assays were performed consecutively at 11°C, 16°C and 21°C in
151 randomised order. Chambers were washed with 96% ethanol and Milli-Q water to thoroughly
152 remove inhibitors and substrates after each assay. To protect fibres from extreme oxygen
153 levels, oxygen concentrations in chambers were kept between 80 nmol O₂ ml⁻¹ (maximum
154 stimulated respiration remained stable down to 80 nmol O₂ ml⁻¹) and air saturation (270-340
155 nmol O₂ ml⁻¹) by re-oxygenation with pure oxygen gas.

156 To assess mitochondrial function, respirometry on permeabilised skinned heart fibres was
157 combined with a substrate-inhibitor protocol as follows. Mitochondria were first fuelled
158 successively and in excess with the amino acid proline (5 mmol l⁻¹), ADP (2.5 mmol l⁻¹) as
159 well as pyruvate (5 mmol l⁻¹) and succinate (10 mmol l⁻¹) in order to reach maximum coupled
160 oxidative phosphorylation (i.e. state 3 respiration tied to ATP production). The choice of
161 substrates corresponds to own tests and to previous studies that showed high rates of
162 oxidation for proline, pyruvate and succinate in squid hearts (Ballantyne et al., 1981;
163 Mommsen and Hochachka, 1981). In situ, cephalopod hearts derive pyruvate most evidently
164 from blood glucose or oxidised octopine and proline from high intracellular stores (> 12 μmol
165 g⁻¹). Succinate can originate from both pyruvate derivatives and amino acids (e.g. ornithine,
166 arginine, glutamate and proline) with the latter feeding into the Krebs cycle at the level of
167 alpha-ketoglutarate (Ballantyne et al., 1981; Mommsen and Hochachka, 1981; Hochachka
168 and Fields, 1982).

169 Integrity of mitochondrial membranes was tested by the addition of cytochrome c (0.01
170 mmol l⁻¹) that would stimulate respiration in case of damaged outer membranes. Proline
171 stimulated respiration without ADP (state 2) provided an estimate of mitochondrial proton
172 leak (Iftikar et al., 2010). Uncoupling of respiration from ATP production by
173 carbonylcyanide-p-(trifluoromethyl) phenylhydrazone (FCCP, up to 2.5 μmol l⁻¹) denoted
174 maximum capacity of the electron transport chain. The loss of activity with rotenone (2.5
175 μmol l⁻¹) indicated NADH dehydrogenase (complex I) activity. The loss of activity with
176 antimycin A (2.5 μmol l⁻¹) quantified the non-mitochondrial background respiration.
177 Cytochrome c oxidase (complex IV) activity was tested using the redox pair ascorbate (2
178 mmol l⁻¹) and N,N,N',N'- tetramethyl-p-phenylenediamine dihydrochloride (TMPD, 0.5
179 mmol l⁻¹, see list in supplementary table S1). Background auto-oxidation of the redox pair
180 was determined for all experimental temperatures and subtracted from final results. Suitability
181 and concentrations of substrates were tested beforehand. All chemicals were purchased from
182 Sigma-Aldrich (Schnelldorf, Germany)

183 **Enzyme assays**

184 *Protein extraction and sample preparation*

185 To extract the protein, frozen heart tissue was ground by hand in a mortar filled with
186 liquid nitrogen. The frozen tissue powder was then weighed and suspended in 10 volumes
187 (w:v) of ice cold extraction buffer (50 mmol l⁻¹ Tris-HCl (pH 7.4 at 16°C), 1 mmol l⁻¹ EDTA,
188 0.1% Triton X-100) and sonicated for 90 s at 0°C in a Branson Sonifier 450 (output control 8,
189 Duty cycle 50%). Following two centrifugations for 10 min at 6000g and 4°C, supernatants
190 were used for protein and enzyme assays. Pellets remaining after centrifugation were
191 resuspended and re-extracted in ice cold extraction buffer to test for residual protein and
192 citrate synthase activity.

193 For enzyme assays, samples were diluted 1:10 (v:v) with 75 mmol l⁻¹ Tris-HCl (pH 8.1 at
194 16°C) and equilibrated to approximately 16°C using a temperature controlled metal block
195 connected to a thermostat (Haake C25, Thermo Scientific, Karlsruhe, Germany) prior to each
196 measurement. Absorbance was measured in triplicates using 96 well F-bottom microplates
197 (Nunc GmbH & Co. KG, Wiesbaden, Germany), a multiplate reader (Fluostar Galaxy, BMG
198 Labtechnologies, Offenburg, Germany) and analyzing software (FLUO 32, version 4.31 R5).

199 *Enzyme activities and protein content*

200 Citrate synthase (EC 4.1.3.7) activity was determined after Sidell et al. (1987) as the
201 increase of absorbance at 412 nm by means of DTNB (5,5'-dithio-bis-(2-nitrobenzoic acid)).
202 First, background deacylase activity was measured with 20 µl of diluted sample supernatant
203 and 160 µl reaction mixture (in mmol l⁻¹, 75 Tris-HCl, 0.25 DTNB, 0.4 acetyl~CoA, pH 8.1)
204 followed by the addition of 20 µl 0.5 mmol l⁻¹ oxaloacetate to start the reaction. As DTNB
205 reacts with sulphur groups (SH), dithiothreitol (DTT) was used as standard for the calculation
206 of SH group turn-over.

207 Activities of octopine dehydrogenase (ODH, EC 1.5.1.11) and lactate dehydrogenase
208 (LDH, EC 1.1.1.27) indicate capacities to form or recycle anaerobic end products in
209 cephalopods (Grieshaber and Gäde, 1976; Storey, 1977; Storey and Storey, 1979). ODH and
210 LDH activities were determined after Storey (1977) and Driedzic et al. (1990) as the decrease
211 of absorbance at 340 nm. First, background activity was measured by combining 20 µl of
212 diluted sample supernatant and 160 µl reaction mixture (in mmol l⁻¹, 100 Tris-HCl (pH 7.0),
213 0.45 NADH, 1 KCN, 10 arginine were added for quantifying the ODH reaction) followed by
214 the addition of 20 µl 4 mmol l⁻¹ pyruvate to start the reaction. Final ODH activity was

215 determined by subtracting the LDH activity, which was measured simultaneously. NADH
216 was used as a standard.

217 Enzyme activities were standardised to protein content determined after Bradford (1976).
218 Sample supernatant and pellet homogenate were diluted 1:10 (v:v) with 0.9% NaCl. 5 µl were
219 transferred to 250 µl Bradford dye reagent (0.1 mg ml⁻¹ Coomassie Brilliant Blue G-250, 5%
220 ethanol, 8.5% H₃PO₄). After mixing and incubation for 10 min at room temperature (21°C),
221 absorbance was recorded at 595 nm. Bovine albumin serum (BSA) was used as protein
222 standard.

223 **Statistical analysis**

224 Statistical analysis was performed to spot significant differences ($P < 0.05$) using SPSS
225 (SPSS© Inc., version 14.0.1) by employing tests as followed: analysis of variance (ANOVA)
226 or non-parametric tests; additional post-hoc Tukey or Hochberg's GT2 test for equal or
227 unequal sample sizes respectively to compare acclimation treatments or assay temperatures;
228 Pearson's correlation analysis to test for allometric effects; Kolmogorov-Smirnov and
229 Levene's test assessed normality and homogeneity of variances, respectively. Data were
230 expressed as means and the range of their 95% confidence interval if not stated otherwise.
231 Outliers were detected using Nalimov's test ($P < 0.01$) and excluded if justified. The
232 temperature coefficient Q_{10} was calculated as follows:

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{\frac{10}{(T_2 - T_1)}}$$

233 R denotes the respiratory rate at a higher (T_2) or lower temperature (T_1). Mitochondrial
234 respiration was expressed per mg wet weight of blotted heart fibres.

235 RESULTS

236 Evolutionary Adaptation

237 Mitochondrial respiration differed clearly between the two genetically distinct cuttlefish
238 populations. On average, cardiac mitochondria, fully fuelled by substrates and ADP (state 3),
239 showed significantly higher respiration rates in Adriatic cuttlefish with $41.1 \text{ pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$
240 (95% CI range from 37.3 to 44.9 $\text{pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$) compared to only $32.9 \text{ pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$
241 (30.0-35.9) as seen in English Channel cuttlefish (ANOVA, $F(1, 128) = 11.92$, $P < 0.01$,
242 calculations based on entire pooled data set). These 20% higher cardiac aerobic capacities in
243 cuttlefish from warmer waters were most pronounced at lower assay temperatures, and were
244 mainly due to a higher contribution of pyruvate to overall respiration (average contribution of
245 $15.2 \text{ pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$, (13.1-17.3)) for Adriatic cuttlefish compared to $9.0 \text{ pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$,
246 (7.8-10.2) for English Channel cuttlefish, Figure 1). Consequently, fractions of pyruvate
247 dependent respiration constituted 35.4% (32.8-38.0) in systemic hearts of Adriatic Sea but
248 only 26.2% (24.1-28.3) in English Channel cuttlefish. Conversely, proline dependent fractions
249 were higher (45.2% (42.5-47.8)) in English Channel- than in Adriatic Sea cuttlefish (35.2%
250 (32.4-38.0)). Succinate dependent fractions were similar in the two populations.

251 While body weights did not differ between populations (ANOVA, $F(1, 64) = 2.36$, $P =$
252 0.13), English Channel cuttlefish contained 43% heavier systemic and 38% heavier branchial
253 hearts than Adriatic cuttlefish (systemic hearts: Kruskal-Wallis, $H(1) = 13.69$, $P < 0.01$;
254 branchial hearts: Kruskal-Wallis, $H(1) = 10.22$, $P < 0.01$, for raw data see supplementary
255 table S2). Relative weights of systemic hearts constituted 0.083% (0.078-0.088) in English
256 Channel and 0.059% (0.056-0.062) in Adriatic cuttlefish. Similarly, relative weights of
257 branchial hearts constituted 0.047% (0.044-0.050) in English Channel and 0.039% (0.036-
258 0.042) in Adriatic cuttlefish. As a result, respiration calculated for the whole systemic heart
259 (state 3 respiration x heart weight) did not differ between populations (ANOVA, $F(1, 41) =$
260 0.03 , $P = 0.86$), due to the larger hearts in English Channel cuttlefish.

261 Further, systemic hearts of Adriatic cuttlefish showed partly higher protein specific
262 enzyme activities compared to English Channel cuttlefish, which was most apparent for
263 octopine dehydrogenase (Table 1). Enzyme activities of branchial hearts were similar between
264 populations except for enhanced lactate dehydrogenase activities in English Channel
265 cuttlefish acclimated to 11°C and 16°C (Table 1).

266 **Thermal acclimation**

267 Acclimation of cuttlefish to 11°C, 16°C and 21°C did not affect overall maximum state 3
 268 respiration in the English Channel (ANOVA, $F(2, 68) = 0.04$, $P = 0.96$) and in the Adriatic
 269 Sea populations (Kruskal-Wallis, $H(2) = 0.94$, $P = 0.63$) but caused minor shifts in substrate
 270 dependent fractions (Figure 1) and slight changes in cardiac enzyme activities of English
 271 Channel cuttlefish, shown by enhanced citrate synthase activities at 11°C and lower lactate
 272 dehydrogenase activities at 21°C compared to animals acclimated to 16°C (Table 1). Also,
 273 relative heart weights did not change with thermal acclimation for systemic- (ANOVA,
 274 English Channel $F(2, 29) = 0.50$, $P = 0.61$, Adriatic Sea $F(2, 30) = 0.95$, $P = 0.40$) and
 275 branchial hearts (ANOVA, English Channel $F(2, 29) = 0.46$, $P = 0.64$, Adriatic Sea $F(2, 30)$
 276 $= 3.0$, $P = 0.07$).

277 Thermal acclimation affected the activity of mitochondrial complexes in systemic heart
 278 fibres of cuttlefish. In this regard, Adriatic cuttlefish acclimated to 21°C showed between 10-
 279 19% lower complex I activity compared to cuttlefish acclimated to 11°C and 16°C whereas
 280 complex I activity in English Channel cuttlefish did not respond to thermal acclimation
 281 (Figure 2A). In contrast, systemic hearts of English Channel cuttlefish contained
 282 mitochondria that displayed increased complex IV activity following acclimation to 21°C, at
 283 assay temperatures of 11°C and 21°C (Figure 2B). In Adriatic cuttlefish hearts, however,
 284 complex IV activity remained unaffected by thermal acclimation but showed between 8-30%
 285 higher enzyme activity at lower acclimation temperatures compared to English Channel
 286 cuttlefish (Figure 2B).

287 **Thermal sensitivity of cardiac mitochondria**

288 In both cuttlefish populations cardiac mitochondria responded similarly to acute
 289 temperature changes ranging from 11°C to 21°C. State 3 respiration increased from 20.4
 290 (18.1-22.8) to 46.4 pmol O₂ s⁻¹ mg⁻¹ (42.9-49.9) in English Channel and from 27.8 (24.8-30.8)
 291 to 51.2 pmol O₂ s⁻¹ mg⁻¹ (45.3-57.1) in Adriatic cuttlefish, with average Q₁₀ of 2.3 in English
 292 Channel and 2.0 in Adriatic animals (Figure 3). Respiration resolved for substrates revealed a
 293 high thermal sensitivity of proline- and pyruvate stimulated respiration but a low thermal
 294 sensitivity of succinate stimulated respiration (Figure 3). This caused succinate to contribute
 295 most to overall state 3 respiration at 11°C assay temperature (English Channel 40.4% (37.8-
 296 43.0); Adriatic Sea 43.4% (39.6-47.2)) but to be a minor substrate at 21°C (English Channel
 297 18.7% (15.9-21.5); Adriatic Sea 19.0% (16.0-22.0)). Conversely, proline and pyruvate

298 prevailed as oxidative substrates at 21°C assay temperature, whereas their fractions declined
299 upon cooling towards 11°C (Figure 1).

300 Mitochondrial complexes displayed different temperature dependencies upon acute
301 exposure. Similar to succinate stimulated respiration (which indicates activity of complex II)
302 the activity of complex IV (indicated by respiration fuelled with ascorbate and TMPD)
303 showed low thermal sensitivity (Figure 3).

304 Interestingly, although absolute leak rates increased with warmer assay temperatures up to
305 21°C, mitochondrial proton leak relative to ADP stimulated respiration decreased in systemic
306 hearts of English Channel cuttlefish and by trend in Adriatic cuttlefish upon warming (Figure
307 4A, B). Outer membrane integrity, tested by cytochrome c addition, decreased with increasing
308 assay temperatures in both cuttlefish populations (Figure 5). Further, English Channel
309 cuttlefish showed more defective outer membranes and higher relative proton leak at 11°C
310 assay temperature compared to Adriatic cuttlefish (Figure 5).

311 **DISCUSSION**

312 **Evolutionary adaptation**

313 There is strong evidence for evolutionary changes of heart function between the
314 genetically distinct temperate and subtropical cuttlefish populations. Systemic hearts of
315 temperate (English Channel) cuttlefish contained lower aerobic and partly lower anaerobic
316 capacities (Figure 1 & Table 1) that may reduce their cardiac output and hence lower their
317 ability to increase heart rates during exercise or elevated temperatures. Whether mitochondrial
318 density parameters, enzyme numbers or specific enzyme activities accounted for the
319 differential aerobic capacities remains to be resolved. Nevertheless, similar total heart
320 respiration between populations showed that English Channel cuttlefish compensate for
321 reduced respiratory capacities by larger hearts. Compensatory increases of heart weight are
322 also common in teleost fish living at cooler temperatures (Foster et al., 1993; Driedzic et al.,
323 1996). Larger hearts support pumping of larger blood volumes per stroke, thereby, they may
324 compensate for low energetic capacities but also for rising blood viscosities at colder
325 temperatures (Goolish, 1987; Driedzic et al., 1996). Thus, cuttlefish heart function in the cold
326 may rather be sustained through an increase in organ size, than through an increase of cellular
327 energetic capacities.

328 Furthermore, the findings of predominant oxidation of proline in systemic hearts of
329 temperate cuttlefish but enhanced pyruvate oxidation in subtropical cuttlefish are well in line
330 with findings for cold adapted fish that show suppressed carbohydrate metabolism but
331 enhanced lipid oxidation (Crockett and Sidell, 1990; Sidell et al., 1995). Fuels like lipids and
332 proline are less oxygen efficient than carbohydrates, which are generally favoured in
333 cephalopods (Hochachka, 1994). Carbohydrate oxidation reduces oxygen consumption of the
334 heart and threats from tissue hypoxia due to higher ATP yields per molecule of oxygen
335 consumed (Higgins et al., 1980; Kahles et al., 1982; Hochachka, 1994). In cephalopod hearts,
336 a mole of proline requires half a mole of dioxygen to oxidize it to glutamate before entering
337 the Krebs cycle, thus decreasing its oxygen efficiency (Mommensen and Hochachka, 1981).
338 Therefore, impaired oxygen supply at higher temperatures or environmental hypoxia, which
339 cuttlefish face often in northern Adriatic lagoons (Diaz, 2001; Sorokin and Dallochio, 2008),
340 may favour carbohydrates and thus, pyruvate as the more oxygen efficient substrate.

341 **Thermal acclimation**

342 *Substrate oxidation*

343 In cuttlefish hearts, thermal acclimation did not affect aerobic capacities (i.e. state 3
344 respiration, Figure 1) and caused only minor changes of citrate synthase and lactate
345 dehydrogenase activities in temperate cuttlefish (Table 1). At first glance, this contrasts with
346 the thermal compensation found for English Channel cuttlefish. Here, routine metabolic rates
347 of cuttlefish acclimated to 20°C fell below routine metabolic rates of individuals acclimated
348 to 15°C, once above 20°C experimental temperature, thereby supporting an upward shift of
349 limiting temperatures beyond 23°C (Melzner, 2005). This pattern of thermal acclimation may
350 well be explained by a suppression of oxygen consumption rates in organs other than the
351 hearts (e.g. hepatopancreas, mantle). Such a one-sided reduction of oxygen consumption in
352 some tissues but concomitant maintenance of cardiac capacities may then free aerobic scope
353 necessary to shift thermal tolerance upwards. On the other hand, in parallel to cardiac
354 capacities, routine metabolic rates decline by 2-3.5 times from 20°C to 8°C to equally low
355 levels irrespective of the acclimation mode (Melzner, 2005). As a result, cardiac as well as
356 whole animal energetic capacities decline with decreasing temperatures and thus match a
357 reduction of energy turnover during the cold season, marked by reduced growth rates of
358 cuttlefish during winter (Le Goff and Daguzan, 1991).

359 *Mitochondrial complexes*

360 Effects of warm acclimation on mitochondrial complexes I and IV in systemic hearts of
361 cuttlefish suggest modifications that delay heat-induced tissue hypoxia and formation of
362 reactive oxygen species (ROS). Due to constraints on their systemic (i.e. low blood oxygen
363 carrying capacity) and intracellular oxygen transport system (e.g. lack of intracellular oxygen
364 delivering protein like myoglobin, (Hochachka, 1994)), cephalopods are chronically prone to
365 facing critically low intracellular oxygen concentrations, particularly when high temperatures
366 induce tissue hypoxia (Pörtner, 2001). Gnaiger et al. (1998) argued that an excess capacity of
367 complex IV sustains high affinities for oxygen in mitochondria. Therefore, the increase of
368 complex IV capacity in systemic hearts of English Channel cuttlefish following warm
369 acclimation to 21°C (Figure 2B) likely enhanced oxygen affinity, thus supporting oxygen
370 diffusion to mitochondria at high and hypoxia inducing temperatures. This response is in
371 agreement with fishes that are capable to increase cardiac complex IV activity following
372 warm-acclimation as seen in the highly hypoxia tolerant carp (Cai and Adelman, 1990) but
373 also in cod (Foster et al., 1993) or in the liver of Antarctic eelpout (Windisch et al., 2011).

374 ROS formation in response to heat stress and environmental hypoxia is common among
375 marine ectotherms (Abele et al., 2007), and may occur more often in the warm and often
376 hypoxic lagoons of the northern Adriatic Sea during summer (Diaz, 2001; Sorokin et al.,
377 2002) where cuttlefish are common (Rossetto, 2001). Complex I is one of the major sites of
378 ROS production and has been related to cardiac failure (Ide et al., 1999; Sorescu and
379 Griendling, 2002), therefore, a suppression of complex I capacity following warm-
380 acclimation in Adriatic cuttlefish (Figure 2A) may reduce heat and hypoxia related ROS
381 formation to preserve mitochondrial function. As a corollary, changed activities of
382 mitochondrial complexes following warm acclimation may serve cuttlefish to reduce
383 temperature induced hypoxia or harmful oxygen stress in cardiac tissues.

384 **Thermal sensitivity of cardiac mitochondria**

385 *Substrate oxidation*

386 During summer, cuttlefish pass a steep thermocline during their daily vertical migration
387 with possible temperature changes of up to 2°C per m (e.g. Adriatic Sea: 23°C to 14°C down
388 to 80 m; English Channel: 19°C to 11°C down to 40 m) (Artegiani et al., 1997; Sharples et al.,
389 2001). Our findings demonstrate that noticeable increases of mitochondrial respiration (state
390 3) from 11°C to 21°C support systemic hearts of cuttlefish to operate aerobically over this
391 range of temperatures. Interestingly, various substrates contributed differently to this thermal
392 response (Figure 3). The substrate dependent effect on thermal sensitivity of oxidative
393 pathways may allow mitochondria to produce aerobic energy over a broader range of
394 temperatures. While less temperature sensitive pathways (e.g. that augment succinate)
395 attenuate a rapid decline of mitochondrial respiration (i.e. ATP provision) at low
396 temperatures, thermally more sensitive pathways (e.g. fuelled by pyruvate or proline)
397 complement less responsive pathways at higher temperatures to match enhanced ATP
398 demands of the heart. Although pyruvate and proline oxidation declines at low temperatures,
399 *in situ* supply of succinate may be sustained by amino acids such as glutamate, ornithine or
400 arginine that are well oxidized and readily available from the blood or intracellular stores
401 (Ballantyne et al., 1981; Mommsen and Hochachka, 1981; Hochachka and Fields, 1982;
402 Mommsen et al., 1983). Observations in fish hearts support this pattern, as fatty acid
403 oxidation has mostly lower or at least different thermal sensitivities than carbohydrate
404 oxidation, which kicks in in the warmth (Sephton et al., 1990; Sephton and Driedzic, 1991).

405 The predominance of pyruvate and proline oxidation at higher temperatures (Figure 1)
406 improves oxygen efficient production of ATP and likely reduces the threat of tissue hypoxia.

407 After entering mitochondria, proline is oxidized to glutamate and then enters the Krebs cycle
408 via alpha-ketoglutarate and yields 4-5 NADH and 1 FADH₂ (i.e. assuming no diffusive loss of
409 products), similar to pyruvate that yields 4 NADH and 1 FADH₂ during its oxidation via the
410 citric acid cycle (Storey and Storey, 1983). As a consequence, proline and pyruvate feed about
411 80% of their electrons into complex I and thus produce more ATP per mole of oxygen
412 consumed. In contrast, succinate yields 1 NADH and 1 FADH₂ and thus, diverts only 50% of
413 its electrons to complex I when fully catabolized to oxaloacetate. Oxygen efficient ATP
414 production may be advantageous at high temperatures when oxygen becomes limiting
415 ((Pörtner, 2010) for recent review) and is well in line with the general trend of modifications
416 in substrate use by mitochondria that enhance oxygen efficiency in cephalopods (Hochachka,
417 1994).

418 *Membrane leakiness and integrity*

419 Surprisingly, in cuttlefish hearts, relative proton leak decreased and outer membrane
420 integrity increased (i.e. shown by decreased stimulation of respiration by cytochrome c) up to
421 a temperature close to the whole animal critical temperature of 23°C ((Melzner et al., 2007b),
422 Figure 4, 5). In contrast to the present observation, relative proton leak increases with
423 temperature in mitochondria isolated from hearts, red muscle and liver of fish (Hardewig et
424 al., 1999; Fangue et al., 2009; Hilton et al., 2010), gills of Antarctic bivalves (Pörtner et al.,
425 1999) or the body wall of a lugworm (Sommer and Pörtner, 2002). Our finding thus contrasts
426 with the general view that higher temperatures increase membrane fluidity and thus proton
427 leak (Hazel, 1995; Pörtner, 2001). This could be explained by a decrease of mitochondrial
428 membrane potential via increased ATP synthase activity relative to electron flux through the
429 electron transport system, which would induce reduced proton gradients over the membrane
430 leading to reduced proton leak (Nicholls, 2004). Alternatively, structural changes of
431 mitochondrial membranes that reduce membrane permeability may have occurred, as
432 indicated by reduced respiratory stimulation by cytochrome c (Figure 5), which passes the
433 outer membrane by diffusion only (Gellerich et al., 2000). Such evident changes of the outer
434 membrane permeability likely affected the inner mitochondrial membrane too and hence
435 supported the observed decline of relative proton leak with rising temperatures. Even though
436 the underlying mechanisms remain unclear, decreasing relative proton leak with acutely rising
437 temperature aids cuttlefish hearts to be more oxygen efficient and thereby shift temperatures
438 that entail oxygen deficiency to higher tolerated values. These functional characteristics may
439 be adaptive in supporting eurythermy of this species.

440 Interestingly, temperate English Channel cuttlefish mitochondria displayed lower
441 respiratory capacities and contained more permeable and therefore less efficient membranes at
442 11°C assay temperature than the subtropical Adriatic cuttlefish (Figure 4, 5). Their
443 mitochondria operate thus less efficient and more “costly” at cool temperatures. Conversely, a
444 higher “futile cycling” through proton leak would make them more responsive to sudden
445 increases in workloads. This may be needed less in warm acclimated hearts.

446 **Conclusion**

447 Our study about cuttlefish heart fibres and their mitochondria revealed inherent potential
448 to cope with thermal challenges faced during an individual’s lifetime, but also genetic
449 plasticity between populations relevant for adaptation to long-term environmental change.
450 Most modifications improve cardiac efficiency and extend tolerance to high temperatures and
451 associated hypoxemic conditions. Cold compensation on evolutionary time scales occurs via a
452 shift in substrate and enhanced proton leak as a sign of enhanced futile cycling in
453 mitochondrial metabolism. Lack of cold compensation by adjusting mitochondrial or enzyme
454 capacities during cold acclimation, on the other hand, suggest decreased cardiac energetic
455 capacities during the cold season. Overall, we conclude that the observed flexibility of cardiac
456 function, based on specific cephalopod-type characteristics, assures cardiac power output and
457 systemic oxygen delivery at various temperatures. It thus conforms to the capacity of *Sepia*
458 *officinalis* to tolerate a broad range of temperatures and supports their ability to contend with
459 fishes - their prime competitors - in rapidly changing environments.

460 **LIST OF SYMBOLS AND ABBREVIATIONS**

461	ATP	Adenosin-5'-triphosphate
462	BSA	Bovine Serum Albumin
463	Complex I	NADH dehydrogenase
464	Complex IV	Cytochrome c oxidase
465	DTNB	5,5'-dithio-bis-(2-nitro-benzoic acid)
466	DTT	Dithiothreitol
467	EDTA	Ethylenediaminetetraacetic acid
468	EGTA	Ethylene glycol tetraacetic acid
469	HEPES	N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid)
470	LDH	Lactate dehydrogenase
471	ODH	Octopine dehydrogenase
472	Q ₁₀	Temperature coefficient
473	ROS	Reactive oxygen species
474	TMPD	N,N,N',N'-tetramethyl-p-phenylenediamine dihydrochloride

475 **ACKNOWLEDGEMENTS**

476 We would like to thank Guido Krieten, Fredy Veliz Moraleda, Timo Hirse and Nils
477 Koschnick for technical support, Anne Thonig for conducting parts of the experiments as well
478 as colleagues for constant support and advice. Special thanks to Anthony J. R. Hickey
479 (University of Auckland) for his technical and theoretical training to apply the skinned fibre
480 technique. We further would like to thank the two anonymous reviewers for their many
481 helpful comments and suggestions on an earlier version of this manuscript.

482 **FUNDING**

483 This study was supported by the Alfred Wegener Institute for Polar and Marine Research
484 and is a contribution to Natural Environment Research Council Grant
485 NERC/A/S/2002/00812.

486 FIGURES LEGENDS

487 Figure 1: Maximum state 3 respiration ($\text{pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$) resolved for each substrate,
488 after addition of ADP, at 11°C, 16°C and 21°C assay temperature in comparison
489 between acclimation treatments ($N = 5-11$) as well as between cuttlefish from (A) the
490 English Channel and (B) Adriatic Sea. Significant differences ($P < 0.05$) between
491 acclimation groups are marked by asterisks and for total state 3 respiration between
492 populations by roofs. Values are means \pm 95% C.I..

493
494 Figure 2: Effects of thermal acclimation on mitochondrial complex I / NADH
495 dehydrogenase (A, relative decrease (%) of uncoupled respiration following rotenone
496 addition) and complex IV / cytochrome c oxidase activity (B, respiration ($\text{pmol O}_2 \text{ s}^{-1}$
497 mg^{-1}) after addition of ascorbate and TMPD) in systemic hearts of cuttlefish from the
498 English Channel and Adriatic Sea. Significant differences ($P < 0.05$) between
499 acclimation temperatures ($N=5-11$) are marked by asterisks and between populations
500 by roofs.

501
502 Figure 3: Temperature coefficients (Q_{10} , means \pm 95% C.I.) of cardiac mitochondrial
503 respiration from 11°C to 21°C, resolved for each substrate and for total state 3 and
504 complex IV respiration. As there were no differences between acclimations, data
505 were pooled for each case. Q_{10} of pyruvate and succinate are based on stimulating
506 respiration.

507
508 Figure 4: Change of proline stimulated respiration without ADP (denoted as proton
509 leak) and with ADP ($\text{pmol O}_2 \text{ s}^{-1} \text{ mg}^{-1}$) and relative proton leak (% , calculated as
510 fraction of leak rates relative to coupled respiration with ADP) from 11°C to 21°C
511 assay temperature in systemic hearts, in comparison between (A) English Channel
512 ($N=24$) and (B) Adriatic ($N=19-20$) cuttlefish. Due to the lack of differences between
513 acclimation treatments, data were pooled for each assay temperature and
514 population. Values are means \pm 95% C.I. Significant differences ($P < 0.05$) between
515 assay temperatures for relative proton leak are marked by unequal letters of the
516 same case and between populations by roofs.

517

518 Figure 5: Relative change of maximum state 3 respiration (%) following cytochrome c
519 addition, from 11°C to 21°C assay temperature in systemic hearts, in comparison
520 between English Channel ($N=23-24$) and Adriatic ($N=19-20$) cuttlefish. Due to the
521 lack of differences between acclimation treatments, data were pooled for each assay
522 temperature and population. Values are means \pm 95% C.I.. Significant differences (P
523 < 0.05) between assay temperatures are marked by unequal letters of the same case
524 and between populations by roofs.

525

526 Table 1: Enzyme activities ($\mu\text{mol min}^{-1} \text{mg protein}^{-1}$, at 16°C) of (A) citrate synthase,
527 (B) lactate dehydrogenase and (C) octopine dehydrogenase of cuttlefish systemic
528 and branchial hearts, in comparison between thermal acclimations, as well as
529 between English Channel and Adriatic Sea populations.

530 **REFERENCES**

- 531 **Abele, D., Philipp, E., Gonzalez, P. and Puntarulo, S.** (2007). Marine invertebrate
532 mitochondria and oxidative stress. *Front. Biosci.* **12**, 933-946.
- 533 **Agnisola, C., Driedzic, W. R., Foster, A. R., Houlihan, D. F. and Stewart, J. M.** (1991).
534 Oxygen consumption, carbon dioxide production and enzyme activities of isolated
535 working Octopus heart. *J. Exp. Biol.* **157**, 543-549.
- 536 **Artegiani, A., Paschini, E., Russo, A., Bregant, D., Raicich, F. and Pinardi, N.** (1997).
537 The Adriatic Sea general circulation. Part I: Air-sea interactions and water mass
538 structure. *J. Phys. Oceanogr.* **27**, 1492-1514.
- 539 **Ballantyne, J., Hochachka, P. and Mommsen, T.** (1981). Studies on the metabolism of the
540 migratory squid, *Loligo opalescens*: enzymes of tissues and heart mitochondria. *Mar.*
541 *Biol. Lett* **2**, 75-85.
- 542 **Boucaud-Camou, E. and Boismery, J.** (1991). The migrations of the cuttlefish (*Sepia*
543 *officinalis* L.) in the English Channel. In *Acta of the 1st International Symposium on*
544 *the Cuttlefish Sepia*, (ed. E. Boucaud-Camou), pp. 179-189. Caen: Université de Caen
545 Publications.
- 546 **Bradford, M.** (1976). A rapid and sensitive method for the quantitation of microgram
547 quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**,
548 248-254.
- 549 **Cai, Y. J. and Adelman, I. R.** (1990). Temperature acclimation in respiratory and
550 cytochrome c oxidase activity in common carp (*Cyprinus carpio*). *Comp. Biochem.*
551 *Physiol. A Physiol.* **95**, 139-144.
- 552 **Crockett, E. and Sidell, B.** (1990). Some pathways of energy metabolism are cold adapted in
553 Antarctic fishes. *Physiol. Zool.* **63**, 472-488.
- 554 **Diaz, R. J.** (2001). Overview of hypoxia around the world. *J. Environ. Qual.* **30**, 275-281.
- 555 **Driedzic, W., Bailey, J. and Sephton, D.** (1996). Cardiac adaptations to low temperature in
556 non-polar teleost fish. *J. Exp. Zool.* **275**, 186-195.
- 557 **Driedzic, W., Sidell, B., Stewart, J. and Johnston, I.** (1990). Maximal activities of enzymes
558 of energy metabolism in cephalopod systemic and branchial hearts. *Physiol. Zool.* **63**,
559 615-629.
- 560 **Fangue, N., Richards, J. and Schulte, P.** (2009). Do mitochondrial properties explain
561 intraspecific variation in thermal tolerance? *J. Exp. Biol.* **212**, 514-522.

- 562 **Farrell, A. P.** (2002). Cardiorespiratory performance in salmonids during exercise at high
563 temperature: insights into cardiovascular design limitations in fishes. *Comp. Biochem.*
564 *Physiol. Part A Mol. Integr. Physiol.* **132**, 797-810.
- 565 **Foster, A., Hall, S. and Houlihan, D.** (1993). The effects of temperature acclimation on
566 organ/tissue mass and cytochrome c oxidase activity in juvenile cod (*Gadus morhua*).
567 *J. Fish Biol.* **42**, 947-957.
- 568 **Frederich, M. and Pörtner, H. O.** (2000). Oxygen limitation of thermal tolerance defined by
569 cardiac and ventilatory performance in spider crab, *Maja squinado*. *Am. J. Physiol.*
570 *Regul. Integr. Comp. Physiol.* **279**, R1531-R1538
- 571 **Gellerich, F., Trumbeckaite, S., Opalka, J., Seppet, E., Rasmussen, H. N., Neuhoff, C.**
572 **and Zierz, S.** (2000). Function of the mitochondrial outer membrane as a diffusion
573 barrier in health and diseases. *Biochem. Soc. Trans.* **28**, 164-169.
- 574 **Gnaiger, E., Lassnig, B., Kuznetsov, A., Rieger, G. and Margreiter, R.** (1998).
575 Mitochondrial oxygen affinity, respiratory flux control and excess capacity of
576 cytochrome c oxidase. *J. Exp. Biol.* **201**, 1129-1139.
- 577 **Goolish, E.** (1987). Cold-acclimation increases the ventricle size of carp, *Cyprinus carpio*. *J.*
578 *Therm. Biol.* **12**, 203-205.
- 579 **Grieshaber, M. and Gäde, G.** (1976). The biological role of octopine in the squid, *Loligo*
580 *vulgaris* (Lamarck). *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **108**, 225-
581 232.
- 582 **Hanley, J. S., Shashar, N., Smolowitz, R., Bullis, R. A., Mebane, W. N., Gabr, H. R. and**
583 **Hanlon, R. T.** (1998). Modified laboratory culture techniques for the European
584 cuttlefish *Sepia officinalis*. *Biol. Bull. (Woods Hole)* **195**, 223-225.
- 585 **Hardewig, I., Pörtner, H. O. and Peck, L. S.** (1999). Thermal sensitivity of mitochondrial
586 function in the Antarctic Notothenioid *Lepidonotothen nudifrons*. *J. Comp. Physiol. B*
587 *Biochem. Syst. Environ. Physiol.* **169**, 597-604.
- 588 **Hazel, J. R.** (1995). Thermal adaptation in biological membranes: is homeoviscous
589 adaptation the explanation? *Annu. Rev. Physiol.* **57**, 19-42.
- 590 **Higgins, A., Morville, M., Burges, R., Gardiner, D., Page, M. and Blackburn, K.** (1980).
591 Oxfenicine diverts rat muscle metabolism from fatty acid to carbohydrate oxidation
592 and protects the ischaemic rat heart. *Life Sci.* **27**, 963-970.
- 593 **Hilton, Z., Clements, K. D. and Hickey, A. J. R.** (2010). Temperature sensitivity of cardiac
594 mitochondria in intertidal and subtidal triplefin fishes. *J. Comp. Physiol. B Biochem.*
595 *Syst. Environ. Physiol.* **180**, 979-990.

- 596 **Hochachka, P.** (1994). Oxygen efficient design of cephalopod muscle metabolism. *Mar.*
597 *Freshw. Behav. Physiol.* **25**, 61-67.
- 598 **Hochachka, P. W. and Fields, J. H. A.** (1982). Arginine, glutamate, and proline as
599 substrates for oxidation and for glycogenesis in cephalopod tissues. *Pac. Sci.* **36**, 325-
600 335.
- 601 **Ide, T., Tsutsui, H., Kinugawa, S., Utsumi, H., Kang, D., Hattori, N., Uchida, K.,**
602 **Arimura, K., Egashira, K. and Takeshita, A.** (1999). Mitochondrial electron
603 transport complex I is a potential source of oxygen free radicals in the failing
604 myocardium. *Circ. Res.* **85**, 357-363.
- 605 **Iftikar, F. I., MacDonald, J. and Hickey, A. J. R.** (2010). Thermal limits of portunid crab
606 heart mitochondria: Could more thermo-stable mitochondria advantage invasive
607 species? *J. Exp. Mar. Biol. Ecol.* **395**, 232-239.
- 608 **Jereb, P. and Roper, C. F. E.** (2005). Cephalopods of the World. An annotated and
609 illustrated catalogue of species known to date. In *FAO Species Catalogue for Fishery*
610 *Purposes* vol. 1 (ed. F.A.O.), pp. 1-277. Rome.
- 611 **Johnston, I. and Harrison, P.** (1987). Morphometrics and ultrastructure of myocardial tissue
612 in Notothenioid fishes. *Fish Physiol. Biochem.* **3**, 1-6.
- 613 **Kahles, H., Hellige, G., Hunneman, D., Mezger, V. A. and Bretschneider, H.** (1982).
614 Influence of myocardial substrate utilization on the oxygen consumption of the heart.
615 *Clin. Cardiol.* **5**, 286-293.
- 616 **Kent, J., Koban, M. and Prosser, C.** (1988). Cold-acclimation-induced protein hypertrophy
617 in channel catfish and green sunfish. *J. Comp. Physiol. B Biochem. Syst. Environ.*
618 *Physiol.* **158**, 185-198.
- 619 **Kleckner, N. and Sidell, B.** (1985). Comparison of maximal activities of enzymes from
620 tissues of thermally acclimated and naturally acclimatized chain pickerel (*Esox niger*).
621 *Physiol. Zool.* **58**, 18-28.
- 622 **Kolok, A. S.** (1991). Temperature compensation in two centrarchid fishes: Do winter-
623 quiescent fish undergo cellular temperature compensation? *Trans. Am. Fish. Soc.* **120**,
624 52-57.
- 625 **Kuznetsov, A. V., Veksler, V., Gellerich, F. N., Saks, V., Margreiter, R. and Kunz, W. S.**
626 (2008). Analysis of mitochondrial function in situ in permeabilized muscle fibers,
627 tissues and cells. *Nat. Protoc.* **3**, 965-976.
- 628 **Le Goff, R. and Daguzan, J.** (1991). Growth and life cycles of the cuttlefish *Sepia officinalis*
629 L. (Mollusca: Cephalopoda) in south Brittany (France). *Bull. Mar. Sci.* **49**, 341-348.

- 630 **Melzner, F.** (2005). Systemic investigations on the physiology of temperature tolerance in the
631 common cuttlefish *Sepia officinalis*. In *Alfred Wegener Institute for Polar and Marine*
632 *Research*, vol. PhD, pp. 234. Bremerhaven: University of Bremen.
- 633 **Melzner, F., Bock, C. and Pörtner, H. O.** (2007a). Coordination between ventilatory
634 pressure oscillations and venous return in the cephalopod *Sepia officinalis* under
635 control conditions, spontaneous exercise and recovery. *J. Comp. Physiol. B Biochem.*
636 *Syst. Environ. Physiol.* **177**, 1-17.
- 637 **Melzner, F., Mark, F. C. and Pörtner, H. O.** (2007b). Role of blood-oxygen transport in
638 thermal tolerance of the cuttlefish, *Sepia officinalis*. *Integr. Comp. Biol.* **47**, 645-655.
- 639 **Mommsen, T. P. and Hochachka, P. W.** (1981). Respiratory and enzymatic properties of
640 squid heart mitochondria. *Eur. J. Biochem.* **120**, 345-350.
- 641 **Mommsen, T. P., Hochachka, P. W. and French, C. J.** (1983). Metabolism of arginine,
642 proline, and ornithine in tissues of the squid, *Illex illecebrosus*. *Can. J. Zool.* **61**, 1835-
643 1846.
- 644 **Nicholls, D. G.** (2004). Mitochondrial membrane potential and aging. *Aging Cell* **3**, 35-40.
- 645 **O'Dor, R. K. and Webber, D. M.** (1986). The constraints on cephalopods: why squid aren't
646 fish. *Can. J. Zool.* **64**, 1591-1605.
- 647 **O'Dor, R. K. and Webber, D. M.** (1991). Invertebrate athletes: trade-offs between transport
648 efficiency and power density in cephalopod evolution. *J. Exp. Biol.* **160**, 93-112.
- 649 **Perez-Losada, M., Nolte, M. J., Crandall, K. A. and Shaw, P. W.** (2007). Testing
650 hypotheses of population structuring in the Northeast Atlantic Ocean and
651 Mediterranean Sea using the common cuttlefish *Sepia officinalis*. *Mol. Ecol.* **16**, 2667-
652 2679.
- 653 **Podrabsky, J. E., Javillonar, C., Hand, S. C. and Crawford, D. L.** (2000). Intraspecific
654 variation in aerobic metabolism and glycolytic enzyme expression in heart ventricles.
655 *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **279**, R2344-R2348
- 656 **Pörtner, H. O.** (1994). Coordination of metabolism, acid-base regulation and haemocyanin
657 function in cephalopods. *Mar. Behav. Physiol.* **25**, 131-148.
- 658 **Pörtner, H. O.** (2001). Climate change and temperature-dependent biogeography: oxygen
659 limitation of thermal tolerance in animals. *Naturwissenschaften* **88**, 137-146.
- 660 **Pörtner, H. O.** (2002a). Climate variations and the physiological basis of temperature
661 dependent biogeography: systemic to molecular hierarchy of thermal tolerance in
662 animals. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **132**, 739-761.

- 663 **Pörtner, H. O.** (2002b). Environmental and functional limits to muscular exercise and body
664 size in marine invertebrate athletes. *Comp. Biochem. Physiol. Part A Mol. Integr.*
665 *Physiol.* **133**, 303-321.
- 666 **Pörtner, H. O.** (2010). Oxygen-and capacity-limitation of thermal tolerance: a matrix for
667 integrating climate-related stressor effects in marine ecosystems. *J. Exp. Biol.* **213**,
668 881.
- 669 **Pörtner, H. O. and Zielinski, S.** (1998). Environmental constraints and the physiology of
670 performance in squids. *S. Afr. J. Mar. Sci.* **20**, 207-221.
- 671 **Pörtner, H. O. and Knust, R.** (2007). Climate Change Affects Marine Fishes Through the
672 Oxygen Limitation of Thermal Tolerance. *Science* **315**, 95-97.
- 673 **Pörtner, H. O. and Farrell, A. P.** (2008). Ecology. Physiology and climate change. *Science*
674 **322**, 690-692.
- 675 **Pörtner, H. O., Hardewig, I. and Peck, L. S.** (1999). Mitochondrial function and critical
676 temperature in the Antarctic bivalve, *Laternula elliptica*. *Comp. Biochem. Physiol.*
677 *Part A Mol. Integr. Physiol.* **124**, 179-189.
- 678 **Rosa, R. and Seibel, B. A.** (2008). Synergistic effects of climate-related variables suggest
679 future physiological impairment in a top oceanic predator. *Proc. Natl. Acad. Sci. USA*
680 **105**, 20776–20780.
- 681 **Rossetto, L.** (2001). The Management of Fishery in the Lagoon of Venice. In *IIFET2000*,
682 pp. 1-10. Corvallis: International Institute of Fisheries Economics and Trade, Oregon
683 State University.
- 684 **Saks, V. A., Veksler, V. I., Kuznetsov, A. V., Kay, L., Sikk, P., Tiivel, T., Tranqui, L.,**
685 **Olivares, J., Winkler, K. and Wiedemann, F.** (1998). Permeabilized cell and
686 skinned fiber techniques in studies of mitochondrial function in vitro. *Mol. Cell.*
687 *Biochem.* **184**, 81-100.
- 688 **Schipp, R.** (1987). General morphological and functional characteristics of the cephalopod
689 circulatory system. An introduction. *Cell. Mol. Life Sci.* **43**, 474-477.
- 690 **Sephton, D. and Driedzic, W.** (1991). Effect of acute and chronic temperature transition on
691 enzymes of cardiac metabolism in white perch (*Morone americana*), yellow perch
692 (*Perca flavescens*), and smallmouth bass (*Micropterus dolomieu*). *Can. J. Zool.* **69**,
693 258-262.
- 694 **Sephton, D., Bailey, J. and Driedzic, W.** (1990). Impact of acute temperature transition on
695 enzyme activity levels, oxygen consumption, and exogenous fuel utilization in sea

696 raven (*Hemipterus americanus*) hearts. *J. Comp. Physiol. B Biochem. Syst. Environ.*
697 *Physiol.* **160**, 511-518.

698 **Sharples, J., Moore, C. M., Rippeth, T. P., Holligan, P. M., Hydes, D. J., Fisher, N. R.**
699 **and Simpson, J. H.** (2001). Phytoplankton distribution and survival in the
700 thermocline. *Limnol. Oceanogr.* **46**, 486-496.

701 **Sidell, B., Crockett, E. and Driedzic, W.** (1995). Antarctic fish tissues preferentially
702 catabolize monoenoic fatty acids. *J. Exp. Zool.* **271**, 73-81.

703 **Sidell, B., Driedzic, W., Stowe, D. and Johnston, I.** (1987). Biochemical correlations of
704 power development and metabolic fuel preferenda in fish hearts. *Physiol. Zool.* **60**,
705 221-232.

706 **Somero, G. N.** (2010). The physiology of climate change: how potentials for acclimatization
707 and genetic adaptation will determine 'winners' and 'losers'. *J. Exp. Biol.* **213**, 912-
708 920.

709 **Sommer, A. M. and Pörtner, H. O.** (2002). Metabolic cold adaptation in the lugworm
710 *Arenicola marina*: comparison of a North Sea and a White Sea population. *Mar. Ecol.*
711 *Prog. Ser.* **240**, 171-182.

712 **Sorescu, D. and Griendling, K. K.** (2002). Reactive oxygen species, mitochondria, and
713 NAD(P)H oxidases in the development and progression of heart failure. *Congest.*
714 *Heart Fail.* **8**, 132-140.

715 **Sorokin, P. Y., Sorokin, Y. I., Zakuskina, O. Y. and Ravagnan, G. P.** (2002). On the
716 changing ecology of Venice lagoon. *Hydrobiologia* **487**, 1-18.

717 **Sorokin, Y. I. and Dallochio, F.** (2008). Dynamics of phosphorus in the Venice lagoon
718 during a picocyanobacteria bloom. *J. Plankton Res.* **30**, 1019-1026.

719 **Storey, K.** (1977). Tissue specific isozymes of octopine dehydrogenase in the cuttlefish,
720 *Sepia officinalis*. The roles of octopine dehydrogenase and lactate dehydrogenase in
721 *Sepia*. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **115**, 159-169.

722 **Storey, K. B. and Storey, J. M.** (1979). Octopine metabolism in the cuttlefish, *Sepia*
723 *officinalis*, octopine production by muscle and its role as an aerobic substrate for non-
724 muscular tissues. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **131**, 311-319.

725 **Storey, K. B. and Storey, J. M.** (1983). Carbohydrate metabolism in cephalopod molluscs.
726 In *The Mollusca: Metabolic biochemistry and molecular biomechanics*, vol. 1 (ed. P.
727 W. Hochachka), pp. 91-136. New York: Academic Press.

728 **Wang, J., Pierce, G. J., Boyle, P. R., Denis, V., Robin, J. P. and Bellido, J. M.** (2003).
729 Spatial and temporal patterns of cuttlefish (*Sepia officinalis*) abundance and

730 environmental influences - A case study using trawl fishery data in French Atlantic
731 coastal, English Channel, and adjacent waters. *ICES J. Mar. Sci.* **60**, 1149-1158.

732 **Wells, M.** (1992). The cephalopod heart: The evolution of a high-performance invertebrate
733 pump. *Cell. Mol. Life Sci.* **48**, 800-808.

734 **Wells, M. and Smith, P.** (1987). The performance of the octopus circulatory system: a
735 triumph of engineering over design. *Cell. Mol. Life Sci.* **43**, 487-499.

736 **Windisch, H. S., Kathover, R., Portner, H. O., Frickenhaus, S. and Lucassen, M.** (2011).
737 Thermal acclimation in Antarctic fish: transcriptomic profiling of metabolic pathways.
738 *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **301**, R1453-1466.

739 **Wolfram, K., Mark, F. C., John, U., Lucassen, M. and Portner, H. O.** (2006).
740 Microsatellite DNA variation indicates low levels of genetic differentiation among
741 cuttlefish (*Sepia officinalis* L.) populations in the English Channel and the Bay of
742 Biscay. *Comp Biochem Phys D* **1**, 375-383.

743

744

Figures

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Table 1

Enzyme	Heart type	Systemic heart			Branchial heart		
		Acclimation temperature (°C)					
		11	16	21	11	16	21
Citrate synthase	English Channel	0.89 (0.82-0.96) ^a	0.73 (0.65-0.82) ^b	0.74 (0.60-0.88)	0.64 (0.51-0.77)	0.78 (0.67-0.89)	0.81 (0.74-0.88)
	Adriatic Sea	1.07 (0.53-1.63)	1.01 (0.80-1.22) ▲	0.81 (0.67-0.96)	0.63 (0.53-0.74)	0.64 (0.51-0.76)	0.78 (0.63-0.93)
Lactate dehydrogenase	English Channel	0.10 (0.07-0.14)	0.11 (0.09-0.12) ^a	0.05 (0.04-0.07) ^b	0.12 (0.10-0.14)	0.10 (0.09-0.12)	0.08 (0.06-0.10)
	Adriatic Sea	0.13 (0.08-0.19)	0.09 (0.05-0.13)	0.08 (0.07-0.10) ▲	0.07 (0.05-0.10) ▼	0.07 (0.06-0.09) ▼	0.06 (0.06-0.07)
Octopine dehydrogenase	English Channel	0.40 (0.34-0.46) ▲	0.48 (0.43-0.53)	0.39 (0.29-0.48)	0.22 (0.17-0.27)	0.24 (0.20-0.27)	0.23 (0.20-0.26)
	Adriatic Sea	0.56 (0.46-0.66)	0.57 (0.35-0.80)	0.62 (0.49-0.75) ▲	0.26 (0.18-0.33)	0.23 (0.18-0.29)	0.28 (0.23-0.34)

Values are means with 95% C.I.in parentheses. $N = 5-17$

▲/▼ indicate significant differences ($P < 0.05$) between populations and unequal letters between acclimation temperatures

Table S1. Action and concentrations of the agents used to assay mitochondrial respiration of permeabilised heart fibres from the European cuttlefish

Agent	Action	Final concentration
Substrates		
Proline	Amino acid substrate	5 mmol l ⁻¹
Pyruvate	Carbohydrate substrate	5 mmol l ⁻¹
Succinate	Carbohydrate substrate	10 mmol l ⁻¹
ADP	Substrate for ATP generation	2.5 mmol l ⁻¹
Cytochrome <i>c</i>	Electron carrier, tests for outer membrane integrity	0.01 mmol l ⁻¹
TMPD	Complex IV substrate	5 mmol l ⁻¹
Ascorbate	Complex IV substrate	20 mmol l ⁻¹
Inhibitors		
Antimycin	Inhibits complex III, to test non-mitochondrial respiration	2.5 μmol l ⁻¹
Rotenone	Complex I inhibitor	2.5 μmol l ⁻¹
Oligomycin	Inhibits ATP synthase, to test proton leak	2 μg ml ⁻¹
Uncoupler		
FCCP	Uncouples the electron transport system electron flow from ATP generation by relieving the proton gradient	2.5 μmol l ⁻¹

Table S2. Measurements of the European cuttlefish *Sepia officinalis* used in acclimation experiments

Identifier	Origin	Acclimation temperature (°C)	Sex	Date of sampling	Total mass (g)	Mantle length (cm)	Systemic heart mass (mg)	Branchial heart #1 mass (mg)	Branchial heart #2 mass (mg)	Comments
A06	Adriatic	11	Male	7/07/2009	90.22	8.8	58.3	38.5	33.5	
A10	Adriatic	11	Male	14/07/2009	63.34	8.0	39.3	25.4	28.9	
A13	Adriatic	11	Male	16/07/2009	83.75	8.5	49.3	37.0	34.3	Animal was inking during capture
A30	Adriatic	11	Female	21/09/2009	99.20	9.1	58.6	38.0	37.4	Animal was inking during capture
A63	Adriatic	11	Female	14/07/2010	110.53	9.3	58.3	39.7	39.2	Animal was inking during capture
A64	Adriatic	11	Male	19/07/2010	78.47	8.2	40.1	27.8	30.9	Animal was inking during capture
A65	Adriatic	11	Female	20/07/2010	86.40	8.0	43.1	34.9	35.3	Animal was slightly inking during capture
A69*	Adriatic	11	Male	22/11/2010	48.20	–	24.6	19.9	18.4	
A07	Adriatic	16	Female	8/07/2009	46.90	6.8	30.5	23.7	22.4	
A09	Adriatic	16	Male	13/07/2009	40.57	7.0	27.0	11.6	19.5	Animal was inking during capture
A14	Adriatic	16	Male	20/07/2009	72.52	8.2	44.9	24.7	27.5	Animal was inking during capture
A19	Adriatic	16	Female	30/07/2009	77.88	7.9	46.4	42.2	38.5	Animal was inking during capture, very well developed gonads
A24	Adriatic	16	Female	10/08/2009	59.11	7.4	37.1	41.6	42.6	
A46*	Adriatic	16	Female	20/04/2010	50.10	7.1	27.3	18.3	18.7	Animal was inking during capture
A47*	Adriatic	16	Male	21/04/2010	49.87	7.0	27.1	15.0	16.5	Animal was inking during capture
A48*	Adriatic	16	Male	26/04/2010	59.67	7.5	33.7	24.5	21.8	Animal was inking during capture
A49*	Adriatic	16	Female	27/04/2010	62.66	7.2	33.0	20.9	24.7	Animal was inking during capture
A50*	Adriatic	16	Male	28/04/2010	52.05	6.9	29.1	22.1	24.3	Animal was inking during capture
A51*	Adriatic	16	Female	29/04/2010	41.34	6.2	25.6	13.6	16.7	Animal was inking during capture
A61	Adriatic	16	Male	12/07/2010	109.93	9.5	77.4	42.1	44.5	Animal was inking during capture
A62	Adriatic	16	Male	13/07/2010	144.00	10.5	101.1	56.3	56.2	
A52*	Adriatic	21	Female	26/05/2010	148.82	10.5	74.7	40.3	43.8	
A53*	Adriatic	21	Male	31/05/2010	128.71	10.0	68.7	40.0	39.8	
A54*	Adriatic	21	Female	2/06/2010	228.17	11.2	113.2	65.3	78.8	Female had proper developed gonads with yellowish transparent eggs
A55*	Adriatic	21	Male	3/06/2010	252.92	12.7	135.6	74.7	74.3	
A56*	Adriatic	21	Male	7/06/2010	149.30	10.4	75.7	55.3	47.6	
A57*	Adriatic	21	Male	9/06/2010	150.37	10.1	75.6	41.5	45.3	
A58	Adriatic	21	Female	7/07/2010	147.75	10.0	86.7	58.7	56.9	Female had eggs
A59	Adriatic	21	Female	8/07/2010	226.54	11.3	140.7	82.9	79.9	Animal was inking during capture, female had eggs
A60	Adriatic	21	Male	9/07/2010	198.63	11.6	109.0	69.4	74.8	Animal was inking during capture
A66	Adriatic	21	Female	21/07/2010	184.21	10.8	145.4	87.7	87.2	Female already mated as there were sucker marks on the head, had hardly ink in ink sac
A67	Adriatic	21	Female	23/07/2010	144.90	9.5	108.5	57.4	59.1	Female already mated as there were sucker marks on the head, had hardly ink in ink sac
A68	Adriatic	21	Female	26/07/2010	248.32	12.2	161.3	96.0	101.3	
A08	Oosterschelde	11	Male	10/07/2009	59.09	7.3	42.8	23.9	27.5	
A11	Oosterschelde	11	Male	15/07/2009	79.46	7.9	63.5	41.2	38.3	
A12	Oosterschelde	11	Female	21/07/2009	89.19	8.3	63.2	37.5	39.1	
A23	Oosterschelde	11	Male	6/08/2009	96.64	8.9	81.2	51.3	48.5	Animal was inking during capture
A25	Oosterschelde	11	Male	12/08/2009	112.12	9.5	70.1	47.0	42.0	
A26	Oosterschelde	11	Male	13/08/2009	77.84	8.5	60.1	37.0	38.5	
A27	Oosterschelde	11	Male	18/08/2009	92.99	8.6	73.0	43.4	45.8	Animal was inking during capture
A28	Oosterschelde	11	Male	8/09/2009	79.98	8.1	65.1	38.1	36.0	
A29	Oosterschelde	11	Male	17/09/2009	83.31	8.7	83.6	53.9	50.6	
A04	Oosterschelde	16	Female	4/06/2009	72.46	8.5	–	–	–	

A05	Oosterschelde	16	Female	17/06/2009	84.50	8.6	57.3	31.6	31.4	Animal was inking during capture
A15	Oosterschelde	16	Female	22/07/2009	97.70	8.9	61.6	38.4		
A18	Oosterschelde	16	Male	29/07/2009	108.29	9.1	111.4	46.4	43.8	Animal was inking during capture
A31	Oosterschelde	16	Male	22/09/2009	125.10	9.3	114.5	50.4	59.3	
A32	Oosterschelde	16	Male	23/09/2009	79.75	7.7	73.1	34.4	33.8	
A33	Oosterschelde	16	Female	24/09/2009	253.83	12.5	315.7	130.0	134.3	Female had proper developed gonads with yellowish transparent and white eggs
A34*	Oosterschelde	16	Female	28/08/2009	164.90	9.3	133.6	93.5	86.7	
A35	Oosterschelde	16	Male	29/09/2009	155.89	10.2	131.5	65.4	68.0	Animal was inking during capture
A36	Oosterschelde	16	Female	30/09/2009	128.38	9.1	97.3	54.7	56.1	Animal was inking during capture
A37	Oosterschelde	16	Female	1/10/2009	165.94	10.3	183.0	90.1	87.0	
A38	Oosterschelde	16	Male	5/10/2009	164.39	11.4	167.9	79.0	70.0	
A39*	Oosterschelde	16	Male	6/10/2009	278.93	12.6	214.8	127.8	133.0	
A40*	Oosterschelde	16	Male	7/10/2009	177.30	11.0	142.2	87.4	80.5	
A41*	Oosterschelde	16	Male	8/10/2009	178.59	10.5	123.9	80.8	89.8	Animal was slightly inking during capture
A42*	Oosterschelde	16	Female	12/10/2009	225.04	10.9	183.3	108.6	116.9	
A43*	Oosterschelde	16	Male	13/10/2009	258.39	12.3	210.0	131.8	138.1	Animal was slightly inking during capture
A44*	Oosterschelde	16	Female	14/10/2009	299.15	12.8	194.7	161.2	133.2	Female had proper developed gonads with yellowish transparent eggs
A45*	Oosterschelde	16	Female	15/10/2009	276.24	12.3	184.6	146.7	155.6	Female had proper developed gonads with yellowish transparent eggs
A16	Oosterschelde	21	Female	27/07/2009	71.33	7.9	56.8	36.2	32.7	
A17	Oosterschelde	21	Male	28/07/2009	83.59	8.1	77.7	41.4	39.5	Heart muscle was relatively tough
A20	Oosterschelde	21	Female	3/08/2009	102.13	8.5	75.5	45.7	44.5	
A21	Oosterschelde	21	Male	4/08/2009	124.66	9.9	106.3	62.3	67.9	
A22	Oosterschelde	21	Male	5/08/2009	123.77	8.9	110.7	65.3	60.4	

Individuals marked with asterisks were used for morphometric statistics but not for measurement of mitochondrial respiration.