

Changes of loggerhead turtle (*Caretta caretta*) dive behavior associated with tropical storm passage during the inter-nesting period

Maria Wilson^{1,*}, Anton D. Tucker^{2,3}, Kristian Beedholm⁴ and David A Mann^{2,5}

1) Institute of Biology, University of Southern Denmark, Odense Denmark.

2) Mote Marine Laboratory and Aquarium, Sarasota, FL 34236, USA.

3) Department of Parks and Wildlife, Marine Science Program, 17 Dick

Perry Ave, Kensington, WA 6151, Australia.

4) Zoophysiology, Department of Bioscience, Aarhus University, Denmark.

5) Loggerhead Instruments, Sarasota, FL 34238, USA.

*Email of corresponding author: im.mariawilson@gmail.com

Keywords: activity level; climate change; animal motion tags; satellite tags; loggerhead turtle; tropical storm

Summary statement

Data retrieved from motion dataloggers and satellite tags showed that a tropical storm can have a large effect on swimming energetics of a sea turtle, but with little effect on nesting.

Abstract

To improve conservation strategies for threatened sea turtles more knowledge on their ecology, behavior, and how they cope with severe and changing weather conditions is needed. Satellite and animal motion datalogging tags were used to study the inter-nesting behavior of two female loggerhead turtles in the Gulf of Mexico, which regularly has hurricanes and tropical storms during nesting season. We contrast the behavioral patterns and swimming energetics of two turtles, the first tracked in calm weather and a second tracked before, during, and after a tropical storm. Turtle #1 was highly active and swam at the surface or submerged 95% of the time during the entire inter-nesting period with high estimated specific oxygen consumption rate ($0.95 \text{ ml min}^{-1} \text{ kg}^{-0.83}$). Turtle #2 was inactive for most of the first nine days of the inter-nesting period where she rested at the bottom (80% of the time) with low estimated oxygen consumption ($0.62 \text{ ml min}^{-1} \text{ kg}^{-0.83}$). Midway through the inter-nesting period turtle #2 encountered a tropical storm and became highly active (swimming 88% of the time during and 95% after the storm). Her oxygen consumption increased significantly to $0.97 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ during and $0.98 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ after the storm. However, despite of the tropical storm turtle #2 returned to the nesting beach, where she successfully re-nested 75 meters from her previous nest. Thus, the tropical storm had a minor effect on this female's individual nesting success, even though the storm caused 90% loss of Casey Key nests.

Introduction

The rising temperatures of the oceans caused by global warming are expected to increase the intensity and frequency of tropical storms and hurricanes (Mann and Emanuel 2006). Understanding how increased storm activity may affect marine animals is important to improve conservation strategies for threatened species and it has been identified as a main key questing for the marine megafauna (Hays et al. 2016). Hurricanes can cause significant destruction on coral reefs with corresponding changes in the reef fish population (Woodley et al. 1981). In an estuarine environment a significant change in fish assemblages was observed after the passage of a cyclone with reduction in species diversity and variation in the seasonal pattern of abundance (Sudeshna et al. 2012). A satellite tracking study on manatees in southwest Florida showed no significant effect on movement patterns before and during hurricane passages, and it was therefore concluded that the hurricanes had a minor effect on this species (Langtimm et al. 2006). Juvenile blacktip sharks left an estuary during barometric pressure drops from an impending hurricane (Heupel et al. 2003). Thus, marine vertebrates can respond differently to storm passages.

The loggerhead turtle (*Caretta caretta*) is listed as threatened under the US Federal Endangered and Threatened Species Act of 1977 and the North West Atlantic subpopulation is IUCN red-listed as Least Concern (Ceriani and Meylan 2015). Some of the biggest threats against sea turtles are human activities and fishery bycatch, which might have played a significant role in declines in the loggerhead population (Finkbeiner et al., 2011; McDaniel et al., 2000; Witherington et al., 2009). Other major threats are loss of eggs due to nest predation, and human disturbances (Engeman et al., 2016). Naturally occurring threats like tropical storms and hurricanes may also have a major damaging effect on the nests (Hillis and Phillips, 1995; Milton and Leone, 1994; Starbird et al., 1992), but our knowledge about how juvenile and adult sea turtles are affected by severe weather conditions is limited (Limpus and Reed, 1985), mainly because of difficulties in

studying sea turtles after they leave the nesting ground. Storms may have profound impacts on the oceanic stages of juvenile loggerhead turtles, blowing them to unexpected locations with potential impact on their fitness (Monzon-Arguello et al. 2012). After hatching males never return to land. Thus, tagging studies involving male sea turtles are difficult and must be done by capture at sea (Schofield et al., 2010). Mature female loggerheads only return to the beach every second to seven years to nest (Plotkin, 2003). Loggerheads deposit multiple clutches of eggs at 10-21 day inter-nesting intervals across a nesting season (Hays et al., 2002; Sato et al., 1998; Schroeder et al., 2003). When a female has found a nesting beach she often shows strong nest site fidelity and will tend to nest within 5 km of the previous nest (Tucker 2010). However, a small percentage of turtles have weak nest site fidelity and will utilize more distant nesting sites in the general area (Bjorndal et al., 1983; Schofield et al., 2010). Because of the strong site fidelity, loggerhead sea turtles are susceptible to negative impact from development and destruction of beach areas, but it also gives opportunities to study the inter-nesting behavior of female loggerhead sea turtles.

The use of advanced technical equipment (archival and satellite tags) makes it possible to collect important information about sea turtle diving behavior, their ecology and habitat use (Eckert and Martins, 1989; Eckert et al., 1986; Hays et al., 2004a; Hays et al., 1991; Houghton et al., 2002; Minamikawa et al., 2000; Minamikawa et al., 1997; Sakamoto et al., 1990a; Sakamoto et al., 1990b; Sato et al., 1995; Wilson et al., 2006). Studies show that loggerhead turtles exhibit plasticity in behavior during the inter-nesting period and their behavior can be linked to the local environment and how close they are to the next nesting event (Tucker et al. 1996; Houghton et al., 2002; Schofield et al. 2009; Fossette et al. 2012). In areas where food is abundant both green (*Chelonia mydas*) (Hochscheid et al., 2010) and loggerhead turtles may opt to forage during the inter-nesting period (Sakamoto et al., 1990b), whereas if food is limited, they may save energy for reproduction and rest on the seabed (Hays et al., 1999; Minamikawa et al., 1997). At the end of an inter-nesting

period both green and loggerhead turtles spend less time resting on the seabed and more time near the surface (Hays et al., 1991; Hays et al., 1999; Houghton et al., 2002). It would therefore seem that sea turtles tend to optimize energy reserves in a way best suited to local environmental conditions (Houghton et al., 2002), and when inter-nesting sea turtles are exposed to storms or hurricanes it could be expected to cause changes in their behavior to cope with the severe oceanographic conditions. Two studies have examined the effect of severe weather conditions on the behavior of a loggerhead turtle (Sakamoto et al., 1990b) and a hawksbill turtle (Storch et al., 2006). Both studies found changed swimming behavior by turtles during the storm passage. The hawksbill encountering hurricane George in the Caribbean, made shorter dives, and spent less time at the surface (Storch et al., 2006). A loggerhead turtle encountering a typhoon made more dives and increased the dive depth and time spent at depth to avoid the wave action (Sakamoto et al., 1990b). However, after the passage of the severe weather both turtles resumed their normal behavior.

Florida's coasts are significant nesting grounds for loggerhead sea turtles with recent years ranging from 77,975 to 122,706 annual nests (FFWCC 2017). Florida is annually hit by hurricanes and tropical storms and there is close overlap of the tropical storm/hurricane season and turtle nesting season. Hurricanes and tropical storms are therefore potential threats to the loggerhead turtle population because of beach erosion and nest losses. We used Argos satellite tags and high-speed multi-channel animal motion datalogging tags to study the behaviors of inter-nesting female loggerhead turtles. One deployment took place during the passage of a tropical storm in the Gulf of Mexico which gave us a unique opportunity to conduct a detailed analysis of how the inter-nesting behavior of a loggerhead sea turtle is altered during a severe weather conditions.

Materials and methods

Animals

We attached Argos satellite tags (Wildlife Computers, SPOT5) to six female loggerhead turtles during the 2012 nesting season between May and July at Casey Key, Florida. Five females were co-instrumented with an animal motion tag (OpenTag, Loggerhead Instruments, Sarasota, FL). Transmitters were glued to the carapace using a 2-part epoxy resin and covered in antifouling paint. The Argos tag transmitted approximate location (accuracy from 0.1 to 2.0 km), whereas data from the motion tags were stored to a microSD card (4 GB Amazon Basics) and retrieval of the motion tags was necessary to access the data. Four of the six satellite tracked females spread their subsequent nests widely, and two departed without laying an additional nest. However motion tag recoveries were only possible if the turtles returned to a cooperative tagging project for the SW Florida coast at either Casey Key, Manasota Key, or Keewaydin Island. One turtle bearing a motion tag was intercepted within the 6 km patrolled area at Casey Key and a second turtle's motion tag was recovered at Keewaydin Island 140 km to the south. The two recovered motion tags provided continuous recordings from 31 May 2012 to 14 June 2012 (tag#1) and from 14 June 2012 to 29 June 2012 (tag#2). Turtle #1 was 95.7 cm curved carapace length (estimated weight 102 kg based on Ehrhart, 1976)) and carried PTT 115649; Turtle #2 was 109.0 cm curved carapace length (estimated weight 136 kg, based on Ehrhart, 1976)) and carried PTT 115650.

Argos satellite tag

The Argos satellite tags were programmed to be continuously on and a salt water switch prevented signal transmission during submergence. Locations were retrieved and analyzed using the Satellite Tracking and Analysis Tool (Coyne and Godley, 2004). Locations used for turtle

movements were Argos Location Classes for expected accuracy: 3, 2, 1, 0, A, B and filtered to remove unrealistic swimming speeds exceeding 10 km/h and positions inland (Witt et al., 2011).

Animal motion tag

The animal motion tags contain a 3-axis accelerometer, 3-axis gyroscope, 3-axis magnetometer (each sampled at 100 Hz), and temperature and pressure sensors (each sampled at 1 Hz). The gyroscope data of rotational velocity around the vertical plane were used to calculate flipper beats/min. The gyroscope signal was high-pass filtered to remove DC-offset (defined as the mean amplitude displacement from zero). The flipper beats were estimated based on a 8192-point FFT (fast fourier transform) analysis of the high-pass filtered gyroscope signal with a calculated rate for every 2 seconds (Fig. 2). Wave action close to the surface interfered with the flipper beat signal and therefore the flipper beat rate was only estimated at depths > 0.5 meters. Based on accelerometer data in the three orthogonal planes (heave: dorso-ventral acceleration, sway: lateral acceleration, and surge: anterior-posterior acceleration) the dynamic body acceleration (DBA) can be calculated (Shepard et al., 2008; Wilson et al., 2006). The influence of gravitational acceleration was reduced by subtracting the running mean over one second from the raw accelerometer signal independently in all three dimensions. A vector of the dynamic body acceleration (VDBA) was then calculated from the remaining dynamic x, y, and z acceleration values as the norm $(x^2 + y^2 + z^2)^{1/2}$ in 2 s time intervals. DBA can be used as a proxy for energy consumption (Enstipp et al., 2011; Fossette et al., 2012). A recent study on captive green turtles found a strong correlation between PDBA (partial dynamic body acceleration) and oxygen consumption with (Enstipp et al., 2011):

$$sVo_2 = 12.17PDBA + 0.03T_w - 0.46$$

where sVo_2 is in $ml\ min^{-1}\ kg^{-0.83}$, PDBA is in g and T_w is in $^{\circ}C$. In addition to VDBA we therefore also calculated PDBA from heave and sway accelerometer signals ($PDBA = |x| + |y|$)

to estimate the daily oxygen consumption of loggerheads. The use of PDBA to estimate oxygen consumption has to our knowledge only been conducted on green turtles and no calibration exists on loggerhead. However, the values we calculated with this equation are similar to the values measured for green turtles (Hays et 2000). Even if the calculated estimate is off with loggerhead turtles, the values are still useful for comparing estimated energy expenditure of swimming versus energy required for egg production. The estimated oxygen consumption for a green turtle of a comparable size (150 kg) to turtle #2 was $0.63 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ during resting dives (Hays et al 2000; Enstripp et al 2011). The estimated oxygen consumption for turtle #2 was $0.62 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ during resting dives.

Dive classification

Depth and temperature data were retrieved from the accelerometer tag and analyzed. Dives had to be longer than 60 s and deeper than 3 m to be classified as a dive or else it was categorized as swimming or resting close to the surface. Classification of the different dives followed the classification of Minamikawa et al. (1997, 2000). Dives were assigned into five different categories based on their profile characteristics (Fig. 2A). Type 1 dives consist of a descent phase, a flat bottom phase (longer than 60 s) and an ascent phase. The bottom phase is often associated with a resting period (Hochscheid, 2014). Type 2 dives are often a short dive, consisting of only a descent and an ascent phase. These dives are observed when turtles are travelling, for example during the first 24 hours after a nesting event away from shore (Hochscheid, 2014). Type 3 dives consist of three phases: a descent, a gradual ascent (longer than 60 s) and a final ascent. Type 4 dives consist of four phases: a rapid descent to maximum depth, followed immediately by ascent to a certain depth (often where the turtles are believed to be neutrally buoyant), and lastly a final rapid ascent.

The last categories of dives are type 5. When a dive did not have a profile that fit into one of the first four categories it was placed in the type 5 category.

Tropical storm Debby

Tag #2 was deployed just before tropical storm Debby passed through the Gulf of Mexico on June 23-27, 2012. The tropical storm caused extensive flooding in Florida after it developed from a low pressure cell in the central Gulf of Mexico on June 23. The storm slowly strengthened to peak intensity with maximum sustained winds of 65 mph (100 km/h) at 1800 UTC on June 25. On June 26 at 2100 UTC, the storm made landfall near Steinhatchee, Florida with winds of 40 mph (65 km/h). Once inland, the system weakened and crossed Florida to the Atlantic on June 27 (Kimberlain, 2012).

Statistics

Statistical analyses were performed using MATLAB 8.2 (The MathWorks, Inc.) and PAST3 (version 1.0.0.0) (Hammer, 2001). We used a non-parametric Mann-Whitney-test for matched pairs to test for differences in the temperature experienced by the two turtles, because temperature data were not normal distributed. A non-parametric Spearman's rank correlation was used to test for the relation between the two estimated activity measures: flipper beats/min and VDBA (*g*). To evaluate the effect on the tropical storm on the swim behaviour of turtle #2, a non-parametric Kruskal-Wallis test was performed to test for significance in the difference in medians between the daily energy estimate, amount of time swimming, amount of time submerged, and kilometres travelled before, during and after the tropical storm. If a significant difference in median values was found, a Dunn's non-parametric multiple comparisons test was conducted to test for pairwise differences ($\alpha = 0.05$ for the tests). Non-parametric tests were used, because data was not normally distributed (Bagdonavicius et al. 2011).

Results

The two recovered animal motion tag (OpenTags) contained continuous recordings for 15 days, a total of 347 hours and 57 minutes (tag #1: turtle #1), and 16 days, a total of 379 hours and 42 minutes (tag #2: turtle #2). Tag #1 was operating during a period of relatively calm weather from May 31, 2012 to June 14, 2012, whereas tag #2 was operating from June 14 – 29, 2012, while tropical storm Debby passed through the Gulf of Mexico (June 23-27, 2012). Both tags were recovered as the two turtles re-nested; turtle #1 at Keewaydin Island on June 18, 2012 (inter-nesting period: 18 days) >140 km from her previous nest and turtle #2 at Casey Key on July 5, 2012 75 meters from her previous nest (inter-nesting period: 21 days).

The total distance traveled for turtle #1 was 675 km (45 km/day), for turtle #2 it was 613 km (38 km/day). Before the tropical storm turtle #2 traveled a total of 177 km (20 km/day), during the storm 285 km (57 km/day), after the storm 151 km (75 km/day) (Fig. 1B).

Turtle #1 spent 94% of the entire time actively swimming, whereas turtle #2 spent 54% of the time actively swimming. An overview of the dive activities, distance travelled per day and amount of time swimming per day are given in Table 1A and Fig. 3A (turtle #1) and Table 1B and Fig. 3B (turtle #2).

Turtle #1 spent 42% (146 hours) of the total time at depths deeper than 3 meters (6% conducting type 1, 0% type 2 dives, 19% type 3 dives, 7% type 4 dives, and 9% type 5 dives (Fig. 4B)). During 7 out of the 37 type 1 dives turtle #1 was resting at the bottom (defined as no flipper beat activity in at least 75% of the bottom phase). Mean duration of type 1 dives was 50 min \pm 22 min, max 87 min, type 2 dives was 7 min \pm 2 min, max 10 min, type 3 dives was 15 min \pm 9 min, max 38 min and type 4 dives was 22 min \pm 13 min, max 57 min.

Turtle #2 spent 69% (262 hours) at depths deeper than 3 meters (42% conducting type 1, 4% type 2 dives, 5% type 3 dives, 17% type 4 dives, and 15% type 5 dives (Fig. 4C)). In contrast to turtle #1, turtle #2 spent 135 out of 176 of the type 1 resting at the bottom. Mean duration of type 1 dives was $33 \text{ min} \pm 20 \text{ min}$, max 90 min, type 2 dives was $5 \text{ min} \pm 4 \text{ min}$, max 15 min, type 3 dives was $19 \text{ min} \pm 8 \text{ min}$, max 43 min and type 4 dives was $23 \text{ min} \pm 10 \text{ min}$, max 45 min.

Figure 4 gives typical examples of type 3 dives (A, B: turtle #1, F, G: turtle #2) and type 4 dives (C, D: turtle #1, H, I: turtle #2). Both turtles swam during the gradual ascent phase (Fig 4 E and J). For turtle #2, Type 3 and 4 dives was mainly conducted during and after the tropical storm (type 3; 35 dives out of a 56 (63%) and type 4; 71 of 97 (73%)).

VDBA and oxygen consumption

We used data from the rotational velocity gyroscope signal in the vertical plane to estimate the direct flipper beat rate. Furthermore we used data from the tri-axial accelerometer to calculate the VDBA (Enstipp et al., 2011). Both measures reflect the activity level of the turtles and a Spearman's correlation was performed to determine the relationship between the flipper beat rate and VDBA values. There was a strong, positive correlation between flipper beat rate and VDBA ($r_s = 0.64$, $n = 2301$, $p < 0.001$) (Fig. S7).

The daily oxygen consumption was estimated for both turtles and the average daily oxygen consumption (for the entire period) was slightly higher for turtle #1, with a median of $0.953 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ (lower quartile: 0.899; upper quartile: 1.029) compared to turtle #2 with a median of $0.803 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ (lower quartile: 0.614; upper quartile: 0.991). However, there was no significant difference in the overall daily amount of oxygen used for the two turtles during the tagged periods (Mann-Whitney signed-ranked test, $|z| = 1.70$, $n_{\text{turtle\#1}} = 15$, $n_{\text{turtle\#2}} = 16$, $p = 0.088$) (Fig. 6).

Temperature

The two turtles were exposed to relatively high water temperatures, with a median of 28.1°C (min: 23.1°C, max: 31.1°C) for turtle #1, and with a median of 27°C (min: 25.3°C max: 30.5°C) for turtle #2 during the entire deployment periods (Fig. 6). There was a significant difference between the median temperatures (Mann-Whitney signed-ranked test, ($|z| = -3.76$, $n=15$, $p<0.001$).

Tropical storm Debby

We obtained storm tracking data from a NASA summary report on Tropical Debby (Tables 1 and 2 in Kimberlain, 2013). There was a significant change in the behavior of turtle #2 when she encountered the tropical storm. Table 2 summarizes changes in daily median oxygen consumption, amount of time spent swimming, amount of time submerged and distance travelled before, during and after the storm with supportive statistical tests. Oxygen consumption was significantly higher during the storm ($0.97 \text{ ml min}^{-1} \text{ kg}^{-0.83}$) compared to before the storm ($0.62 \text{ ml min}^{-1} \text{ kg}^{-0.83}$). After the storm the oxygen consumption was also higher compared to before, but not significantly higher ($0.99 \text{ ml min}^{-1} \text{ kg}^{-0.83}$) (Figure 6B). Amount of time swimming was significantly higher during the storm (91%) compared to before the storm (20 %) and also after the storm (86%) versus before. Amount of time submerged (86 %) was significantly higher before the storm compared to during the storm (44 %), but there was no significant difference before the storm compared to after the storm (57 %) Daily distance travelled was significantly higher during the storm (57 km) compared to before the storm (17 km) and also after the storm (70 km) compared to before.

Discussion

Sea turtles exhibit plasticity in behavior during inter-nesting periods (Hochscheid, 2014) and the present study supports previous findings. We documented different behaviors by two female loggerheads nesting at the same rookery. Both turtles were instrumented early in the nesting season. The entire period where turtle #1 was tagged and the first nine days where turtle # 2 was tagged were periods with calm and warm weather. Despite the same weather conditions the turtles displayed different inter-nesting strategies, this complicates a direct comparison between females even without the tropical storm.

A fundamental factor affecting inter-nesting behavior is water temperature (Fossette et al., 2012; Hays et al., 2002; Sato et al., 1998; Schofield et al., 2009). Sea turtles are ectotherms and the maturation of eggs is therefore dependent on the surrounding water temperature (Schofield et al., 2009). Active maintenance of a high and stable body temperature is a clear benefit, however, both of the turtles in the present study experienced water temperature above 23°C and water temperature therefore seems an unlikely reason why different inter-nesting strategies are observed.

Another explanation for higher activity by turtle #1 could be that food was available. Instead of resting and saving energy, females may invest energy into foraging to supplement their body reserves and maximize reproduction outcomes. This type of behaviour has been observed in both a Greek loggerhead population (Schofield et al., 2009) and in Japan, where pelagic feeding took place during the inter-nesting period (Narazaki et al., 2013). Pelagic feeding events were mainly observed during the gradual ascent phase of type 3 and type 4 dives (Narazaki et al., 2013). Of the 335 dives turtle #1 conducted during the inter-nesting period, 276 of the dives were either type 3 or type 4 , and it is therefore possible that turtle #1 encountered waters with a high concentration of gelatinous

food items and that she was foraging. There were no video corroborations of feeding in the present study of neritic Gulf of Mexico loggerheads to uncover if she was feeding or not. However, decades of systematic necropsies find negligible or empty gastrointestinal tracts in gravid female loggerheads during Florida's nesting season (A. Foley-FWC- pers. comm.; G. Lovewell-Mote Marine Lab-pers. comm.).

Dive types and estimated aerobic dive limit

Both turtles conducted relatively long type 1 dives with maximum durations of 90 min and 87 min for turtle number #1 and #2 respectively. By using the estimated resting oxygen consumption of $0.62 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ and the approximately oxygen store of a loggerhead turtle of $22.2 \text{ ml O}_2/\text{kg}$ (Hochscheid et al. 2005) the aerobic dive limit would be 89 min corresponding to the maximum length of type 1 dives in the present study. Our study supports previous findings, that loggerheads very rarely make anaerobic dives (Hochscheid et al. 2005).

Both turtles conducted type 3 and type 4 dives with a gradual ascent phase to between 10-20 meters, where turtles are neutrally buoyant (Hays et al., 2004b; Minamikawa et al., 2000). Studies on loggerhead turtles in Japan (Minamikawa et al., 1997; Minamikawa et al., 2000) and in Cyprus (Houghton et al., 2002) found that type 3 and 4 dives are used for midwater resting by females during the inter-nesting period. However, loggerhead dive types might also have different purposes than in green turtles that travel by swimming or gliding during type 3 and 4 dives (Hochscheid et al., 1999; Rice and Balazs, 2008) or pelagic foraging on gelatinous prey during these dives (Narazaki et al., 2013). For both turtles in the present study we found that they were swimming during most of type 3 and type 4 dives (Fig. 5), and not resting as previously observed in other female loggerheads during the inter-nesting period. Turtle #2 conducted the main part of the type 3 and 4 dives during or after the tropical storm (Fig. 4), where she also moved relatively long

distances (Fig. 3). Swimming at a depth of neutral buoyancy is energetically efficient, since turtles would not have to allocate energy to remain at a certain depth nor struggle with surface waves. Some of the type 4 dives were quite long with maximum duration of 57 min and 45 min for turtle #1 and turtle # 2, respectively. By using the estimated active oxygen consumption of $0.95 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ for turtle #1 and $0.97 \text{ ml min}^{-1} \text{ kg}^{-0.83}$ for turtle #2 active dives will become anaerobic after 58 and 56 min for the two turtles. None of the active dives exceeded the estimated aerobic dive limit.

Tropical storm Debby

Turtle #2 encountered changing weather conditions during the sixteen days she was instrumented. Before the storm she was resting, which is common for sea turtle between nesting events and agrees with previous data recorded for loggerheads (Sakamoto et al., 1993; Minamikawa et al., 1997; Houghton et al., 2002), green turtles (Hochscheid et al., 1999; Rice and Balazs 2008; Cheng 2009), and hawksbill turtles (Storch et al., 2006). During the storm the behavior of turtle #2 changed significantly and she became highly active moving in a northern direction, consistent with surface currents generated by the storm (Kimberlain, 2013). We interpreted the displacement as passive storm-generated drift rather than active directed movement by the turtle (Fig. 1). During the storm she spent more time close to the surface as opposed to the two former studies that found that turtles spend less time at the surface when encountering severe weather (Sakamoto et al., 1990b; Storch et al., 2006). The dominant dive type during the storm was type 4 (Figs. 2), in which the turtle descended to the bottom but shortly afterwards ascended to the neutral buoyancy zone between 10-20 meters (Hays et al., 2004b; Minamikawa et al., 2000) where she swam during the gradual ascent phase. After the storm most dives were short dives while she travelled south back to the nesting beach (Figs. 1 and 4G). She returned to the same nesting beach (Casey Key) where she successfully re-nested only 75 meters from her previous nest. According to Sato et al., (1998) there

is a negative correlation between the time span between nesting attempts and the temperature of the surrounding water in loggerhead sea turtles. A surrounding water temperature of 22°C, will cause an inter-nesting period of approximately 21.7 days, whereas if the temperature is 27°C, the inter-nesting period decreases to approximately 14.9 days. Turtle #2 was exposed to an average water temperature of 27°C, thus according to the study by Sato et al (1998), a predicted inter-nesting period would be fifteen days. The actual inter-nesting period was twenty-one days, six days longer than the predicted value. Twenty-one days is still within the normal range of loggerheads and the longer inter-nesting interval could be explained by individual variation. An equally parsimonious explanation is displacement from the tropical storm.

The question is how much energy did the tropical storm cost? If we assume the turtle is metabolizing fat (Schmit-Nielsen 1992) the daily energy expenditure based on the oxygen consumption estimate would be 1029 kJ before the tropical storm, whereas during and after the storm the daily energy consumption would be 1608 kJ, more than a 50% increase. Is it much compared to the energy used during a nesting event? Energy expenditure for a nesting event is very high (Jackson 1979). Hays and Speakman (1991) estimated the mean oxygen consumption by nesting loggerhead turtles on the beach to be 0.23 O₂ kg⁻¹ h⁻¹ which correspond to an energy expenditure of 4.52 kJ O₂ kg⁻¹ h⁻¹. The egg production is, however by far the most energy consuming process in the nesting event. Assuming the volume-specific energy content of loggerhead eggs is the same as for green turtles, the energy content of a loggerhead egg would be 165 kJ (Hays and Speakman, 1991). Clutch size depends on turtle size and her carrying capacity. For turtle #2 the estimated clutch size would be approximately 150 eggs (Hays and Speakman 1991) corresponding to 24,750 kJ per clutch. A nesting event depositing a clutch size of 150 eggs takes approximately 100 min (Hays and Speakman 1991). Based on these values the total energy expenditure of the entire nesting event for turtle #2 would be 25,777 kJ. If turtle #2 did not

encounter the tropical storm and she rested during the entire inter-nesting period, she would expend 1,994 kJ less energy, corresponding to the energy content in 12 eggs. Therefore the significant change in turtle #2's behavior during the tropical storm would likely have had a minor effect on the overall energy budget.

The overall estimated oxygen consumption for the tracked time span was actually lower for turtle #2 compared to turtle #1, which encountered calm weather. Consequently, the tropical storm effects on a single sea turtle appear to have a negligible effect on site fidelity of the turtle and her ability to nest, despite any behavioral changes at sea for the dive profile. The same tropical storm had a more severe effect on the beach itself, where almost 90% of the incubating nests at Casey Key were destroyed (Tucker et al. 2012). Thus, in terms of conservation priorities focus should be on securing the incubating nest from beach erosion.

Acknowledgements

The authors thank the sea turtle team at Mote Marine Laboratory for help with sea turtle tagging and retrieval of animal motion tags. Dave Addison of the Conservancy of Southwest Florida recovered the tag on Keewaydin Island. Thanks go to Magnus Wahlberg, Sabrina Fossette, Allen Foley, Brian Stacey, and Gretchen Lovewell for helpful comments on earlier versions of the manuscript. Tagging and instrument attachment was conducted with animal ethics approval and permits from Florida Fish and Wildlife Conservation Commission Permits MTP126 and MTP155 and IACUC permits at Mote Marine Laboratory.

Competing interests

D. Mann is President of Loggerhead Instruments, which designed and manufactured the open source tags used in the project. The other authors declare no competing interests.

Author contributions

Conceived and designed the experiments: MW, DM. Performed the experiments: MW, AT, DM.

Analyzed the data: MW. Contributed analysis tools: KB. Wrote the paper: MW, AT, KB, DM

Performed statistical analyses: MW.

Funding

Satellite tags were funded by the Florida Sea Turtle Grants Program supported by the Sea Turtle License Plate Fund (www.helpingseaturtles.org). MW was funded by the Danish Council for Independent Research, Natural Science and Carlsberg Foundation.

References

- Bagdonavicius, V., Kruopis, J., Nikulin, M.S.** (2011). Non-parametric tests for complete data
ISTE & WILEY: London & Hoboken.
- Bjorndal, K. A., Meylan, A. B. and Turner, B. J.** (1982). Sea turtles nesting at Melbourne Beach,
Florida, I. Size, growth and reproductive biology. *Biological Conservation* **26**, 65-77.
- Ceriani, S. A. and Meylan, A. B.** 2015. *Caretta caretta* (North West Atlantic subpopulation).
<http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T84131194A84131608.en>
- Cheng, I. J., Bentivegna, F. and Hochscheid, S.** (2013). The behavioural choices of green turtles
nesting at two environmentally different islands in Taiwan. *Journal of Experimental Marine
Biology and Ecology* **440**, 141–148.
- Coyne, M.S. and Godley B.J.** (2005) Satellite Tracking and Analysis Tool (STAT): an integrated
system for archiving, analyzing and mapping animal tracking data. *Marine Ecology Progress Series*
301:1-7.
- Eckert, S. A. and Martins, H. R.** (1989). Transatlantic travel by juvenile loggerhead turtle. *Marine
Turtle Newsletter* **45**, 15.
- Eckert, S., Nellis, D. W., Eckert, K. L. and Kooyman, G. L.** (1986). Diving Patterns of Two
Leatherback Sea Turtles (*Dermochelys coriacea*) during Internesting Intervals at Sandy Point, St.
Croix, U.S. Virgin Islands. *Herpetologica* **42**, 381–388.
- Engeman, R. M., Addison, D. and Griffin, J. C.** (2016). Defending against disparate marine
turtle nest predators: nesting success benefits from eradicating invasive feral swine and caging nests
from raccoons. *Oryx* **50**, 289-295.

Enstipp, M. R., Ciccione, S. and Gineste, B. (2011). Energy expenditure of freely swimming adult green turtles (*Chelonia mydas*) and its link with body acceleration. *Journal of Experimental Biology* **214**, 4010–4020.

Finkbeiner, E. M., Wallace, B. P. and Moore, J. E. (2011). Cumulative estimates of sea turtle bycatch and mortality in USA fisheries between 1990 and 2007. *Biological Conservation* **144**, 2719–2727.

Florida Fish and Wildlife Conservation Commission. (2017). Statewide nesting beach survey program- loggerhead nesting data, 2012-2016.

Fossette, S., Schofield, G., Lilley, M. K. S., Gleiss, A. and Hays, G. C. (2012). Acceleration data reveal the energy management strategy of a marine ectotherm during reproduction. *Functional Ecology* **26**, 324–333.

Hammer, Ø. Harper, D.T. & Ryan, P.D. (2001). *Past: Paleontological statistics software package for education and data analysis.*

Hays, G. C., Broderick, A. C., Glen, F., Godley, B. J., Houghton, J. D. R. and Metcalfe, J. D. (2002). Water temperature and internesting intervals for loggerhead (*Caretta caretta*) and green (*Chelonia mydas*) sea turtles. *Journal of Thermal Biology* **27**, 429–432.

Hays, G. C., Ferreira, L. C., Sequeira, A. M.M., Meekan, M. G., Duarte, C. M., et al. (2016). Key Questions in Marine Megafauna Movement Ecology. *Trends in Ecology & Evolution* **31**, 463–475.

Hays, G. C., Luschi, P., Papi, F., del Seppia, C. and Marsh, R. (1999). Changes in behaviour during the inter-nesting period and post-nesting migration for Ascension Island green turtles.

Marine Ecology Progress Series **189**, 263–273.

Hays, G. C., Marshall, A. and Seminoff, J. A. (2007). Flipper beat frequency and amplitude changes in diving green turtles, *Chelonia mydas*. *Marine Biology* **150**, 1003–1009.

Hays, G. C., Metcalfe, J. D., Walne, A. W. and Wilson, R. P. (2004a). First records of flipper beat frequency during sea turtle diving. *Journal of Experimental Marine Biology and Ecology* **303**, 243–260.

Hays, G. C., Metcalfe, J. D. and Walne, A. W. (2004b). The implication of lung-regulated buoyancy control for dive depth and duration. *Ecology* **85**, 1137–1145.

Hays, G.C. and Speakman, J. R. (1991). Reproductive investment and optimum clutch size of loggerhead sea turtles (*Caretta caretta*). *Journal of Animal Ecology* **60**, 455–462.

Hays, G. C., Webb, P. I. and Hayes, J. P. (1991). Satellite tracking of a loggerhead turtle (*Caretta caretta*) in the Mediterranean. *Journal of the Marine Biology and Ecology* **71**, 743–746.

Heupel, M. R., Simpfendorfer, C. A., and Hueter, R. E. (2003). Running before the storm: blacktip sharks respond to falling barometric pressure associated with Tropical Storm Gabrielle. *Journal of Fish Biology* **63**, 1357–1363.

Hillis, Z. and Phillips, B. (1995). *The Hurricane Season of the Century, Buck Island Reef National Monument, St. Croix, Virgin Islands*.

Hochscheid, S. (2014). Why we mind sea turtles' underwater business: A review on the study of diving behavior. *Journal of Experimental Marine Biology and Ecology* **450**, 118–136.

Hochscheid, S., Bentivegna, F. and Hays, G. C. (2005). First records of dive durations in a hibernating sea turtle. *Biology Letters* **1**, 82-87. doi: 10.1098/rsbl.2004.0250.

Hochscheid, S., Bentivegna, F. and Hays, G. C. (2010). When surfacers do not dive: multiple significance of extended surface times in marine turtles. *The Journal of Experimental Biology* **213**, 1328–1337.

Hochscheid, S., Godley, B. J. and Broderick, A. C. (1999). Reptilian diving: highly variable dive patterns in the green turtle *Chelonia mydas*. *Marine Ecology Progress Series* **185**, 101-112.

Hooker, S. K. and Miller, P. (2005). Ascent exhalations of Antarctic fur seals: a behavioural adaptation for breath-hold diving? *Proceedings of the Royal Society B: Biological Sciences* **272**, 355-363.

Houghton, J. D. R., Broderick, A. C., Godley, B. J., Metcalfe, J. D. and Hays, G. C. (2002). Diving behaviour during the internesting interval for loggerhead turtles *Caretta caretta* nesting in Cyprus. *Marine Ecology Progress Series* **227**, 63–70.

Jackson, D.C. and Prange, H.D. (1979). Ventilation and gas exchange during rest and exercise in adult green sea turtles. *Journal of Comparative Physiology* **134**, 315-319.

Kimberlain, T. B. (2013). Tropical Cyclone Report: Tropical Storm Debby. *National Hurricane Center* (Report). Miami, Florida: National Oceanic and Atmospheric Administration.

Kimberlain, T. B. (2012) Post-tropical cyclone Debby discussion number 18 (TXT). *National Hurricane Center* (Report). Miami, Florida: National Oceanic and Atmospheric Administration

Limpus, C. J. and Reed, P. C. (1985) Green sea turtles stranded by Cyclone Kathy on the South-Western coast of the Gulf of Carpentaria. *Australian Wildlife Research* **12**: 523-533.

- McDaniel, C. J., Crowder, L. B. and Priddy, J. A.** (2000). Spatial dynamics of sea turtle abundance and shrimping intensity in the US Gulf of Mexico. *Conservation Ecology* **4**(1), 15
- Miller, P., Johnson, M. P., Tyack, P. L., and Terry, E. A.** (2004). Swimming gaits, passive drag and buoyancy of diving sperm whales *Physeter macrocephalus*. *Journal of Experimental Biology* **207**, 1953-1967.
- Milton, S. L. and Leone, S.** (1994). Effects of hurricane Andrew on the sea turtle nesting beaches of South Florida. *Bulletin of Marine Science - Miami* **54**(3), 974-981.
- Minamikawa, S., Naito, Y., Sato, K., Matsuzawa, Y., Bando, T. and Sakamoto, W.** (2000). Maintenance of neutral buoyancy by depth selection in the loggerhead turtle *Caretta caretta*. *The Journal of Experimental Biology* **203**, 2967–2975.
- Minamikawa, S., Naito, Y. and Uchida, I.** (1997). Buoyancy control in diving behavior of the loggerhead turtle *Caretta caretta*. *Journal of Ethology* **15**, 109–118.
- Monzon-Arguello, C., F. Dell'Amico, F., Morinière, P., Marco, A., López-Jurado L. F., et al.** (2012). Lost at sea: genetic, oceanographic and meteorological evidence for storm-forced dispersal. *Journal of the Royal Society Interface* **9**, 1725-32.
- Narazaki, T., Sato, K., Abernathy, K. J., Marshall, G. J. and Miyaxaki, N.** (2013). Loggerhead turtles (*Caretta caretta*) use vision to forage on gelatinous prey in mid-water. *PLOS one* **8**(6).
- Plotkin, P.** (2003). Adult migrations and habitat use. In *The Biology of Sea Turtles II* (ed. P. L. Lutz, J. A. Musick and J. Wyneken) pp. 225-233. **CRC Press.**

Rice, M. R. and Balazs, G. H. (2008). Diving behavior of the Hawaiian green turtle (*Chelonia mydas*) during oceanic migrations. *Journal of Experimental Marine Biology and Ecology* **356**, 121-127.

Sakamoto, W., Naito, Y., Uchida, I. and Kureha, K. (1990a). Circadian rhythm on diving motion of the loggerhead turtle *Caretta caretta* during inter-nesting and its fluctuations induced by the oceanic environmental events. *Nippon Suisan Gakkaishi* **56**, 263-272.

Sakamoto, W., Sato, K., Tanaka, H. and Naito, Y. (1993). Diving patterns and swimming environment of two loggerhead turtles [*Caretta caretta*] during internesting. *Nippon Suisan Gakkaishi* **59**, 1129-1137.

Sakamoto, W., Uchida, I., Naito, Y., Kureha, K., Tujimura, M. and Sato, K. (1990b). Deep diving behavior of the loggerhead turtle near the frontal zone. *Nippon Suisan Gakkaishi* **56**, 1435–1443.

Sato, K., Matsuzawa, Y., Tanaka, H., Bando, T., Minamikawa, S., Sakamoto, W. and Naito, Y. (1998). Internesting intervals for loggerhead turtles, *Caretta caretta*, and green turtles, *Chelonia mydas*, are affected by temperature. *Canadian Journal of Zoology* **76**, 1651–1662.

Sato, K., Sakamoto, W., Matsuzawa, Y., Tanaka, H., Minamikawa, S. and Naito, Y. (1995). Body temperature independence of solar radiation in free-ranging loggerhead turtles, *Caretta caretta*, during internesting periods. *Marine Biology* **123**, 197–205.

Schmidt-Nielsen, K. (1997) Animal physiology: adaptation and environment. New York, NY, USA, Cambridge University Press.

Schofield, G., Bishop, C. M., Katselidis, K. A., Dimopoulos, P., Pantis, J. D. and Hays, G. C.

(2009). Microhabitat selection by sea turtles in a dynamic thermal marine environment. *Journal of Animal Ecology* **78**, 14–21.

Schofield, G., Hobson, V. J., Lilley, M. K. S., Katselidis, K. A., Bishop, C. M., Brown, P. K.

and Hays, G. C. (2010). Inter-annual variability in the home range of breeding turtles: Implications for current and future conservation management. *Biological Conservation* **143**, 722–730.

Schroeder, B. A., Foley, A. M. and Bagley, D. A. (2003). Nesting patterns, reproductive

migrations, and adult foraging areas of loggerhead turtles. In *Loggerhead sea turtles*. (ed. A. B., Bolten and B. E. Witherington) pp. 114-124. Washington, DC: Smithsonian Books.

Shepard, E. L. C., Wilson, R. P., Quintana, F., Laich, A. G., Liebsch, N., Albareda, D. A.,

Halsey, L. G., Gleiss, A., Morgan, D. T., Myers, A. E., et al. (2008). Identification of animal movement patterns using tri-axial accelerometry. *Endangered Species Research* **10**, 47–60.

Starbird, C., Hillis, Z. M., Salmon, M. and Wyneken, J. (1992). *The effects of Hurricane Hugo on the nesting behavior of hawksbill sea turtles on Buck Island National Monument, United States Virgin Island. US Dep.*

Storch, S., Hays, G. C., Hillis-Starr, Z. and Wilson, D. P. (2006). The behaviour of a hawksbill turtle data-logged during the passage of hurricane Georges through the Caribbean. *Marine and Freshwater Behavior and Physiology* **39**, 307–313.

Tucker, A.D., Fitz-Simmons, N. N. and Limpus, C. J. (1996). Conservation implications of inter-nesting habitat use by loggerhead turtles (*Caretta caretta*) in Woongarra Marine Park, Queensland, Australia. *Pacific Conservation Biology* **2**: 157-166.

- Tucker, A. D.** (2010) Nest site fidelity and clutch frequency of loggerhead turtles are better elucidated by satellite telemetry than by nocturnal tagging efforts. *Journal of Experimental Marine Biology and Ecology* **383**, 48-55.
- Tucker, A. D., Mazzarella, K, Hirsch, S., and Klingensmith, K.** (2012). Sea turtle Monitoring, nest evaluation and protection measures for Siesta Key and Casey Key 2012. *Mote Marine Laboratory Technical Report* **1647**.
- Williams, T. M., Davis, R. W., Fuiman, L. A. and Francis, J.** (2000). Sink or swim: strategies for cost-efficient diving by marine mammals. *Science* **288**, 133.
- Wilson, R. P., White, C. R., Quintana, F., Halsey, L. G., Liebsch, N., Martin, G. R. and Butler, P. J.** (2006). Moving towards acceleration for estimates of activity-specific metabolic rate in free-living animals: the case of the cormorant. *Journal of Animal Ecology* **75**, 1081–1090.
- Witt MJ, Åkesson S, Broderick AC, Coyune MS, Ellick J, Formia A, Hays GC, Luschi P, Stroud S, Godley BJ** (2010) Assessing accuracy and utility of satellite-tracking data using Argos-linked FAsTloc-GPS. *Animal Behaviour* **80**, 571-581.
- Witherington, B., Kubilis, P. and Brost, B.** (2009). Decreasing annual nest counts in a globally important loggerhead sea turtle population. *Ecological Applications* **19**, 30-54.

Figures




Figure 1: Swimming tracks of the two loggerhead turtles equipped with an animal motion tag and a satellite tag based on satellite positions from 2012-05-31 to 2012-06-4 (turtle #1: blue circles unaffected by storm) and 2012-06-14 to 2012-07-04 (turtle #2: black/red/green circles delineate

before/during/after storm passage). Red stars indicate the track of tropical storm Debby passage in the Gulf of Mexico. The strongest winds are on the right side of the storm in counterclockwise circulation of hurricanes in the Northern Hemisphere. Despite the fact that Debby never achieved hurricane strength, the slow movement of the system and prolonged period of onshore flow allowed a moderate storm surge to move into the Florida Big Bend. Maximum wind speed was 55 knots from a southerly direction over the period while the turtle drifted northward (red circles).




Figure 2: Illustration of the fluke rate estimation based on an 8192 point FFT analysis of the gyroscope signal with a calculated rate for every 2 seconds. A) FFT of the gyro signal in the vertical plane. Red circle indicates peak energy of the signal and gives the flipper beats/second. B) The high-pass filtered gyro signal in the vertical plane. C) Corresponding dive profile.




Figure 3: Activity budget of the two loggerhead sea turtles. Overview of distance traveled per day, amount of time spent submerged per day and amount of time spent swimming (A: Turtle #1, B: Turtle #2). Red circles are distance traveled (km/day), black triangles are % of the time submerged and black squares are % of time swimming. Gray shaded area in B indicates the passage of the tropical storm.




Figure 4: Different dive categories. A) Classic dive categories and illustration of the dive profiles of each dive type conducted by the female loggerhead turtles during the inter-nesting period (modified after Minamikawa et al. 1997). B-C) Histograms illustrating the proportion of time spent conducting different dive types during the inter-nesting period for B) turtle #1 and C) turtle #2. Gray shaded area in C indicates the passage of the tropical storm.









Figure 5: Type 3 and type 4 dives. Examples of a type 3 (A: turtle #1, F: turtle #2) and a type 4 (C: turtle #1, H: turtle #2). Color code gives flipper beats/min (FB/min). B, D, G and I gives the flipper beats of the dives in A, C, F and H, respectively. Histogram is number of type 3 and type 4 dives where the turtles are swimming more than 95% of time at the gradual ascent phase (red) or resting (blue) (E: turtle #1, J: turtle #2).




Figure 6: Oxygen consumption and water temperature. Mean daily estimated oxygen consumption in ml min⁻¹ kg^{-0.83} (black line), and mean daily temperature in °C (red: mean temperature, turquoise: maximum temperature and green: minimum temperature). (A: Turtle #1, B: Turtle #2). Gray shaded area in B indicates the passage of the tropical storm.

Tables

A	Number of dives	Time submerged (%)	Time swimming (%)	Distance traveled (km)	Mean temp (°C)	Min temp (°C)	Max temp (°C)	Max depth (m)	Mean VDBA (g)	O ₂ C (ml min ⁻¹ kg ^{-0.83})
Date										
2012-05-31	51	55	96	20	28.6	25.7	30.3	12.8	0.04	0.87
2012-06-01	49	68	83	28	27.9	27.6	28.2	14.7	0.03	0.79
2012-06-02	42	54	99	39	28.0	27.3	29.7	20.5	0.04	0.87
2012-06-03	26	63	90	38	27.0	24.7	29.2	40.0	0.04	0.83
2012-06-04	26	66	80	57	26.3	24.0	29.1	50.9	0.04	0.84
2012-06-05	17	30	97	40	27.0	23.1	28.2	57.0	0.05	0.97
2012-06-06	7	10	99	39	27.1	24.5	27.5	47.4	0.06	1.11
2012-06-07	8	13	99	59	27.2	25.4	27.8	41.0	0.06	1.08
2012-06-08	22	28	98	55	27.4	24.9	30.2	45.6	0.05	0.98
2012-06-09	16	36	85	41	27.1	24.3	28.9	47.1	0.05	0.95
2012-06-10	41	67	99	43	28.2	27.2	30.2	31.2	0.04	0.89
2012-06-11	43	51	97	51	29.1	28.1	30.9	16.8	0.04	0.92
2012-06-12	37	49	97	60	28.9	28.3	30.8	21.1	0.04	0.95
2012-06-13	6	5	100	54	29.2	28.5	31.1	23.1	0.05	1.06
2012-06-14	23	36	100	51	29.5	28.7	31.1	22.8	0.05	1.05

B	Number of dives	Time submerged (%)	Time swimming (%)	Distance traveled (km)	Mean temp (°C)	Min temp (°C)	Max temp (°C)	Max depth (m)	Mean VDBA (g)	O ₂ C (ml min ⁻¹ kg ^{-0.83})
Date										
2012-06-14	73	75	85	40	28.7	26.7	30.5	29.1	0.05	0.99
2012-06-15	38	94	24	25	26.8	25.9	29.3	33.7	0.02	0.56
2012-06-16	24	94	12	17	26.2	25.9	29.4	34.8	0.01	0.51
2012-06-17	21	92	11	12	26.2	25.7	28.8	36.1	0.01	0.50
2012-06-18	22	88	13	12	26.4	25.8	29.2	36.9	0.02	0.53
2012-06-19	24	82	27	16	26.8	26.1	28.2	35.9	0.02	0.64
2012-06-20	20	75	24	15	26.8	26.0	28.3	36.4	0.02	0.64
2012-06-21	20	83	16	18	27.4	26.3	28.2	35.6	0.02	0.61
2012-06-22	26	76	30	22	27.5	26.7	28.1	37.5	0.03	0.71
2012-06-23	19	50	72	35	27.1	25.3	27.5	48.9	0.04	0.89
2012-06-24	14	41	88	71	26.8	25.4	27.3	49.5	0.06	1.07
2012-06-25	16	38	90	69	26.7	25.5	27.1	45.3	0.06	1.02
2012-06-26	18	46	89	52	26.9	26.4	27.7	40.8	0.05	0.99
2012-06-27	52	57	98	58	27.2	26.6	28.1	31.4	0.05	0.95
2012-06-28	60	47	92	94	27.5	26.7	29.9	22.8	0.05	0.91
2012-06-29	85	66	95	57	27.4	26.8	28.5	22.1	0.04	0.87

Table 1: Overview of the daily activity for turtle #1 (A) and turtle #2 (B) given as: number of dives, % of the time the turtles are submerged, % of the time the turtles are swimming, distance travelled, mean, minimum and maximum temperature, mean VDBA and oxygen consumption.

	Before	During	After	Kruskal Wallis test	Dunn's multiple comparison test		
	n=9	n=3	n=4	dr=2	before vs during	before vs after	During vs after
Median daily O_2 C ($ml\ min^{-1}\ kg^{-0.83}$)	0.62	0.97	0.99	$p<0.05^*$ Hc=10.76	Q=2.78 $p<0.05^*$	Q=1.61 $p>0.05$	Q=0.78 $p>0.05$
Daily amount of time swimming (%)	20	91	86	$p<0.01^{**}$ Hc=10.5	Q=2.74 $p<0.05^*$	Q=2.48 $p<0.05^*$	Q=0.0 $p>0.05$
Daily amount of time submerged (%)	84	44	57	$p<0.001^{***}$ Hc=11.23	Q=3.20 $p<0.05^*$	Q=1.93 $p>0.05$	Q=0.85 $p<0.05$
Daily distance travelled (km)	17	57	70	$p<0.01^{**}$ Hc=10.5	Q=2.70 $p<0.05^*$	Q=1.56 $p<0.05^*$	0.34 $p>0.05$

Table 2: The effect of the tropical storm on the behaviour of turtle #2 provided as estimated daily median oxygen consumption, median daily amount of time swimming and amount submerged and median distance travelled. n=number of days. Kruskal Wallis and Dunn's multiple comparison tests were conducted a * significant where $p<0.05$ and ** significant where $p<0.01$.

Supplemental figures




Figure S1. (A-D) Four examples of diving behaviour for turtle #1: A) 5/31 just after the turtle nested, B) 6/4; 4 days after turtle #1 nested, C) 6/6; 6 days after turtle #1 nested and D) 6/10; 8 days before turtle #1 renested. (E-H) Four examples of diving behaviour for turtle #2: E) 6/14; just after the turtles nested, F) 6/16; 2 days after turtle #2 nested, G) 6/24; 10 days after turtle #2 nested and during the tropical storm and H) 6/28; 6 days before turtle #2 renested and after tropical storm had passed the Gulf of Mexico. Color code gives flipper beats/min




Figure S2: Median flipper beat of the different dive phases. Turtle #1: A) Dive type 1. B) Dive type 2. C) Dive type 3. D) Dive type 4. Turtle #2: E) Dive type 1. F) Dive type 2. G) Dive type 3. H) Dive type 4. Middle line in box represents the median, lower box bounds the first quartile, upper box bounds the third quartile, whiskers are 1.5 times the interquartile range, + are outliers. n.s is no significant difference. *significant where $p < 0.05$, ** significant where $p < 0.01$, and *** significant where $p < 0.001$ based on a Mann-Whitney test.




Figure S3: Instantaneous VDBA in (g) plotted against flipper beat/min using all data from the two turtles (Blue: Turtle #1, Red: Turtle #2).

Supplemental figures




Figure S1. (A-D) Four examples of diving behaviour for turtle #1: A) 5/31 just after the turtle nested, B) 6/4; 4 days after turtle #1 nested, C) 6/6; 6 days after turtle #1 nested and D) 6/10; 8 days before turtle #1 renested. (E-H) Four examples of diving behaviour for turtle #2: E) 6/14; just after the turtles nested, F) 6/16; 2 days after turtle #2 nested, G) 6/24; 10 days after turtle #2 nested and during the tropical storm and H) 6/28; 6 days before turtle #2 renested and after tropical storm had passed the Gulf of Mexico. Color code gives flipper beats/min




Figure S2: Median flipper beat of the different dive phases. Turtle #1: A) Dive type 1. B) Dive type 2. C) Dive type 3. D) Dive type 4. Turtle #2: E) Dive type 1. F) Dive type 2. G) Dive type 3. H) Dive type 4. Middle line in box represents the median, lower box bounds the first quartile, upper box bounds the third quartile, whiskers are 1.5 times the interquartile range, + are outliers. n.s is no significant difference. *significant where $p < 0.05$, ** significant where $p < 0.01$, and *** significant where $p < 0.001$ based on a Mann-Whitney test.




Figure S3: Instantaneous VDBA in (g) plotted against flipper beat/min using all data from the two turtles (Blue: Turtle #1, Red: Turtle #2).