

REVIEW

Linking environmental salinity to respiratory phenotypes and metabolic rate in fishes: a data mining and modelling approach

Till S. Harter¹, Christian Damsgaard^{2,3} and Matthew D. Regan^{4,*}

ABSTRACT

The gill is the primary site of ionoregulation and gas exchange in adult teleost fishes. However, those characteristics that benefit diffusive gas exchange (large, thin gills) may also enhance the passive equilibration of ions and water that threaten osmotic homeostasis. Our literature review revealed that gill surface area and thickness were similar in freshwater (FW) and seawater (SW) species; however, the diffusive oxygen (O_2) conductance (G_d) of the gill was lower in FW species. While a lower G_d may reduce ion losses, it also limits O_2 uptake capacity and possibly aerobic performance in situations of high O_2 demand (e.g. exercise) or low O_2 availability (e.g. environmental hypoxia). We also found that FW fishes had significantly higher haemoglobin (Hb)– O_2 binding affinities than SW species, which will increase the O_2 diffusion gradient across the gills. Therefore, we hypothesized that the higher Hb– O_2 affinity of FW fishes compensates, in part, for their lower G_d . Using a combined literature review and modelling approach, our results show that a higher Hb– O_2 affinity in FW fishes increases the flux of O_2 across their low- G_d gills. In addition, FW and SW teleosts can achieve similar maximal rates of O_2 consumption ($\dot{M}_{O_2,max}$) and hypoxia tolerance (P_{crit}) through different combinations of Hb– O_2 affinity and G_d . Our combined data identified novel patterns in gill and Hb characteristics between FW and SW fishes and our modelling approach provides mechanistic insight into the relationship between aerobic performance and species distribution ranges, generating novel hypotheses at the intersection of cardiorespiratory and ionoregulatory fish physiology.

KEY WORDS: Teleost, Osmorespiratory compromise, Ionoregulation, Aerobic metabolism, Exercise, Hypoxia

Introduction

Seawater (SW) and freshwater (FW) environments present different osmoregulatory challenges to resident species. Teleosts in the two environments have similar internal ion concentrations (Holmes and Donaldson, 1969) and, therefore, are hypo-osmotic compared with ion-rich SW, where they risk losing water to the environment, and hyper-osmotic compared with ion-poor FW, where they risk losing ions (Evans et al., 2005). The fish gill is the primary site of gas exchange and presents a large, thin epithelium to the surrounding water that facilitates the uptake of oxygen (O_2) and the excretion of the metabolic by-products carbon dioxide and ammonia. However, those gill morphometrics that maximize diffusive gas exchange may

also increase passive ion and water fluxes that, if unopposed, would lead to osmotic disturbances. These conflicting requirements for ionoregulation and O_2 uptake at the fish gill were recognized over 100 years ago by August Krogh (Ege and Krogh, 1914; Krogh, 1937) and later substantiated in seminal work by Randall et al. (1972), Gonzalez and McDonald (1992) and others. Today, the effect is known as the osmorespiratory compromise, and it has been treated in detailed reviews (Gilmour and Perry, 2018; Gonzalez, 2011; Nilsson et al., 2012; Perry, 1998; Sardella and Brauner, 2007; Wood and Eom, 2021). The present Review builds on previous work to uncover general patterns among teleost fishes that link environmental salinity to gill and blood phenotypes and to provide mechanistic insight into the physiology underlying these observations by using a modelling approach.

FW and SW teleosts have evolved divergent strategies to overcome the osmotic challenges imposed by their respective environments, involving anatomical and physiological adaptations at different organ systems. Briefly, FW fishes take up ions at the gills and excrete dilute urine via the kidneys, whereas SW fishes take up water and ions via their digestive tracts and excrete ions via the gills and kidneys, and with their faeces (Evans et al., 2005; Grosell, 2010; Marshall and Grosell, 2006). These divergent mechanisms enable FW and SW fishes to maintain water and ion homeostasis in their respective environments and are well studied, but their effects on gill morphometrics and, thus, O_2 uptake capacity, have not been studied broadly. To address this knowledge gap, we conducted a systematic literature review of gill, blood and metabolic characteristics in 355 teleost species in relation to their aquatic habitat (155 FW, 200 SW species; see Appendix for details on the literature review and Table S2 for raw data).

Gill oxygen conductance and Hb– O_2 binding affinity are associated with salinity

Our initial analysis indicated that FW fishes, on average, had significantly smaller mass-specific gill surface area (GSA) than SW fishes ($P=0.019$; Fig. 1A), consistent with what has long been reported in the literature. However, we also observed differences in body mass between the investigated FW and SW fishes, which biases this analysis because GSA scales allometrically with body mass. Furthermore, these interspecific comparisons are dominated by a few hyper-diverse taxa (Ostariophysi in FW and Percomorpha in SW; Vega and Wiens, 2012), which may introduce an additional phylogenetic bias. Therefore, to test whether salinity affected GSA while taking into account allometric scaling and phylogenetic non-independence of species, we fitted a phylogenetically corrected linear model to the \log_{10} -transformed GSA and body mass data (Fig. 1B), and then calculated the vertical residuals that provided a body mass-independent measure of GSA (Garland et al., 1992). These residuals were independent of salinity ($P=0.672$; Fig. 1C), indicating that, once corrected for mass and phylogenetic relatedness, teleosts in FW and SW had similar GSAs. These

¹Marine Biology Research Division, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92093, USA. ²Zoophysiology, Department of Biology, Aarhus University, 8000 Aarhus, Denmark. ³Aarhus Institute of Advanced Studies, Aarhus University, 8000 Aarhus, Denmark. ⁴Département de sciences biologiques, Université de Montréal, Montréal, QC, Canada, H3T 1J4.

*Author for correspondence (matthew.regan@umontreal.ca)

© T.S.H., 0000-0003-1712-1370; C.D., 0000-0002-5722-4246; M.D.R., 0000-0001-9341-5747




Fig. 1. See next page for legend.

Fig. 1. Systematic differences in the gill and blood characteristics of freshwater (FW, green) and seawater (SW, blue) teleost fishes.

(A) The effect of salinity on mass-specific gill surface area (GSA). (B) There was a significant effect of body mass on GSA, as determined by a generalized linear mixed model that accounted for phylogenetic non-independence of species. Regression line: $\log_{10}(\text{gill surface area}) = 0.63 + 0.93 \log_{10}(\text{body mass})$, where GSA is in mm^2 and body mass is in g. (C) The residuals of this model provide a mass-independent measure of GSA and the effect of salinity was tested by a phylogenetic ANOVA simulation that revealed no differences between FW and SW species. (D) The blood–water diffusion distance (gill thickness) in FW and SW species. (E) Gill thickness was independent of body mass and (F) the phylogenetically corrected residuals were not significantly different between FW and SW species. (G) Mass-specific gill oxygen (O_2) conductance (G_d) was calculated from GSA and thickness (and the parameters reported in Table S1). (H) G_d was dependent on body mass. Regression line: $\log_{10}(G_d) = 0.86 + 0.92 \log_{10}(\text{body mass})$, where G_d is in $\mu\text{mol mmHg}^{-1} \text{h}^{-1}$ and body mass is in g. (I) The residuals of $\log_{10}G_d$ were lower in FW than in SW species. (J) Haemoglobin (Hb) O_2 affinity (expressed as P_{50} of Hb – the P_{O_2} at which Hb is 50% saturated with O_2). (K) Hb P_{50} was dependent on temperature. Regression line: $\log_{10}(P_{50}) = 6.332 - 1534T^{-1}$, where P_{50} is in mmHg and temperature (T) is in K. (L) The phylogenetically and temperature-corrected residuals of $\log P_{50}$ were significantly lower in FW than in SW fishes. All values were mined from the literature according to the search protocols described in the Appendix. Points are values for individual fish species.

findings illustrate the importance of accounting for body mass and species relatedness in interspecific comparisons of traits that scale allometrically with body mass. In contrast, the blood–water diffusion distance (hereafter, gill thickness; Fig. 1D) was unaffected by body mass ($P=0.789$; Fig. 1E) and, thus, these data were only corrected for the phylogenetic relatedness of the species. As with GSA, we found no significant effect of salinity on gill thickness ($P=0.140$; Fig. 1F), indicating that other factors must determine the observed variability.

To explore the interacting effects of GSA and thickness, we calculated the O_2 diffusive conductance of the gills (G_d) in FW and SW species, which describes the amount of O_2 that can diffuse across the gills for a given P_{O_2} gradient. In our study, G_d was calculated from GSA and gill thickness values from the literature, and from the gill characteristics described in Table S1 (Dejours, 1981). We found 33 species with reported values for both GSA and thickness (Fig. 1G) and, after correcting for body mass and phylogenetic relatedness (Fig. 1H), FW fishes had significantly lower G_d values than SW fishes ($P=0.017$; Fig. 1I). This result should be interpreted in light of the relatively few species with thickness measurements and the fact that some species with extreme gill characteristics may disproportionately influence the analysis (such as the highly active Scombridae in SW and some facultative air-breathing fishes in FW). We tried expanding our analysis to all actinopterygian fishes to obtain additional values for G_d but found that gill thickness values were also largely unavailable. Additional measurements of gill thickness in other FW and SW fishes are needed to strengthen the relationships between G_d and environmental salinity that we found in our analysis. Nevertheless, the data that are currently available indicate that, even after mass and phylogenetic corrections, FW teleosts have lower G_d than SW teleosts, and if substantiated more broadly, this finding may change our understanding of the physiological implications of environmental salinity on gill phenotypes.

Smaller and thicker gills can benefit fishes by reducing the rate of passive ion and water fluxes that can disrupt osmotic homeostasis (Greco et al., 1996; Henriksson et al., 2008; Perry, 1998; Sollid et al., 2003). However, the gill is also the primary site for gas exchange and smaller and thicker gills will result in lower G_d . In FW fishes, lower G_d may limit O_2 -uptake capacity and could impair their performance in situations of high O_2 demand (e.g. exercise) or

low O_2 availability (e.g. environmental hypoxia). Interestingly, our data revealed a potential compensatory mechanism, as FW species had a ~ 2 -fold higher haemoglobin (Hb) O_2 affinity than SW species (expressed as lower P_{50} values of Hb – the P_{O_2} at which Hb is 50% saturated with O_2 ; $P<0.001$; Fig. 1L). Hb P_{50} values were corrected for phylogenetic relatedness and a significant temperature effect using a Van't Hoff plot ($P=0.001$; Fig. 1K). Because Hb is the principal O_2 carrier in the blood and its O_2 -binding characteristics determine the kinetics of O_2 loading at the gills and unloading at the tissues, the higher Hb– O_2 affinity of FW species could enhance the diffusion gradient of O_2 across the fish gill and theoretically support higher rates of metabolic O_2 consumption (\dot{M}_{O_2} ; a proxy for metabolic rate). These observations led us to hypothesize that the high Hb– O_2 affinities of FW species compensate for their lower G_d , allowing FW species to maintain similar \dot{M}_{O_2} as SW species during aerobic challenges, such as exercise and hypoxia.

To test this hypothesis, we explored the interacting effects of G_d and Hb P_{50} using a mathematical model (based on Malte and Weber, 1985) that predicts the maximal \dot{M}_{O_2} ($\dot{M}_{\text{O}_2, \text{max}}$) that a fish can support over a range of gill and blood characteristics, as well as environmental conditions (see Appendix for detailed model description). Specifically, we tested three predictions derived from our hypothesis: all else being equal, (1) a higher Hb– O_2 affinity will compensate for a lower G_d to maintain arterial O_2 transport; (2) FW and SW teleosts will achieve similar $\dot{M}_{\text{O}_2, \text{max}}$ using different combinations of G_d and Hb– O_2 affinity; and (3) FW and SW teleosts will achieve similar hypoxia tolerance (represented by P_{crit} – the lowest water P_{O_2} at which a fish can maintain its standard \dot{M}_{O_2} ; $\dot{M}_{\text{O}_2, \text{std}}$) using different combinations of G_d and Hb– O_2 affinity. We then compared the modelled results with empirical values mined from the literature that revealed novel patterns in the divergent gill and Hb phenotypes of FW and SW teleosts in relation to environmental salinity.

Prediction 1: higher Hb– O_2 affinity will compensate for lower G_d to maintain arterial O_2 transport

First, we used our $\dot{M}_{\text{O}_2, \text{max}}$ model to investigate the effects of gill and blood characteristics on the equilibration of P_{O_2} across the gill epithelium using the lowest, median and highest G_d values for FW and SW fishes that we observed in the literature (using only values that were based on $N>1$), over the range of Hb– O_2 affinities reported in Fig. 1. FW fishes with the lowest G_d were severely diffusion-limited, reflected in a poor equilibration of P_{O_2} between the counter-current flows of water and blood (Fig. 2A). $\dot{M}_{\text{O}_2, \text{max}}$ in these fishes was generally low, but the highest values were achieved at the lowest Hb P_{50} . Therefore, when all else is equal, a higher Hb– O_2 affinity can partially compensate for a low G_d by increasing the P_{O_2} gradient across the epithelium. This finding is consistent with previous work that modelled the diffusion of O_2 across the fish gill and found an increased O_2 extraction from the water and lower ventilatory requirements at high Hb– O_2 affinities (Malte and Weber, 1987). In addition, a reduction in Hb– O_2 affinity caused a steep decline in the arterial O_2 saturation of Hb (S_{aO_2}), which determines the maximum capacity for blood O_2 transport and $\dot{M}_{\text{O}_2, \text{max}}$ (Gallaughier et al., 2001). In SW species, the lowest, median and highest values for G_d were higher than in FW fishes. Consequently, O_2 uptake at the gills in SW was generally more efficient, as indicated by a nearly complete equilibration of P_{O_2} across the gill epithelium, especially at the lowest Hb P_{50} values (Fig. 2D).

An increase in G_d to median values increased $\dot{M}_{\text{O}_2, \text{max}}$ in both FW and SW fishes and shifted the optimal Hb P_{50} to higher values (Fig. 2B,E). Higher G_d also reduced the effect of Hb P_{50} on




Fig. 2. Simulations modelling the outcome of counter-current oxygen uptake at the fish gill. The partial pressure of O_2 (P_{O_2}) in blood (red) and water is shown along the length of the gill, where blood is oxygenated in the 0 to 1 direction and water is deoxygenated from 1 to 0. (A–C) O_2 uptake in FW fishes (green) for the values of G_d shown in Fig. 1 (0.02, 0.13 and $0.36 \mu\text{mol mmHg}^{-1} \text{g}^{-1} \text{h}^{-1}$ for lowest, median and highest G_d , respectively). (D–F) O_2 uptake in SW fishes (blue) for their range of G_d values (0.10 , 1.2 and $11.4 \mu\text{mol mmHg}^{-1} \text{g}^{-1} \text{h}^{-1}$ for lowest, median and highest G_d , respectively). Haemoglobin (Hb) O_2 affinity (Hb P_{50} , the partial pressure at which Hb is 50% saturated with O_2 ; mmHg) was set to cover the ranges shown in Fig. 1 for both groups. The $\dot{M}_{O_2, \text{max}}$ ($\mu\text{mol g}^{-1} \text{h}^{-1}$) and arterial Hb saturation (Sa_{O_2} ; %) values resulting from these simulations are listed in each panel.

Sa_{O_2} because O_2 uptake became less dependent on the P_{O_2} diffusion gradient. The fact that a high Hb- O_2 affinity exerts the greatest benefits in fish with the lowest G_d is consistent with our literature-mined data. Combined, these findings indicate that the higher Hb- O_2 affinities observed in FW fishes may have adaptive significance to compensate for their low- G_d gills, an idea that could be substantiated by investigating G_d in additional species.

SW fishes with median G_d achieved full O_2 saturation during blood transit through the gill, suggesting that $\dot{M}_{O_2, \max}$ is limited by maximal cardiac output (\dot{Q}_{\max}), rather than the diffusion characteristics of the gill (Fig. 2D), and the same was true for those FW and SW species with the highest G_d (Fig. 2C,F). These findings generally agree with experimental work, indicating that O_2 uptake at the gill is not diffusion-limited in normoxia, but rather is a function of maximal perfusion and/or ventilation rates (Daxboeck et al., 1982; Malte and Weber, 1985; Randall and Daxboeck, 1984). However, in >50% of the FW species in our mined dataset, $\dot{M}_{O_2, \max}$ may be limited by gill diffusion characteristics, where increases in \dot{Q} are inconsequential. These results must be interpreted with respect to the \dot{Q}_{\max} value for rainbow trout that we used in our simulations, which may be higher than those of many low- G_d FW species. Nevertheless, our results highlight systematic differences in cardiorespiratory phenotypes between FW and SW teleosts that have ecological consequences. Those species with the highest G_d may achieve increases in $\dot{M}_{O_2, \max}$ through increases in \dot{Q} , enabling more active lifestyles that are not available to low- G_d species. A broader assessment of G_d , $\dot{M}_{O_2, \max}$ and \dot{Q} in relation to environmental salinity in teleosts may shed light on the mechanistic basis for our observations and their ecological significance.

Finally, our simulations also showed that the Hb P_{50} values that achieved the highest Sa_{O_2} did not necessarily produce the highest $\dot{M}_{O_2, \max}$. Lower Hb P_{50} values can safeguard Sa_{O_2} and improve $\dot{M}_{O_2, \max}$ in those fish with the lowest G_d . However, the higher G_d we observed in SW species would lessen the benefits of a lower Hb P_{50} , potentially enabling them to exploit a higher range of Hb P_{50} values that promote the offloading of O_2 at the tissues, which we explore in more detail below.

Prediction 2: FW and SW teleosts will achieve similar $\dot{M}_{O_2, \max}$ using different combinations of G_d and Hb- O_2 affinity

The optimal Hb P_{50} is a compromise between the physiological requirements for O_2 loading at the gas exchange surface and unloading at the tissue capillaries (Brauner and Wang, 1997; Harter and Brauner, 2017; Wang and Malte, 2011). To highlight this trade-off in more detail, Fig. 3 shows the $\dot{M}_{O_2, \max}$ that can theoretically be attained over the range of Hb P_{50} and G_d values in FW and SW teleosts. Generally, those species with the lowest G_d achieved the lowest $\dot{M}_{O_2, \max}$ and increasing G_d improved $\dot{M}_{O_2, \max}$, consistent with a diffusion limitation on O_2 uptake. However, $\dot{M}_{O_2, \max}$ in species with median G_d matched those of species with the highest G_d , indicating a progressive perfusion limitation, which is in line with the results shown in Fig. 2. As the upper boundary for $\dot{M}_{O_2, \max}$ was set by perfusion, the two groups achieved similar $\dot{M}_{O_2, \max}$ of $\sim 13 \mu\text{mol g}^{-1} \text{h}^{-1}$ at the \dot{Q}_{\max} for rainbow trout ($53 \text{ ml kg}^{-1} \text{min}^{-1}$).

Increasing Hb- O_2 affinity also had beneficial effects on $\dot{M}_{O_2, \max}$, but only until an optimum value was reached, after which further increases in Hb- O_2 affinity severely decreased $\dot{M}_{O_2, \max}$. The effect of Hb- O_2 affinity on $\dot{M}_{O_2, \max}$ depended on G_d , where a reduced G_d shifted the optimal P_{50} for $\dot{M}_{O_2, \max}$ to lower values. For example, when G_d was reduced from median to lowest values, the optimal P_{50} shifted from 31 to 3 mmHg in FW fishes (Fig. 3A) and from 39 to 25 mmHg in SW fishes (Fig. 3B). These ranges of optimal P_{50} map well onto the mined P_{50} values for FW and SW fishes, respectively (Fig. 1).

Our simulations are also in line with data on rainbow trout acclimated to soft water that thickened their gills from 3 to 6 μm as a result of ionocyte proliferation and compensated for the impaired G_d by decreasing P_{50} from 18 to 12 mmHg (Greco et al., 1996; Perry, 1998; Perry et al., 1996). Gill morphology and P_{50} are plastic traits, and it seems that they may respond in concert to overcome physiological challenges to O_2 uptake in a way that depends on environmental salinity. The situation is complicated as the diffusive pathways for O_2 , water and ions may differ and many species exert some control over these 'passive' fluxes, such as observed in extremely hypoxia-tolerant species (Wood et al., 2009), active




Fig. 3. Maximal rate of oxygen consumption ($\dot{M}_{O_2, \max}$) as a function of haemoglobin O_2 affinity (Hb P_{50}) and gill O_2 conductance (G_d). (A,B) Results are for FW (green) and SW (blue) fishes, respectively. G_d and Hb P_{50} values were chosen to cover the ranges reported in Fig. 1 for both groups (see Fig. 2 caption for details).

species (Gonzalez and McDonald, 1994), and after exercise training (Gallaughan et al., 2001; Postlethwaite and McDonald, 1995). A salinity-specific effect on gill permeability has not been studied broadly, but data for a few species exist. FW-acclimated killifish have higher branchial water permeability than SW conspecifics, but they display similar diffusive ion fluxes (in opposite directions) during hypoxic hyperventilation (Giacomin et al., 2019; Wood et al., 2019). However, while FW-acclimated killifish do not alter ionocyte density, they do decrease their overall GSA (Giacomin et al., 2019), which is consistent with our observation of lower G_d in FW fishes that may perhaps occur via different mechanisms in different species and conditions. If this effect is substantiated more broadly, one may hypothesize that the hyper-osmoregulatory strategy of FW fishes has led not only to lower G_d but also to higher Hb–O₂ affinities that balance the requirements for O₂ uptake and tissue O₂ extraction at the prevailing gill diffusion characteristics. Likewise, the hypo-osmoregulatory strategy of SW fishes may be permissive of high- G_d gills that are best matched by higher Hb P_{50} and that enable increases in \dot{Q} and $\dot{M}_{O_2,max}$ in species with active lifestyles where exercise performance is linked to evolutionary fitness.

To further explore the interacting effects of G_d and Hb–O₂ affinity on $\dot{M}_{O_2,max}$, we plotted the upper and lower boundaries for G_d as a function of Hb P_{50} , resulting in areas of theoretical $\dot{M}_{O_2,max}$ that can be achieved by the gill and blood characteristics of FW and SW fishes (Fig. 4A). The lower areas were calculated at the \dot{Q}_{max} for rainbow trout, and because of the perfusion limitation of $\dot{M}_{O_2,max}$, FW and SW fishes with the highest G_d achieved similar $\dot{M}_{O_2,max}$. To corroborate these modelling results, we then plotted empirically measured $\dot{M}_{O_2,max}$ values onto the theoretical areas. Generally, the measured and calculated values agreed well. However, one cluster of FW fishes with high Hb–O₂ affinity fell above the predicted lines, which is likely explained by higher tissue O₂ conductance that maintains tissue O₂ extraction from high-affinity Hbs, and perhaps by lower Hill coefficients that enhance O₂ extraction from the water (Malte and Weber, 1987); both these values were kept constant in our model. Other FW and SW species may achieve \dot{Q}_{max} values that exceed those of rainbow trout. For instance, sockeye salmon swimming in FW reached $\dot{M}_{O_2,max}$ of 27.2 $\mu\text{mol g}^{-1} \text{h}^{-1}$ at a \dot{Q}_{max} of 67.8 $\text{ml kg}^{-1} \text{min}^{-1}$ (Steinhausen et al., 2008) and in some sockeye salmon populations with longer migration routes, \dot{Q}_{max} can exceed 100 $\text{ml kg}^{-1} \text{min}^{-1}$, nearly 2-fold the value of rainbow trout (Eliason et al., 2013). Whether the gill characteristics of these anadromous sockeye salmon are truly representative of a FW species is unclear, but despite their semelparous life history, they appear to exert tight control over ion homeostasis, even during their final FW stage.

We then explored whether a salinity-specific pattern in $\dot{M}_{O_2,max}$ exists across empirically measured values for 130 fish species (51 FW, 79 SW), and we corrected these data for allometric scaling with body mass, temperature and phylogenetic relatedness (Fig. 4B). Across all species, there was no significant effect of salinity on $\dot{M}_{O_2,max}$ ($P < 0.067$; Fig. 4C), consistent with our prediction that FW and SW teleosts may achieve similar $\dot{M}_{O_2,max}$ through different combinations of gill and blood characteristics. However, the low P -value in our analysis on 130 species suggests that future studies, especially those with larger sample sizes, may be able to resolve potential differences in $\dot{M}_{O_2,max}$ between FW and SW teleosts. Finally, $\dot{M}_{O_2,max}$ is a complex physiological trait that is strongly influenced by methodology, animal condition and behaviour (Killen et al., 2017; Norin and Clark, 2016); exploring these sources of variability and accounting for

them by statistical means may be a worthwhile avenue for future analyses.

Some of the highest $\dot{M}_{O_2,max}$ values are found in the fast-swimming, pelagic, SW teleosts (Wegner et al., 2010), including billfishes (Istiophoridae and Xiphiidae), dolphinfishes (Coryphaenidae), jacks (Carangidae) and tunas (Scombridae). Reliable data during maximal exercise in these animals are notoriously difficult to obtain, but a few measurements are available. Based on submaximal swimming trials, the $\dot{M}_{O_2,max}$ in skipjack tuna has been estimated at 68.7 $\mu\text{mol g}^{-1} \text{h}^{-1}$ (Brill, 1987; Dewar and Graham, 1994; Gooding et al., 1981) and therefore exceeds the value for rainbow trout by more than 5-fold. These high $\dot{M}_{O_2,max}$ values are enabled by GSAs of nearly 20 $\text{cm}^2 \text{g}^{-1}$ in yellowfin and skipjack tunas (Brill and Bushnell, 2001), and gill thicknesses as low as 0.5 μm (Hughes, 1970, 1984), resulting in G_d values that are 3-fold higher than the highest values of any FW species. Clearly, a unique physiology in tunas is required to sustain such high $\dot{M}_{O_2,max}$ that cannot be achieved with the cardio-respiratory characteristics of rainbow trout or most other teleosts. Therefore, to determine what physiological adaptations are required in tuna to attain their high $\dot{M}_{O_2,max}$, we re-calibrated our model to the skipjack tuna (Fig. 4A; light-shaded areas). This adjustment revealed that the high $\dot{M}_{O_2,max}$ in tuna is feasible only if \dot{Q}_{max} is increased ~ 6 -fold (to 318 $\text{ml kg}^{-1} \text{min}^{-1}$), blood [Hb] is increased 2.3-fold and O₂ conductance at the tissues is increased ~ 3 -fold over the values in rainbow trout. Whether these values are representative of some tuna species remains to be validated experimentally; however, the available data indicate that they may not be unreasonable estimates. Even in spinally blocked skipjack tuna, \dot{Q}_{max} has been measured at 132 $\text{ml kg}^{-1} \text{min}^{-1}$ (Brill, 1987; Bushnell, 1988), blood [Hb] can reach 2.3 mmol l^{-1} (Brill and Bushnell, 1991) and tissue O₂ conductance is undoubtedly increased by extremely high capillary density (Hulbert et al., 1979) and myoglobin concentrations (George and Stevens, 1978; Stevens and Carey, 1981). Theoretically, high \dot{Q}_{max} , [Hb] and tissue O₂ conductance may also be attainable by FW teleosts; but matching high- G_d gills may not. Our simulations show that even the highest G_d values in FW fishes are insufficient to produce tuna-like $\dot{M}_{O_2,max}$ and the upper boundary levels off at $\sim 40 \mu\text{mol g}^{-1} \text{h}^{-1}$. These values are still higher than any $\dot{M}_{O_2,max}$ recorded for a FW teleost, indicating that factors other than G_d may set the actual limit. Regardless, these fundamental relationships may explain why no FW fishes match the high $\dot{M}_{O_2,max}$ found in some SW species. Whether FW tuna-like fishes are, in fact, absent from the fossil record is a worthwhile avenue for future investigations. However, based on our findings it would seem that FW environments should remain devoid of fast-swimming pelagic analogues to the tunas.

Prediction 3: FW and SW teleosts will achieve similar P_{crit} using different combinations of GSA and Hb–O₂ affinity

Another important driver for variation in gill characteristics and Hb–O₂ affinity among fishes is the availability of environmental O₂ (Mandic et al., 2009). Under conditions of low environmental O₂ (hypoxia), a fish's indefinite survival depends on its ability to sustain $\dot{M}_{O_2,std}$ (Hughes, 1973). The lowest water P_{O_2} at which a fish can sustain $\dot{M}_{O_2,std}$ is termed the critical O₂ tension (P_{crit}), and at P_{O_2} below P_{crit} , a fish's survival becomes dependent on some combination of anaerobic glycolysis and metabolic depression, both of which are unsustainable in the long term (Beamish, 1964; Regan et al., 2017; Ultsch and Regan, 2019; Ultsch et al., 1978). P_{crit} is therefore a common metric of hypoxia tolerance that represents the




Fig. 4. $\dot{M}_{O_2,max}$ of FW and SW teleosts. (A) $Hb P_{50}$ was chosen to span the ranges reported in Fig. 1 for both groups. $\dot{M}_{O_2,max}$ was calculated for the upper and lower boundaries of gill G_d observed in FW (green) and SW (blue) fishes (see Fig. 2 caption for details). The dark shaded areas represent the model results using the cardio-respiratory characteristics of rainbow trout (see Table S1). The light shaded areas represent simulations after the model was adjusted to the high cardiac output (Q), $[Hb]$ and tissue G_d of tunas (respectively, 6-fold, 2.3-fold and 3-fold higher values than in rainbow trout). Empirically measured values for $Hb P_{50}$ and $\dot{M}_{O_2,max}$ from the literature were overlaid for FW and SW species whenever both parameters were available (1 rainbow trout, *Oncorhynchus mykiss*; 2 skipjack tuna, *Katsuwonus pelamis*; see search protocols described in the Appendix). (B) There were significant effects of body mass and temperature on $\dot{M}_{O_2,max}$, as determined by a generalized linear mixed model that accounted for phylogenetic non-independence of species. Regression line: $\log_{10}(\dot{M}_{O_2,max}) = 0.84 + 0.019T + 0.88 \log_{10}(\text{body mass})$, where $\dot{M}_{O_2,max}$ is in $\mu\text{mol h}^{-1}$, temperature (T) is in $^{\circ}\text{C}$ and body mass is in g. (C) The residuals of this model provide mass- and temperature-independent measures of $\dot{M}_{O_2,max}$ and the effect of salinity was tested by a phylogenetic ANOVA simulation that revealed no significant differences between FW and SW species.

suite of aerobic contributions to hypoxia survival in a single value (Regan et al., 2019); thus, we explored this metric in our simulations.

To estimate how differences in G_d and $Hb-O_2$ affinity may influence P_{crit} , we adjusted our model to predict $\dot{M}_{O_2,max}$ at progressively decreasing inspired water P_{O_2} (from 150 to 0 mmHg) and then fixed a horizontal line representing $\dot{M}_{O_2,std}$ at $1.5 \mu\text{mol g}^{-1} \text{h}^{-1}$, which is the value reported for resting rainbow trout (Kiceniuk and Jones, 1977). The P_{O_2} at which the $\dot{M}_{O_2,max}$ and $\dot{M}_{O_2,std}$ lines intersect represents the theoretical P_{crit} for a given set of

gill and blood characteristics. Our approach assumed that $\dot{M}_{O_2,max}$ and $\dot{M}_{O_2,std}$ respond similarly to reductions in P_{O_2} below P_{crit} and that the typical P_{crit} curve is biphasic. The former assumption is consistent with previous theoretical work (Esbaugh et al., 2021), empirical measurements (Claireaux et al., 2000) and the fact that aerobic scope for activity at sub- P_{crit} P_{O_2} is zero (Claireaux and Chabot, 2016; Ern et al., 2016). The latter assumption is true for some fish species, but not all (Wood, 2018; Farrell et al., 2021). However, most important for our simulations was not the shape of the $\dot{M}_{O_2,std}$ curve at P_{O_2} below P_{crit} , but rather the presence of a

benchmark $\dot{M}_{O_2, \text{std}}$ value to which the sub- P_{crit} $\dot{M}_{O_2, \text{max}}$ line could be compared. A biphasic P_{crit} curve enabled this.

The outcomes of these simulations are shown in Fig. 5 for the lowest, median and highest G_d that we observed in FW and SW fishes (Fig. 1). In all cases, increasing Hb–O₂ affinity or G_d led to lower P_{crit} values, representing a higher hypoxia tolerance of the fish. SW fishes achieved lower P_{crit} than FW fishes when G_d was at the lowest or median values. However, the lowest overall P_{crit} values were achieved by FW fishes with high G_d and high Hb–O₂ affinity. These and our previous results show that an increased Hb–O₂ affinity can generally compensate for impaired diffusion characteristics at the gill caused by either low G_d or low environmental O₂. In both cases, \dot{M}_{O_2} is limited by the reduction in SaO₂ that can be improved by increasing Hb–O₂ affinity, and these results are in close agreement with previous work (Wang and Malte, 2011).

To validate our simulations, we plotted the P_{crit} values for high and low G_d as a function of Hb–O₂ affinity, spanning the area of theoretical P_{crit} that were predicted by the model. We then overlaid empirically measured P_{crit} values from the literature for FW and SW fishes (Fig. 6A). For FW fishes, we found an excellent agreement between the model predictions and measured P_{crit} values. In addition, P_{crit} values in FW fishes were highly variable, spanning an order of magnitude. GSA and gill thickness are plastic traits that respond to hypoxia, environmental ion concentration and temperature. The underlying mechanisms include dynamic changes in the redistribution of branchial blood flow and cellular responses that reversibly alter the morphology of the gill (Wood and Eom, 2021). For example, the crucian carp can increase GSA by ~7.5-fold after an acclimation period to hypoxia, largely through the receding of an interlamellar cell mass (Sollid et al., 2003). These acclimation responses can occur quickly, as in goldfish, where GSA increases ~2-fold in just hours (Regan et al., 2017). High GSA plasticity is often found in hypoxia-tolerant rather than -intolerant species (Dhillon et al., 2013), and in more FW than SW species (Gilmour and Perry, 2018). This may be because many FW environments are particularly hypoxia-prone (Diaz and Breitburg, 2009) or simply because relatively few SW species have been studied in this respect.

In contrast, SW fishes occupied a narrow band in P_{crit} values over a broad range of Hb P_{50} , which is supported by the modelled and empirical data (Fig. 6A). However, the modelled data were generally offset to lower P_{crit} values. The reasons for this discrepancy may relate to our model assumptions not accurately representing some of the *in vivo* characteristics of SW fishes; for example: (i) many SW species may have higher $\dot{M}_{O_2, \text{std}}$ than rainbow trout, and/or (ii) the boundaries for G_d in our dataset may not match those of SW species for which P_{crit} and Hb P_{50} data are available, both representing only a small subset of SW species. Regardless, the combined data revealed that FW and SW fishes occupy different quadrants in the P_{crit} versus Hb P_{50} relationship, which indicates that they achieve their hypoxia tolerance by different mechanisms. SW fishes rely on their higher G_d to maintain branchial O₂ uptake in hypoxia (Fig. 5), resulting in P_{crit} values that are largely independent of Hb P_{50} and thus fall within a narrow range (Fig. 6A). In contrast, FW fishes rely on their high Hb–O₂ affinity to maintain branchial O₂ uptake in hypoxia and, when combined with high G_d values, achieve the lowest P_{crit} observed in teleosts.

To test whether there is a systematic difference in P_{crit} between FW and SW teleosts, we mined literature values for 142 species (46 FW and 96 SW). After correcting for phylogenetic relatedness

of species, we found no significant difference in P_{crit} between FW and SW teleosts ($P=0.891$; Fig. 6C). This stands in contrast with our dataset when not corrected for phylogeny, which showed significantly lower P_{crit} in FW than in SW fish, and to the metanalysis of Rogers et al. (2016) and intraspecific salinity trials of Haney and Nordlie (1997), which reported lower P_{crit} in FW fishes under some, but not all, conditions. Therefore, the variation in P_{crit} across salinity environments appears to be driven by a few teleost clades with extreme P_{crit} values, again highlighting the importance of applying appropriate corrections of the raw data. Relatedly, most SW species on which P_{crit} experiments have been performed are native to environments that regularly experience hypoxia, such as coral reefs and intertidal zones. The mean SW P_{crit} value may therefore be skewed towards these hypoxia-adapted SW species and away from pelagic species that are less likely to regularly encounter hypoxia. Fewer P_{crit} values for pelagic species exist, perhaps because these fishes are not easily caught and experimented on. However, their addition to this comparison could further distinguish the salinity groups and is a worthwhile direction for future studies. Finally, there is substantial variability in P_{crit} values in the literature and, as with other complex performance metrics, P_{crit} should be interpreted with some caution regarding differences in methodology, animal condition and behaviour (Wood, 2018). Part of this variability may be due to the plasticity in gill morphology and blood characteristics in teleosts, perhaps coupled with different methodologies regarding the rates of hypoxia onset (Rogers et al., 2016). Nevertheless, the close agreement of the empirical and modelling data indicates that the general patterns in P_{crit} between FW and SW fishes are largely driven by factors that are manipulated in the model. Therefore, studying the interspecific variability in G_d , Hb–O₂ affinity and their interactions may provide a roadmap for future investigations into the mechanistic basis for P_{crit} in fish.

Conclusion and perspectives

Our analysis provides compelling evidence for systematic differences in gill and blood phenotypes between FW and SW teleosts that may be related to the osmotic characteristics of their aquatic environments. However, aerobic performance, assessed as $\dot{M}_{O_2, \text{max}}$ and P_{crit} , did not differ between FW and SW fishes; therefore, the two groups may use different combinations of gill and blood characteristics to support similar aerobic capacities. FW fishes generally had lower G_d , which has consequences for aerobic metabolism and may reflect a physiological constraint by their hyper-osmoregulatory strategy. The higher Hb–O₂ affinity of FW fishes may thus have adaptive significance, as it increases the diffusion gradient for O₂ across the gills. The benefits of a higher Hb–O₂ affinity may not fully compensate for the disadvantages of a low G_d , but rather, will enable a higher $\dot{M}_{O_2, \text{max}}$ within this constraint, by balancing the conflicting requirements for O₂ loading at the gills and unloading at the tissues. In contrast, the hypo-osmoregulatory strategy of SW teleosts is clearly permissive of higher G_d , which in turn may have lifted the brakes on the evolution of higher \dot{Q}_{max} and [Hb]. In tuna, these coordinated adaptations of their cardio-respiratory systems ultimately enabled extremely high $\dot{M}_{O_2, \text{max}}$ that are not attainable even by FW fishes with the highest G_d .

Gill morphology and Hb–O₂ binding characteristics are highly plastic traits in teleosts, and many species with euryhaline, anadromous or catadromous life cycles routinely transition between the FW and SW environments and similar transitions occurred repeatedly in the course of teleost evolution (Betancur-R




Fig. 5. Calculated critical oxygen tension (P_{crit}) for FW and SW teleost fishes. Simulations were run for the lowest, median and highest gill G_d in FW (green; A–C) and SW (blue; D–F) fishes, respectively (see Fig. 2 for details). The rate of O_2 consumption (\dot{M}_{O_2}) at decreasing water P_{O_2} (mmHg) was calculated with an $\dot{M}_{O_2,max}$ model (see Appendix), and standard \dot{M}_{O_2} ($\dot{M}_{O_2,std}$) was set to the value reported for rainbow trout, of $1.5 \mu\text{mol g}^{-1} \text{h}^{-1}$ (Kiceniuk and Jones, 1977). P_{crit} (mmHg) was determined as the intersection between the \dot{M}_{O_2} and $\dot{M}_{O_2,std}$ curves over the range of Hb P_{50} that we observed in FW and SW fishes (Fig. 1).




Fig. 6. Modelled and empirically measured P_{crit} for FW and SW teleosts. (A) Modelling results for P_{crit} of FW (green) and SW (blue) fishes as a function of $Hb P_{50}$ (mmHg). The solid symbols and lines represent the calculated P_{crit} values in FW and SW fish, as shown in Fig. 5, and the shaded areas represent the combinations of P_{crit} and $Hb P_{50}$ that are possible based on the gill and blood characteristics observed in both groups (Fig. 1). Empirically measured values for $Hb P_{50}$ and P_{crit} from the literature were overlaid for FW and SW species whenever both parameters were available. Dashed lines and shading indicate linear regressions through the empirical datasets for FW and SW fishes and their respective 95% confidence intervals. (B) There were no significant effects of body mass on P_{crit} , as determined by a generalized linear mixed model that accounted for phylogenetic non-independence of species. (C) The residuals of this model provide a mass-independent measure of P_{crit} and the effect of salinity was tested by a phylogenetic ANOVA simulation that revealed no significant differences between FW and SW species.

et al., 2015). The evolutionary dynamics that govern these transitions are worthy of further investigation and fish clades that include species that have recently transitioned between FW and SW may be well suited for this purpose.

The osmorepiratory compromise describes the conflicting requirements for O_2 uptake and ion/water flux at the gills and has typically been studied with respect to GSA. Our data indicate that gill thickness and G_d are important factors that determine $\dot{M}_{O_2,max}$ in fishes and should be considered in future studies on the osmorepiratory compromise by measuring O_2 uptake and unidirectional ion/water flux in more species in both FW and SW. Ideally, future studies should use standardized protocols for determining gill morphometrics, unidirectional ion/water flux, $\dot{M}_{O_2,max}$, P_{crit} and Hb characteristics in animals acclimated to a clearly defined set of environmental conditions, allowing for more reproducible interspecific

comparisons. These future studies may lay the foundation to test our hypothesis that invasions of the FW environment and the divergent selective pressures of the osmorepiratory compromise gave rise to adaptations that reduced G_d and $Hb P_{50}$, thus profoundly shaping the aerobic pathway of FW fishes.

Appendix

Literature review

We conducted literature reviews of five cardio-respiratory variables in teleost fishes: gill surface area (GSA), blood–water diffusion distance (gill thickness), $Hb-O_2$ affinity (expressed as P_{50} , the P_{O_2} at which Hb is 50% saturated with O_2), maximal O_2 consumption rate ($\dot{M}_{O_2,max}$) and critical O_2 tension (P_{crit} , the lowest P_{O_2} at which a fish can maintain standard \dot{M}_{O_2}). We included only teleost species in our datasets, which are by far the most diverse

group of fishes and account for ~95% of all extant species (Nelson et al., 2006). The reasons for excluding other groups, such as the Chondrichthyes, were based on their osmo-conforming physiology that changes the relationship between respiratory and ionoregulatory demands at the gill, and this group is almost exclusively found in the marine environment. We also excluded amphibious and obligatory air-breathing fishes (Damsgaard et al., 2020), as these traits fundamentally change the relationship between osmoregulation and respiration that we sought to examine (Graham, 1997). Additionally, the values we used were limited to those from individuals in juvenile or adult developmental stages, measured under control conditions (i.e. untreated and non-acclimated individuals).

Commenting on the methodological approaches used in literature studies was outside the scope of this Review and, generally, we considered data generated with different methods, while being aware that our variables of interest are sensitive to different experimental protocols. However, we did limit our search to Hb P_{50} measured on whole blood, and excluded those on haemolysates, which are less representative of *in vivo* conditions even when standardizing for allosteric effectors (Berenbrink, 2006). For GSA, we limited our search to measurements of total GSA or lamellar surface area.

All literature searches were performed on Clarivate's Web of Science®, in 2020 and 2021, using the following search terms: GSA (gill AND surface area AND fish; lamellar surface area AND fish); Hb P_{50} (oxygen* binding affinity AND fish; hemoglobin* AND P_{50} AND fish); P_{crit} (critical oxygen* AND fish; P_{crit} +fish; P_{crit} +hypoxia+fish); $\dot{M}_{O_2,max}$ (maximum metabolic rate+fish; MO_2max +fish). In instances where the mined studies cited other studies that our search did not uncover, we included these data if they were appropriate. Two recent reviews were particularly valuable in our search, providing many of the mined data points: Killen et al. (2017) and Rogers et al. (2016). Values, references and URLs for all cited studies can be found in Table S2.

We analysed our data in R Studio (v.1.3.1056) using a generalized linear mixed model (GLMM) to identify correlations between continuous variables using the *MCMCglmm()* function in the *MCMCglmm* package in R (Hadfield, 2010; Hadfield and Nakagawa, 2010). This method treats individual studies on the same species as a random effect (i.e. taking into account intraspecific variation) and accounted for phylogenetic non-independence of species. Here, we used a maximum clade credibility tree generated from 100 Bayesian posterior probability trees from Rabosky et al. (2018) using the *maxCladeCred()* function in the *phangorn* package in R (Schliep, 2011).

To test the effect of salinity on GSA and G_d , while taking into account the hypo-allometric scaling with body mass, we fitted a GLMM model to the \log_{10} -transformed data and calculated the residuals. These residuals provide a mass-independent measure of GSA and G_d (Garland et al., 1992). We then tested for an effect of salinity on the residuals with a phylogenetic analysis of variance simulation (Garland et al., 1993), using the *phylANOVA()* function in the *phytools* package in R (Revell, 2012). Similar approaches were used for the other variables, as detailed below.

To test for the effect of salinity on $\dot{M}_{O_2,max}$, while taking into account temperature and body mass, we fitted three GLMM models to the data:

$$\begin{aligned} \log_{10}(\dot{M}_{O_2,max}) &\sim \log_{10}(\text{body mass}) + \text{temperature} \quad (\text{DIC} = -124) \\ \log_{10}(\dot{M}_{O_2,max}) &\sim \text{temperature} \quad (\text{DIC} = -81.6) \\ \log_{10}(\dot{M}_{O_2,max}) &\sim \log_{10}(\text{body mass}) \quad (\text{DIC} = 205) \end{aligned}$$

We selected the full model based on the lowest deviance information criterion value (DIC), calculated the residuals, and tested for an effect of salinity on the $\dot{M}_{O_2,max}$ residuals, as described above. To test for the effect of salinity on Hb P_{50} we first fitted a GLMM model to $\log_{10}(P_{50})$ versus absolute temperature⁻¹ and calculated the residuals that we tested for an effect of salinity. P_{crit} and gill thickness were body mass independent; thus, we used the raw values in a phylogenetic ANOVA to test for an effect of salinity.

Mathematical model of gas exchange

To study the effects of Hb P_{50} and G_d on gas exchange at the fish gills, we implemented the previously published mathematical model of Malte and Weber (1985) using R v.3.6.2. in RStudio v.1.2.5033. Briefly, the model solves a system of differential equations to calculate the changes in water and blood P_{O_2} during counter-current gas exchange along the length of a model fish gill. In R, the differential equations were solved as a boundary value problem with venous P_{O_2} and inspired water P_{O_2} as the input parameters, using the *bvpsolve* package (Mazzia et al., 2014), with *bvptwp()* that uses a mono-implicit Runge–Kutta (MIRK) method with deferred corrections (Cash and Mazzia, 2005; Cash and Wright, 1991) and a continuation method. The output from these gas exchange simulations gives the calculated values for expired water P_{O_2} and arterial P_{O_2} , which is then used to calculate other arterial blood parameters. Next, the model predicts the outcome of gas exchange at the tissues to calculate venous P_{O_2} and other venous blood parameters that are then used for the next iteration of gas exchange at the gills; the cycle is repeated until the final calculated venous P_{O_2} is in equilibrium with the gas exchange systems at the gills and at the tissues (Malte and Weber, 1985, 1987, 1989, 2011). Our model focused only on the exchange of O_2 at the fish gill and did not take into account changes in CO_2 excretion; however, the model does simulate a pH shift at the tissues that leads to an increase in Hb P_{50} by the Bohr effect and recovery of Hb P_{50} during venous transit.

The model was 'calibrated' against the cardio-respiratory characteristics measured in rainbow trout (*Oncorhynchus mykiss*) at 10°C by Kiceniuk and Jones (1977). Specifically, the diffusion coefficient for O_2 at the gills (D_{O_2}) and the tissue O_2 conductance, two values for which no reliable measurements exist, were adjusted so that the model output matched the empirically measured $\dot{M}_{O_2,max}$, arterial and venous P_{O_2} , and tissue O_2 extraction in rainbow trout. All other input parameters for the model calibration are listed in Table S1 with their respective references.

In all simulations, [Hb] was set to 1 mmol l⁻¹ based on the summary data by Gallagher and Farrell (1998) and the cooperativity of Hb– O_2 binding (the Hill coefficient) was set to $n=2$. While this represents a simplistic approach, the effects of cooperativity on gas exchange at the fish gill have been addressed previously (Malte and Weber, 1987) and we observed no systematic differences in cooperativity between FW and SW teleosts that would warrant a more detailed investigation within the context of our study. In addition, some groups (e.g. Cottidae) display a wide range of Hb P_{50} at rather constant Hill coefficients, indicating that phylogeny may be a better predictor of Hill coefficients than, for example, Hb P_{50} (Mandic et al., 2009). Finally, to test our hypotheses, we adjusted the parameters of G_d and Hb P_{50} based on the values for FW and SW fishes that resulted from our review of the literature.

Acknowledgements

We thank Nick Wegner and Rod Wilson for insightful discussions and help with our literature review, Tommy Norin for critical feedback on our manuscript, and the two anonymous reviewers for their insightful comments.

Competing interests

The authors declare no competing or financial interests.

Funding

This study was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant (NSERC RGPIN-2021-03109) to M.D.R. C.D. is supported by the Carlsberg Foundation (Carlsbergfondet, CF18-0658), the Lundbeck Foundation (Lundbeckfonden, R346-2020-1210), the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement (754513), and the Aarhus Universitets Forskningsfond. T.S.H. is supported by a National Science Foundation (NSF) grant to Martin Tresguerres (1754994).

Data availability

All R code is publicly available from GitHub ($\dot{M}_{O_2,max}$ model and data analyses: github.com/tillharter/Fish_MO2max_models_in_R).

References

- Beamish, F. W. H.** (1964). Respiration of fishes with special emphasis on standard oxygen consumption: II. Influence of weight and temperature on respiration of several species. *Can. J. Zool.* **42**, 177-188. doi:10.1139/z64-016
- Berenbrink, M.** (2006). Evolution of vertebrate haemoglobins: histidine side chains, specific buffer value and Bohr effect. *Respir. Physiol. Neurobiol.* **154**, 165-184. doi:10.1016/j.resp.2006.01.002
- Betancur-R, R., Ortí, G. and Pyron, R. A.** (2015). Fossil-based comparative analyses reveal ancient marine ancestry erased by extinction in ray-finned fishes. *Ecol. Lett.* **18**, 441-450. doi:10.1111/ele.12423
- Brauner, C. J. and Wang, T.** (1997). The optimal oxygen equilibrium curve: a comparison between environmental hypoxia and anemia. *Am. Zool.* **37**, 101-108. doi:10.1093/icb/37.1.101
- Brill, R. W.** (1987). On the standard metabolic rates of tropical tunas, including the effect of body size and acute temperature change. *Fish. Bull.* **85**, 25-35.
- Brill, R. W. and Bushnell, P. G.** (1991). Effects of open- and closed-system temperature changes on blood oxygen dissociation curves of skipjack tuna, *Katsuwonus pelamis*, and yellowfin tuna, *Thunnus albacores*. *Can. J. Zool.* **69**, 1814-1821. doi:10.1139/z91-250
- Brill, R. W. and Bushnell, P. G.** (2001). The cardiovascular system of tunas. In *Fish Physiology* (ed. B. A. Block and E. D. Stevens), pp. 79-120. San Diego: Academic Press.
- Bushnell, P. G.** (1988). Cardiovascular and respiratory responses to hypoxia in three species of obligate ram ventilating fishes, skipjack tuna (*Katsuwonus pelamis*), yellowfin tuna (*Thunnus albacares*), and bigeye tuna (*T. obesus*). *PhD thesis*, Department of Physiology, University of Hawaii.
- Cash, J. R. and Mazzia, F.** (2005). A new mesh selection algorithm, based on conditioning, for two-point boundary value codes. *J. Comput. Appl. Math.* **184**, 362-381.
- Cash, J. R. and Wright, M. H.** (1991). A Deferred Correction Method for Nonlinear Two-Point Boundary Value Problems: Implementation and Numerical Evaluation. *SIAM J. Sci. and Stat. Comput.* **12**, 971-989.
- Claireaux, G. and Chabot, D.** (2016). Responses by fishes to environmental hypoxia: integration through Fry's concept of aerobic metabolic scope. *J. Fish Biol.* **88**, 232-251. doi:10.1111/jfb.12833
- Claireaux, G., Webber, D. M., Lagardère, J.-P. and Kerr, S. R.** (2000). Influence of water temperature and oxygenation on the aerobic metabolic scope of Atlantic cod (*Gadus morhua*). *J. Sea Res.* **44**, 257-265. doi:10.1016/S1385-1101(00)00053-8
- Damsgaard, C., Baliga, V. B., Bates, E., Burggren, W., McKenzie, D. J., Taylor, E. and Wright, P. A.** (2020). Evolutionary and cardio-respiratory physiology of air-breathing and amphibious fishes. *Acta Physiologica* **228**, e13406. doi:10.1111/apha.13406
- Daxboeck, C., Davie, P. S., Perry, S. F. and Randall, D. J.** (1982). Oxygen uptake in a spontaneously ventilating, blood-perfused trout preparation. *J. Exp. Biol.* **101**, 35-45. doi:10.1242/jeb.101.1.35
- Dejours, P.** (1981). *Principles of Comparative Respiratory Physiology*. Amsterdam: Elsevier/North-Holland Biomedical Press.
- Dewar, H. and Graham, J.** (1994). Studies of tropical tuna swimming performance in a large water tunnel - energetics. *J. Exp. Biol.* **192**, 13-31. doi:10.1242/jeb.192.1.13
- Dhillon, R. S., Yao, L., Matey, V., Chen, B.-J., Zhang, A.-J., Cao, Z.-D., Fu, S.-J., Brauner, C. J., Wang, Y. S. and Richards, J. G.** (2013). Interspecific differences in hypoxia-induced gill remodeling in carp. *Physiol. Biochem. Zool.* **86**, 727-739. doi:10.1086/673180
- Diaz, R. J. and Breitburg, D. L.** (2009). Chapter 1 The hypoxic environment. In *Fish Physiology, Hypoxia* (ed. J. Richards, A. Farrell and C. Brauner), pp. 1-23. Academic Press.
- Ege, R. and Krogh, A.** (1914). On the relation between the temperature and the respiratory exchange in fishes. *Int. Rev. Gesamten Hydrobiol. Hydrogr.* **7**, 48-55. doi:10.1002/iroh.19140070105
- Eliason, E. J., Clark, T. D., Hinch, S. G. and Farrell, A. P.** (2013). Cardiorespiratory performance and blood chemistry during swimming and recovery in three populations of elite swimmers: adult sockeye salmon. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **166**, 385-397. doi:10.1016/j.cbpa.2013.07.020
- Ern, R., Norin, T., Gamperl, A. K. and Esbaugh, A. J.** (2016). Oxygen dependence of upper thermal limits in fishes. *J. Exp. Biol.* **219**, 3376-3383.
- Esbaugh, A. J., Ackerly, K. L., Dichiera, A. M. and Negrete, B., Jr** (2021). Is hypoxia vulnerability in fishes a by-product of maximum metabolic rate? *J. Exp. Biol.* **224**, jeb232520. doi:10.1242/jeb.232520
- Evans, D. H., Piermarini, P. M. and Choe, K. P.** (2005). The multifunctional fish gill: dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiol. Rev.* **85**, 97-177. doi:10.1152/physrev.00050.2003
- Farrell, A. P., Mueller, C. A. and Seymour, R. S.** (2021). Coming up for air. *J. Exp. Biol.* **224**, jeb243101. doi:10.1242/jeb.243101
- Gallaugher, P. and Farrell, A. P.** (1998). Hematocrit and blood oxygen-carrying capacity. In *Fish Physiology* (ed. S. F. Perry, II and B. L. Tufts), pp. 185-227. New York: Academic press.
- Gallaugher, P. E., Thorarensen, H., Kiessling, A. and Farrell, A. P.** (2001). Effects of high intensity exercise training on cardiovascular function, oxygen uptake, internal oxygen transport and osmotic balance in chinook salmon (*Oncorhynchus tshawytscha*) during critical speed swimming. *J. Exp. Biol.* **204**, 2861-2872. doi:10.1242/jeb.204.16.2861
- Garland, T., Jr, Harvey, P. H. and Ives, A. R.** (1992). Procedures for the analysis of comparative data using phylogenetically independent contrasts. *Syst. Biol.* **41**, 18-32. doi:10.1093/sysbio/41.1.18
- Garland, T., Jr., Dickerman, A. W., Janis, C. M. and Jones, J. A.** (1993). Phylogenetic analysis of covariance by computer simulation. *Syst. Biol.* **42**, 265-292. doi:10.1093/sysbio/42.3.265
- George, J. C. and Stevens, E. D.** (1978). Fine structure and metabolic adaptation of red and white muscles in tuna. *Environ. Biol. Fish* **3**, 185-191. doi:10.1007/BF00691942
- Giacomin, M., Bryant, H. J., Val, A. L., Schulte, P. M. and Wood, C. M.** (2019). The osmorepiratory compromise: physiological responses and tolerance to hypoxia are affected by salinity acclimation in the euryhaline Atlantic killifish (*Fundulus heteroclitus*). *J. Exp. Biol.* **222**, jeb206599. doi:10.1242/jeb.206599
- Gilmour, K. M. and Perry, S. F.** (2018). Conflict and compromise: using reversible remodeling to manage competing physiological demands at the fish gill. *Physiology* **33**, 412-422. doi:10.1152/physiol.00031.2018
- Gonzalez, R. J.** (2011). Role of the gills | the osmorepiratory compromise. In *Encyclopedia of Fish Physiology* (ed. A. P. Farrell), pp. 1389-1394. San Diego: Academic Press.
- Gonzalez, R. J. and McDonald, D. G.** (1992). The relationship between oxygen consumption and ion loss in a freshwater fish. *J. Exp. Biol.* **163**, 317-332. doi:10.1242/jeb.163.1.317
- Gonzalez, R. and McDonald, D.** (1994). The relationship between oxygen uptake and ion loss in fish from diverse habitats. *J. Exp. Biol.* **190**, 95-108. doi:10.1242/jeb.190.1.95
- Gooding, R. M., Neill, W. H. and Dizon, A. E.** (1981). Respiration rates and low-oxygen tolerance limits in skipjack tuna, *Katsuwonus pelamis*. *Fish. Bull.* **79**, 31-48.
- Graham, J. B.** (1997). *Air-breathing fishes: Evolution, Diversity, and Adaptation*. New York: Academic Press.
- Greco, A. M., Fenwick, J. C. and Perry, S. F.** (1996). The effects of soft-water acclimation on gill structure in the rainbow trout *Oncorhynchus mykiss*. *Cell Tissue Res.* **285**, 75-82. doi:10.1007/s004410050622
- Grosell, M.** (2010). The role of the gastrointestinal tract in salt and water balance. In *Fish Physiology. The Multifunctional Gut of Fish* (ed. M. Grosell, A. P. Farrell and C. J. Brauner), pp. 135-164. London, UK: Academic Press.
- Hadfield, J. D.** (2010). MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R Package. *J. Stat. Soft.* **33**, 1-22. doi:10.18637/jss.v033.i02
- Hadfield, J. D. and Nakagawa, S.** (2010). General quantitative genetic methods for comparative biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *J. Evol. Biol.* **23**, 494-508. doi:10.1111/j.1420-9101.2009.01915.x
- Haney, D. and Nordlie, F.** (1997). Influence of environmental salinity on routine metabolic rate and critical tension of *Cyprinodon variegatus*. *Physiol. Zool.* **70**, 511-518. doi:10.1086/515867
- Harter, T. S. and Brauner, C. J.** (2017). The O₂ and CO₂ transport system in teleosts and the specialized mechanisms that enhance Hb-O₂ unloading to tissues. In *Fish Physiology* (ed. A. K. Gamperl, T. E. Gillis, A. P. Farrell and C. J. Brauner), pp. 1-107. New York: Academic Press.
- Henriksson, P., Mandic, M. and Richards, J. G.** (2008). The osmorepiratory compromise in sculpins: Impaired gas exchange is associated with freshwater tolerance. *Physiol. Biochem. Zool.* **81**, 310-319. doi:10.1086/587092
- Holmes, W. N. and Donaldson, E. M.** (1969). The Body Compartments and the Distribution of Electrolytes. In *Fish Physiology* (ed. Hoar, W. S.) and Randall, D. J.), pp. 1-90. New York: Academic Press.

- Hughes, G. M. (1970). Morphological measurements on the gills of fishes in relation to their respiratory function. *Folia Morphol (Praha)* **18**, 78-95.
- Hughes, G. M. (1973). Respiratory responses to hypoxia in fish. *Am. Zool.* **13**, 475-489. doi:10.1093/icb/13.2.475
- Hughes, G. M. (1984). General anatomy of the gills. In *Fish Physiology* (ed. W. S. Hoar and D. J. Randall), pp. 1-63. New York: Academic Press.
- Hulbert, W. C., Guppy, M., Murphy, B. and Hochachka, P. W. (1979). Metabolic sources of heat and power in tuna muscles: I. Muscle fine structure. *J. Exp. Biol.* **82**, 289-301. doi:10.1242/jeb.82.1.289
- Kiceniuk, J. W. and Jones, D. R. (1977). The oxygen transport system in trout (*Salmo gairdneri*) during sustained exercise. *J. Exp. Biol.* **69**, 247-260. doi:10.1242/jeb.69.1.247
- Killen, S. S., Norin, T. and Halsey, L. G. (2017). Do method and species lifestyle affect measures of maximum metabolic rate in fishes? *J. Fish Biol.* **90**, 1037-1046. doi:10.1111/jfb.13195
- Krogh, A. (1937). Osmotic regulation in fresh water fishes by active absorption of chloride ions. *J. Comp. Physiol. A* **24**, 656-666. doi:10.1007/BF00592303
- Malte, H. and Weber, R. E. (1985). A mathematical model for gas exchange in the fish gill based on non-linear blood gas equilibrium curves. *Resp. Physiol.* **62**, 359-374. doi:10.1016/0034-5687(85)90091-X
- Malte, H. and Weber, R. E. (1987). The effect of shape and position of the oxygen equilibrium curve on extraction and ventilation requirement in fishes. *Resp. Physiol.* **70**, 221-228. doi:10.1016/S0034-5687(87)80045-2
- Malte, H. and Weber, R. E. (1989). Gas exchange in fish gills with parallel inhomogeneities. *Resp. Physiol.* **76**, 129-137. doi:10.1016/0034-5687(89)90023-6
- Mandic, M., Todgham, A. E. and Richards, J. G. (2009). Mechanisms and evolution of hypoxia tolerance in fish. *Proc. R. Soc. Lond. B Biol. Sci.* **276**, 735-744.
- Marshall, W. S. and Grosell, M. (2006). Ion transport, osmoregulation and acid-base balance. In *The Physiology of Fishes* (ed. D. Evans and J. B. Claiborne), pp. 177-230. Boca Raton: CRC Press.
- Mazza, F., Cash, J. R. and Soetaert, K. (2014). Solving boundary value problems in the open source software R: package bvpSolve. *Opuscula Math.* **34**, 387-403. doi:10.7494/OpMath.2014.34.2.387
- Nelson, J. S., Grande, T. C. and Wilson, M. V. (2006). *Fishes of the World*, 4th edn. Hoboken, NJ: John Wiley & Sons.
- Nilsson, G. E., Dymowska, A. and Stecyk, J. A. W. (2012). New insights into the plasticity of gill structure. *Resp. Physiol. Neurobiol.* **184**, 214-222. doi:10.1016/j.resp.2012.07.012
- Norin, T. and Clark, T. D. (2016). Measurement and relevance of maximum metabolic rate in fishes. *J. Fish Biol.* **88**, 122-151. doi:10.1111/jfb.12796
- Perry, S. F. (1998). Relationships between branchial chloride cells and gas transfer in freshwater fish. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **119**, 9-16. doi:10.1016/S1095-6433(97)00411-X
- Perry, S. F., Reid, S. G., Wankiewicz, E., Iyer, V. and Gilmour, K. M. (1996). Physiological responses of rainbow trout (*Oncorhynchus mykiss*) to prolonged exposure to soft water. *Physiol. Zool.* **69**, 1419-1441. doi:10.1086/physzool.69.6.30164267
- Postlethwaite, E. K. and McDonald, D. G. (1995). Mechanisms of Na⁺ and Cl⁻ regulation in freshwater-adapted rainbow trout (*Oncorhynchus mykiss*) during exercise and stress. *J. Exp. Biol.* **198**, 295-304. doi:10.1242/jeb.198.2.295
- Rabosky, D. L., Chang, J., Tittle, P. O., Cowman, P. F., Sallan, L., Friedman, M., Kaschner, K., Garilao, C., Near, T. J., Coll, M. et al. (2018). An inverse latitudinal gradient in speciation rate for marine fishes. *Nature* **559**, 392-395. doi:10.1038/s41586-018-0273-1
- Randall, D. J., Daxboeck, C. (1984). Oxygen and carbon dioxide transfer across fish gills. In *Fish Physiology* (ed. W. S. Hoar and D. J. Randall), pp. 263-314. New York: Academic Press.
- Randall, D. J., Baumgarten, D. and Malyusz, M. (1972). The relationship between gas and ion transfer across the gills of fishes. *Comp. Biochem. Physiol. A Comp. Physiol.* **41**, 629-637. doi:10.1016/0300-9629(72)90017-5
- Regan, M. D., Gill, I. and Richards, J. G. (2016). Calorespirometry reveals that goldfish prioritize aerobic metabolism over metabolic rate depression in all but near-anoxic environments. *J. Exp. Biol.* **220**, 564-572. doi:10.1242/jeb.145169
- Regan, M. D., Gill, I. S. and Richards, J. G. (2017). Metabolic depression and the evolution of hypoxia tolerance in threespine stickleback, *Gasterosteus aculeatus*. *Biol. Lett.* **13**, 20170392. doi:10.1098/rsbl.2017.0392
- Regan, M. D., Mandic, M., Dhillon, R. S., Lau, G. Y., Farrell, A. P., Schulte, P. M., Seibel, B. A., Speers-Roesch, B., Utsch, G. R. and Richards, J. G. (2019). Don't throw the fish out with the respirometry water. *J. Exp. Biol.* **222**, jeb200253. doi:10.1242/jeb.200253
- Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217-223. doi:10.1111/j.2041-210X.2011.00169.x
- Rogers, N. J., Urbina, M. A., Reardon, E. E., McKenzie, D. J. and Wilson, R. W. (2016). A new analysis of hypoxia tolerance in fishes using a database of critical oxygen level (P_{crit}). *Conserv Physiol* **4**, cow012. doi:10.1093/conphys/cow012
- Sardella, B. A. and Brauner, C. (2007). The Osmo-respiratory Compromise in Fish. In *Fish Respiration and Environment* (ed. M. N. Fernandes, F. T. Rantin, M. L. Glass and B. G. Kapoor), pp. 147-165. Enfield: Science Publishers.
- Schliep, K. P. (2011). phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592-593. doi:10.1093/bioinformatics/btq706
- Sollid, J., De Angelis, P., Gundersen, K. and Nilsson, G. E. (2003). Hypoxia induces adaptive and reversible gross morphological changes in crucian carp gills. *J. Exp. Biol.* **206**, 3667-3673. doi:10.1242/jeb.00594
- Steinhausen, M. F., Sandblom, E., Eliason, E. J., Verhille, C. and Farrell, A. P. (2008). The effect of acute temperature increases on the cardiorespiratory performance of resting and swimming sockeye salmon (*Oncorhynchus nerka*). *J. Exp. Biol.* **211**, 3915-3926. doi:10.1242/jeb.019281
- Stevens, E. D. and Carey, F. G. (1981). One why of the warmth of warm-bodied fish. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **240**, R151-R155. doi:10.1152/ajpregu.1981.240.3.R151
- Utsch, G. R. and Regan, M. D. (2019). The utility and determination of P_{crit} in fishes. *J. Exp. Biol.* **222**, jeb203646. doi:10.1242/jeb.203646
- Utsch, G. R., Boschung, H. and Ross, M. J. (1978). Metabolism, Critical Oxygen Tension, and Habitat Selection in Darters (*Etheostoma*). *Ecology* **59**, 99-107. doi:10.2307/1936635
- Vega, G. C. and Wiens, J. J. (2012). Why are there so few fish in the sea? *Proc. R. Soc. Lond. B Biol. Sci.* **279**, 2323-2329.
- Wang, T. and Malte, H. (2011). O₂ uptake and transport: the optimal P50. In *Encyclopedia of Fish Physiology: From Genome to Environment* (ed. A. P. Farrell, E. D. Stevens, J. R. Chech and J. G. Richards), pp. 893-898. Elsevier.
- Wegner, N. C., Sepulveda, C. A., Bull, K. B. and Graham, J. B. (2010). Gill morphometrics in relation to gas transfer and ram ventilation in high-energy demand teleosts: Scombrids and billfishes. *J. Morphol.* **271**, 36-49. doi:10.1002/jmor.10777
- Wood, C. M. (2018). The fallacy of the Pcrit – are there more useful alternatives? *J. Exp. Biol.* **221**, jeb163717. doi:10.1242/jeb.163717
- Wood, C. M. and Eom, J. (2021). The osmorepiratory compromise in the fish gill. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **254**, 110895. doi:10.1016/j.cbpa.2021.110895
- Wood, C. M., Iftikar, F. I., Scott, G. R., De Boeck, G., Sloman, K. A., Matey, V., Domingos, F. X. V., Duarte, R. M., Almeida-Val, V. M. and Val, A. L. (2009). Regulation of gill transcellular permeability and renal function during acute hypoxia in the Amazonian oscar (*Astronotus ocellatus*): new angles to the osmorepiratory compromise. *J. Exp. Biol.* **212**, 1949-1964. doi:10.1242/jeb.028464
- Wood, C. M., Ruhr, I. M., Schauer, K. L., Wang, Y., Mager, E. M., McDonald, M. D., Stanton, B. and Grosell, M. (2019). The osmorepiratory compromise in the euryhaline killifish: water regulation during hypoxia. *J. Exp. Biol.* **222**, jeb204818. doi:10.1242/jeb.204818

Table S1. Input parameters used for the calibration of the mathematical model of $\dot{M}O_{2\max}$ using reported data for rainbow trout at $10\pm 1^\circ\text{C}$

Parameters	Input	Units	Reference
Gill surface area	2.4	$\text{cm}^2 \text{g}^{-1}$	(Hughes, 1972)
Gill thickness	4.92	μm	(Greco et al., 1996) average of control and softwater acclimated fish
beta epithelium	1.77×10^{-3}	$\mu\text{mol cm}^{-3} \text{mmHg}^{-1}$	(Boutilier et al., 1984; Jensen, 2017)
Diffusion coefficient (D_{O_2})	4.4×10^{-6}	$\text{cm}^2 \text{s}^{-1}$	Based on (Dejours, 1981; Hills and Hughes, 1970; Piiper and Baumgarten-Schumann, 1968) and adjusted to fit (Kiceniuk and Jones, 1977)
Gill diffusive conductance (G_d)	2.3	$\mu\text{mol mmHg}^{-1} \text{min}^{-1} \text{kg}^{-1}$	Calculated from above
Tissue diffusive conductance (G_{dt})	6.4	$\mu\text{mol mmHg}^{-1} \text{min}^{-1} \text{kg}^{-1}$	Based on (Wang and Malte, 2011) and adjusted to fit (Kiceniuk and Jones, 1977)
O_2 solubility coefficient (α) in water	2.24	$\mu\text{mol L}^{-1} \text{mmHg}^{-1}$	(Boutilier et al., 1984)
α in blood plasma	1.99	$\mu\text{mol L}^{-1} \text{mmHg}^{-1}$	(Boutilier et al., 1984)
Water PO_2	150	mmHg	(Wang and Malte, 2011)
Haemoglobin P_{50}	22.2	mmHg	(Vorger, 1985)
Hill coefficient	2.09		(Vorger, 1985)
Bohr coefficient	-0.82		(Vorger, 1985)
Max. arterial-venous pH shift	-0.12		(Kiceniuk and Jones, 1977)
Haemoglobin concentration	1.05	mM	Based on (Nikinmaa et al., 1981) and adjusted to fit (Kiceniuk and Jones, 1977)
Max. cardiac output (\dot{Q}_{\max})	53	$\text{mL kg}^{-1} \text{min}^{-1}$	(Kiceniuk and Jones, 1977)
Ventilation rate (\dot{V}_w)	1700	$\text{mL kg}^{-1} \text{min}^{-1}$	(Kiceniuk and Jones, 1977)

Table S2.

[Click here to download Table S2](#)

References

- Boutilier, R. G., Heming, T. A. and Iwama, G. K.** (1984). Physicochemical parameters for use in fish respiratory physiology. In *Fish Physiology* (ed. Hoar W. S. and Randall, D. J.), pp. 403–426. New York: Academic Press.
- Dejours, P.** (1981). *Principles of comparative respiratory physiology*. Amsterdam: Elsevier/North-Holland Biomedical Press.
- Greco, A. M., Fenwick, J. C. and Perry, S. F.** (1996). The effects of soft-water acclimation on gill structure in the rainbow trout *Oncorhynchus mykiss*. *Cell Tissue Res.* **285**, 75–82.
- Hills, B. A. and Hughes, G. M.** (1970). A dimensional analysis of oxygen transfer in the fish gill. *Resp. Physiol.* **9**, 126–140.
- Hughes, G. M.** (1972). Morphometrics of fish gills. *Respir Physiol* **14**, 1–25.
- Jensen, P.** (2017). A finite element model of oxygen diffusion in the fish gill.
- Kiceniuk, J. W. and Jones, D. R.** (1977). The oxygen transport system in trout (*Salmo gairdneri*) during sustained exercise. *J. Exp. Biol.* **69**, 247–260.
- Nikinmaa, M., Soivio, A. and Railo, E.** (1981). Blood volume of *Salmo gairdneri*: Influence of ambient temperature. *Comp. Biochem. Physiol., A: Physiol.* **69**, 767–769.
- Piiper, J. and Baumgarten-Schumann, D.** (1968). Effectiveness of O₂ and CO₂ exchange in the gills of the dogfish (*Scyliorhinus stellaris*). *Resp. Physiol.* **5**, 338–349.
- Vorger, P.** (1985). The Bohr effect of the blood in rainbow trout (*Salmo gairdneri*). A comparative study with human blood, using precise oxygen equilibrium curves and the Adair model. *Comp. Biochem. Physiol.* **82A**, 915–924.
- Wang, T. and Malte, H.** (2011). O₂ uptake and transport: the optimal P50. In *Encyclopedia of Fish Physiology: From Genome to Environment* (ed. Farrell, A. P.), Stevens, E. D.), Chech, J. R.), and Richards, J. G.), pp. 893–898. Elsevier.