

SHORT COMMUNICATION

Weanling gut microbiota composition of a mouse model selectively bred for high voluntary wheel-running behavior

Monica P. McNamara¹, Emily M. Venable², Marcell D. Cadney³, Alberto A. Castro¹, Margaret P. Schmill^{4,5}, Lawrence Kazzazi¹, Rachel N. Carmody² and Theodore Garland, Jr^{1,*}

ABSTRACT

We compared the fecal microbial community composition and diversity of four replicate lines of mice selectively bred for high wheel-running activity over 81 generations (HR lines) and four non-selected control lines. We performed 16S rRNA gene sequencing on fecal samples taken 24 h after weaning, identifying a total of 2074 bacterial operational taxonomic units. HR and control mice did not significantly differ for measures of alpha diversity, but HR mice had a higher relative abundance of the family Clostridiaceae. These results differ from a study of rats, where a line bred for high forced-treadmill endurance and that also ran more on wheels had lower relative abundance of Clostridiaceae, as compared with a line bred for low endurance that ran less on wheels. Within the HR and control groups, replicate lines had unique microbiomes based on unweighted UniFrac beta diversity, indicating random genetic drift and/or multiple adaptive responses to selection.

KEY WORDS: 16S rRNA, Behavior, Exercise, Gut microbiota, Microbiome, Selection experiment

INTRODUCTION

The mammalian gut microbiome plays an essential role in host biology, including immune system function, energy extraction and protection from pathogens (Gilbert et al., 2018; Kohl and Carey, 2016). Within an individual, the microbiome is shaped by both host genetics and environmental factors (Benson et al., 2010; Carmody et al., 2015; Tamburini et al., 2016). Both acute and chronic voluntary exercise can affect the gut microbiome in rodents and humans (Campbell and Wisniewski, 2017; Mailing et al., 2019; Mohr et al., 2020). For example, adult rats given wheels for 5 weeks had more butyrate-producing bacteria in their ceca compared with sedentary controls, and an increased amount of cecal n-butyrate, a short-chain fatty acid essential for intestinal epithelial cell health (Matsumoto et al., 2008). In human marathon runners, fecal samples taken before and after a run demonstrated rapid changes to their microbiome (Scheiman et al., 2019).

Conversely, the gut microbiome can affect both exercise ability and motivation to engage in exercise (Dohnalová et al., 2022). Among adult male C57BL/6N mice treated with antibiotics then

gavaged with cecal microbial communities harvested from sedentary versus exercised mice, recipients of microbiome from exercised animals ran more on wheels compared with recipients of the microbiome from sedentary mice (Oyanagi et al., 2018). With respect to exercise ability, 5 h after mice were gavaged with a lactate-metabolizing strain of *Veillonella atypica* cultured from post-marathon athletes, they had significantly longer run times to 'exhaustion' versus those gavaged with *Lactobacillus bulgaricus*, a common bacterial symbiont that cannot metabolize lactate (Scheiman et al., 2019). However, the method used to motivate mice to run was not stated and *L. bulgaricus* can synthesize lactate, thus complicating interpretation (Carmody and Baggish, 2019).

We compared the weanling gut microbiota of mice that differ in both exercise ability and motivation: four replicate high runner (HR) lines selectively bred for high voluntary wheel-running behavior over 81 generations, and 4 non-selected control lines (Swallow et al., 1998). The selection criterion is revolutions on days 5 and 6 of a 6-day access period between 6–8 weeks of age. HR and control mice differ in several ways that potentially correlate with unique gut microbial communities. First, HR mice run ~2.5- to 3-fold more revolutions per day (Careau et al., 2013; Copes et al., 2015) and also have higher levels of activity when housed individually without wheels (Copes et al., 2015; Malisch et al., 2009). High activity is accompanied by elevated food consumption relative to body size (Copes et al., 2015; Hiramatsu and Garland, 2018; Swallow et al., 2001), which could directly affect gut microbial community composition via changes in luminal resources (Alcock et al., 2014). HR and control mice have not been found to differ in small or large intestine mass or length, thus suggesting a faster transit time in HR mice (Kelly et al., 2017). HR mice also have higher body temperatures when active (Rhodes et al., 2000), altered hormone levels (Garland et al., 2016; Malisch et al., 2009), and tend to be smaller at weaning (Cadney et al., 2021; McNamara et al., 2022; Swallow et al., 1999).

Previously, we reported that adult male HR and control mice differ in gut microbial community composition, regardless of diet or exercise manipulation during early life (McNamara et al., 2021). In addition, when the microbiome is reduced with oral antibiotics, running behavior of control mice is largely unaffected, whereas HR mice run significantly fewer revolutions per day (McNamara et al., 2022), suggesting the gut microbiome is an important component of the high wheel-running phenotype.

Unique gut microbial phenotypes in HR mice could result from acute effects of differences in HR and control traits and/or changes in the selective regime experienced by the microbiota. Our two previous studies cannot distinguish between these possibilities because adults will have experienced many weeks of differences in physical activity, food consumption, and other physiological differences that could acutely affect the microbiome. Although physical activity of HR and control pups prior to weaning has not been quantified, related aspects

¹Department of Evolution, Ecology, and Organismal Biology, University of California, Riverside, CA 91521, USA. ²Department of Human Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA. ³Neuroscience Research Institute, University of California, Santa Barbara, CA 93106, USA. ⁴Neuroscience Graduate Program, University of California, Riverside, CA 92521, USA. ⁵Medpace, 717th St, Suite 500, Denver, CO 80202, USA.

*Author for correspondence (tgarland@ucr.edu)

DOI: 10.1242/jeb.245081

of pre-weaning behavior do not significantly differ, including first day of eye opening, moving, and feeding on solid food, and locomotor play (see fig. 2A–C in Hiramatsu et al., 2017, 10–20 days old; N. N. D. Whitehead and T.G., unpublished results; 15 days old). Therefore, we compared the gut microbiota at weaning to better discriminate whether microbial signatures are specific to the HR lines or a secondary response to acute phenotypes, such as wheel running, differences in physiology, etc.

MATERIALS AND METHODS

All experiments and methods were approved by the Institutional Animal Use and Care Committee of the University of California, Riverside.

Experimental animals

Females were sampled from generation 81 of the high runner selection experiment, which began in 1993 with a population of 224 outbred Hsd:ICR mice (Swallow et al., 1998). Briefly, mice are weaned at 21 days of age and housed 4 per cage separated by line and sex. At ~6–8 weeks of age, they are housed individually for 6 days in cages attached to a 1.12 m circumference wheel. Each generation, within each of the 4 replicate HR lines (lab designations 3,6,7,8), the highest-running male and female from within each of 10 families are chosen as breeders, based on the average revolutions on days 5 and 6. For the 4 replicate control lines (1,2,4,5), one male and one female are taken from each family without regard to running. Mice are paired within their line, and no sibling matings are allowed. Following weaning, mice are provided with Standard Laboratory Rodent Diet from Harlan Teklad (Envigo) (W-8604), which contains 24.3% kJ from protein, 4% kJ from fat and 40.2% kJ from carbohydrate. Pregnant dams are given Harlan Teklad (Envigo) Lab Mouse Breeder Diet [S-2335] 7004 through weaning.

In the present study, mice were weaned at 21 days of age and housed individually for 24 h prior to fecal sampling. Mice were checked for signs of distress (e.g. hunched posture) while individually housed and no such signs were observed. Each was from a different litter, with six exceptions. From an initial sample of 100, 5 mice were excluded owing to cage issues or premature death post-weaning, resulting in $N=95$ [11–12 from each line, except line 6 (polymorphic for the mini-muscle phenotype, described below), which had 14].

Fecal sampling

Fecal samples were collected from mice 20–24 h after weaning. Mice were grasped at the nape until defecation occurred into a sterile tube, which was immediately placed on dry ice and stored at -80°C . Fecal samples were shipped on dry ice to the Nutritional and Microbial Ecology Lab at Harvard University and stored at -80°C until processing.

DNA extraction

We used an established 16S ribosomal RNA (rRNA) gene sequencing pipeline to assess gut microbial community composition in each sample (Carmody et al., 2015, 2019). Briefly, we isolated DNA from fecal samples using the Qiagen PowerSoil DNA Isolation Kit (cat. no. 12888). Next, we PCR amplified the hypervariable V4 region of the 16S rRNA gene using custom barcoded 515F and 806R primers, which target conserved regions around the V4 region (Caporaso et al., 2011, 2012; Shahi et al., 2017). PCR amplification was performed in triplicate using the following reaction mix: 11 μl nuclease-free H_2O , 1 μl 25 mmol l^{-1} MgCl_2 , 10 μl Quantabio 5Prime Hot MasterMix

(cat. no. 2200410), 2 μl primers (1 μl of forward primer and 1 μl of reverse primer) and 1 μl template DNA. We included a negative control reaction per sample to ensure that primers and reagents were not contaminated. PCR was performed using BioRad T100 thermocyclers and the following protocol: 94°C for 3 min; 35 cycles of 94°C for 45 s, 50°C for 30 s and 72°C for 90 s; and 10 min at 72°C . PCR amplicons were checked by running recombinant triplicate reactions, negative controls and a 100 bp DNA ladder on a 1.5% agarose gel in an electrophoresis chamber. Amplicons were purified using Agencourt AMPure XP solution (cat. no. A63880) and resuspended in 40 μl of $1\times$ TE buffer. Cleaned amplicons were quantified using the Quant-iT PicoGreen dsDNA Assay Kit (cat. no. P11495), with fluorescence measured with a Spectramax Gemini XS Plate Reader set to 480 nm excitation and 520 nm emission. Cleaned amplicons were pooled at sample-specific volumes to obtain 80 ng DNA per sample. We purified 100 μl of the pooled solution using the Qiaquick MinElute kit (cat. no. 28004). The eluted DNA was then gel-purified by 1.5% agarose gel electrophoresis. Band size was compared against a 100 bp DNA ladder, and the targeted 381 bp band was cut from the gel with a sterile razor and resuspended using the Qiaquick PCR Purification kit (cat. no. 28104). The pool was diluted to 10 nmol l^{-1} and submitted for sequencing on one lane of an Illumina HiSeq rapid flow cell (1×150 bp) at the Harvard Bauer Core.

Analysis of 16S rRNA gene sequences

Raw sequences were processed using the Quantitative insights into microbial ecology (QIIME) package v. 1.8 (Caporaso et al., 2010). After quality filtering, we obtained a mean sequencing depth of $123,557\pm 50,871$ (s.e.m.) reads per sample. Operational taxonomic units (OTUs) were picked at 97% similarity (Caporaso et al., 2010). Bacterial relative abundances at taxonomic levels from phylum to genus were generated using the summarize_taxa.py script. Prior to alpha diversity analysis, we rarefied the dataset at 25,000 reads per sample. Alpha diversity (Shannon diversity index, Chao1, unique OTUs and Faith's phylogenetic diversity) was analyzed using the alpha_diversity.py script. Prior to beta diversity analysis, we subsampled the dataset at 33,600 reads and used the beta_diversity_through_plots.py script to generate Bray–Curtis, unweighted UniFrac and weighted UniFrac distance matrices and associated principal coordinates. The bacterial 16S rRNA sequences have been deposited in the NCBI Sequence Read Archive as BioProject PRJNA911624 (<https://www.ncbi.nlm.nih.gov/sra/PRJNA911624>).

Statistical analyses

We used mixed models to analyze alpha diversity metrics and taxonomic relative abundances in SAS 9.4 Procedure Mixed (SAS Institute, Cary, NC, USA). Our models tested the effects of linetype (HR versus control lines) against the variance among replicate lines, nested as a random effect within linetype, with 1 and 6 d.f. We also tested the effect of mini-muscle status, a phenotype currently present in two of the four HR lines that is caused by a single base pair change in the myosin heavy polypeptide 4 gene *Myh4* (Kelly et al., 2013). The mini-muscle phenotype is characterized by a ~50% reduction in hindlimb muscle mass, larger internal organs and a variety of other differences compared with mice with normal muscle (e.g. see Garland et al., 2002; Swallow et al., 2009; Wallace and Garland, 2016). In the present study, 26 of 95 mice had the phenotype (all 12 in HR line 3, and 6 of 14 in HR line 6).

Beta diversity of the gut microbiome was assessed by calculating unweighted UniFrac, weighted UniFrac and Bray–Curtis distance

matrices and performing principal coordinate analyses (PCoA) to visualize the microbial community clustering based on distance. We used the `adonis` function within the `vegan` package in R (<https://CRAN.R-project.org/package=vegan>) to perform permutational analysis of variance (PERMANOVA) to determine significant clustering within the dataset (Anderson, 2001, 2017). We permuted the distance matrix over line type and mini-muscle status 999 times. We also permuted the distance matrix over line 999 times for separate analyses of the 4 replicate HR and 4 replicate control lines. Replicate line was not treated as a nested random effect because this feature is not available in the `vegan` package. One sample was removed from the unweighted UniFrac distance matrix based on its appearance as a strong outlier in the initial PCoA.

Bacterial relative abundances were log or arcsine square-root transformed to improve normality of residuals (Brown et al., 2020; Kohl et al., 2016). To focus on taxa present widely in our population, we limited analyses to taxa found with at least 50% prevalence among samples. We then used a targeted approach to test for differentially abundant bacteria between the HR and control line types. Specifically, we analyzed taxa previously associated with exercise in rodents (Campbell et al., 2016; Clarke et al., 2014; Codella et al., 2018; Hughes, 2020; Mach and Fuster-Botella, 2017; Munukka et al., 2018; Queipo-Ortuño et al., 2013), including the phyla: Proteobacteria, Bacteroidetes, Firmicutes, Tenericutes and Actinobacteria; families: Rikenellaceae, Lactobacillaceae and Clostridiaceae; genera: *Clostridium* (genus within the family Clostridiaceae), *Lactobacillus*, *Bifidobacterium*, *Akkermansia* and *Oscillospira*. (Note that the genus *Clostridium* presently includes OTUs that are nested within multiple different families, reflecting a taxonomy that is not fully phylogenetic.) We were also interested in the genus *Veillonella* for its prior association with endurance exercise in humans (Scheiman et al., 2019), but *Veillonella* was not present. Statistical significance was judged at the $P=0.05$ level. For completeness, analyses of additional taxa with at least 50% prevalence among samples are presented in Table S1. A measure of effect size (Pearson's r) was calculated for all main and interactive effects for bacterial relative abundance and alpha diversity (Sullivan and Feinn, 2012).

RESULTS AND DISCUSSION

Alpha diversity of the weanling gut microbiome

Based on mixed models, the average number of unique OTUs per mouse and other alpha diversity metrics did not statistically differ between weanling HR and control mice (unique OTUs, $F_{1,6}=0.01$, $P=0.7611$; Shannon index, $F_{1,6}=0.28$, $P=0.6131$; Chao1, $F_{1,6}=0$, $P=0.9884$; Faith's phylogenetic diversity, $F_{1,6}=0.09$, $P=0.7777$) or between mini-muscle and normal muscle mice (unique OTUs, $F_{1,86}=0.52$, $P=0.4727$; Shannon index, $F_{1,86}=0.0$, $P=0.9621$; Chao1, $F_{1,86}=0.81$, $P=0.3704$; Faith's phylogenetic diversity, $F_{1,86}=0.5$, $P=0.4797$) (Table S1). Within the two line types, the 4 individual replicate control lines differed in the number of unique OTUs ($F_{3,41}=3.14$, $P=0.0353$) and Faith's phylogenetic diversity metric ($F_{3,41}=3.06$, $P=0.0386$), but not Chao1 ($F_{3,41}=2.53$, $P=0.0707$) or Shannon index ($F_{3,41}=1.28$, $P=0.2954$). The 4 individual replicate HR lines did not differ in the number of unique OTUs ($F_{3,45}=0.93$, $P=0.4356$), Faith's phylogenetic diversity ($F_{3,45}=0.49$, $P=0.6938$), Chao 1 ($F_{3,45}=0.96$, $P=0.4220$), or Shannon index ($F_{3,45}=0.62$, $P=0.6081$).

The gut microbiota typically becomes more diverse with age in both humans and rodents (Koenig et al., 2011; Schloss et al., 2012; Yatsunen et al., 2012). Consistent with this pattern, our weanling mice had fewer average OTUs ($N=373$) compared with our previous

study of adult HR and control mice ($N=430$), although it is important to note that these numbers cannot be strictly compared because they were based on different marker genes (McNamara et al., 2021).

Two prior rodent studies examined the gut microbiome in response to selective breeding for aspects of exercise capacity, although not in weanlings. Two lines of rats bred for either high or low endurance capacity during forced treadmill exercise did not differ in alpha diversity metrics at either 7 or 40 weeks of age (Pekkala et al., 2017; weaning occurs at 4 weeks of age). Four replicate lines of bank voles bred for oxygen consumption during swimming exercise at a temperature below the thermal neutral zone also did not differ in alpha diversity as compared with four control lines (mean age: 166 days) (Kohl et al., 2016).

Dominant phyla of the weanling gut microbiome

Across the entire sample of 95 mice, 2074 OTUs were identified, representing 11 phyla, 21 classes, 38 orders, 108 families and 218 genera of microbes. Typical for mice, composition was dominated by the phyla Firmicutes (48.4±14%; mean±s.d.) and Bacteroidetes (37.5±17.1%) (Fig. S1). Based on mixed models, neither line type (HR vs control) nor mini-muscle status statistically affected the relative abundance of the phyla Bacteroidetes, Firmicutes, Proteobacteria, Tenericutes or Actinobacteria (Table S1).

Previously (and as consistently observed in murine studies), we reported that the adult gut microbiota in these mice was dominated by Bacteroidetes (~68%), with Firmicutes (~28%) being second most abundant (McNamara et al., 2021). For the present sample from weanlings, the phylum Proteobacteria constitutes a much larger portion of the weanling (11.8±8.7%) compared to the adult (~1%) gut microbiome. Although our weanling and adult datasets were based on different sequencing methods, these trends are consistent with previous studies in lab rodents showing that the weanling gut microbiome is initially dominated by Firmicutes (Cox et al., 2014; Pantoja-Feliciano et al., 2013) followed by a shift towards Bacteroidetes (Cox et al., 2014; Nagpal et al., 2018).

Targeted taxonomic comparisons

HR mice had significantly higher relative abundance of family Clostridiaceae compared with control mice (least squares means±standard errors: HR=0.0698±0.0082; control=0.0279±0.0117) ($F_{1,6}=10.54$, $P=0.0175$, Fig. 1), with no statistical difference for the families Rikenellaceae and Lactobacillaceae or genera *Clostridium*, *Bifidobacterium*, *Lactobacillus*, *Akkermansia* and *Oscillospira* (Table S1). In our previous study of adults, HR mice also had a higher relative abundance of the family Clostridiaceae compared with controls, although the difference was not statistically significant ($P=0.0750$: Table S2 in McNamara et al., 2021).

The HCR (high endurance capacity) and LCR (low endurance capacity) lines of rats (see above) also differed in gut microbiome community composition (Liu et al., 2015; Pekkala et al., 2017). HCR rats have higher maximal aerobic capacity ($\dot{V}_{O_{2,max}}$) and higher voluntary wheel running compared with the LCR line (Karvinen et al., 2015; Park et al., 2016; Swallow et al., 2010), paralleling the elevated endurance capacity (Meek et al., 2009) and $\dot{V}_{O_{2,max}}$ (Cadney et al., 2021) of HR mice. In addition, HCR are smaller than LCR rats (Pekkala et al., 2017; Wisløff et al., 2005) and HR are smaller than control mice (Dumke et al., 2001; Kelly et al., 2017). Thus, we anticipated some similar patterns of differentiation in the gut microbiome. However, whereas adult HCR rats had significantly lower relative abundance of Clostridiaceae compared with LCR (Liu et al., 2015), HR mice had a higher relative




Fig. 1. Relative abundance of the family Clostridiaceae in the mouse microbiome. Based on mixed models simultaneously comparing high running (HR) and control lines, as well as mini-muscle and normal muscle mice, HR mice had significantly higher relative abundance of Clostridiaceae compared with control mice (ANOVA, $F_{1,6}=10.54$, $P=0.0175$, $r=0.7983$), with no effect of mini-muscle status ($F_{1,86}=3.34$, $P=0.0712$, $r=0.1934$). Shown are least squares means \pm standard errors for arcsine square-root transformed values. Pearson's r is a measure of effect size. $N=90$ mice in total.

abundance of Clostridiaceae compared with the non-selected control lines for both adults (McNamara et al., 2021) and weanlings (present study). One obvious explanation for this difference is that they are different species, and rats and mice are known to differ in various ways with respect to exercise physiology and responses to exercise training (e.g. see Dumke et al., 2001; Kowalski and Bruce, 2014). Future studies in rodents bred for exercise-related traits, including our own HR mice, should examine the gut microbiome community across various timepoints and generations to illuminate if the abundance of Clostridiaceae changes during development and/or across generations.

Beta diversity of the weanling gut microbiome

Principal coordinate analysis based on unweighted UniFrac distances did not indicate separation between HR and control mice along the first two principal coordinate axes (Fig. 2A: although together they account for only 14% of the total variance in the data). PERMANOVA of the unweighted UniFrac distance matrix also indicated no significant differentiation between HR and control mice ($R^2=0.013$, $P=0.080$), nor did reanalysis based on Bray–Curtis or weighted UniFrac distances (Fig. S2). This lack of differentiation contrasts with our previous results for adults, where HR and control mice clustered separately, regardless of diet and/or exercise treatment during early life, based on unweighted UniFrac distances (PERMANOVA $P=0.009$; McNamara et al., 2021).

Mini-muscle and normal muscle mice separated somewhat on the third PCoA axis for unweighted UniFrac distances, and PERMANOVA indicates statistically significant separation for unweighted UniFrac (Fig. 2B; $R^2=0.014$, $P=0.047$). Thus, weanling mice have distinct bacterial communities based on mini-muscle status. Mini-muscle individuals differ from normal muscle individuals in several ways that might affect the microbiome fairly directly, including higher mass-adjusted food consumption (Meek et al., 2014), larger stomachs, a trend for longer small intestines, and a trend for heavier caecum dry masses (Kelly et al., 2017). In addition, their smaller muscles but larger internal organs (heart, soleus, spleen, liver, kidney, lung; Kelly et al., 2017) could impact




Fig. 2. Principal coordinate analysis of unweighted UniFrac distances between the 16S rRNA gene sequencing-based profiles of weanling fecal microbiomes. PCoA in relation to (A) line type and (B) mini-muscle status (one outlier removed). PERMANOVAs indicate significant separation based on mini-muscle status mice ($R^2=0.014$, $P=0.047$), but not between line types ($R^2=0.013$, $P=0.080$).

energetic demands and also the microbiome. However, no significant separation was detected when measured by Bray–Curtis or weighted UniFrac distances, suggesting that the effects of mini muscle status may be limited to differences in phylogenetic representation in the microbiome and not differences in relative abundance (Fig. S2).

Finally, we considered potential separation among the four replicate HR lines and among the four replicate control lines (Fig. 3). Differentiation was significant within both line types based on PERMANOVAs on unweighted UniFrac distances (HR lines

with one outlier removed: $R^2=0.028$, $P=0.038$; control lines: $R^2=0.037$, $P=0.006$). Based on Bray–Curtis distances, separation among control lines was statistically significant, but not among HR lines, and weighted UniFrac distances indicated no separation among the replicate lines for either linetype (Fig. S3). Jointly, these data suggest that drift has been a stronger force on the gut microbiome in control lines than in HR lines, where, as expected, selection has been more important in HR lines.

When measured as adults (37 weeks of age after 11 weeks of individual housing with or without wheel access), the HCR and

LCR lines of rats mentioned above clustered separately, based on unweighted and weighted UniFrac distances (Liu et al., 2015). In the bank vole selection experiment, a total of 16 lines were included: 4 bred for aerobic capacity, 4 for ability to maintain body weight when fed a low-quality diet for 4 days, and 4 for predatory behavior towards crickets. As adults, none of the sets of selected lines had gut microbial profiles that differed from those of the control lines using weighted UniFrac distances (Kohl et al., 2016). However, unweighted UniFrac distances indicated the herbivorous lines differed in gut microbial profiles compared with the other groups.

Concluding remarks and future directions

Previously, we reported that selective breeding for voluntary wheel running resulted in higher relative abundance of Clostridiaceae ($P=0.0750$) among adults in 4 replicate HR lines as compared with their 4 non-selected control lines, in addition to substantial differentiation in beta diversity. In our previous study, we sequenced the internal transcribed spacer region. Here, using 16S rRNA sequencing, we show that the HR lines also have higher relative abundance of Clostridiaceae at weaning. Notably, our study only examines how gut microbial community structure differs between mouse lines at weaning and does not examine function. Future studies should test for differences in levels of physical activity between the HR and control lines before weaning, as that could contribute to microbiome differences. In addition, transplantation of the HR microbiome into control mice, and vice versa, following ablation via antibiotics (McNamara et al., 2022), or into germ-free animals, would shed further light on the contribution of the gut microbiome to increased aerobically supported activity among HR mice. Reanalysis of data from previous studies in other systems could indicate if Clostridiaceae may be associated with exercise behavior and/or physiology.

Acknowledgements

We thank the UCR vivarium staff for help maintaining the mouse colony.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: M.P.M., R.N.C., T.G.; Methodology: M.P.M., R.N.C., T.G.; Software: R.N.C.; Validation: R.N.C., T.G.; Formal analysis: M.P.M., E.M.V., R.N.C., T.G.; Investigation: M.P.M., E.M.V., M.D.C., A.A.C., M.P.S., L.K., R.N.C., T.G.; Resources: R.N.C., T.G.; Data curation: M.P.M., E.M.V., R.N.C., T.G.; Writing - original draft: M.P.M., E.M.V., R.N.C., T.G.; Writing - review & editing: M.P.M., E.M.V., M.D.C., A.A.C., M.P.S., L.K., R.N.C., T.G.; Visualization: M.P.M., E.M.V., R.N.C., T.G.; Supervision: R.N.C., T.G.; Project administration: T.G.; Funding acquisition: R.N.C., T.G.

Funding

This work was supported by U.S. National Science Foundation grant IOS-2038528 to T.G. and N. C. Holt and by a grant from the Harvard University Dean's Competitive Fund for Promising Scholarship to R.N.C.

Data availability

The bacterial 16S rRNA sequences are available in the NCBI Sequence Read Archive as BioProject PRJNA911624 (<https://www.ncbi.nlm.nih.gov/sra/PRJNA911624>).

References

- Alcock, J., Maley, C. C. and Aktipis, C. A. (2014). Is eating behavior manipulated by the gastrointestinal microbiota? Evolutionary pressures and potential mechanisms. *BioEssays* **36**, 940–949. doi:10.1002/bies.201400071
- Anderson, M. J. (2017). Permutational multivariate analysis of variance (PERMANOVA). In *Wiley StatsRef: Statistics Reference Online* (ed. N. Balakrishnan, T. Colton, B. Everitt, W. Piegorisch, F. Ruggeri and J. L. Teugels), pp. 1–15. Wiley.




Fig. 3. Principal coordinate analysis of unweighted UniFrac distances between the 16S rRNA gene sequencing-based profiles. PCoA was performed for each line type separately on (A) 4 HR lines (3,6,7,8) and (B) 4 control lines (1,2,4,5). Values are means \pm s.e. for scores on PCoA axes. PERMANOVAs indicate significant separation among the 4 HR lines and among the 4 C lines (HR lines: $R^2=0.028$, $P=0.038$; control lines: $R^2=0.037$, $P=0.006$).

- Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral. Ecol.* **26**, 32–46. doi:10.1046/j.1442-9993.2001.01070.x
- Anderson, M. J. (2017). Permutational Multivariate Analysis of Variance (PERMANOVA). In *Wiley StatsRef: Statistics Reference Online*, pp. 1–15. John Wiley & Sons, Ltd.
- Benson, A. K., Kelly, S. A., Legge, R., Ma, F., Low, S. J., Kim, J., Zhang, M., Oh, P. L., Nehrenberg, D., Hua, K. et al. (2010). Individuality in gut microbiota composition is a complex polygenic trait shaped by multiple environmental and host genetic factors. *Proc. Natl. Acad. Sci. USA* **107**, 18933–18938. doi:10.1073/pnas.1007028107
- Brown, T. A., Tashiro, H., Kasahara, D. I., Cho, Y. and Shore, S. A. (2020). Early life microbiome perturbation alters pulmonary responses to ozone in male mice. *Physiol. Rep.* **8**, e14290. doi:10.14814/phy2.14290
- Cadney, M. D., Hiramatsu, L., Thompson, Z., Zhao, M., Kay, J. C., Singleton, J. M., de Albuquerque, R. L., Schmill, M. P., Saltzman, W. and Garland, T., Jr. (2021). Effects of early-life exposure to Western diet and voluntary exercise on adult activity levels, exercise physiology, and associated traits in selectively bred High Runner mice. *Physiol. Behav.* **234**, 113389. doi:10.1016/j.physbeh.2021.113389
- Campbell, S. C. and Wisniewski, P. J. (2017). Exercise is a novel promoter of intestinal health and microbial diversity. *Exerc. Sport Sci. Rev.* **45**, 41–47. doi:10.1249/JES.0000000000000096
- Campbell, S. C., Wisniewski, P. J., Noji, M., McGuinness, L. R., Häggblom, M. M., Lightfoot, S. A., Joseph, L. B. and Kerkhof, L. J. (2016). The effect of diet and exercise on intestinal integrity and microbial diversity in mice. *PLoS ONE* **11**, e0150502. doi:10.1371/journal.pone.0150502
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Peña, A. G., Goodrich, J. K., Gordon, J. I. et al. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* **7**, 335–336. doi:10.1038/nmeth.f.303
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., Fierer, N. and Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. USA* **108**, 4516–4522. doi:10.1073/pnas.1000080107
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M., Betley, J., Fraser, L., Bauer, M. et al. (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J.* **6**, 1621–1624. doi:10.1038/ismej.2012.8
- Careau, V., Wolak, M. E., Carter, P. A. and Garland, T., Jr. (2013). Limits to behavioral evolution: the quantitative genetics of a complex trait under directional selection: quantitative genetics of a selection limit. *Evolution* **67**, 3102–3119. doi:10.1111/evo.12200
- Cardo, R. N. and Baggish, A. L. (2019). Working out the bugs: microbial modulation of athletic performance. *Nat. Metab.* **1**, 658–659. doi:10.1038/s42255-019-0092-1
- Cardo, R. N., Gerber, G. K., Luevano, J. M., Gatti, D. M., Somes, L., Svenson, K. L. and Turnbaugh, P. J. (2015). Diet dominates host genotype in shaping the murine gut microbiota. *Cell Host Microbe* **17**, 72–84. doi:10.1016/j.chom.2014.11.010
- Cardo, R. N., Bisanz, J. E., Bowen, B. P., Maurice, C. F., Lyalina, S., Louie, K. B., Treen, D., Chadaideh, K. S., Maini Rekhal, V., Bess, E. N. et al. (2019). Cooking shapes the structure and function of the gut microbiome. *Nat. Microbiol.* **4**, 2052–2063. doi:10.1038/s41564-019-0569-4
- Clarke, S. F., Murphy, E. F., O'Sullivan, O., Lucey, A. J., Humphreys, M., Hogan, A., Hayes, P., O'Reilly, M., Jeffery, I. B., Wood-Martin, R. et al. (2014). Exercise and associated dietary extremes impact on gut microbial diversity. *Gut* **63**, 1913–1920. doi:10.1136/gutjnl-2013-306541
- Codella, R., Luzzi, L. and Terruzzi, I. (2018). Exercise has the guts: how physical activity may positively modulate gut microbiota in chronic and immune-based diseases. *Dig. Liver Dis.* **50**, 331–341. doi:10.1016/j.dld.2017.11.016
- Copes, L. E., Schutz, H., Dlugosz, E. M., Acosta, W., Chappell, M. A. and Garland, T., Jr. (2015). Effects of voluntary exercise on spontaneous physical activity and food consumption in mice: Results from an artificial selection experiment. *Physiol. Behav.* **149**, 86–94. doi:10.1016/j.physbeh.2015.05.025
- Cox, L. M., Yamanishi, S., Sohn, J., Alekseyenko, A. V., Leung, J. M., Cho, I., Kim, S. G., Li, H., Gao, Z., Mahana, D. et al. (2014). Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell* **158**, 705–721. doi:10.1016/j.cell.2014.05.052
- Dohnalová, L., Lundgren, P., Carty, J. R. E., Goldstein, N., Wenski, S. L., Nanudorn, P., Thiengmag, S., Huang, K.-P., Litichevskiy, L., Descamps, H. C. et al. (2022). A microbiome-dependent gut–brain pathway regulates motivation for exercise. *Nature* **612**, 739–747. doi:10.1038/s41586-022-05525-z
- Dumke, C. L., Rhodes, J. S., Garland, T., Jr, Maslowski, E., Swallow, J. G., Wetter, A. C. and Cartee, G. D. (2001). Genetic selection of mice for high voluntary wheel running: effect on skeletal muscle glucose uptake. *J. Appl. Physiol.* **91**, 1289–1297. doi:10.1152/jappl.2001.91.3.1289
- Garland, T., Jr, Morgan, M. T., Swallow, J. G., Rhodes, J. S., Girard, I., Belter, J. G. and Carter, P. A. (2002). Evolution of a small-muscle polymorphism in lines of house mice selected for high activity levels. *Evolution* **56**, 1267–1275. doi:10.1111/j.0014-3820.2002.tb01437.x
- Garland, T., Jr, Zhao, M. and Saltzman, W. (2016). Hormones and the evolution of complex traits: insights from artificial selection on behavior. *Integr. Comp. Biol.* **56**, 207–224. doi:10.1093/icb/icw040
- Gilbert, J. A., Blaser, M. J., Caporaso, J. G., Jansson, J. K., Lynch, S. V. and Knight, R. (2018). Current understanding of the human microbiome. *Nat. Med.* **24**, 392–400. doi:10.1038/nm.4517
- Hiramatsu, L. and Garland, T., Jr. (2018). Mice selectively bred for high voluntary wheel-running behavior conserve more fat despite increased exercise. *Physiol. Behav.* **194**, 1–8. doi:10.1016/j.physbeh.2018.04.010
- Hiramatsu, L., Kay, J., Thompson, Z., Singleton, J., Claghorn, G., Albuquerque, R. L., Ho, B., Ho, B., Sanchez, G. and Garland, T., Jr. (2017). Maternal exposure to Western diet affects adult body composition and voluntary wheel running in a genotype-specific manner in mice. *Physiol. Behav.* **179**, 235–245. doi:10.1016/j.physbeh.2017.06.008
- Hughes, R. L. (2020). A review of the role of the gut microbiome in personalized sports nutrition. *Front. Nutr.* **6**, 191. doi:10.3389/fnut.2019.00191
- Karvonen, S., Waller, K., Silvennoinen, M., Koch, L. G., Britton, S. L., Kaprio, J., Kainulainen, H. and Kujala, U. M. (2015). Physical activity in adulthood: genes and mortality. *Sci. Rep.* **5**, 18259. doi:10.1038/srep18259
- Kelly, S. A., Bell, T. A., Selitsky, S. R., Buus, R. J., Hua, K., Weinstock, G. M., Garland, T., Jr, Pardo-Manuel de Villena, F. and Pomp, D. (2013). A novel intronic single nucleotide polymorphism in the *myosin heavy polypeptide 4* gene is responsible for the mini-muscle phenotype characterized by major reduction in hind-limb muscle mass in mice. *Genetics* **195**, 1385–1395. doi:10.1534/genetics.113.154476
- Kelly, S. A., Gomes, F. R., Kolb, E. M., Malisch, J. L. and Garland, T., Jr. (2017). Effects of activity, genetic selection and their interaction on muscle metabolic capacities and organ masses in mice. *J. Exp. Biol.* **220**, 1038–1047. doi:10.1242/jeb.148759
- Koenig, J. E., Spor, A., Scalfone, N., Fricker, A. D., Stombaugh, J., Knight, R., Angenent, L. T. and Ley, R. E. (2011). Succession of microbial consortia in the developing infant gut microbiome. *Proc. Natl. Acad. Sci. USA* **108** Suppl. 1, 4578–4585. doi:10.1073/pnas.1000081107
- Kohl, K. D. and Carey, H. V. (2016). A place for host-microbe symbiosis in the comparative physiologist's toolbox. *J. Exp. Biol.* **219**, 3496–3504. doi:10.1242/jeb.136325
- Kohl, K. D., Sadowska, E. T., Rudolf, A. M., Dearing, M. D. and Koteja, P. (2016). Experimental evolution on a wild mammal species results in modifications of gut microbial communities. *Front. Microbiol.* **7**, 634. doi:10.3389/fmicb.2016.00634
- Kowalski, G. M. and Bruce, C. R. (2014). The regulation of glucose metabolism: implications and considerations for the assessment of glucose homeostasis in rodents. *Am. J. Physiol. Endocrinol. Metab.* **307**, E859–E871. doi:10.1152/ajpendo.00165.2014
- Liu, T.-W., Park, Y.-M., Holscher, H. D., Padilla, J., Scroggins, R. J., Welly, R., Britton, S. L., Koch, L. G., Vieira-Potter, V. J. and Swanson, K. S. (2015). Physical activity differentially affects the cecal microbiota of ovariectomized female rats selectively bred for high and low aerobic capacity. *PLoS ONE* **10**, e0136150. doi:10.1371/journal.pone.0136150
- Mach, N. and Fuster-Botella, D. (2017). Endurance exercise and gut microbiota: a review. *J. Sport Health Sci.* **6**, 179–197. doi:10.1016/j.jshs.2016.05.001
- Mailing, L. J., Allen, J. M., Buford, T. W., Fields, C. J. and Woods, J. A. (2019). Exercise and the gut microbiome: a review of the evidence, potential mechanisms, and implications for human health. *Exerc. Sport Sci. Rev.* **47**, 75–85. doi:10.1249/JES.0000000000000183
- Malisch, J. L., Breuner, C. W., Kolb, E. M., Wada, H., Hannon, R. M., Chappell, M. A., Middleton, K. M. and Garland, T., Jr. (2009). Behavioral despair and home-cage activity in mice with chronically elevated baseline corticosterone concentrations. *Behav. Genet.* **39**, 192–201. doi:10.1007/s10519-008-9246-8
- Matsumoto, M., Inoue, R., Tsukahara, T., Ushida, K., Chiji, H., Matsubara, N. and Hara, H. (2008). Voluntary running exercise alters microbiota composition and increases n-butyrate concentration in the rat cecum. *Biosci. Biotechnol. Biochem.* **72**, 572–576. doi:10.1271/bbb.70474
- McNamara, M. P., Singleton, J. M., Cadney, M. D., Ruegger, P. M., Borneman, J. and Garland, T., Jr. (2021). Early-life effects of juvenile Western diet and exercise on adult gut microbiome composition in mice. *J. Exp. Biol.* **224**, jeb239699. doi:10.1242/jeb.239699
- McNamara, M. P., Cadney, M. D., Castro, A. A., Hillis, D. A., Kallini, K. M., Macbeth, J. C., Schmill, M. P., Schwartz, N. E., Hsiao, A. and Garland, T., Jr. (2022). Oral antibiotics reduce voluntary exercise behavior in athletic mice. *Behav. Processes* **199**, 104650. doi:10.1016/j.beproc.2022.104650
- Meek, T. H., Lonquich, B. P., Hannon, R. M. and Garland, T., Jr. (2009). Endurance capacity of mice selectively bred for high voluntary wheel running. *J. Exp. Biol.* **212**, 2908–2917. doi:10.1242/jeb.028886
- Meek, T. H., Eisenmann, J. C., Keeney, B. K., Hannon, R. M., Dlugosz, E. M. and Garland, T., Jr. (2014). Effects of early-life exposure to Western diet and wheel access on metabolic syndrome profiles in mice bred for high voluntary exercise. *Genes Brain Behav.* **13**, 322–332. doi:10.1111/gbb.12098
- Mohr, A. E., Jäger, R., Carpenter, K. C., Kerksick, C. M., Purpura, M., Townsend, J. R., West, N. P., Black, K., Gleeson, M., Pyne, D. B. et al. (2020). The athletic gut microbiota. *J. Int. Soc. Sports Nutr.* **17**, 24. doi:10.1186/s12970-020-00353-w

- Munukka, E., Ahtiainen, J. P., Puigbó, P., Jalkanen, S., Pakkala, K., Keskitalo, A., Kujala, U. M., Pietilä, S., Hollmén, M., Elo, L. et al. (2018). Six-week endurance exercise alters gut metagenome that is not reflected in systemic metabolism in over-weight women. *Front. Microbiol.* **9**, 2323. doi:10.3389/fmicb.2018.02323
- Nagpal, R., Wang, S., Solberg Woods, L. C., Seshie, O., Chung, S. T., Shively, C. A., Register, T. C., Craft, S., McClain, D. A. and Yadav, H. (2018). Comparative microbiome signatures and short-chain fatty acids in mouse, rat, non-human primate, and human feces. *Front. Microbiol.* **9**, 2897. doi:10.3389/fmicb.2018.02897
- Oyanagi, E., Uchida, M., Kremenik, M. J. and Yano, H. (2018). Altered gut microbiota by voluntary exercise induces high physical activity in high-fat diet mice. *J. Phys. Fit. Sports Med.* **7**, 81-85. doi:10.7600/jpfsm.7.81
- Pantoja-Feliciano, I. G., Clemente, J. C., Costello, E. K., Perez, M. E., Blaser, M. J., Knight, R. and Dominguez-Bello, M. G. (2013). Biphasic assembly of the murine intestinal microbiota during early development. *ISME J.* **7**, 1112-1115. doi:10.1038/ismej.2013.15
- Park, Y.-M., Kanaley, J. A., Padilla, J., Zidon, T., Welly, R. J., Will, M. J., Britton, S. L., Koch, L. G., Rueggsegger, G. N., Booth, F. W. et al. (2016). Effects of intrinsic aerobic capacity and ovariectomy on voluntary wheel running and nucleus accumbens dopamine receptor gene expression. *Physiol. Behav.* **164**, 383-389. doi:10.1016/j.physbeh.2016.06.006
- Pekkala, S., Lensu, S., Nokia, M., Vanhatalo, S., Koch, L. G., Britton, S. L. and Kainulainen, H. (2017). Intrinsic aerobic capacity governs the associations between gut microbiota composition and fat metabolism age-dependently in rat siblings. *Physiol. Genomics* **49**, 733-746. doi:10.1152/physiolgenomics.00081.2017
- Queipo-Ortuño, M. I., Seoane, L. M., Murri, M., Pardo, M., Gomez-Zumaquero, J. M., Cardona, F., Casanueva, F. and Tinahones, F. J. (2013). Gut microbiota composition in male rat models under different nutritional status and physical activity and its association with serum leptin and ghrelin levels. *PLoS ONE* **8**, e65465. doi:10.1371/journal.pone.0065465
- Rhodes, J. S., Koteja, P., Swallow, J. G., Carter, P. A. and Garland, T., Jr. (2000). Body temperatures of house mice artificially selected for high voluntary wheel-running behavior: repeatability and effect of genetic selection. *J. Therm. Biol.* **25**, 391-400. doi:10.1016/S0306-4565(99)00112-6
- Scheiman, J., Lubner, J. M., Chavkin, T. A., MacDonald, T., Tung, A., Pham, L.-D., Wibowo, M. C., Wurth, R. C., Punthambaker, S., Tierney, B. T. et al. (2019). Meta-omics analysis of elite athletes identifies a performance-enhancing microbe that functions via lactate metabolism. *Nat. Med.* **25**, 1104-1109. doi:10.1038/s41591-019-0485-4
- Schloss, P. D., Schubert, A. M., Zackular, J. P., Iverson, K. D., Young, V. B. and Petrosino, J. F. (2012). Stabilization of the murine gut microbiome following weaning. *Gut Microbes* **3**, 383-393. doi:10.4161/gmic.21008
- Shahi, S. K., Freedman, S. N. and Mangalam, A. K. (2017). Gut microbiome in multiple sclerosis: the players involved and the roles they play. *Gut Microbes* **8**, 607-615. doi:10.1080/19490976.2017.1349041
- Sullivan, G. M. and Feinn, R. (2012). Using effect size—or why the p value is not enough. *J. Grad. Med. Educ.* **4**, 279-282. doi:10.4300/JGME-D-12-00156.1
- Swallow, J. G., Carter, P. A. and Garland, T., Jr. (1998). Artificial selection for increased wheel-running behavior in house mice. *Behav. Genet.* **28**, 227-237. doi:10.1023/A:1021479331779
- Swallow, J. G., Koteja, P., Carter, P. A. and Garland, T., Jr. (1999). Artificial selection for increased wheel-running activity in house mice results in decreased body mass at maturity. *J. Exp. Biol.* **202**, 2513-2520. doi:10.1242/jeb.202.18.2513
- Swallow, J. G., Koteja, P., Carter, P. A. and Garland, T., Jr. (2001). Food consumption and body composition in mice selected for high wheel-running activity. *J. Comp. Physiol. [B]* **171**, 651-659. doi:10.1007/s003600100216
- Swallow, J. G., Hayes, J. P., Koteja, P. and Garland, T., Jr. (2009). Selection experiments and experimental evolution of performance and physiology. In *Experimental Evolution: Concepts, Methods, and Applications of Selection Experiments* (ed. T. Garland, Jr. and M. R. Rose), pp. 301-351. University of California Press.
- Swallow, J. G., Wroblewska, A. K., Waters, R. P., Renner, K. J., Britton, S. L. and Koch, L. G. (2010). Phenotypic and evolutionary plasticity of body composition in rats selectively bred for high endurance capacity. *J. Appl. Physiol.* **109**, 778-785. doi:10.1152/jappphysiol.01026.2009
- Tamburini, S., Shen, N., Wu, H. C. and Clemente, J. C. (2016). The microbiome in early life: implications for health outcomes. *Nat. Med.* **22**, 713-722. doi:10.1038/nm.4142
- Wallace, I. J. and Garland, T., Jr. (2016). Mobility as an emergent property of biological organization: Insights from experimental evolution. *Evol. Anthropol. Issues News Rev.* **25**, 98-104. doi:10.1002/evan.21481
- Wisløff, U., Najjar, S. M., Ellingsen, Ø., Haram, P. M., Swoap, S., Al-Share, Q., Fernström, M., Rezaei, K., Lee, S. J., Koch, L. G. et al. (2005). Cardiovascular risk factors emerge after artificial selection for low aerobic capacity. *Science* **307**, 418-420. doi:10.1126/science.1108177
- Yatsunenko, T., Rey, F. E., Manary, M. J., Trehan, I., Dominguez-Bello, M. G., Contreras, M., Magris, M., Hidalgo, G., Baldassano, R. N., Anokhin, A. P. et al. (2012). Human gut microbiome viewed across age and geography. *Nature* **486**, 222. doi:10.1038/nature11053

Figure S1




Fig. S1. Community composition of the weanling gut microbiome for all experimental mice (N=95) was dominated by Firmicutes ($48.4 \pm 14\%$) (mean \pm S.D.) and Bacteroidetes ($37.5 \pm 17.1\%$), with additional phyla being much less abundant: Proteobacteria ($11.8 \pm 8.7\%$), Tenericutes ($1.6 \pm 2\%$), Cyanobacteria ($0.28 \pm 0.53\%$), Verrucomicrobia ($0.24 \pm 1.1\%$), Actinobacteria ($0.09 \pm 0.16\%$), Deferribacteres ($0.08 \pm 0.24\%$), Fusobacteria ($0.0002 \pm 0.0006\%$), and TM7 ($0.0001 \pm 0.0004\%$).

Figure S2




Fig. S2. Beta diversity (among experimental groups) of the weanling fecal microbiome based on 16S rRNA sequence data. A and B are PCoA plots based on Bray-Curtis distances, which consider bacterial OTU sequence relative abundances. C and D are PCoA plots based on weighted UniFrac distances, which consider both bacterial OTU sequence relative abundances and phylogenetic distances. PERMANOVAs based on Bray-Curtis or weighted UniFrac distance matrices indicated no statistically significant separation based on either linetype (Bray-Curtis: $R^2=0.0147$, $P=0.108$, weighted UniFrac: $R^2=0.0116$, $P=0.307$) or mini-muscle status (Bray-Curtis: $R^2=0.009$, $P=0.638$, weighted UniFrac: $R^2=0.005$, $P=0.773$).

Figure S3




Fig. S3. PCoA plots from separate weighted UniFrac and Bray-Curtis analyses of the 4 HR lines (lab designations 3,6,7,8: A and C) and of the 4 C lines (lab designations 1,2,4,5: B and D). Values are means and standard errors for scores on PCoA axes. Separately, we used PERMANOVAs to test for significant separation among the 4 replicate HR lines and among the 4 non-selected C lines. Differences among the 4 replicate C lines were statistically significant based on Bray-Curtis ($R^2=0.051$, $P=0.005$), but not among the 4 replicate HR lines ($R^2=0.014$, $P=0.839$). Weighted UniFrac indicated no significant separation among replicate lines for either linetype (HR lines: $R^2=0.005$, $P=0.957$; C lines: $R^2=0.030$, $P=0.228$).