

Planar cell polarity controls directional Notch signaling in the *Drosophila* leg

Amalia Capilla^{1,2,*}, Ruth Johnson^{3,*}, Maki Daniels³, María Benavente¹, Sarah J. Bray^{3,§} and Máximo Ibo Galindo^{1,2,§}

SUMMARY

The generation of functional structures during development requires tight spatial regulation of signaling pathways. Thus, in *Drosophila* legs, in which Notch pathway activity is required to specify joints, only cells distal to ligand-producing cells are capable of responding. Here, we show that the asymmetric distribution of planar cell polarity (PCP) proteins correlates with this spatial restriction of Notch activation. Frizzled and Dishevelled are enriched at distal sides of each cell and hence localize at the interface with ligand-expressing cells in the non-responding cells. Elimination of PCP gene function in cells proximal to ligand-expressing cells is sufficient to alleviate the repression, resulting in ectopic Notch activity and ectopic joint formation. Mutations that compromise a direct interaction between Dishevelled and Notch reduce the efficacy of repression. Likewise, increased Rab5 levels or dominant-negative Deltex can suppress the ectopic joints. Together, these results suggest that PCP coordinates the spatial activity of the Notch pathway by regulating endocytic trafficking of the receptor.

KEY WORDS: Planar cell polarity, Notch, Dishevelled, *Drosophila*

INTRODUCTION

The development and physiology of all multicellular organisms requires cell communication through well-defined signaling pathways that each consist of distinct canonical components. Cross-talk between pathways is required for the relatively limited number of pathways to match the anatomical and functional complexity that cell signaling has to regulate. The mechanisms that enable the pathway cross-talk are thus of major importance for generation and maintenance of complex structures.

One highly conserved signaling pathway important for coordinating many developmental processes is mediated by Notch transmembrane receptors (Fiuza and Arias, 2007; Fortini, 2009). Besides Notch (N), core members of this pathway in *Drosophila* include two transmembrane ligands: Serrate (Ser; Jagged in vertebrates) and Delta (DI). Upon ligand binding, N suffers two consecutive proteolytic cleavages and releases its cytoplasmic portion, which enters the nucleus and mediates a transcriptional response by binding to CSL transcription factors. Behind this apparent simplicity, a wide variety of biological functions and modes of action are made possible by context-dependent accessory mechanisms that help regulate the activation of N (Andersson et al., 2011; Bray, 2006). These include post-translational modifications, such as glycosylation and ubiquitinylation, that affect endocytic sorting of both N and its ligands.

N activity is also modulated by key aspects of tissue organization, including planar cell polarity (PCP). PCP was first characterized in *Drosophila* epithelial cells, where it establishes a polarity axis in the tissue plane, orthogonal to the apical-basal axis (Goodrich and Strutt, 2011; Vladar et al., 2009). Its relevance is evident in the orientation of cell projections, such as hairs or microvilli, and it is also important in coordinating behavior in fields of cells, ensuring that they respond in a homogeneous directional fashion, including convergent extension in vertebrate embryos and ommatidial rotation in insect eyes. The latter is one example for which PCP and N are known to converge (Cooper and Bray, 1999; Fanto and Mlodzik, 1999; Tomlinson and Struhl, 1999).

The proteins of one of the main PCP pathways (the core, Fz or Stan system) associate in complexes at the cell membrane. They include the transmembrane proteins Van Gogh (Vang; also known as Strabismus, Stbm) (Taylor et al., 1998; Wolff and Rubin, 1998), Frizzled (Fz) (Vinson et al., 1989) and Flamingo (Fmi; also known as Starry Night, Stan) (Chae et al., 1999; Usui et al., 1999) as well as the cytoplasmic proteins Prickle (Pk) (Gubb et al., 1999), Dishevelled (Dsh) (Klingensmith et al., 1994; Theisen et al., 1994) and Diego (Dgo) (Feiguin et al., 2001). In the wing epithelium, PCP protein complexes acquire an asymmetric proximal-distal localization (Strutt and Strutt, 2009). A Fz-Dsh complex localizes to the distal side of cells, together with Dgo, whereas a Stbm-Pk complex is localized to the proximal domain. These two complexes repel each other within the cell and both require Fmi and other proteins for their correct localization. Most of these core PCP proteins function in other planar polarized systems in *Drosophila* and in vertebrates, although the details of their localization or cellular actions might differ (Seifert and Mlodzik, 2007). Besides this role in PCP, a non-canonical Wnt pathway, Fz and Dsh are also required in canonical Wnt signaling, for which they trigger nuclear accumulation of β -catenin upon Wnt activation (MacDonald et al., 2009). Most mutations in Fz affect its role in both PCP and Wnt signaling (Povelones et al., 2005), whereas PCP-specific mutations of Dsh affect protein localization (Axelrod et al., 1998). In addition,

¹Developmental Cell Biology Unit, Instituto de Biomedicina de Valencia (IBV-CSIC), Jaime Roig 11, 46010 Valencia, Spain. ²Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Alvaro de Bazán 10, 46010 Valencia, Spain.

³Department of Physiology Development and Neuroscience, University of Cambridge, Cambridge CB2 3DY, UK.

*These authors contributed equally to this work

[§]Present address: Biology Department, 52 Lawn Avenue, Room 212, Hall-Atwater Laboratories, Wesleyan University, Middletown, CT 06459, USA

[§]Authors for correspondence (sjb32@cam.ac.uk; ibo@ibv.csic.es)

the interaction partners also influence the outcome as the association with Dgo produces a bias towards PCP to the detriment of Wnt (Wu et al., 2008).

A striking feature of flies mutant for core PCP members is that they have supernumerary joints in the tarsal region of the leg (Held et al., 1986). Normally composed of five segments (T1 to T5) separated by joints with a ball and socket structure, tarsi mutant for core PCP genes contain ectopic joints in segments T2, T3 and T4 and, less frequently, T1. Joints are determined at the end of larval development, when a stripe of Ser-expressing cells is specified within each segment and activates the receptor in distal cells triggering transcription of several N targets that control different aspects of joint differentiation (Bishop et al., 1999; de Celis et al., 1998; Rauskolb and Irvine, 1999). Ser appears to be the functional N-ligand in this process, as joints are absent in Ser mutants although other aspects of leg morphology appear normal, and in PCP mutant legs the ectopic joints correlate with ectopic Notch activity although the mechanism is unknown (Bishop et al., 1999).

The ectopic joint phenotype in PCP mutant flies implies that the PCP system has a role in regulating N signaling (Bishop et al., 1999). The likely scenario is that, when PCP is disrupted, N becomes activated in cells proximal to Ser-expressing cells, as well as those distal. How this regulation occurs is, however, unknown. In the eye, where the R3/R4 photoreceptor fate choice is crucial for ommatidial polarity, Fz activity in the presumptive R3 is essential for polarizing N activity (Cooper and Bray, 1999; Fanto and Mlodzik, 1999; Tomlinson and Struhl, 1999). It does so via a combination of mechanisms including effects on *Dl* transcription and activation as well as on endocytic regulation of N (Cho and Fischer, 2011; del Alamo and Mlodzik, 2006; Strutt et al., 2002) that may be amplified via Fmi upregulation (Das et al., 2002). Whether these mechanisms operate during other processes, such as joint development, remains to be established. Given the prevalence of PCP in many tissues, understanding how it can influence the ability of a cell to send or receive signals is of widespread relevance.

MATERIALS AND METHODS

Fly stocks

The following mutant alleles were used, either homozygous or in mitotic clones: *pk^{sple1}*, *dsh¹*, *dsh^{v26}*, *fmi¹⁹²*, *fz²²*, *Drok²* (described in FlyBase, <http://flybase.org>). To monitor N activation we used the following reporter lines: *bib-lacZ*, *disco-lacZ*, *E(spl)mβ1.5-CD2* and *E(spl)mβ1.5-lacZ*. Subcellular localization of proteins was analyzed using GFP fusions: *Ac5-Vang-GFP* (Strutt et al., 2002), *arm-fz-GFP* (Strutt, 2001), *dsh-GFP* (Axelrod, 2001). For directed expression *ap-Gal4* or the flip-out cassette *Ac>CD2>Gal4* were used to drive the expression from UAS constructs: *UAS-GFP*, *UAS-dsh-myc*, *UAS-Rab5*, *UAS-Rab7*, *UAS-Rab11* (<http://flybase.org>), *UAS-dx*, *UAS-dx^{APRM}*, *UAS-dx^{mRZF}*, *UAS-dx^{ΔNBS}* (Matsuno et al., 2002).

The FLP/FRT technique was used to generate mutant clones (Xu and Rubin, 1993) with appropriate recombinant chromosomes. To induce the FLPase, 48–72 hours after egg laying (AEL) (second instar) larvae were heat-shocked at 37°C in a water bath for one hour.

Histology, immunofluorescence and microscopy

Prepupal leg discs were dissected in PBS and fixed in 4% paraformaldehyde in PBS. Primary antibodies were: rabbit anti-β-galactosidase (Life Technologies, Grand Island, NY, USA), mouse anti-β-galactosidase (Promega, Madison, WI, USA), rabbit anti-GFP (Rockland, Gilbertsville, PA, USA), mouse anti-CD2 (AbD Serotec, Kidlington, UK), rabbit anti-Serrate (gift of Ken Irvine, Waksman Institute of Microbiology, NJ, USA) and rabbit anti-Rab5 (Abcam, Cambridge, UK). In addition, mouse anti-Arm (β-catenin), mouse anti-Fmi, mouse anti-Nicd, mouse

anti-Necd and rat anti-E-cadherin were obtained from the Developmental Studies Hybridoma Bank (University of Iowa, USA). Actin cytoskeleton was labeled with Phalloidin-TRITC (Sigma, St Louis, MO, USA).

Samples were examined using a Leica DM RXA2 microscope and Leica TCS SP Confocal system (Leica Microsystems, Wetzlar, Germany). Images were processed and analyzed with Leica Confocal Software, ImageJ suite and Adobe Photoshop.

dsh genomic constructs

Specific mutations were introduced into the coding sequence of the *dsh-GFP* genomic fragment (Axelrod, 2001) using site-directed mutagenesis. In brief, a 1.4 kb fragment containing part of the promoter and the coding region for the DIX and PDZ domains was subcloned into pKS for mutagenesis using the QuikChange Site-Directed Mutagenesis Kit (Agilent Technologies, Santa Clara, CA, USA) to introduce the K46V and Q47A mutations. The mutated region was then substituted into the full length *dsh-GFP* by replacing the 400 bp *MluI-KpnI* fragment encompassing the DIX domain. The entire *dsh-GFP* mutant genomic fragment was introduced into the transformation vector pWhiteRabbit. Single copy insertions of the resulting plasmid were generated using conventional P-element-mediated transformation and multiple insertion lines were mapped to chromosomes and analyzed for expression. Suitable lines were then crossed into *dsh¹* and *dsh^{v26}* backgrounds to generate *w¹¹⁴ dsh^[x] / FM7*; *dsh^{mut6} (w⁺) / dsh^{mut6} (w⁺)* and the phenotypes analyzed in *dsh^[x] / Y* males. Over 100 legs were scored for ectopic joint phenotype for each one of the independent constructs.

Biochemistry

To map the domains of Dsh interacting with the intracellular domain of Notch (NIC), glutathione s-transferase (GST) pull-down experiments were performed as described (Djiane et al., 2005); by incubating GST-NIC fusion with ³⁵S-labeled Dsh fragments (primers are listed in supplementary material Tables S1, S2). Specific amino acids were mutated by site-directed mutagenesis with the QuikChange Kit (Agilent Technologies, Santa Clara, CA, USA). The relative intensity of the bands was calculated using the gel analysis application of the ImageJ suite, by plotting the lane profile and calculating of the resulting peak areas. Two gels were analyzed for each pair of bands.

RESULTS

The double joint planar cell polarity phenotype correlates with ectopic N activity

Mutations affecting core PCP genes result in supernumerary joints in the tarsal region, most commonly in tarsal segments 2–4 (Bishop et al., 1999; Held et al., 1986). Ectopic joints are located proximal to the normal joint and have reversed polarity (Fig. 1A,B). Such ectopic joints appear to be a bona fide PCP phenotype, as the defect occurs with mutations affecting any core PCP gene: phenotypes of *pk*, *fz*, *dsh*, *Vang*, *fmi* and *dgo* are consistent and differ from other mutant conditions affecting joint development (Held et al., 1986; Wolff and Rubin, 1998) (<http://flybase.org>; our unpublished data).

The implication of this phenotype is that PCP controls the directionality of cell signaling mediated by the N pathway, the main agent in joint determination (Bishop et al., 1999; de Celis et al., 1998; Rauskolb and Irvine, 1999). To ascertain whether this interpretation is correct, we have analyzed expression of three reporters of N activation in different viable mutants of core PCP genes. Reporter expression is fully established by 2–6 hours after puparium formation (APF) when the leg disc starts to evert. The best characterized direct N targets are the *E(spl)* genes. An *E(spl)mβ1.5-lacZ* reporter has previously been shown to respond to N in the leg (Cooper et al., 2000; de Celis Ibeas and Bray, 2003), where it is expressed in a stripe distal to, and partially overlapping, the Ser-expressing cells. In prepupal legs of *dsh¹*, a PCP-specific




Fig. 1. N is ectopically activated in *Drosophila* PCP mutants. (A,B) Wild-type leg with normal joints (A) and *pk^{sple1}* leg with ectopic joints with inverted polarity (B). In these and subsequent panels arrowheads indicate normal joints terminating tarsal segments 2, 3 and 4, and the corresponding domains of expression. (C,D) *E(spl)mβ1.5-lacZ* expression (X-gal, blue) in wild-type prepupal leg (C) and *dsh¹* mutant (D) with ectopic stripes. (E,F) *bib-lacZ* expression (green) in a single row of cells (anti-β-catenin, purple, cell contours) at the segment boundary in wild type (E) is duplicated in a *pk^{sple1}* (F). (G,H) *disco-lacZ* (green) is expressed in a wider domain in wild type (G); normal and duplicated domains merge in a single broad territory in *pk^{sple1}* (H). (I,I') *bib-lacZ* (green) is expressed distally adjacent to Ser (I, purple; I', gray). (J,J') In *pk^{sple1}* the ectopic stripe (green) appears proximally adjacent to Ser (J, purple; J', gray); a sub-apical confocal section is shown to capture the nuclear β-galactosidase. (K,K') Large *fz^{J22}* clone marked by absence of GFP (green); expression of Ser (K, purple; K', gray) is not altered.

allele, an ectopic domain appeared, confirming that disruptions in core PCP gene activity result in ectopic N activity (Fig. 1C,D). Equivalent ectopic domains of *E(spl)mβ1.5-lacZ* were seen with alleles affecting other core PCP genes (see below).

Similar results were obtained with a *lac-Z* insertion into *big brain* (*bib-lacZ*), expression of which is also regulated by N (de Celis et al., 1998; Pueyo and Couso, 2011). *bib-lacZ* is expressed in a narrow band one or two cells wide just distal to the domain of Ser expression (Fig. 1E,I). In prepupal legs of *pk^{sple1}*, a strong

hypomorphic allele of *pk* (Gubb et al., 1999), there was an ectopic *bib-lacZ* stripe proximal to the Ser-expressing cells (Fig. 1F,J). In addition, the levels of *bib-lacZ* were sometimes reduced or discontinuous (see below). A further marker of leg joints (although not a known direct target of N) is *disco-lacZ* (Bishop et al., 1999), an insertion into *disconnected*, which is expressed in a broader domain spanning three or four cell diameters (Fig. 1G). In a *pk^{sple1}* mutant, the *disco-lacZ* domain was duplicated and, owing to its larger territory, the ectopic domain merged with the endogenous one from the preceding segment, resulting in a continuous domain of expression in most of the tarsal region (Fig. 1H).

Despite the evidence for ectopic N activity, there was no change in expression of the Ser ligand in *pk^{sple1}* legs or in mitotic clones of *fz^{J22}* (a strong *fz* hypomorph; Fig. 1I-K) or of Df present at later stages (Bishop et al., 1999). Neither was there a clear alteration in the expression profile of the N receptor itself (supplementary material Fig. S1). Therefore, the extra stripes of N activation are a general feature of mutations affecting PCP but are unlikely to be the result of a simple change in the expression of N or its ligands.

Cell-autonomous effects of PCP alleles suggest a requirement in the signal-receiving cell

In order to know whether the mechanism of action of PCP on N is direct or indirect, or whether it is likely to act in ligand-sending or signal-receiving cells, we analyzed defects caused by clones of mutant cells. If the effects are direct in signal-receiving cells, activation of N targets should only be detected autonomously within mutant cells in clones located proximal to the site of ligand expression. Conversely, if PCP regulation affects the ligand, some non-autonomous defects would be seen. It is important to note that not all mutant alleles would be useful for this analysis, because several of them show a directional domineering non-autonomy owing to reorganization in PCP over neighboring cells. Therefore, we used only alleles reported to show autonomous polarization phenotypes: *pk^{sple1}*, *fmi¹⁹²*, *dsh¹* and *fz^{J22}* (Chae et al., 1999; Jones et al., 1996; Lee and Adler, 2002; Strutt and Strutt, 2007).

We first examined effects on the *E(spl)mβ1.5-CD2* reporter [containing the same regulatory element as *E(spl)mβ1.5-lacZ*], a direct target of N pathway. In clones of *fz^{J22}*, expression of *E(spl)mβ1.5-CD2* is de-repressed autonomously within the mutant cells (Fig. 2A). Furthermore, in several examples the mutant cells were juxtaposed with putative ligand-producing cells that were wild type. These results argue that the effect of *fz^{J22}* on the N pathway is autonomous and is most likely to occur in the signal-receiving cells. *E(spl)mβ1.5-CD2* is also autonomously de-repressed in *dsh¹* mutant clones (not shown).

The behavior of *disco-lacZ* in *pk^{sple1}* and *fmi¹⁹²* clones was identical to that of *E(spl)mβ1.5-CD2*. In both genotypes, *disco-lacZ* was de-repressed in a completely autonomous manner (Fig. 2B,C). Again, expression of *lacZ* in cells at clone edges suggested a requirement for polarization in the signal-receiving cell rather than on the ligand.

The effects of mutations on *bib-lacZ* were, however, slightly different. In all three of the mutants tested (*dsh¹*, *pk^{sple1}*, *fmi¹⁹²*) de-repression of the reporter was detected only in mutant cells and not in adjacent wild-type cells (Fig. 2D,E; supplementary material Fig. S2). However, unlike *E(spl)mβ1.5-CD2* and *disco-lacZ*, ectopic activation of *bib-lacZ* was variable in intensity (weak to similar to endogenous *bib-lacZ*) and in extent. In some clones, only a few cells showed ectopic expression whereas in others it filled the whole width of the mutant clone. Nevertheless, the fact that *bib-lacZ* can be de-repressed at the clone boundaries is consistent with




Fig. 2. Proximal de-repression of N in *Drosophila* PCP mutants is cell autonomous. In all panels, mutant clones are revealed by absence of GFP (green); ectopic expression of reporters (purple; single channel on the right) is indicated by arrowheads. (A) Mutant *fzJ22* clone; ectopic expression of *E(spl)mβ1.5-CD2* coincides with clone border despite presence of adjacent wild-type (ligand-expressing) cells. (B) Mutant *fmi192* clones; autonomous expression of *disco-lacZ* in cells adjacent to wild-type GFP-positive cells. (C) Mutant clones of *pksp1*; several ectopic domains of *disco-lacZ* occur autonomously within the mutant clones. (D) Large clone of cells mutant for *dsh1*; some ectopic expression of *bib-lacZ* appears in tarsal segment 3, but not in segment 2. (E) Elongated clones of cells mutant for *pksp1*; ectopic activation of *bib-lacZ* that can occupy the full width of the clone. Occasional downregulation of the normal stripe of *bib-lacZ* is also detected (arrows).

a requirement for PCP in signal-receiving cells, rather than through effects on Ser. We note also that in some *fmi* and *pk* clones the endogenous domain of *bib-lacZ* is weakened within the clone (Fig. 2E). This observation also correlates with the fact that in whole legs mutant for core PCP genes, normal *bib-lacZ* expression can be weakened compared with the ectopic expression (Fig. 1F; supplementary material Fig. S2).

Both the variability of *bib-lacZ* de-repression within clones and the weakening of endogenous expression in mutant cells might reflect a difference in the threshold of N activity required for *bib-lacZ* activation compared with *disco-lacZ* and *E(spl)mβ1.5-CD2* (see Discussion). Nevertheless, N reporters were de-repressed autonomously in all the core PCP mutant genotypes when mutant cells were located proximal to the Ser domain. No ectopic expression of the reporters was observed in wild-type tissue adjacent to the mutant cells, which argues against an effect of PCP

on Ser or on a second signaling pathway. These results were replicated in earlier stages (third instar leg discs) and in smaller clones (supplementary material Fig. S3).

Asymmetric distribution of core PCP proteins in the developing leg

Core PCP proteins adopt a polarized distribution in *Drosophila* pupal wing cells, which are arranged in a regular hexagonal lattice (between 20 and 24 hours APF). Dsh and Fz become localized to the distal edge of each cell, Pk and Vang to the proximal edge (Strutt and Strutt, 2009). Some degree of polarization is also evident in prepupal stages (Aigouy et al., 2010; Strutt et al., 2011). Cells in the prepupal leg epithelium are mostly irregular in shape and size, unlike the wing, making it difficult to detect clear organization in cell morphology or protein distribution. To investigate whether there was any asymmetry in core PCP protein distribution, we generated patches of cells expressing Fz::GFP, Vang::GFP and Dsh::GFP and examined protein distributions at clone borders, as was done previously in the wing to investigate protein asymmetries (Axelrod, 2001; Strutt et al., 2002; Strutt, 2001). Fz::GFP and Dsh::GFP levels were higher at the distal side of cells compared with proximal (Fig. 3A,B). Conversely, Vang::GFP levels were highest on the proximal sides of each cell (Fig. 3C). Some differences in Fmi localization were also evident: the protein was more enriched at proximal-distal boundaries than at dorsal-ventral (Fig. 3D), a characteristic that was most obvious in the tarsus-pretarsus boundary where cells have a more regular morphology (Fig. 3D').

Cells receiving the N signal also exhibited distinct morphology. Detection of β-catenin, localized to sub-apical adherens junctions, revealed that the *bib-lacZ*-expressing cells were roughly quadrangular. Their distal edges formed a straight line, probably marking the boundary between adjacent tarsal segments (Fig. 3E). These features were also observed in the ectopic domain of *bib-lacZ* in the PCP mutants (Fig. 3F). In addition, the intensity of *bib-lacZ* expression was correlated with cell morphology, both in the normal and ectopic domains of expression. This suggests that high levels of N activation result in ordered alignment of the leg epithelial cells.

Direct interaction of N and Dsh

Drok (Rok – FlyBase) is one of the main mediators of the cytoskeletal response to PCP: it is important for restricting wing hair generation and for ommatidial rotation (Winter et al., 2001). We tested whether Drok had any effect on the *bib-lacZ* and *E(spl)mβ1.5-CD2* reporters. Although *Drok* mutant clones showed defects in ommatidial rotation and in tissue morphology (data not shown), none resulted in ectopic expression of the *bib-lacZ* or *E(spl)mβ1.5-CD2* reporters (Fig. 4A,B). Therefore, this function of the core PCP pathway does not seem to be mediated by the actin cytoskeleton.

Previous studies have shown a physical interaction between Dsh and N, which contributes to inhibition of N signaling in the wing margin (Axelrod et al., 1996; Munoz-Descalzo et al., 2010). We questioned whether a similar mechanism operates in joint regulation. Overexpression of Dsh driven by *ap-Gal4* in T4 and proximal T5 had two clear effects (Fig. 4C,D). First, there was a disruption of PCP that has already described for Dsh overexpression in the wing (Axelrod et al., 1998). Second, formation of the joint between these tarsi was repressed, as would be expected if Dsh were capable of repressing N. A possible caveat to this interpretation is that Dsh overexpression could produce




Fig. 3. Cell biology of PCP in the *Drosophila* prepupal leg.

(A-D') Localization of PCP proteins, proximal-distal orientation is top left to bottom right; cells outlined with anti-E-cadherin (A-C, purple). Arrows indicate proximal cell boundaries and arrowheads indicate distal cell boundaries. (A) Cells at the border of Fz::GFP-expressing clones reveal that the fusion protein is enriched at the distal side. (B) Expression of Dsh::GFP is comparatively weak, but also accumulates distally (arrowhead). (C) Vang::GFP localizes to the proximal side of cells (arrow). (D,D') Fmi is absent from cell borders oriented along the dorsal-ventral axis in T2 (D) and in the tarsus-pretarsus boundary (D'). (E) Late prepupal leg, cells expressing *bib-lacZ* (green, dotted) have a larger sub-apical diameter (β -catenin, purple; single channel on the right), a more regular shape and their borders align to form a straight line. (F) Late prepupal *pk^{sp}le1* leg, ectopic rows of *bib-lacZ* expression (green, dotted) have the same features; intensity of *bib-lacZ* appears to correlate with cell size, shape and alignment both in normal and ectopic domains.

patterning defects causing a secondary effect on joints. However, the effects of expressing Dsh in clones do not support this possibility. For example, a large dorsal clone of Dsh-expressing cells produced autonomous repression of *bib-lacZ* (Fig. 4E,E'). Although this clone contained two small putative axis duplications (ectopic leg tips in the form of circular domains of *bib-lacZ* expression in tarsi 2 and 5; Fig. 4C', arrowhead), segmentation was largely unaltered.

To test functional relevance of the direct interaction of Dsh and N in leg segmentation, we set out to find a mutant form of Dsh that had reduced ability to interact physically with N. To map the interacting regions we used glutathione-s-transferase (GST) pull-down experiments. From a set of deletions spanning different regions of Dsh, only constructs with an intact DIX domain were successfully retained by the intracellular portion of N (GST-NIC; Fig. 5A,B). Deletion of the N-terminal region of this DIX domain was sufficient to abolish this interaction (compare constructs Dsh5 and Dsh6). Unfortunately, as the DIX domain is also required for Axin recruitment (Julius et al., 2000; Kishida et al., 1999), these deletions also prevented Axin binding (Fig. 5B). It was therefore important to identify mutations that would affect binding of NIC but not Axin. We first tested effects




Fig. 4. N downregulation by Dsh. (A,A') Clone of cells homozygous for *Drok2* (marked by the absence of GFP, green); no ectopic expression of *bib-lacZ* is detected (purple, single channel in A'). (B,B') In clones of the same genotype, expression of *E(spl)mβ1.5-CD2* is also unaffected. (C) *ap-Gal4* expression domain revealed with UAS-GFP (green), includes T4 and proximal part of T5 (anti-E-cadherin, purple). (D) *ap-Gal4*, UAS-*dsh* causes defects in planar polarity and the joint between T4 and T5 is absent (bracket indicates *ap-Gal4* territory). (E,E') Large dorsal clone expressing Dsh::myc (anti-myc, green) results in autonomous repression of *bib-lacZ* (purple, single channel in E') and putative axis duplications (ectopic leg tips, arrowheads). All tarsal segments are present with segmental grooves detected as normal.

of mutations in residues that are conserved between Dsh and the mammalian Dvl proteins. Of the three mutations tested, only V43E (present in *mut1* and *mut4*) abolished interaction with NIC and this also affected Axin binding (Fig. 5C-E; data not shown). Substitution of two adjacent hydrophilic residues that are specific to *Drosophila* Dsh, K46V+Q47A, showed some specificity for NIC. Quantitative analysis of the band intensities revealed that this mutation (*mut6*) reduced binding to NIC by 95%, but interaction with Axin was reduced by only 65%. This mutation was therefore a candidate to test relevance for N regulation.

To investigate whether the *mut6* form of Dsh was compromised for N regulation in the leg, we introduced the K46V+Q47A mutation into a *dsh* genomic rescue construct that had been used previously (Axelrod, 2001). Multiple insertions of the mutant protein were tested for their ability to rescue *dsh^l* and *dsh^{v26}* mutant phenotypes. The former only affects PCP function; the latter is a null allele affecting both PCP and Wnt signaling. All three constructs could rescue the embryonic lethality of *dsh^{v26}*, so they can function in the canonical Wnt pathway. As expected, the PCP ectopic leg joint phenotype was rescued by the wild-type *dsh* construct but not by *dsh^l* (Fig. 6A). Importantly, the *mut6* constructs showed a reduced ability to rescue the ectopic leg joint phenotype (Fig. 6A,B), although the proximal-distal and dorsal-ventral patterns were wild type, including the distal tip, which is the part of the leg most sensitive to alterations in Wnt signaling (Galindo et al., 2002). In addition, when combined with *dsh^{v26}*, the resulting adult flies had largely wild-type wings and both leg bristles and wing hairs




Fig. 5. Direct interaction of Dsh and N. (A) Domain structure of Dsh protein, with the different deletion constructs employed in the GST pull-down assays depicted below: constructs retained by GST::NIC are ticked. (B) Autoradiograph of ^{35}S -labeled Dsh constructs pulled down with GST::NIC and GST::Axin. Input proteins and a negative control pull-down using an unrelated GST construct (–) are also shown. (C) Alignment of the N- and Axin-interacting region of *Drosophila* Dsh with two human Dishevelled (Dvl) proteins and *Drosophila* and human Axin. Residues mutated are in bold: D39 and V43 are conserved in all Dsh proteins but not in Axins, and K46 and Q47 are present only in *Drosophila* Dsh. (D) Six different site-directed mutants were generated with single or double mutations. (E) GST pull-down assays with the different ^{35}S -labeled Dsh mutants. V43E prevents interaction with NIC and Axin; K46V + Q47A (mut6) hinders the interaction with NIC but retains some interaction with Axin. As a control, GST::Grh fails to bind to any Dsh derivatives.

exhibited normal planar polarization (Fig. 6B-E). Expression levels of the GFP-tagged Dsh proteins produced by *mut6* were similar to those from the wild-type *dsh* construct (Fig. 6B,C). These results, therefore, are consistent with the hypothesis that a direct interaction between Dsh and N is important for suppressing N activity in the domain proximal to Ser stripe. However, we cannot rule out the alternative possibility that the inability of *mut6* to rescue leg joints reflects a difference in the threshold levels of Dsh activity required for this process compared with others.

The antagonistic effect of Dsh on N signaling has been described previously in the wing, where N is also important for patterning sensory organs at the wing margin. However, although overexpression of known N repressors such as Numb (Nb) and Hairless (H) (Frise et al., 1996; Nagel et al., 2005) produced nicks in the wing margin and repression of the Notch-target Cut (supplementary material Fig. S4), ectopic Dsh was sufficient neither to produce nicks in the adult wing nor to repress Cut at larval stages (supplementary material Fig. S4). Although the




Fig. 6. Rescue with genomic constructs.

(A) Presence of ectopic joints in tarsal segments 2,3 and 4 in *dsh¹* and *dsh^{v26}* mutant backgrounds rescued with genomic constructs encoding different forms of *dsh*: *dsh¹* (one insertion), wild-type *dsh* (five different insertions) and *dsh^{mut6}* (three insertions); *yw* is used as a wild-type control. (B,C) Representative legs of *dsh^{v26}* rescued with the genomic constructs for wild-type *dsh* (B) and for *dsh^{mut6}* (C). Arrowheads point to partial ectopic joints; insets show similar expression levels of Dsh::GFP and Dsh^{mut6}::GFP in leg imaginal discs. (D,E) Wings of *dsh^{v26}*; *dsh^{mut6}* (D, trichomes are forked) and *dsh¹*; *dsh^{mut6}* rescued (E) flies with largely normal wing margin and PCP.

interpretation of these results might be confounded by the fact that overexpression of Dsh at the margin can also lead to expression of N ligands and to N activation, possibly explaining the ectopic bristles observed in the wing blade (supplementary material Fig. S4), the effects of overexpression of N are nevertheless relatively minor compared with the overexpression of Nb and H. The results suggest, therefore, that the ability of Dsh to suppress N is restricted to certain contexts, making it less likely that it antagonizes the cleaved, active form of N (N_{icd}).

To investigate further whether Dsh could antagonize N_{icd}, we assayed the effects on an N-responsive reporter (NRE-luciferase) of co-expressing Dsh and N_{icd} in transient transfection assays. Expression of N_{icd} alone resulted in strong induction of NRE-luciferase that was little altered by co-expression with Dsh (supplementary material Fig. S4). Therefore, although Dsh binds to the intracellular domain of N, it does not inhibit N_{icd} trans-activation function, suggesting that it is likely to regulate the receptor prior to cleavage.

Role of endocytic regulators

It has been suggested that Dsh influences endocytic trafficking of proteins including N (Chen et al., 2003; Munoz-Descalzo et al., 2010; Yu et al., 2007). One model therefore is that recruitment of Dsh to the distal edge of the cells would result in downregulation of N by endocytosis. To investigate this, we tested the consequences of expressing several different endocytic regulators in the T4 segment with *ap-Gal4* (Fig. 4C) to determine their effect on the ectopic joint present in *dsh¹* mutants.

First, we tested consequences of expressing three different Rab GTPases, Rab5, Rab7 and Rab11, which regulate vesicle trafficking to early endosome, late endosome and recycling endosome compartments, respectively (Stenmark, 2009). In previous studies, overexpression of Rab5-GFP and Rab7-GFP were able to suppress the ectopic N activation seen in *lethal giant discs* [*lgd; l(2)gd1* – FlyBase] mutants (Jaekel and Klein, 2006). Overexpression of Rab5 in a wild-type background had no effect (Fig. 7A,C), but in a *dsh¹* mutant background it was able to modify the ectopic joint phenotype in 80% of the legs examined, resulting in a partial suppression (Fig. 7B,D). By contrast, neither Rab11 nor Rab7 had any effect in this assay (data not shown), suggesting that the defect is linked to transit to early endosomes. The implication is that *dsh¹* leads to a defect in endocytic transport of N from the plasma membrane, preventing its activation by Ser, and that this can be compensated for by increasing the levels of Rab5. To test this hypothesis, we examined whether overexpression of Dsh::myc, which resulted in lack of joints and downregulation of *bib-lacZ* (Fig. 4C-E), had any impact on N protein distribution. In *ap-Gal4, UAS-dsh-myc* legs there was reduced N at the apical membrane and a large fraction of N colocalized with Dsh in intracellular puncta (Fig. 8A). Many of the N-containing puncta appeared to correspond to early endosomes based on their colocalization with Rab5 (Fig. 8B). Furthermore, N depletion from the cell surface and accumulation in puncta was evident using antibodies against either the extracellular or the intracellular portion of N (Fig. 8C,D). This implies that a significant fraction of the endocytosed N is uncleaved and, therefore, that the Dsh-mediated change in localization occurs independently of ligand binding or γ -secretase cleavage.

Rab5 has also been found to inhibit the ectopic N signaling caused by increased levels of the Deltex (Dx) E3 ubiquitin ligase (Hori et al., 2004; Matsuno et al., 2002). The effect of Dx is complex, as it can result in ligand-independent activation or in downregulation of N signaling, depending on the context




Fig. 7. Endocytic control of N signaling in the leg. Endogenous (black arrows/arrowheads) or ectopic (white arrows/arrowheads) joints in the tarsus 4/5 region detected in wild-type (A,C,E,G) and *dsh¹* (B,D,F,H) backgrounds after expression of different endocytic regulators driven by *ap-Gal4*. Arrows are complete joints, arrowheads partial joints. (A,B) Normal and ectopic joints. (C,D) Expression of Rab5 has no effect on wild type (C) but partially suppresses ectopic joints in *dsh¹* (D, white arrowhead; note only legs with ectopic T3 joint were scored for rescue of ectopic T4 joint). (E,F) Expression of Dx elicits ectopic joint in wild type resembling *dsh¹* (E) and fails to modify *dsh¹* (F). (G,H) By contrast, expression of Dx^{NBS} partially inhibits normal joints in wild type (G) and partially suppresses normal and ectopic joint in *dsh¹* (H).

(Mukherjee et al., 2005; Wilkin et al., 2008; Yamada et al., 2011). Both modes of regulation require Rab5-mediated endocytosis of N to early endosomes. To test whether the *dsh¹* phenotype in T4 could be modified by expression of Dx and derivatives, we assayed full-length Dx and mutations affecting some of its functional domains (Matsuno et al., 2002): proline-rich region (Dx^{Δpro}), ring-H2 domain (Dx^{mRZF}) and N-binding region (Dx^{ΔNBS}).

Expression of Dx and Dx^{Δpro} phenocopied the defects caused by PCP mutations (Fig. 7E), arguing that N activity at these ectopic sites might involve modifications to its trafficking, and neither was able to modify the *dsh¹* phenotype (Fig. 7F). Dx^{mRZF} had no effect in wild type or in *dsh¹* (not shown). By contrast, expression of Dx^{ΔNBS} resulted in a striking phenotype of joint fusion (Fig. 7G,H) in both wild type and *dsh¹*, resembling consequences of *dx* null alleles in certain conditions of altered N activity (Gorman and Girton, 1992). Dx^{ΔNBS} thus appears to have a dominant-negative effect, blocking the endogenous T4-T5 joint as well as the ectopic joint in *dsh¹*. This contrasts with the wing, in which Dx^{ΔNBS} exhibits little residual activity (Matsuno et al., 2002) and suggests that the Dx context-dependent effects (Wilkin et al., 2008) are likely to rely on other factors that could be titrated by Dx^{ΔNBS}.

Our results reveal that two endocytic regulators, Rab5 and Dx, can alter N activation at the site of ectopic joint formation, although there are mechanistic differences between them. Overexpression of Rab5 had no effect on wild-type tarsus but could rescue the mutant phenotype of *dsh¹*. This suggests that ectopic N activity in PCP mutants is associated with a change in N trafficking that can be




Fig. 8. Ectopic expression of Dsh::myc in the ap territory. (A) Overexpressed Dsh::myc (anti-myc, purple) appears in a vesicular pattern and colocalizes with N (anti-Necd, green). (B) Some N vesicles (Necd, green), coincide with Rab5-positive puncta (anti-Rab5, purple; e.g. arrows). (C,D) Sub-cellular localization of N (green) is altered in Dsh-expressing cells (purple). Immunofluorescence associated with apical cell membrane is decreased, internal puncta are increased. Similar results are obtained with antibodies against extracellular (C) or intracellular (D) portions of N.

suppressed by Rab5. By contrast, overexpression of Dx induced ectopic joints even in the wild-type background arguing that it is sufficient to overcome repression of N mediated by PCP.

DISCUSSION

Spatially coordinated regulation of signaling pathways is essential to generate correct anatomical and functional structures, as exemplified by the *Drosophila* leg, in which activity of the N pathway is required to specify leg joints (Bishop et al., 1999; de Celis et al., 1998; Rauskolb and Irvine, 1999). In this case, only cells distal to the stripe of Ser expression appear to be capable of responding to the ligand. Here, we show that activity of the core PCP pathway is required in those cells proximal to the domain of Ser expression to prevent them from responding to this N ligand. This regulation correlates with the asymmetric distribution of the core PCP proteins, as we show that Fz and Dsh are enriched at the distal side of each cell, which in the non-responding cells faces the neighboring Ser-expressing cells. Conversely, in those cells distal to Ser, Fz and Dsh are depleted from the proximal side, leaving N free to interact with its ligand to promote joint formation. It appears that elimination of core PCP gene function in cells proximal to the Ser-expressing cells is sufficient to alleviate the repression resulting in ectopic N activity and ectopic joint formation. Such regulation of the membrane availability of Notch could equally affect D1-mediated activation, although Ser appears to be the major ligand responsible in the joints (Bishop et al., 1999). Other factors are likely to influence proximal repression of N because ectopic joints are also observed in alterations of the EGFR pathway (Galindo et al., 2005) and mutants of *defective proventriculus* (Shirai et al., 2007).

We note also that the domains of N activation (both normal and ectopic) extend beyond the cells at the interface with Ser. We have not sought to investigate this additional level of regulation here, but our results indicate that it is unlikely to be due to a secondary signal emanating from the Ser-interfacing cells because the loss of function clones show complete autonomy, without any ‘shadow’ of activation adjacent to the clone. An alternative possibility is that the cells make more extensive contacts, as has been seen in other tissues (Cohen et al., 2010; De Jussineau et al., 2003; Demontis and Dahmann, 2007).

PCP regulation of N has been observed in other developmental processes, most notably in photoreceptor fate choice in the *Drosophila* eye (Cooper and Bray, 1999; del Alamo and Mlodzik, 2006; Fanto and Mlodzik, 1999; Strutt et al., 2002). There, much of the regulation is via effects on levels and activity of the ligand.

However, we detected no change in the pattern of N or Ser expression in PCP mutants. Instead, our evidence suggests that regulation involves direct interaction between Dsh and N and that this interaction has consequences on the endocytic trafficking of N, resulting in its inactivation. The interaction requires the amino-terminal portion of the Dsh DIX domain, which is also required for Axin binding in the canonical Wnt pathway (Julius et al., 2000; Kishida et al., 1999), making it difficult to dissect its role in the PCP-mediated N inhibition. Nevertheless, we were able to generate one mutation that reduced interactions with N with minor consequences on Axin binding. Rescue experiments with this mutant form of Dsh indicated that it was less effective in PCP function in the leg joints compared with others (e.g. polarity of leg bristles). These results support the model that a direct interaction between Dsh and N is relevant in the context of joint determination. However, we cannot fully rule out the possibility that the mutation has more generalized effects on Dsh, if the joints are particularly sensitive to the levels of Dsh activity.

Several studies indicate that endocytic sorting of N is involved in its regulation, with either positive or negative effects depending on the particular context (Fortini and Bilder, 2009; Furthauer and Gonzalez-Gaitan, 2009). Our findings suggest that regulation of N by PCP in the leg is mediated by interaction with Dsh, and probably involves the control of N endocytic trafficking. This suggests a model whereby the interaction between Dsh and N results in increased endocytosis of the N receptor, so reducing its capability to interact with ligands on neighboring cell. Removal of Fz or Dsh compromises this endocytic trafficking, allowing N to be activated. The interaction between Dsh and N is thus only likely to be relevant under circumstances in which there is a strong localization of Dsh co-incident with an interface between N and ligand-expressing cells.

Previous studies have also suggested a role for Dsh in regulating N and on promoting its endocytosis (Axelrod et al., 1996; Munoz-Descalzo et al., 2010). In both instances, these effects were linked to Wg signaling, rather than to the core PCP pathway as here. Nevertheless several aspects are consistent with our results, most notably the direct binding between Dsh and N. Additionally it has been argued that Dsh specifically antagonizes Dx-mediated effects of N (Romain et al., 2001), which is compatible with their complementary effects on joint formation. However, it is also evident that the ability of Dsh to inhibit N depends on the developmental context. For example, whereas overexpression of Dsh in the leg is sufficient to inhibit N activation at presumptive joints, overexpression of Dsh at the wing margin is not sufficient

to repress N signaling: there are no nicks and *cut* expression is not inhibited. Interestingly, differences in Dx behavior are also evident in these two contexts. At the wing margin (Matsuno et al., 2002), Dx^{Δpro} acts as a dominant-negative form of Dx, whereas Dx^{ΔNBS} is inactive. By contrast, in the leg joints Dx^{Δpro} behaves as wild-type Dx, whereas Dx^{ΔNBS} is a dominant negative. We postulate, therefore, that the subcellular localization of Dsh and the availability of Dx are important for determining the regulation of N trafficking at joints.

The autonomous effect of core PCP mutants was clear when we used the *E(spl)mβ1.5-CD2* N reporter and *disco-lacZ*. However, the consequences on *bib-lacZ* were more complex. Although larger clones of mutant cells always exhibited autonomous ectopic expression, similar to *E(spl)mβ1.5-CD2*, some narrow clones exhibited no ectopic expression. We suggest that this might be due to *bib-lacZ* having a higher threshold of response, so it would need stronger N activation. The domain of *bib-lacZ* is narrower than that of the other known reporters, in agreement with this model. Furthermore, we found some cases in which there was a reduction of the normal *bib-lacZ* expression in the mutant cells, in addition to ectopic expression. This suggests that PCP-mediated distal localization of Dsh would be required not only for inhibition of N in proximal cells, but also for efficient activation of N in distal ones.

Acknowledgements

We thank Emma Harrison and Franziska Aurich for technical assistance; Jeff Axelrod for the pCasper-dsh-GFP and for sharing wild-type insertion lines with us; David Strutt for flies; Ken Irvine for the anti-Ser antibody.

Funding

This work was funded by a programme grant from the UK Medical Research Council [G0300034 to S.J.B.]; a Spanish Ministry of Science and Innovation project grant [BFU2009-07949 to M.I.G.]; and a 'Ramón y Cajal' fellowship [to M.I.G.]. M.B. is supported by a Spanish Research Council (CSIC) JAE-Tec contract. Deposited in PMC for release after 6 months.

Competing interests statement

The authors declare no competing financial interests.

Supplementary material

Supplementary material available online at
<http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.077446/-DC1>

References

- Aigouy, B., Farhadifar, R., Staple, D. B., Sagner, A., Roper, J. C., Julicher, F. and Eaton, S. (2010). Cell flow reorients the axis of planar polarity in the wing epithelium of *Drosophila*. *Cell* **142**, 773-786.
- Andersson, E. R., Sandberg, R. and Lendahl, U. (2011). Notch signaling: simplicity in design, versatility in function. *Development* **138**, 3593-3612.
- Axelrod, J. D. (2001). Unipolar membrane association of Dishevelled mediates Frizzled planar cell polarity signaling. *Genes Dev.* **15**, 1182-1187.
- Axelrod, J. D., Matsuno, K., Artavanis-Tsakonas, S. and Perrimon, N. (1996). Interaction between Wingless and Notch signaling pathways mediated by dishevelled. *Science* **271**, 1826-1832.
- Axelrod, J. D., Miller, J. R., Shulman, J. M., Moon, R. T. and Perrimon, N. (1998). Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and Wingless signaling pathways. *Genes Dev.* **12**, 2610-2622.
- Bishop, S. A., Klein, T., Arias, A. M. and Couso, J. P. (1999). Composite signalling from Serrate and Delta establishes leg segments in *Drosophila* through Notch. *Development* **126**, 2993-3003.
- Bray, S. J. (2006). Notch signalling: a simple pathway becomes complex. *Nat. Rev. Mol. Cell Biol.* **7**, 678-689.
- Chae, J., Kim, M. J., Goo, J. H., Collier, S., Gubb, D., Charlton, J., Adler, P. N. and Park, W. J. (1999). The *Drosophila* tissue polarity gene starry night encodes a member of the protocadherin family. *Development* **126**, 5421-5429.
- Chen, W., ten Berge, D., Brown, J., Ahn, S., Hu, L. A., Miller, W. E., Caron, M. G., Barak, L. S., Nusse, R. and Lefkowitz, R. J. (2003). Dishevelled 2 recruits beta-arrestin 2 to mediate Wnt5A-stimulated endocytosis of Frizzled 4. *Science* **301**, 1391-1394.
- Cho, B. and Fischer, J. A. (2011). Ral GTPase promotes asymmetric Notch activation in the *Drosophila* eye in response to Frizzled/PCP signaling by repressing ligand-independent receptor activation. *Development* **138**, 1349-1359.
- Cohen, M., Georgiou, M., Stevenson, N. L., Miodownik, M. and Baum, B. (2010). Dynamic filopodia transmit intermittent Delta-Notch signaling to drive pattern refinement during lateral inhibition. *Dev. Cell* **19**, 78-89.
- Cooper, M. T. and Bray, S. J. (1999). Frizzled regulation of Notch signalling polarizes cell fate in the *Drosophila* eye. *Nature* **397**, 526-530.
- Cooper, M. T., Tyler, D. M., Furriols, M., Chalkiadaki, A., Delidakis, C. and Bray, S. (2000). Spatially restricted factors cooperate with notch in the regulation of Enhancer of split genes. *Dev. Biol.* **221**, 390-403.
- de Celis Ibeas, J. M. and Bray, S. J. (2003). Bowl is required downstream of Notch for elaboration of distal limb patterning. *Development* **130**, 5943-5952.
- de Celis, J. F., Tyler, D. M., de Celis, J. and Bray, S. J. (1998). Notch signalling mediates segmentation of the *Drosophila* leg. *Development* **125**, 4617-4626.
- Das, G., Reynolds-Kenneally, J. and Mlodzik, M. (2002). The atypical cadherin Flamingo links Frizzled and Notch signaling in planar polarity establishment in the *Drosophila* eye. *Dev. Cell* **2**, 655-666.
- De Jossineau, C., Soule, J., Martin, M., Anguille, C., Montcourrier, P. and Alexandre, D. (2003). Delta-promoted filopodia mediate long-range lateral inhibition in *Drosophila*. *Nature* **426**, 555-559.
- del Alamo, D. and Mlodzik, M. (2006). Frizzled/PCP-dependent asymmetric neuralized expression determines R3/R4 fates in the *Drosophila* eye. *Dev. Cell* **11**, 887-894.
- Demontis, F. and Dahmann, C. (2007). Apical and lateral cell protrusions interconnect epithelial cells in live *Drosophila* wing imaginal discs. *Dev. Dyn.* **236**, 3408-3418.
- Djiane, A., Yogev, S. and Mlodzik, M. (2005). The apical determinants aPKC and dPati regulate Frizzled-dependent planar cell polarity in the *Drosophila* eye. *Cell* **121**, 621-631.
- Fanto, M. and Mlodzik, M. (1999). Asymmetric Notch activation specifies photoreceptors R3 and R4 and planar polarity in the *Drosophila* eye. *Nature* **397**, 523-526.
- Feiguin, F., Hannus, M., Mlodzik, M. and Eaton, S. (2001). The ankyrin repeat protein Diego mediates Frizzled-dependent planar polarization. *Dev. Cell* **1**, 93-101.
- Fiuzza, U. M. and Arias, A. M. (2007). Cell and molecular biology of Notch. *J. Endocrinol.* **194**, 459-474.
- Fortini, M. E. (2009). Notch signaling: the core pathway and its posttranslational regulation. *Dev. Cell* **16**, 633-647.
- Fortini, M. E. and Bilder, D. (2009). Endocytic regulation of Notch signaling. *Curr. Opin. Genet. Dev.* **19**, 323-328.
- Frise, E., Knoblich, J. A., Younger-Shepherd, S., Jan, L. Y. and Jan, Y. N. (1996). The *Drosophila* Numb protein inhibits signaling of the Notch receptor during cell-cell interaction in sensory organ lineage. *Proc. Natl. Acad. Sci. USA* **93**, 11925-11932.
- Furthauer, M. and Gonzalez-Gaitan, M. (2009). Endocytic regulation of notch signalling during development. *Traffic* **10**, 792-802.
- Galindo, M. I., Bishop, S. A., Greig, S. and Couso, J. P. (2002). Leg patterning driven by proximal-distal interactions and EGFR signaling. *Science* **297**, 256-259.
- Galindo, M. I., Bishop, S. A. and Couso, J. P. (2005). Dynamic EGFR-Ras signalling in *Drosophila* leg development. *Dev. Dyn.* **233**, 1496-1508.
- Goodrich, L. V. and Strutt, D. (2011). Principles of planar polarity in animal development. *Development* **138**, 1877-1892.
- Gorman, M. J. and Girton, J. R. (1992). A genetic analysis of *deltex* and its interaction with the Notch locus in *Drosophila melanogaster*. *Genetics* **131**, 99-112.
- Gubb, D., Green, C., Huen, D., Coulson, D., Johnson, G., Tree, D., Collier, S. and Roote, J. (1999). The balance between isoforms of the prickle LIM domain protein is critical for planar polarity in *Drosophila* imaginal discs. *Genes Dev.* **13**, 2315-2327.
- Held, L., Duarte, C. and Derakhshanian, K. (1986). Extra tarsal joints and abnormal cuticular polarities in various mutants of *Drosophila melanogaster*. *Roux's Arch. Dev. Biol.* **195**, 145-157.
- Hori, K., Fostier, M., Ito, M., Fuwa, T. J., Go, M. J., Okano, H., Baron, M. and Matsuno, K. (2004). *Drosophila* *deltex* mediates suppressor of Hairless-independent and late-endosomal activation of Notch signaling. *Development* **131**, 5527-5537.
- Jaekel, R. and Klein, T. (2006). The *Drosophila* Notch inhibitor and tumor suppressor gene lethal (2) giant discs encodes a conserved regulator of endosomal trafficking. *Dev. Cell* **11**, 655-669.
- Jones, K. H., Liu, J. and Adler, P. N. (1996). Molecular analysis of EMS-induced frizzled mutations in *Drosophila melanogaster*. *Genetics* **142**, 205-215.
- Julius, M. A., Schelbert, B., Hsu, W., Fitzpatrick, E., Jho, E., Fagotto, F., Costantini, F. and Kitajewski, J. (2000). Domains of axin and dishevelled required for interaction and function in wnt signaling. *Biochem. Biophys. Res. Commun.* **276**, 1162-1169.

- Kishida, S., Yamamoto, H., Hino, S., Ikeda, S., Kishida, M. and Kikuchi, A. (1999). DIX domains of Dvl and axin are necessary for protein interactions and their ability to regulate beta-catenin stability. *Mol. Cell. Biol.* **19**, 4414-4422.
- Klingensmith, J., Nusse, R. and Perrimon, N. (1994). The *Drosophila* segment polarity gene *dishevelled* encodes a novel protein required for response to the wingless signal. *Genes Dev.* **8**, 118-130.
- Lee, H. and Adler, P. N. (2002). The function of the frizzled pathway in the *Drosophila* wing is dependent on inturnd and fuzzy. *Genetics* **160**, 1535-1547.
- MacDonald, B. T., Tamai, K. and He, X. (2009). Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Dev. Cell* **17**, 9-26.
- Matsuno, K., Ito, M., Hori, K., Miyashita, F., Suzuki, S., Kishi, N., Artavanis-Tsakonas, S. and Okano, H. (2002). Involvement of a proline-rich motif and RING-H2 finger of Deltex in the regulation of Notch signaling. *Development* **129**, 1049-1059.
- Mukherjee, A., Veraksa, A., Bauer, A., Rosse, C., Camonis, J. and Artavanis-Tsakonas, S. (2005). Regulation of Notch signalling by non-visual beta-arrestin. *Nat. Cell Biol.* **7**, 1191-1201.
- Munoz-Descalzo, S., Sanders, P. G., Montagne, C., Johnson, R. I., Balayo, T. and Arias, A. M. (2010). Wingless modulates the ligand independent traffic of Notch through Dishevelled. *Fly (Austin)* **4**.
- Nagel, A. C., Krejci, A., Tenin, G., Bravo-Patino, A., Bray, S., Maier, D. and Preiss, A. (2005). Hairless-mediated repression of notch target genes requires the combined activity of Groucho and CtBP corepressors. *Mol. Cell. Biol.* **25**, 10433-10441.
- Povelones, M., Howes, R., Fish, M. and Nusse, R. (2005). Genetic evidence that *Drosophila* frizzled controls planar cell polarity and Armadillo signaling by a common mechanism. *Genetics* **171**, 1643-1654.
- Pueyo, J. I. and Couso, J. P. (2011). Tarsal-less peptides control Notch signalling through the Shavenbaby transcription factor. *Dev. Biol.* **355**, 183-193.
- Ramain, P., Khechumian, K., Seugnet, L., Arbogast, N., Ackermann, C. and Heitzler, P. (2001). Novel Notch alleles reveal a Deltex-dependent pathway repressing neural fate. *Curr. Biol.* **11**, 1729-1738.
- Rauskolb, C. and Irvine, K. D. (1999). Notch-mediated segmentation and growth control of the *Drosophila* leg. *Dev. Biol.* **210**, 339-350.
- Seifert, J. R. and Mlodzik, M. (2007). Frizzled/PCP signalling: a conserved mechanism regulating cell polarity and directed motility. *Nat. Rev. Genet.* **8**, 126-138.
- Shirai, T., Yorimitsu, T., Kiritooshi, N., Matsuzaki, F. and Nakagoshi, H. (2007). Notch signaling relieves the joint-suppressive activity of Defective proventriculus in the *Drosophila* leg. *Dev. Biol.* **312**, 147-156.
- Stenmark, H. (2009). Rab GTPases as coordinators of vesicle traffic. *Nat. Rev. Mol. Cell Biol.* **10**, 513-525.
- Strutt, D. and Strutt, H. (2007). Differential activities of the core planar polarity proteins during *Drosophila* wing patterning. *Dev. Biol.* **302**, 181-194.
- Strutt, D., Johnson, R., Cooper, K. and Bray, S. (2002). Asymmetric localization of frizzled and the determination of Notch-dependent cell fate in the *Drosophila* eye. *Curr. Biol.* **12**, 813-824.
- Strutt, D. I. (2001). Asymmetric localization of frizzled and the establishment of cell polarity in the *Drosophila* wing. *Mol. Cell* **7**, 367-375.
- Strutt, H. and Strutt, D. (2009). Asymmetric localisation of planar polarity proteins: Mechanisms and consequences. *Semin. Cell Dev. Biol.* **20**, 957-963.
- Strutt, H., Warrington, S. J. and Strutt, D. (2011). Dynamics of core planar polarity protein turnover and stable assembly into discrete membrane subdomains. *Dev. Cell* **20**, 511-525.
- Taylor, J., Abramova, N., Charlton, J. and Adler, P. N. (1998). Van Gogh: a new *Drosophila* tissue polarity gene. *Genetics* **150**, 199-210.
- Theisen, H., Purcell, J., Bennett, M., Kansagara, D., Syed, A. and Marsh, J. L. (1994). *dishevelled* is required during wingless signaling to establish both cell polarity and cell identity. *Development* **120**, 347-360.
- Tomlinson, A. and Struhl, G. (1999). Decoding vectorial information from a gradient: sequential roles of the receptors Frizzled and Notch in establishing planar polarity in the *Drosophila* eye. *Development* **126**, 5725-5738.
- Usui, T., Shima, Y., Shimada, Y., Hirano, S., Burgess, R. W., Schwarz, T. L., Takeichi, M. and Uemura, T. (1999). Flamingo, a seven-pass transmembrane cadherin, regulates planar cell polarity under the control of Frizzled. *Cell* **98**, 585-595.
- Vinson, C. R., Conover, S. and Adler, P. N. (1989). A *Drosophila* tissue polarity locus encodes a protein containing seven potential transmembrane domains. *Nature* **338**, 263-264.
- Vladar, E. K., Antic, D. and Axelrod, J. D. (2009). Planar cell polarity signaling: the developing cell's compass. *Cold Spring Harb. Perspect. Biol.* **1**, a002964.
- Wilkin, M., Tongngok, P., Gensch, N., Clemence, S., Motoki, M., Yamada, K., Hori, K., Taniguchi-Kanai, M., Franklin, E., Matsuno, K. et al. (2008). *Drosophila* HOPS and AP-3 complex genes are required for a Deltex-regulated activation of notch in the endosomal trafficking pathway. *Dev. Cell* **15**, 762-772.
- Winter, C. G., Wang, B., Ballew, A., Royou, A., Karess, R., Axelrod, J. D. and Luo, L. (2001). *Drosophila* Rho-associated kinase (Drok) links Frizzled-mediated planar cell polarity signaling to the actin cytoskeleton. *Cell* **105**, 81-91.
- Wolff, T. and Rubin, G. M. (1998). Strabismus, a novel gene that regulates tissue polarity and cell fate decisions in *Drosophila*. *Development* **125**, 1149-1159.
- Wu, J., Jenny, A., Mirkovic, I. and Mlodzik, M. (2008). Frizzled-Dishevelled signaling specificity outcome can be modulated by Diego in *Drosophila*. *Mech. Dev.* **125**, 30-42.
- Xu, T. and Rubin, G. M. (1993). Analysis of genetic mosaics in developing and adult *Drosophila* tissues. *Development* **117**, 1223-1237.
- Yamada, K., Fuwa, T. J., Ayukawa, T., Tanaka, T., Nakamura, A., Wilkin, M. B., Baron, M. and Matsuno, K. (2011). Roles of *Drosophila* *deltex* in Notch receptor endocytic trafficking and activation. *Genes Cells* **16**, 261-272.
- Yu, A., Rual, J. F., Tamai, K., Harada, Y., Vidal, M., He, X. and Kirchhausen, T. (2007). Association of Dishevelled with the clathrin AP-2 adaptor is required for Frizzled endocytosis and planar cell polarity signaling. *Dev. Cell* **12**, 129-141.

Table S1. Sequence of the forward and reverse primers and primer combinations used to generate the fragments that were cloned into pGEX for the pull-down experiments

Forward		Ds h1	Ds h2	Ds h3	Ds h4	Ds h5	Ds h6	Ds h7
dsh5\A	AGGAATTCACCATGGACGCGGACAGGGGC	X		X				
dsh5\B	AGGAATTCACCATGGGCAATCCGCTGCTG		X		X			
dsh5\C	AGGAATTCACCATGCCATCCGCCCAGGTGACG					X		
dsh5\6	AGGAATTCACCATGGACGCCGATTTTCGGT						X	
dsh5\7	AGGAATTCACCATGGGTCTGACCAACAGG							X
Reverse								
dsh3\L	AAGCGGCCCGCCTAAGTGTGCGCCACCCAAGC	X	X			X	X	X
dsh3\S	AAGCGGCCCGCCTACGAGGAGGTGCGCGACATG			X	X			

Table S2. Aligned sequences of the primers used for the site-directed mutagenesis of *dsh*

mut1 (D39S+V43E)	GGTGACGCTGCGAT <u>TCG</u> TTCAAGCTC <u>GAG</u> CTGAACAAGCAG
mut2 (D39S+V43Q)	GGTGACGCTGCGAT <u>TCG</u> TTCAAGCTC <u>GAG</u> CTGAACAAGCAG
mut3 (D39S)	CAGGTGACGCTGCGAT <u>TCG</u> TTCAAGCTGGTGCTG
mut4 (V43E)	CGCGATTTCAAGCTC <u>GAG</u> CTGAACAAGCAGAAC
mut5 (V43E)	CGCGATTTCAAGCTC <u>GAG</u> CTGAACAAGCAGAAC
mut6 (K46V+Q47A)	CAAGCTGGTGCTGAACGTT <u>GCG</u> AACAACAACCTAC

Codons where the mutations were introduced are underlined.