


## Transcriptional regulatory networks in epiblast cells and during anterior neural plate development as modeled in epiblast stem cells

**Makiko Iwafuchi-Doi, Kazunari Matsuda, Kazuhiro Murakami, Hitoshi Niwa, Paul J. Tesar, Jun Aruga, Isao Matsuo and Hisato Kondoh**

There was an error published in *Development* **139**, 3926-3937.

Fig. S6 was incorrect. The correct Fig. S6, as shown below, now appears in the supplementary data.

We apologise to the authors and readers for this mistake.



Development 139, 3926–3937 (2012) doi:10.1242/dev.085936  
 © 2012. Published by The Company of Biologists Ltd

# Transcriptional regulatory networks in epiblast cells and during anterior neural plate development as modeled in epiblast stem cells

Makiko Iwafuchi-Doi<sup>1,\*</sup>, Kazunari Matsuda<sup>1</sup>, Kazuhiro Murakami<sup>2,†</sup>, Hitoshi Niwa<sup>2</sup>, Paul J. Tesar<sup>3</sup>, Jun Aruga<sup>4</sup>, Isao Matsuo<sup>5</sup> and Hisato Kondoh<sup>1,§</sup>

## SUMMARY

Somatic development initiates from the epiblast in post-implantation mammalian embryos. Recent establishment of epiblast stem cell (EpiSC) lines has opened up new avenues of investigation of the mechanisms that regulate the epiblast state and initiate lineage-specific somatic development. Here, we investigated the role of cell-intrinsic core transcriptional regulation in the epiblast and during derivation of the anterior neural plate (ANP) using a mouse EpiSC model. Cells that developed from EpiSCs in one day in the absence of extrinsic signals were found to represent the ANP of ~E7.5 embryos. We focused on transcription factors that are uniformly expressed in the E6.5 epiblast but in a localized fashion within or external to the ANP at E7.5, as these are likely to regulate the epiblast state and ANP development depending on their balance. Analyses of the effects of knockdown and overexpression of these factors in EpiSCs on the levels of downstream transcription factors identified the following regulatory functions: cross-regulation among *Zic*, *Otx2*, *Sox2* and *Pou* factors stabilizes the epiblastic state; *Zic*, *Otx2* and *Pou* factors in combination repress mesodermal development; *Zic* and *Sox2* factors repress endodermal development; and *Otx2* represses posterior neural plate development. All of these factors variably activate genes responsible for neural plate development. The direct interaction of these factors with enhancers of *Otx2*, *Hesx1* and *Sox2* genes was demonstrated. Thus, a combination of regulatory processes that suppresses non-ANP lineages and promotes neural plate development determines the ANP.

**KEY WORDS:** Epiblast stem cells, Anterior neural plate, *Zic* factors, *Pou* factors, *Otx2*, *Sox2*

## INTRODUCTION

The epiblast serves as the primordium of all somatic lineages in amniotes, where the earliest derivative is the anterior neural plate (ANP). Despite profound interest in the process by which somatic lineages are generated, cell-intrinsic and transcription factor-dependent regulatory mechanisms remain poorly understood, primarily because of the difficulty in accessing the epiblast in post-implantation mammalian embryos. The cell-extrinsic mechanisms that regulate the fate of the epiblast are better understood. It has been shown, for example, that Nodal antagonists secreted from the anterior visceral endoderm disrupt Nodal signaling, which otherwise stabilizes the epiblast state and elicits ANP development (Camus et al., 2006; Perea-Gomez et al., 2002).

The ANP and the posterior neural plate (PNP) are generated via different mechanisms. The ANP is derived directly from the epiblast (Iwafuchi-Doi et al., 2011), whereas the development of the PNP from the epiblast passes through an intermediate state of axial stem cells, which are common precursors for the PNP and

paraxial mesoderm (Kondoh and Takemoto, 2012; Takemoto et al., 2011; Tzouanacou et al., 2009).

Recent success in establishing cell lines directly from the egg cylinder epiblast [epiblast stem cells (EpiSCs)] (Brons et al., 2007; Tesar et al., 2007) has opened up new avenues to investigate the cell-intrinsic mechanisms in the epiblast and its derivatives. The epiblast state of EpiSCs is maintained by activin (Nodal substitute) and Fgf2 signaling, and the interruption of these pathways elicits the development of neural plate cells (NPCs), partly mimicking the action of Nodal antagonists during the derivation of the ANP in embryos.

In our present study, we first characterized EpiSCs and their immediately derived NPCs under culture conditions without supply of extrinsic signals. The expression profiles of transcription factor genes under these conditions were compared with those in embryonic tissues. Our findings indicate that the immediate derivatives of EpiSCs under the NPC culture conditions represent the ANP cell state in ~E7.5 embryos. Taking advantage of this observation, we then systematically analyzed the effects on downstream genes of the knockdown and overexpression in EpiSCs of several transcription factors.

These analyses of the transcriptional regulatory networks indicated intricate cross-regulation among the factors to stabilize the epiblast state and to derive the ANP. During the derivation of ANP, inhibitory mechanisms were found to play key roles to suppress the development of mesodermal, endodermal and PNP cells, confining the developmental pathway of epiblast derivatives to ANP. This analysis also highlights the crucial involvement of *Zic2/3*, *Otx2* and *Sox2* in these processes. This study thus highlights the advantages of using EpiSCs to examine in detail the regulatory mechanisms that underlie the derivation of somatic lineages.

<sup>1</sup>Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita, Osaka 565-0871, Japan. <sup>2</sup>RIKEN Center for Developmental Biology (CDB), Chuo-ku, Kobe, Hyogo 650-0047, Japan. <sup>3</sup>Department of Genetics and Center for Stem Cell and Regenerative Medicine, Case Western Reserve University School of Medicine, Cleveland, OH 44106, USA. <sup>4</sup>RIKEN Brain Science Institute, Wako, Saitama 351-0198, Japan. <sup>5</sup>Osaka Medical Center and Research Institute for Maternal and Child Health, Murodo-cho, Izumi, Osaka 594-1101, Japan.

\*Present address: Department of Cell and Developmental Biology, University of Pennsylvania School of Medicine, Philadelphia, PA 19104, USA

†Present address: The Wellcome Trust/Cancer Research UK Gurdon Institute, University of Cambridge, Tennis Court Road, Cambridge CB2 1QN, UK

§Author for correspondence (kondohh@fbs.osaka-u.ac.jp)

## MATERIALS AND METHODS

### EpiSC cultures

EpiSCs (Tesar et al., 2007) were cultured under two different sets of conditions: one for epiblastic state maintenance involving the supply of 20 ng/ml activin and 10 ng/ml Fgf2 in N2B27 medium on fibronectin-coated dishes; and another for the promotion of NPC development without the addition of growth factors on gelatin-coated dishes (Iwafuchi-Doi et al., 2011).

### qRT-PCR analysis

Total RNA was extracted from the epiblastic (inner) layer of egg cylinder stage mouse embryos and EpiSCs, and processed for qRT-PCR analysis as described previously (Iwafuchi-Doi et al., 2011). The mRNA levels relative to *Gapdh* were quantified based on standard curves using cloned cDNA sequences. Primer sequences are listed in supplementary material Table S1.

### Microarray analysis

Total RNAs extracted from duplicate epiblastic, NP1 and NP2 cultures were analyzed using an Agilent SurePrint G3 Mouse GE 8×60k Microarray. The normalized data using Agilent Feature Extraction were averaged for duplicate samples, and the data for transcription factor genes, selected by GO filtering and itemized inspection, were analyzed using Excel 2010 (Microsoft) functions. The original data are deposited in the NCBI GEO database with accession number GSE38085.

### Knockdown and overexpression of transcription factor genes

shRNA vectors were constructed in pSilencerU6puro (Ambion) using the sequences listed in supplementary material Table S2. For knockdown analysis, 300 ng shRNA vector DNA was complexed with Lipofectamine 2000 (Invitrogen), added to a suspension of  $4 \times 10^5$  dissociated EpiSCs, and the mixture plated in a well of a 12-well dish (Falcon 3043). Similarly, for overexpression analysis, 50 ng pCAGGS-based expression vector (Sawicki et al., 1998) for each transcription factor was mixed with 300 ng control pSilencerU6puro and used for transfection. After 8 hours, puromycin was added at 5 µg/ml to select transfected cells, and after 24 hours RNAs were extracted for analysis.

### Transactivation assays using 10T1/2 fibroblasts

10T1/2 fibroblasts in a 24-well dish were transfected using Lipofectamine 2000 with DNA mixtures containing (per well) enhancer-δ51-Luciferase constructed in pGL4.10 (120 ng; Promega), pRL-Tk-Renilla luciferase (20 ng; Promega), and varying amounts of pCAGGS-based transcription factor expression vector. The following mouse gene enhancers were used: *Otx2* AN (Kurokawa et al., 2004), *Hesx1* ANP (Spieler et al., 2004) and *Sox2* N2[73bp]<sub>2</sub> (Iwafuchi-Doi et al., 2011). Luciferase activity was measured using the Dual Luciferase Assay Kit (Promega).

### Production of *Otx2*<sup>-/-</sup> embryos transgenic for enhancer N2-lacZ

The enhancer N2-lacZ transgene (Iwafuchi-Doi et al., 2011) was injected into fertilized mouse eggs derived from *Otx2*<sup>+/-</sup> parents, and the genotypes of the implants were determined by PCR analysis of yolk sac DNA.

### Other procedures

Immunofluorescence staining, the enhancer activity assay in EpiSCs and NPCs, and electrophoretic mobility shift assay (EMSA) were performed as described previously (Iwafuchi-Doi et al., 2011).

## RESULTS

### Dynamic changes in transcription factor gene expression during derivation of NPCs from the epiblast in embryos and from EpiSCs in culture


To derive NPCs from EpiSCs, we adopted a culture condition without the supply of exogenous signaling factors, which was employed for neural tissue derivation from embryonic stem (ES) cells in monolayer culturing (Ying et al., 2003). Under this condition, Pax6-immunopositive cells and TuJ1 (Tubb3)-positive cells appeared at NP3 (3 days in the NPC culture condition), and the proportion of these cells increased at NP4 (supplementary

material Fig. S1), indicating progression of neural development. To characterize the initial process by which the NPCs are derived from EpiSCs, we investigated the expression profiles of a panel of transcription factor genes expressed in EpiSCs under the epiblast-maintenance condition and under the NPC condition for 1 (NP1) and 2 (NP2) days using qRT-PCR. In parallel, epiblasts from E6.5 egg cylinders and from the inner cell layer of the anterior and posterior halves of E7.5 embryos were collected (Fig. 1A) and analyzed for the purposes of comparison. The genes included in the panels characterize the epiblast and specific domains of the neural plate (Fig. 1B); references for the expression patterns of these genes in mouse embryos from E6.5 to E8.5 are listed in supplementary material Table S3. *Fgf5* was included in the analysis as a hallmark of the epiblast state, although it is not a transcription factor gene. Transcript levels of the genes were quantified using a standard curve method and are shown relative to  $10^{-3}$  of the *Gapdh* level.

The expression levels of *Fgf5* and *Pou5f1*, which both characterize the epiblast state, are similar between E6.5 epiblast and EpiSC and decrease similarly in the E7.5 inner layers and NP1 cells (Fig. 1C,E). This indicated that EpiSCs in the epiblast maintenance condition share major characteristics of the E6.5 epiblast, and that E7.5 inner layer and NP1 cells have similarly departed the epiblast state. The decrease in the expression of other genes characteristic of epiblast, *Eomes* and *Nanog*, in E7.5 inner layers and NP1 cells (Fig. 1C) also supports this notion, although some differences in the net expression levels might partly reflect differences between the states of cells in embryos and in culture. The NP1 cells were also characterized by activation of *Sip1*, which is characteristic of the neural plate (Chng et al., 2010; Miyoshi et al., 2006). In addition, transcription factor genes that are expressed at high levels exclusively in the ANP (as represented by anterior inner layer), i.e. *Otx2*, *Sox2* and *Pou3f1* (see also Fig. 2, below), were strongly expressed in NP1 cells (Fig. 1D,E). These observations indicate that epiblastic EpiSCs and NP1 cells to a large extent represent the epiblast and ANP states in embryos, respectively. The strong expression of *Sox1* (Uchikawa et al., 2011; Wood and Episkopou, 1999), *Pou3f2* (Bouchard et al., 2005) and *Otx1* (Suda et al., 1999) in the neural plate is known to occur in embryos after E8.0. Here, these genes were activated only in NP2 cells, indicating that NP1 cells represent the neural plate in embryos prior to E8.0 (Fig. 1D-F).

The gradual decrease of *Pou5f1* expression in NP1 and NP2 cells and the steep decrease in *Pou3f1* expression in NP2 cells are consistent with the expression profiles documented for embryonic ANP (Perea-Gómez et al., 1999; Zwart et al., 1996). Genes characteristic of the ANP (*Hesx1*, *Otx1* and *Pax6*) were activated in the NP1/2 cells, whereas expression of genes characteristic of the PNP (*Gbx2* and *Nkx2.1*) remained very low (Fig. 1F,G). Moreover, the expression of brachyury (*T*) and *Sox17*, which mark mesodermal and endodermal precursors, respectively, was strongly inhibited (Fig. 1G). These observations indicate that EpiSCs develop mostly into the ANP under the NPC culture condition.

The NPC culture condition differed from that used for epiblast maintenance by the absence of activin and Fgf2 signaling and by the coating of the culture substrate with gelatin rather than fibronectin. Evaluation of the contribution of these differences to the promotion of ANP development (supplementary material Fig. S2) indicated that the effects of the culture conditions are synergistic, without a simple relationship with a single signaling system.



**Fig. 1. qRT-PCR analysis of transcription factor gene expression in the epiblast and during neural plate derivation.** (A) The regions of the mouse embryo isolated after removal of the endodermal layer and used in the analysis. Scale bar: 200  $\mu$ m. (B) The axial expression domains of transcription factors in the ANP and PNP after E8.0, aligned to E8.0 mouse embryos hybridized for *Sox2* transcripts as shown in lateral (top) and dorsal (bottom) views. (C–G) Gene expression levels (relative to  $10^{-3}$  *Gapdh*) determined by qRT-PCR are indicated for the embryonic regions indicated in A, epiblastic EpiSCs (Epi) and NP1 and NP2 cells ( $n \geq 3$ ). The genes are loosely grouped according to their expression specificity or gene family. Error bars indicate s.e. ANP, anterior neural plate; PNP, posterior neural plate.


### Microarray analysis of transcription factor expression profiles during NPC derivation

Using the same RNAs as used for the qRT-PCR analysis shown in Fig. 1, the expression profiles of a broader range of transcription factor genes were investigated using microarrays. After exclusion

of genes that gave only background signals, including most Hox genes, the expression profiles of 994 transcription factor genes were analyzed during the progression from epiblastic to NP1 and NP2 cells. The results, as summarized in Table 1 and detailed in supplementary material Table S4, were generally consistent with

**Table 1. Classification of transcription factor genes according to microarray expression profiles in epiblastic, NP1 and NP2 cells**

Profile class	Expression profiles			No. assigned among 994 genes	Representative genes
	Epiblastic	NP1	NP2		
A				35	<i>Hesx1, Sip1, Irx3</i>
B				42	<i>Zic2, Gli2</i>
C				74	<i>Nr6a1, Sox1, Sox2, Hes5, Gbx2</i>
D				23	<i>Dmbx1, Sall1</i>
E				290	<i>Otx2, Nanog, Pou3f2, Pbx1</i>
F				265	<i>Sox3, Pou3f4, Zfp521, Pbx2</i>
G				126	<i>Snai2, Gli1, Hes1</i>
H				18	<i>Pou5f1, Mycn, Myc</i>
I				40	<i>Eomes, Klf5</i>
J				31	<i>Sox17, Foxa1</i>
K				18	<i>T, Foxa2, Gata6, Gsc</i>
L				13	<i>Sox15, Pou3f1</i>
M				19	<i>Foxb1, Zfxh2</i>



the qRT-PCR data for the genes presented in Fig. 1. Two-thirds (681) of the genes maintained their expression levels, i.e. any changes did not exceed 2-fold, as compared with the epiblastic state (Table 1, classes E-G), whereas the remaining genes were either activated or downregulated to greater extents. The majority of genes that were strongly activated in NP1 and/or NP2 cells (classes A-D) were associated with neural plate development. The genes that were activated strongly in NP1 cells included *Irx3*, which is activated at E7.5 in the ANP (Houweling et al., 2001), and also those indicated in the qRT-PCR analyses (*Hesx1*, *Sip1*, *Zic2*; Fig. 1). A significant proportion of the genes that were strongly downregulated in NP1 and NP2 cells (classes I-K) were representative of genes involved in endodermal (*Eomes*, *Sox17*, *Foxa1*, *Foxa2*, *Gata6*) or primitive streak-dependent mesodermal (*T*, *Gata6*, *Gsc*) development.

*Nr6a1* (*Gcnf*) plays a major role in the repression of *Pou5f1* during the epiblast-neural plate stages of embryo development (Fuhrmann et al., 2001) and in the retinoic acid-induced differentiation of ES cells (Gu et al., 2005). *Nr6a1* was upregulated during the development of EpiSCs to NP2 cells (class C), accounting for the parallel decrease in *Pou5f1* expression (Fig. 1).

Class L genes, as exemplified by *Sox15*, were sharply repressed in NP2 cells, whereas class M genes, as exemplified by *Foxb1*, were maintained at low expression levels in NP1 and strongly activated in NP2 cells. Strong downregulation of the major Myc genes *Mycn* and *Myc* (class H) was also observed in NP2 cells. These observations indicate the progression of neural plate development from NP1 (<E8.0 ANP) to the less proliferative NP2 (>E8.0 ANP) cells.

### Regional expression of transcription factors in embryos and EpiSC cultures

We noted that *Sox2*, *Pou3f1* and *Otx2* expressed from the E6.5 epiblast stage developed a clear anterior dominance at E7.5 (Fig. 1D,E), consistent with published histological data (Ang et al., 1994; Iwafuchi-Doi et al., 2011; Wood and Episkopou, 1999; Zwart

et al., 1996). It has been reported that *Pou5f1* is downregulated in an inverse spatial relationship with the above three genes (Perea-Gómez et al., 1999). To confirm the expression profiles at the individual cell level, embryos at E6.5 and ~E7.5, as well as EpiSCs under both epiblast and NPC culturing conditions, were immunostained for these transcription factors.


In E6.5 embryos, *Sox2*, *Pou5f1*, *Pou3f1* and *Otx2* proteins were expressed uniformly throughout the epiblast nuclei (Fig. 2A). However, at ~E7.5, a high level of *Sox2* was observed in the nuclei of the developing ANP (Fig. 2Ba), paralleled by a reduction in *Pou5f1* (Fig. 2Bb) and by high expression of *Pou3f1* (Fig. 2Bc) and *Otx2* (Fig. 2Bd) in the same ANP domain.

In the epiblast state, the majority of EpiSC nuclei expressed *Sox2*, *Pou3f1* and *Otx2* with *Pou5f1*, analogous to the E6.5 epiblast (Fig. 2C). However, NP1 cells strongly expressed *Pou3f1* and *Otx2* at the periphery of their colonies (Fig. 2Dbe). Conversely, *Pou5f1* was downregulated at the periphery of colonies but remained highly expressed in the medial portion, with an inverse spatial relationship with *Pou3f1* and *Otx2* (Fig. 2Dcd). This medial high *Pou5f1* expression was reduced in NP2 cells (data not shown). *Sox2* expression also showed a moderate peripheral enhancement (Fig. 2Da). Thus, the development of NPCs from EpiSCs, which initiates at the periphery of the colonies, shares many features with the development of the ANP in embryos.

### Experimental design for transcription factor manipulation in EpiSCs

We reasoned that the transcription factors that are initially expressed throughout the epiblast but shift their expression sites at the beginning of ANP development might play crucial roles in the maintenance of the epiblastic state and in the initiation of ANP development, subject to their balance. We selected *Sox2*, *Pou5f1*, *Pou3f1* and *Otx2* (Fig. 2), as well as *Nanog* and *Zic2/3*, as candidate transcription factors with such functions. *Nanog* is known to stabilize the epiblastic state (Mitsui et al., 2003; Vallier et al., 2009) and to be downregulated in the forming ANP (Hart et





**Fig. 2. Immunofluorescent detection of transcription factors in egg cylinder embryos and EpiSCs.** (A,B) Mouse embryos at (A) E6.5 and (B) E7.5-7.75 were stained for (a) Sox2, (b) Pou5f1, (c) Pou3f1 and (d) Otx2. Whole-mount view (left) and optical sections through the medial plane (right). The ANP regions are outlined in the optical sections. (C) Epiblastic EpiSCs doubly stained for Pou5f1 and (a) Sox2, (b) Pou3f1 or (c) Otx2. (D) Immunofluorescence images of NP1 cells stained for (a) Sox2, (b) Pou3f1, (c) Pou5f1 (the same field as b), (d) a merge of b and c, and (e) Otx2. Scale bars: 100  $\mu$ m.

al., 2004). The involvement of *Zic* factors in the early stages of neural development was first indicated in *Xenopus* (Mizuseki et al., 1998; Nakata et al., 1997) and supported by *Zic3* mutant mouse phenotypes (Inoue et al., 2007; Ware et al., 2006). We expected that knockdown and overexpression of these transcription factors in the EpiSCs would strongly affect the expression of immediate downstream genes, and that analysis of these effects, with the rationale shown in supplementary material Fig. S4, would reveal the transcriptional regulatory networks that regulate epiblast and ANP states.

Expression vectors for shRNAs (supplementary material Table S2) or pCAGGS-based expression vectors for the transcription factor genes were transfected into epiblastic cells, and transfected cells selected by puromycin (supplementary material Table S5). The effects on other genes of manipulating transcription factor levels were evaluated by qRT-PCR after 24 hours. The knockdown of transcription factor genes typically reduced their expression to ~30-40% of normal levels (supplementary material Table S6). Overexpression using pCAGGS-based vectors was very efficient, often reaching levels 100-fold greater than endogenous expression (supplementary material Table S6). However, lower expression levels were sufficient to affect downstream genes (see below; supplementary material Fig. S3).

### The impact of single transcription factor manipulations in epiblastic cells

The results of the transcription factor knockdown and overexpression under the epiblast stabilized culture condition are shown in Fig. 3 and supplementary material Fig. S5. In cases indicative of activation or repression of a gene according to the criteria given in supplementary material Fig. S4, areas are highlighted in yellow or pink, respectively.

Genes characteristic of epiblasts (*Fgf5*, *Eomes* and *Nanog*) were activated by *Zic2/3*. *Fgf5* and *Eomes* were also activated by *Otx2*, indicating analogous regulation. *Fgf5* was also activated by *Pou3f1*, and *Nanog* was activated by *Zic2/3* and *Sox2* rather than by *Otx2*.


The regulation of *Sox2*, *Otx2* and *Pou5f1* suggested cross-regulation in the maintenance of the epiblast state as well as in the promotion of ANP development. *Otx2* and *Sox2* were activated by

*Zic2/3*, and *Otx2* was also activated by *Sox2*. These regulations might maintain the epiblast state in cooperation with *Pou5f1* and promote ANP development once *Pou5f1* is downregulated. It is interesting to note that *Otx2* was activated by *Pou5f1*, whereas *Pou5f1* expression was repressed by overexpression of *Otx2*. This negative-feedback loop might prevent an excess of *Pou5f1* in the epiblast that otherwise would promote neural development through the activation of a set of neural plate genes (*Hesx1*, *Pax6*, *Gbx2* and *Pou3f2/3*). The repression of *Pou5f1* by *Otx2* and *Sox2* might also reinforce the *Nr6a1*-dependent repression of *Pou5f1* (Fuhrmann et al., 2001) during ANP development (Table 1).

The regulation of *Sox1*, *Sip1* and *Otx1*, which are all strongly activated in the neural plate after E8.0 and in NP2 cells (Fig. 1), was similar and involved moderate activation by *Zic2/3* and *Pou5f1/3f1*, and also by *Otx2* in the case of *Sip1* and *Otx1*. Among the *Pou3f* factor genes, *Pou3f1* was uniquely activated by *Zic2/3*, *Sox2* and *Nanog*, which is likely to reflect its strong expression in NP1 cells and downregulation in NP2 cells. The other three *Pou3f* genes responded variably (supplementary material Fig. S5). The regulation of *Zic3* and *Zic2* differed: *Zic3* was activated by *Sox2*, *Pou5f1* and *Nanog*, whereas *Zic2* was activated only by *Pou5f1*.

Regulation of the ANP genes *Hesx1* and *Pax6* was characterized by their activation by *Otx2* and *Pou5f1/3f1*. *Pax6* was also activated by *Sox2* and *Nanog*. This suggested that *Otx2* function is crucial for the neural plate to gain anterior characteristics. The regulation of two genes characteristic of PNP, *Gbx2* and *Nkx1.2*, showed interesting differences. *Gbx2* was strongly activated by *Sox2* and to lesser extent by *Pou5f1/3f1*. By contrast, *Nkx1.2* was activated by *Zic2/3* and *Pou5f1* and repressed by *Otx2* and *Nanog*. The *Otx2*-dependent repression of PNP-specific genes will be elaborated below.

Interestingly, the *T* gene, which represents mesoderm precursors, was strongly repressed by *Otx2* and also inhibited by *Zic2/3* and *Pou5f1/3f1*, showing opposite responses to many of the genes characteristic of neural plate described above. A low level of exogenous *Otx2* expression was sufficient to exert efficient repression of *T* (supplementary material Fig. S3). *Sox17*, which is characteristic of endoderm precursors, was activated by *Otx2* and



**Fig. 3. Effects of knockdown and overexpression of transcription factors on their target genes under the epiblast maintenance culture condition.** Impact of knockdown (K) or overexpression (O) of *Zic2/3* (Z), *Otx2* (O2), *Sox2* (S), *Pou5f1* (P5), *Pou3f1* (P3) and *Nanog* (N) on the expression of the indicated genes as determined by qRT-PCR, compared with control (C) cultures. These effects were judged to have activated (yellow) or repressed (pink) a gene in accordance with the criteria shown in supplementary material Fig. S4. As the effects of *Pou3f1* knockdown on downstream genes were minimal in the initial survey, only overexpression data are shown for *Pou3f1* in order to distinguish between the common and unique functions of *Pou5f1* and *Pou3f1*. Data for genes found to be under analogous regulation are bound by a rectangle. Error bars indicate s.e.

repressed by *Sox2*, and possibly by a high level of *Zic2/3*. The expression of *Zfp521* (Kamiya et al., 2011) was unaffected by these transcription factor manipulations (supplementary material Fig. S5).

Thus, in many cases, *Zic2/3* activity contributed crucially to the activation of epiblast- and neural plate-specific genes and to the repression of the mesodermal gene *T* and endodermal gene *Sox17*. *Sox2* activity was strongly antagonistic to the endodermal genes. A strong association between *Otx2* activity and the activation of ANP genes was also noted. Overexpression of *Pou3f1* elicited analogous effects to that of *Pou5f1*. These and further regulatory interactions are compiled in Table 2 and are classified into lineage regulation and neural plate development categories (see below, Fig. 5).

### Multiple transcription factor manipulations in the epiblastic state

To investigate the possible synergistic effects of altering transcription factor levels, we combined *Pou5f1* knockdown with overexpression of *Otx2* and *Sox2* in the epiblast maintenance culture condition (Fig. 4A) as a means of mimicking the transcription factor changes that take place at the anteriormost part of the E7.0-7.5 embryonic inner layer (Fig. 2A). In most cases, the effects of these combined changes were not synergistic, displaying either one of the effects or their averages (supplementary material Fig. S6). However, in several interesting cases, synergistic or antagonistic effects of changing transcription factor levels were observed.

**Table 2. The regulatory targets of individual transcription factors in cell lineage regulation and neural plate development as deduced from knockdown and overexpression effects**

Transcription factor	Cell lineage regulation	Target genes	Neural plate development	Target genes
Zic2/3	Supporting epiblastic state <b>Inhibition of mesodermal state</b>	<i>Fgf5, Eomes, Nanog, Otx2, Sox2</i> <b>T</b>	Promotion of ANP Promotion of >E8 neural plate	<i>Otx2, Pou3f1, Sox2</i> <i>Sox1, Sip1, Otx1, Gbx2*, Nkx1.2*</i>
'High expression'	<b>Inhibition of endodermal state</b>	<b>Sox17</b>		
Otx2	Supporting epiblastic state Promotion of endodermal state <b>Inhibition of mesodermal state</b>	<i>Fgf5, Eomes, Sox17</i> <b>T</b>	Promotion of ANP Promotion of >E8 neural plate <b>Inhibition of PNP</b>	<i>Hesx1, Pax6, Otx1, Sox2*</i> <i>Sip1</i> <b>Gbx2, Nkx1.2, Sox1*</b>
'High expression'	<b>Preventing excess Pou5f1</b>	<b>Pou5f1</b>	Destabilizing epiblastic state	<i>Pou5f1</i>
Sox2	Supporting epiblastic state <b>Inhibition of endodermal state</b>	<i>Nanog, Zic3, Otx2</i> <b>Sox17, Eomes</b>	Promotion of ANP Promotion of PNP	<i>Otx2, Pou3f1, Pax6</i> <i>Gbx2</i>
'High expression'	<b>Preventing excess Pou5f1</b>	<b>Pou5f1</b>	Destabilizing epiblastic state	<i>Pou5f1</i>
Pou5f1/3f1	<b>Inhibition of mesodermal state</b>	<b>T</b>	Enhancement of some neural plate traits	<i>Hesx1, Pax6, Gbx2</i>
Pou5f1	Supporting epiblastic state	<i>Otx2, Zic2, Zic3, Sox2*</i>	Inhibition of ANP Enhancement of some neural plate traits	<i>Otx1</i> <i>Pou3f2, Pou3f3</i>
Nanog	Supporting epiblastic state	<i>Zic3, Sox3, Pou3f1</i>	Enhancement of some neural plate traits	<i>Pou3f1, Pax6</i>

Inhibitory effects contributing to the ANP state are indicated in bold.

\*Target genes identified by knockdown experiments using NP1 cells (Fig. 4B).

†Regulatory interactions indicated by enhancer analysis (Fig. 6).

*Otx1* expression was augmented by knockdown of *Pou5f1* alone, and further activated by combined overexpression of *Otx2* and *Sox2* (Fig. 4A). *Sox17* was activated by *Otx2* overexpression (Fig. 3), and this was enhanced by combination with *Pou5f1* knockdown. However, this activation of *Sox17* was completely repressed by exogenous *Sox2* expression (Fig. 4A). Similarly, *Eomes*, which is strongly expressed in epiblast cells (Fig. 1) and is implicated in endodermal development (Teo et al., 2011), was activated further by exogenous *Otx2*, but this *Otx2*-dependent activation was cancelled by the coexpression of exogenous *Sox2* (Fig. 5A). These observations indicate that, whereas *Otx2* promotes endodermal development by activating relevant transcription factor genes, *Sox2* strongly antagonizes and overrides this *Otx2* effect.

As indicated above (Fig. 3, supplementary material Fig. S5), *Gbx2*, which is characteristic of PNP, was strongly activated by *Sox2*. However, this *Sox2*-dependent activation was totally suppressed by coexpression of *Otx2* (Fig. 4A). Another PNP gene, *Nkx1.2*, was moderately repressed by *Otx2*, and this effect was enhanced by a combined *Pou5f1* knockdown. These results indicate that *Sox2* expression generally supports neural plate development, regardless of whether it is the ANP or PNP. However, *Otx2* inhibits the posterior characteristics and allows only the development of the ANP while also promoting anterior characteristics such as *Hesx1* expression.

### Transcription factor manipulations that promote ANP development

In the above experiments, specific transcription factor levels were manipulated under epiblast maintenance culture conditions. If the observed regulatory interactions participate in ANP derivation, their proper combination would be expected to accelerate ANP development under the NPC culture condition. To characterize the gene regulatory network in NP1 cells, we individually knocked down *Zic2/3* and *Otx2* in these cells. The knockdown effects of these factors were similar to those in the epiblastic state for many genes, as exemplified by *Fgf5*, *Eomes* and *T* (Fig. 4B), indicating a similarity of the core gene regulatory network between the epiblastic and ANP states. However, *Sox1*, *Nkx1.2* and *Gbx2*,

which are activated in the PNP of embryos after E8.0, responded more sharply to *Zic2/3* knockdown by reducing their expression and to *Otx2* knockdown by enhancing their expression (Fig. 4B). These observations confirmed the involvement of *Zic2/3* in the activation of neural plate genes and the activity of *Otx2* as an antagonist of PNP-dedicated genes.


We next tested the effects of *Otx2* overexpression, *Sox2* overexpression, *Pou5f1* knockdown and combinations of these conditions in NP1 cells. The impact of *Sox2* overexpression alone or in combination with a *Pou5f1* knockdown was no greater than that of single factor manipulations (data not shown). However, the combination of *Otx2* overexpression and *Pou5f1* knockdown synergized their effects, either promoting ANP-specific gene expression or inhibiting PNP-specific genes (Fig. 4C). Expression of *Hesx1* and *Pax6*, which was moderately activated in control NP1 cells, and that of *Otx1*, which was activated only at the NP2 stage (Fig. 1F), was significantly augmented by the combination of *Otx2* overexpression and *Pou5f1* knockdown, in an additive fashion for *Hesx1* and *Otx1* and cooperatively for *Pax6* (Fig. 4C). By contrast, the PNP gene *Nkx1.2* was downregulated by *Otx2* overexpression and *Pou5f1* knockdown. The *Otx2*-dependent repression of *Gbx2* was also confirmed (Fig. 4C).

Immunostaining of NP1 cells with *Pou5f1* knockdown and *Otx2* overexpression confirmed augmented *Pax6* expression in the majority of cells (Fig. 4D), which otherwise occurs only in NP3 cells in control cultures (supplementary material Fig. S1). As the strong expression of *Otx1* and *Pax6* in the ANP occurs only in mouse embryos after E8.0 (Fig. 1), the above observations indicated that the combination of *Pou5f1* knockdown and *Otx2* overexpression accelerated the development of the EpiSC-derived ANP cells to reach stages beyond E8.0 in 1 day of culturing.

### Direct activation of ANP-associated transcription factor gene enhancers by core transcription factors

We sought to establish mechanistic links between the effects of the transcription factor manipulations and the action of the transcription factors themselves on their target genes. The enhancer





**Fig. 4. Effects of combined manipulations of transcription factor levels in epiblastic and NP1 cells.** (A) Effects of overexpression of *Otx2* (O2), *Sox2* (S) or both (O2/S) or in combination with *Pou5f1* knockdown (KD-P5) in epiblastic cells, in comparison with levels in control (–) cells. (B) The effects of *Zic2/3* knockdown (KD-Z) and *Otx2* knockdown (KD-O2) on the expression of six representative transcription factor genes in NP1 cells. The effects on *Fgf5*, *Eomes* and *brachyury* (*T*) were similar to those in the epiblastic state (Fig. 3). However, genes expressed in the PNP of embryos after E8 (*Sox1*, *Gbx2* and *Nkx1.2*) clearly responded to these transcription factor manipulations in NP1 cells, which was not observed in the epiblastic state. (C) Effects of a combination of *Otx2* overexpression and *Pou5f1* knockdown in NP1 cells. (D) Pax6 immunostaining of NP1 control cultures (a) and cultures with *Pou5f1* knockdown and *Otx2* overexpression (b). Nuclei were counterstained with DAPI. Error bars indicate s.e. Scale bar: 100  $\mu$ m.

sequences of the *Otx2*, *Hesx1* and *Sox2* genes that have been shown to be active in the embryonic ANP were cloned in a luciferase reporter vector for transactivation assays. 10T1/2 mouse embryo fibroblasts were chosen as the host cells, as the endogenous expression levels are negligible for those transcription factors to be expressed exogenously (supplementary material Table S7).


The *Otx2* ANP enhancer (Kurokawa et al., 2004) containing a Pou binding site and a putative Zic binding site (supplementary material Fig. S7A) was activated by *Pou3f1* and more strongly by *Pou5f1*, and this activation was augmented further by the coexpression of *Zic2* (supplementary material Fig. S7A). This is consistent with the effects of *Zic2/3* knockdown and Pou factor overexpression on *Otx2* (Fig. 3). The *Hesx1* ANP enhancer, which is located immediately upstream of the promoter and contains three essential *Otx2* binding sites (Chou et al., 2006; Spieler et al., 2004), was activated by exogenous *Otx2* (supplementary material Fig. S7B).

Activation of the N2 core enhancer of the *Sox2* gene depends on the binding of Pou factors to their bipartite binding sites in both epiblast and ANP (Iwafuchi-Doi et al., 2011). However, neither *Pou5f1* nor *Pou3f1* alone activated this enhancer (see below, Fig. 6C). We identified additional binding sites for Zic and Otx factors

in the N2 core sequence (Fig. 6A, supplementary material Fig. S8). Mutations of the Zic site (Mut-ZIC) or Pou sites (Mut-POU) inactivated N2 core enhancer activity in both epiblastic and NP1 cells (Fig. 6A). By contrast, mutation of the Otx factor binding site (Mut-OTX) attenuated N2 enhancer activity only in NP1 cells (Fig. 6Ac) and not in epiblastic cells (Fig. 6Ab), suggesting that *Otx2* activity is essential for N2 enhancer activation in ANP cells only.

That *Otx2* is not required for N2 enhancer activation and N2-dependent *Sox2* expression in the epiblast (E6.75) was confirmed using an *Otx2*<sup>−/−</sup> mutant (Matsuo et al., 1995). When an enhancer N2-*lacZ* transgene was introduced into *Otx2*<sup>−/−</sup> embryos, comparable expression of N2-*lacZ* and *Sox2* at levels equivalent to those in wild-type embryos was observed in the epiblast, whereas N2 activation was totally lost in E7.75 *Otx2*<sup>−/−</sup> embryos, as was *Sox2* expression (Fig. 6B).

We hypothesized that a change in the major Pou factors from *Pou5f1* to the Pou3fs during the derivation of ANP from the epiblast caused the emergence of *Otx2* dependence. To test this model, we transfected 10T1/2 fibroblasts with a luciferase reporter vector carrying the N2 core enhancer (N2[73bp]<sub>2</sub>) together with expression vectors for the transcription factors *Zic2*, *Otx2* and one



**Fig. 5. Regulation of developmental pathways by core transcription factors via their target genes in the EpiSC model.**

Some of these regulatory pathways have also been indicated by studies using ES cells: inhibition of *Sox17* by *Zic2/3* and activation of *Zic3* by *Sox2*, *Pou5f1* and *Nanog* (Lim et al., 2007), and the promotion of neural development by sustained high-level expression of *Pou5f1* (Shimozaki et al., 2003). The function of *Pou5f1* and *Nanog* in stabilizing the epiblastic state has been more clearly indicated in earlier studies (Mitsui et al., 2003; Vallier et al., 2009).

of the Pou factors (*Pou5f1*, *Pou3f1* or *Pou3f4*). None of the Pou factors activated the enhancer (Fig. 6Ca), nor did coexpression of *Otx2* with a Pou factor elicit a substantial level of enhancer activation (Fig. 6Cb). However, when *Zic2* was coexpressed, *Pou5f1* significantly activated the N2 core enhancer, whereas this was not observed with *Pou3f1/4* (Fig. 6Cc). When *Zic2* and *Otx2* were coexpressed, all Pou factors activated the N2 core enhancer efficiently, even at low Pou expression levels (Fig. 6Cd). The activation by *Pou5f1* plus *Zic2* was further augmented by coexpression of *Otx2*. *Zic3* displayed similar effects on N2 enhancer activation (supplementary material Fig. S9).

These results, taken together with our observations of mutant enhancers (Fig. 6A), indicated a model in which (1) the cooperative action of *Zic2/3* is essential for Pou factor-dependent activation of the N2 core enhancer and (2) *Otx2* action is also essential for Pou3f-dependent N2 core enhancer activation in the ANP, but may be dispensable for *Pou5f1*-dependent activation in the epiblast (Fig. 6Ce). This mechanism of N2 enhancer regulation sustains *Sox2* expression during the developmental stages at which the major Pou factor function is shifted from *Pou5f1* to the Pou3fs.

## DISCUSSION

### The core transcriptional networks in the epiblast and during ANP derivation

The establishment of EpiSCs from embryonic epiblast has opened up new avenues for the study of core transcriptional networks that maintain the epiblast state and/or give rise to various somatic lineages. Because the expression profiles of transcription factor genes (Fig. 1, Table 1) indicated that epiblastic and NP1 states of EpiSCs in our culture conditions mimicked ~E6.5 epiblast and

~E7.5 ANP, respectively, we focused our analysis on the epiblast and ANP derivation.


We systematically knocked down or overexpressed several transcription factor genes in the epiblastic or ANP derivation states of EpiSCs and quantitatively analyzed the immediate impact on the expression of downstream transcription factor genes. The genes subjected to manipulation were those that are expressed uniformly in the E6.5 epiblast but shift their expression sites into ANP (*Sox2*, *Otx2* and *Pou3f1*) or that avoid ANP (*Pou5f1* and *Nanog*). *Zic2/3* were also included in the analysis because of their involvement in the early phase of neural plate development (Mizuseki et al., 1998; Nakata et al., 1997). This analysis of short-term effects of altered transcription factor levels should be distinguished from the long-term effects of stably overexpressing exogenous genes using human ES cells and mouse EpiSCs (e.g. Chng et al., 2010; Vallier et al., 2009).

The data shown in Figs 3 and 4 and in supplementary material Figs S6 and S7 and as summarized in Table 2 and Fig. 5 highlight the pivotal roles of *Zic2/3*, *Otx2* and *Sox2* in the epiblast state and in ANP derivation. The data also suggest intricate interactions and cross-regulation among these factors. For instance, *Zic2/3* and *Otx2* support the expression of genes characteristic of epiblast such as *Fgf5* and *Eomes*, whereas *Otx2* and *Sox2* overexpression inhibits *Pou5f1* expression. The function of *Otx2* and *Sox2* in repressing *Pou5f1* expression might be to accelerate the reduction of *Pou5f1* expression triggered by *Nr6a1* activation (Fuhrmann et al., 2001), resulting in the clearance of *Pou5f1* expression from the embryonic ANP (Fig. 2). However, overexpression of *Pou5f1* elicited activation of neural plate-associated genes despite the fact that the normal function of *Pou5f1* is to stabilize the epiblast state. It is thus possible that an appropriate level of *Pou5f1* for the epiblast state is maintained by negative feedback through *Otx2* and *Sox2*.

A remarkable finding is that both the repressive and activating mechanisms function together to generate only ANP from the epiblast (Fig. 5). The repressive mechanisms inhibited mesodermal, endodermal and PNP development to confine the developmental pathway to ANP. *Zic2/3*, *Otx2* and *Pou5f1/3f1* inhibited mesodermal development by repressing *T*. *Zic2/3* and *Sox2* inhibited endodermal development by repressing *Sox17* and *Eomes*. These anti-endodermal effects of *Sox2* dominated over the endoderm-promoting effects of *Otx2* on *Sox17* and *Eomes* (Fig. 4A). Moreover, *Otx2* activity strongly inhibited the development of the PNP transcription factor genes *Nkx1.2*, *Gbx2* and *Sox1* (Fig. 3, Fig. 4A,B).

Along with participating in the repressive mechanisms as described above, *Zic2/3*, *Otx2*, *Sox2* and *Pou3f1* also participate in the activation of various genes characteristic of the neural plate, as exemplified by *Sip1* (Chng et al., 2010; Miyoshi et al., 2006). Among these factors, *Otx2* was found to have a prominent role in the promotion of ANP development. *Hesx1*, which determines the anteriormost ANP subdomain (Andoniadou et al., 2007), is directly activated by *Otx2* through its interaction with the ANP enhancer (Spieler et al., 2004) (supplementary material Fig. S7B). Moreover, *Otx2* overexpression in combination with *Pou5f1* knockdown elicited the accelerated development of a fully Pax6-positive ANP cell population in a 1-day period under the NPC culture condition (Fig. 4D), demonstrating the lead function of *Otx2* in ANP development.

In addition, *Otx2* plays a unique role in sustaining *Sox2* expression during the transitory stages from the epiblast to ANP, when the major Pou function switches from *Pou5f1* to Pou3fs (Fig.



**Fig. 6. Differential regulation of the Sox2 N2 enhancer in the epiblast and ANP.** (A) (a) Transcription factor binding sites in the Sox2 N2 enhancer core sequence as confirmed by EMSA (Iwafuchi-Doi et al., 2011) (supplementary material Fig. S8) and mutation of these binding sites. (b,c) Effects of the binding site mutations on N2 activity as assessed in a luciferase reporter assay in transfected epiblastic (b) and NP1 (c) cells. —, luciferase gene without an enhancer; WT, 176 bp N2 enhancer. Mutations of the POU-2/POU-3 sites also inactivated the enhancer in a manner analogous to the POU-1 site mutation investigated here (Iwafuchi-Doi et al., 2011). (B) Expression of Sox2 (in situ hybridization) and N2-lacZ activity (positive cases out of total indicated in parentheses) in the epiblast at E6.75 were similar in *Otx2*<sup>-/-</sup> and wild-type (*Otx2*<sup>+/+</sup>) embryos (a) but absent from *Otx2*<sup>-/-</sup> at E7.75 (b) (compare black and open arrowheads). Scale bars: 200  $\mu$ m. (C) Transactivation of the N2 core enhancer by the combined action of Zic2, Otx2 and Pou factors in 10T1/2 fibroblasts. N2[73bp]<sub>2</sub>-luciferase expression levels relative to those without exogenous factors are shown. (a) Pou factors only; (b) with Otx2 expression; (c) with Zic2 expression; and (d) with coexpression of Otx2 and Zic2. (e) Model for the differential requirement of Otx2 in the activation of the Sox2 N2 enhancer. Error bars indicate s.e.

6C). In contrast to Pou5f1, Pou3f factors plus Zic2/3 cannot activate the N2 enhancer of *Sox2* without the cooperation of Otx2.

In summary, the following regulatory functions were identified: cross-regulation among Zic, Otx2, Sox2 and Pou factors stabilizes the epiblastic state; Zic, Otx2 and Pou factors in combination repress mesodermal development; Zic and Sox2 factors repress endodermal development; and Otx2 represses PNP development. All of these factors variably activate genes responsible for neural plate development, along with participating in the above-mentioned repression mechanisms.

### In vivo evidence for the deduced functions of Otx2, Zic2/3 and Sox2

The regulatory networks identified in this study using EpiSCs account for some earlier observations in post-implantation

embryos. The repression of *T* by Otx2 is novel but accounts for the previous observation that *T* is upregulated in *Otx2*<sup>-/-</sup> embryos (Kimura et al., 2000). In *Otx2*<sup>-/-</sup>  $\leftrightarrow$  wild-type chimeras, where the visceral endoderm defects of *Otx2*<sup>-/-</sup> embryos were rescued (Rhinn et al., 1999; Rhinn et al., 1998), the ANP tissue that consisted of *Otx2*<sup>-/-</sup> cells somehow acquired morphological features of anterior neural fold at E7.5 but expressed PNP genes after E8, never activating *Hesx1*. This observation is accounted for by the simultaneous loss of PNP inhibition and ANP activation, which are both attributed to Otx2 (Fig. 5).

Consistent with the repression of *Gbx2* by Otx2, the competitive interaction between Otx2 and *Gbx2* in the determination of ANP and PNP territories has been demonstrated by chicken embryo electroporation (Katahira et al., 2000) and in compound knockout mouse embryos (Martinez-Barbera et al., 2001).



The contribution of *Zic2/3* functions to neural plate development has not been fully established using mouse embryos owing to their functional overlap and to the lack of a *Zic2* null allele. However, among the variable phenotypes previously reported for *Zic3* null embryos, the most severe phenotype was in fact characterized by serious neural plate defects (Inoue et al., 2007; Ware et al., 2006).

In contrast to its anti-endodermal activity, Sox2 did not interfere with the expression of the hallmark mesodermal gene *T* (Fig. 3, supplementary material Fig. S6). This observation is likely to reflect the fact that the majority of PNP cells are not directly derived from the epiblast but through neural/mesodermal bipotential intermediates known as axial stem cells, which reside in the caudal lateral epiblast abutting the primitive streak (Kondoh and Takemoto, 2012; Takemoto et al., 2011). The caudal lateral epiblast in chicken embryos and in late streak stage mouse embryos coexpresses low levels of *Sox2* and *T*, consistent with the bipotential nature of axial stem cells (Delfino-Machin et al., 2005; Kondoh and Takemoto, 2012; Perantoni et al., 2005).

Thus, the model summarized in Fig. 5 should provide a new guide for understanding various in vivo phenomena associated with cell lineage selection in post-gastrulation embryos.

### Advantages of EpiSCs in the study of somatic lineage derivation

ES cells are frequently employed as a model system for studying somatic cell derivation. However, to derive a somatic cell lineage with ES cells, the cells must go through the epiblast state, as evidenced by the transient expression of *Fgf5* prior to the activation of genes indicative of somatic development (Kamiya et al., 2011; Thomson et al., 2011). Therefore, the effects of gene manipulations in ES cells can be indirect or ambiguous in terms of determining active stages.

Our present study poses EpiSCs as a more suitable alternative. As EpiSCs serve as the immediate precursor for somatic cells of various lineages, investigations of the short-term effects of manipulating genes will reveal the first-step mechanisms in the derivation of the somatic lineages. Not confined to the epiblast or ANP, as reported in this paper, EpiSCs are useful in the study of various cell lineages. Indeed, it has been demonstrated that EpiSCs can develop into virtually all somatic lineages with appropriate manipulation of culture conditions (Brons et al., 2007; Tesar et al., 2007). In the context of PNP development, it would be interesting if neural/mesodermal bipotential axial stem cells could be produced from EpiSCs.

EpiSCs under various culture conditions can give rise to a critical mass of intermediate state cells during cell lineage derivation. These EpiSC-derived cells can be subjected to a comprehensive analysis of gene regulatory networks, which is not an easy task to undertake in embryonic tissues. In the same way that ES cells have contributed greatly to elucidating transcriptional regulatory networks in pre-implantation embryos (e.g. Masui et al., 2007; Shimosato et al., 2007), EpiSCs will prove to be an excellent tool with which to investigate the regulation of somatic lineage derivation in post-implantation embryos.

### Acknowledgements

We thank Jitsutaro Kawaguchi for technical guidance and members of the H.K. laboratory for discussions.

### Funding

P.J.T. is a Robertson Investigator of the New York Stem Cell Foundation. This study was supported by Grants-in-Aid for Scientific Research from MEXT, Japan [17107005, 22247035 and 231147090 to H.K.].

### Competing interests statement

The authors declare no competing financial interests.

### Supplementary material


Supplementary material available online at <http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.085936/-/DC1>

### References


- Andoniadou, C. L., Signore, M., Sajedi, E., Gaston-Massuet, C., Kelberman, D., Burns, A. J., Itasaki, N., Dattani, M. and Martinez-Barbera, J. P. (2007). Lack of the murine homeobox gene *Hesx1* leads to a posterior transformation of the anterior forebrain. *Development* **134**, 1499-1508.
- Ang, S. L., Conlon, R. A., Jin, O. and Rossant, J. (1994). Positive and negative signals from mesoderm regulate the expression of mouse *Otx2* in ectoderm explants. *Development* **120**, 2979-2989.
- Bouchard, M., Grote, D., Craven, S. E., Sun, Q., Steinlein, P. and Busslinger, M. (2005). Identification of Pax2-regulated genes by expression profiling of the mid-hindbrain organizer region. *Development* **132**, 2633-2643.
- Briata, P., Ilengo, C., Bobola, N. and Corte, G. (1999). Binding properties of the human homeodomain protein OTX2 to a DNA target sequence. *FEBS Lett.* **445**, 160-164.
- Brons, I. G., Smithers, L. E., Trotter, M. W., Rugg-Gunn, P., Sun, B., Chuva de Sousa Lopes, S. M., Howlett, S. K., Clarkson, A., Ahrlund-Richter, L., Pedersen, R. A. et al. (2007). Derivation of pluripotent epiblast stem cells from mammalian embryos. *Nature* **448**, 191-195.
- Camus, A., Perea-Gomez, A., Moreau, A. and Collignon, J. (2006). Absence of Nodal signaling promotes precocious neural differentiation in the mouse embryo. *Dev. Biol.* **295**, 743-755.
- Chng, Z., Teo, A., Pedersen, R. A. and Vallier, L. (2010). SIP1 mediates cell-fate decisions between neuroectoderm and mesendoderm in human pluripotent stem cells. *Cell Stem Cell* **6**, 59-70.
- Chou, S. J., Hermes, E., Hatta, T., Feltner, D., El-Hodiri, H. M., Jamrich, M. and Mahon, K. (2006). Conserved regulatory elements establish the dynamic expression of *Rpx/Hesx1* in early vertebrate development. *Dev. Biol.* **292**, 533-545.
- Ciruna, B. G. and Rossant, J. (1999). Expression of the T-box gene *Eomesodermin* during early mouse development. *Mech. Dev.* **81**, 199-203.
- Delfino-Machin, M., Lunn, J. S., Breitkreuz, D. N., Akai, J. and Storey, K. G. (2005). Specification and maintenance of the spinal cord stem zone. *Development* **132**, 4273-4283.
- Elms, P., Scurry, A., Davies, J., Willoughby, C., Hacker, T., Bogani, D. and Arkell, R. (2004). Overlapping and distinct expression domains of *Zic2* and *Zic3* during mouse gastrulation. *Gene Expr. Patterns* **4**, 505-511.
- Fuhrmann, G., Chung, A. C., Jackson, K. J., Hummelke, G., Baniahmad, A., Sutter, J., Sylvester, I., Schöler, H. R. and Cooney, A. J. (2001). Mouse germline restriction of *Oct4* expression by germ cell nuclear factor. *Dev. Cell* **1**, 377-387.
- Greber, B., Wu, G., Bernemann, C., Joo, J. Y., Han, D. W., Ko, K., Tapia, N., Sabour, D., Sterneckert, J., Tesar, P. and Schöler, H. R. (2010). Conserved and divergent roles of FGF signaling in mouse epiblast stem cells and human embryonic stem cells. *Cell Stem Cell* **6**, 215-226.
- Gu, P., LeMenuet, D., Chung, A. C., Mancini, M., Wheeler, D. A. and Cooney, A. J. (2005). Orphan nuclear receptor GCNF is required for the repression of pluripotency genes during retinoic acid-induced embryonic stem cell differentiation. *Mol. Cell. Biol.* **25**, 8507-8519.
- Hancock, S. N., Agulnik, S. I., Silver, L. M. and Papaioannou, V. E. (1999). Mapping and expression analysis of the mouse ortholog of *Xenopus Eomesodermin*. *Mech. Dev.* **81**, 205-208.
- Hart, A. H., Hartley, L., Ibrahim, M. and Robb, L. (2004). Identification, cloning and expression analysis of the pluripotency promoting *Nanog* genes in mouse and human. *Dev. Dyn.* **230**, 187-198.
- Hebert, J. M., Boyle, M. and Martin, G. R. (1991). mRNA localization studies suggest that murine FGF-5 plays a role in gastrulation. *Development* **112**, 407-415.
- Houweling, A. C., Dildrop, R., Peters, T., Mummehoff, J., Moorman, A. F., Rüther, U. and Christoffels, V. M. (2001). Gene and cluster-specific expression of the Iroquois family members during mouse development. *Mech. Dev.* **107**, 169-174.
- Inoue, T., Nakamura, S. and Osumi, N. (2000). Fate mapping of the mouse prosencephalic neural plate. *Dev. Biol.* **219**, 373-383.
- Inoue, T., Ota, M., Mikoshiba, K. and Aruga, J. (2007). *Zic2* and *Zic3* synergistically control neurulation and segmentation of paraxial mesoderm in mouse embryo. *Dev. Biol.* **306**, 669-684.
- Ivanova, N., Dobrin, R., Lu, R., Kutenko, I., Levorse, J., DeCoste, C., Schafer, X., Lun, Y. and Lemischka, I. R. (2006). Dissecting self-renewal in stem cells with RNA interference. *Nature* **442**, 533-538.
- Iwafuchi-Doi, M., Yoshida, Y., Onichtchouk, D., Leichenring, M., Driever, W., Takemoto, T., Uchikawa, M., Kamachi, Y. and Kondoh, H. (2011). The *Pou5f1/Pou3f*-dependent but *Sox2*-independent regulation of conserved enhancer *N2* initiates *Sox2* expression during epiblast to neural plate stages in vertebrates. *Dev. Biol.* **352**, 354-366.




- Kamiya, D., Banno, S., Sasai, N., Ohgushi, M., Inomata, H., Watanabe, K., Kawada, M., Yakura, R., Kiyonari, H., Nakao, K. et al. (2011). Intrinsic transition of embryonic stem-cell differentiation into neural progenitors. *Nature* **470**, 503-509.
- Kanai-Azuma, M., Kanai, Y., Gad, J. M., Tajima, Y., Taya, C., Kurohmaru, M., Sanai, Y., Yonekawa, H., Yazaki, K., Tam, P. P. et al. (2002). Depletion of definitive gut endoderm in Sox17-null mutant mice. *Development* **129**, 2367-2379.
- Katahira, T., Sato, T., Sugiyama, S., Okafuji, T., Araki, I., Funahashi, J. and Nakamura, H. (2000). Interaction between Otx2 and Gbx2 defines the organizing center for the optic tectum. *Mech. Dev.* **91**, 43-52.
- Kimura, C., Yoshinaga, K., Tian, E., Suzuki, M., Aizawa, S. and Matsuo, I. (2000). Visceral endoderm mediates forebrain development by suppressing posteriorizing signals. *Dev. Biol.* **225**, 304-321.
- Kondoh, H. and Takemoto, T. (2012). Axial stem cells deriving both posterior neural and mesodermal tissues during gastrulation. *Curr. Opin. Genet. Dev.* **22**, 374-380.
- Kurokawa, D., Takasaki, N., Kiyonari, H., Nakayama, R., Kimura-Yoshida, C., Matsuo, I. and Aizawa, S. (2004). Regulation of Otx2 expression and its functions in mouse epiblast and anterior neuroectoderm. *Development* **131**, 3307-3317.
- Lim, L. S., Loh, Y. H., Zhang, W., Li, Y., Chen, X., Wang, Y., Bakre, M., Ng, H. H. and Stanton, L. W. (2007). Zic3 is required for maintenance of pluripotency in embryonic stem cells. *Mol. Biol. Cell* **18**, 1348-1358.
- Martinez-Barbera, J. P., Signore, M., Boyle, P. P., Puelles, E., Acampora, D., Gogoi, R., Schubert, F., Lumsden, A. and Simeone, A. (2001). Regionalisation of anterior neuroectoderm and its competence in responding to forebrain and midbrain inducing activities depend on mutual antagonism between OTX2 and GBX2. *Development* **128**, 4789-4800.
- Masui, S., Nakatake, Y., Toyooka, Y., Shimosato, D., Yagi, R., Takahashi, K., Okochi, H., Okuda, A., Matoba, R., Sharov, A. A. et al. (2007). Pluripotency governed by Sox2 via regulation of Oct3/4 expression in mouse embryonic stem cells. *Nat. Cell Biol.* **9**, 625-635.
- Matsuo, I., Kuratani, S., Kimura, C., Takeda, N. and Aizawa, S. (1995). Mouse Otx2 functions in the formation and patterning of rostral head. *Genes Dev.* **9**, 2646-2658.
- Mitsui, K., Tokuzawa, Y., Itoh, H., Segawa, K., Murakami, M., Takahashi, K., Maruyama, M., Maeda, M. and Yamanaka, S. (2003). The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell* **113**, 631-642.
- Miyoshi, T., Maruhashi, M., Van De Putte, T., Kondoh, H., Huylebroeck, D. and Higashi, Y. (2006). Complementary expression pattern of Zfhx1 genes Sip1 and deltaEF1 in the mouse embryo and their genetic interaction revealed by compound mutants. *Dev. Dyn.* **235**, 1941-1952.
- Mizuseki, K., Kishi, M., Matsui, M., Nakanishi, S. and Sasai, Y. (1998). Xenopus Zic-related-1 and Sox-2, two factors induced by chordin, have distinct activities in the initiation of neural induction. *Development* **125**, 579-587.
- Nakata, K., Nagai, T., Aruga, J. and Mikoshiba, K. (1997). Xenopus Zic3, a primary regulator both in neural and neural crest development. *Proc. Natl. Acad. Sci. USA* **94**, 11980-11985.
- Perantoni, A. O., Timofeeva, O., Naillat, F., Richman, C., Pajni-Underwood, S., Wilson, C., Vainio, S., Dove, L. F. and Lewandoski, M. (2005). Inactivation of FGF8 in early mesoderm reveals an essential role in kidney development. *Development* **132**, 3859-3871.
- Perea-Gómez, A., Shawlot, W., Sasaki, H., Behringer, R. R. and Ang, S. (1999). HNF3beta and Lim1 interact in the visceral endoderm to regulate primitive streak formation and anterior-posterior polarity in the mouse embryo. *Development* **126**, 4499-4511.
- Perea-Gomez, A., Vella, F. D., Shawlot, W., Oulad-Abdelghani, M., Chazaud, C., Meno, C., Pfister, V., Chen, L., Robertson, E., Hamada, H. et al. (2002). Nodal antagonists in the anterior visceral endoderm prevent the formation of multiple primitive streaks. *Dev. Cell* **3**, 745-756.
- Rhinn, M., Dierich, A., Shawlot, W., Behringer, R. R., Le Meur, M. and Ang, S. L. (1998). Sequential roles for Otx2 in visceral endoderm and neuroectoderm for forebrain and midbrain induction and specification. *Development* **125**, 845-856.
- Rhinn, M., Dierich, A., Le Meur, M. and Ang, S. (1999). Cell autonomous and non-cell autonomous functions of Otx2 in patterning the rostral brain. *Development* **126**, 4295-4304.
- Sakurai, Y., Kurokawa, D., Kiyonari, H., Kajikawa, E., Suda, Y. and Aizawa, S. (2010). Otx2 and Otx1 protect diencephalon and mesencephalon from caudalization into metencephalon during early brain regionalization. *Dev. Biol.* **347**, 392-403.
- Sawicki, J. A., Morris, R. J., Monks, B., Sakai, K. and Miyazaki, J. (1998). A composite CMV-IE enhancer/beta-actin promoter is ubiquitously expressed in mouse cutaneous epithelium. *Exp. Cell Res.* **244**, 367-369.
- Shimosato, D., Shiki, M. and Niwa, H. (2007). Extra-embryonic endoderm cells derived from ES cells induced by GATA factors acquire the character of XEN cells. *BMC Dev. Biol.* **7**, 80.
- Shimozaki, K., Nakashima, K., Niwa, H. and Taga, T. (2003). Involvement of Oct3/4 in the enhancement of neuronal differentiation of ES cells in neurogenesis-inducing cultures. *Development* **130**, 2505-2512.
- Spieler, D., Bäumer, N., Stebler, J., Köprunner, M., Reichman-Fried, M., Teichmann, U., Raz, E., Kessel, M. and Witter, L. (2004). Involvement of Pax6 and Otx2 in the forebrain-specific regulation of the vertebrate homeobox gene ANF/Hesx1. *Dev. Biol.* **269**, 567-579.
- Suda, Y., Nakabayashi, J., Matsuo, I. and Aizawa, S. (1999). Functional equivalency between Otx2 and Otx1 in development of the rostral head. *Development* **126**, 743-757.
- Takemoto, T., Uchikawa, M., Yoshida, M., Bell, D. M., Lovell-Badge, R., Papaioannou, V. E. and Kondoh, H. (2011). Tbx6-dependent Sox2 regulation determines neural or mesodermal fate in axial stem cells. *Nature* **470**, 394-398.
- Tamplin, O. J., Kinzel, D., Cox, B. J., Bell, C. E., Rossant, J. and Lickert, H. (2008). Microarray analysis of Foxa2 mutant mouse embryos reveals novel gene expression and inductive roles for the gastrula organizer and its derivatives. *BMC Genomics* **9**, 511.
- Teo, A. K., Arnold, S. J., Trotter, M. W., Brown, S., Ang, L. T., Chng, Z., Robertson, E. J., Dunn, N. R. and Vallier, L. (2011). Pluripotency factors regulate definitive endoderm specification through eomesodermin. *Genes Dev.* **25**, 238-250.
- Tesar, P. J., Chenoweth, J. G., Brook, F. A., Davies, T. J., Evans, E. P., Mack, D. L., Gardner, R. L. and McKay, R. D. (2007). New cell lines from mouse epiblast share defining features with human embryonic stem cells. *Nature* **448**, 196-199.
- Thomson, M., Liu, S. J., Zou, L. N., Smith, Z., Meissner, A. and Ramanathan, S. (2011). Pluripotency factors in embryonic stem cells regulate differentiation into germ layers. *Cell* **145**, 875-889.
- Tzouanacou, E., Wegener, A., Wymeersch, F. J., Wilson, V. and Nicolas, J. F. (2009). Redefining the progression of lineage segregations during mammalian embryogenesis by clonal analysis. *Dev. Cell* **17**, 365-376.
- Uchikawa, M., Yoshida, M., Iwafuchi-Doi, M., Matsuda, K., Ishida, Y., Takemoto, T. and Kondoh, H. (2011). B1 and B2 Sox gene expression during neural plate development in chicken and mouse embryos: universal versus species-dependent features. *Dev. Growth Differ.* **53**, 761-771.
- Vallier, L., Mendjan, S., Brown, S., Chng, Z., Teo, A., Smithers, L. E., Trotter, M. W., Cho, C. H., Martinez, A., Rugg-Gunn, P. et al. (2009). Activin/Nodal signalling maintains pluripotency by controlling Nanog expression. *Development* **136**, 1339-1349.
- Wang, Z. X., Teh, C. H., Kueh, J. L., Lufkin, T., Robson, P. and Stanton, L. W. (2007). Oct4 and Sox2 directly regulate expression of another pluripotency transcription factor, Zfp206, in embryonic stem cells. *J. Biol. Chem.* **282**, 12822-12830.
- Ware, S. M., Harutyunyan, K. G. and Belmont, J. W. (2006). Zic3 is critical for early embryonic patterning during gastrulation. *Dev. Dyn.* **235**, 776-785.
- Waters, S. T., Wilson, C. P. and Lewandoski, M. (2003). Cloning and embryonic expression analysis of the mouse Gbx1 gene. *Gene Expr. Patterns* **3**, 313-317.
- Wilkinson, D. G., Bhatt, S. and Herrmann, B. G. (1990). Expression pattern of the mouse T gene and its role in mesoderm formation. *Nature* **343**, 657-659.
- Wood, H. B. and Episkopou, V. (1999). Comparative expression of the mouse Sox1, Sox2 and Sox3 genes from pre-gastrulation to early somite stages. *Mech. Dev.* **86**, 197-201.
- Yang, Y. P. and Klingensmith, J. (2006). Roles of organizer factors and BMP antagonism in mammalian forebrain establishment. *Dev. Biol.* **296**, 458-475.
- Ying, Q. L., Stavridis, M., Griffiths, D., Li, M. and Smith, A. (2003). Conversion of embryonic stem cells into neuroectodermal precursors in adherent monoculture. *Nat. Biotechnol.* **21**, 183-186.
- Zeineddine, D., Papadimou, E., Chebli, K., Gineste, M., Liu, J., Grey, C., Thurig, S., Behfar, A., Wallace, V. A., Skerjanc, I. S. et al. (2006). Oct-3/4 dose dependently regulates specification of embryonic stem cells toward a cardiac lineage and early heart development. *Dev. Cell* **11**, 535-546.
- Zwart, R., Broos, L., Grosveld, G. and Meijer, D. (1996). The restricted expression pattern of the POU factor Oct-6 during early development of the mouse nervous system. *Mech. Dev.* **54**, 185-194.



**Fig. S1. Immunohistological analysis of NP2 to NP4 cells for expression of Sox2 and Pax6 (double staining) and Otx2 and TuJ1.** Sox2 continued to be expressed in the majority of cells, whereas Otx2 began to be downregulated at NP3, in an inverse relationship with the appearance of Pax6-immunopositive and TuJ1-positive NP3 cells and the increase in NP4 cells. The TuJ1 (neuron-specific class III  $\beta$ -tubulin) staining pattern indicated a wider extension of the neurites in NP4 cells compared with NP3 cells. Scale bars: 100  $\mu$ m.




**Fig. S2. Synergistic contributions of the removal of growth factors and changing the culture substrate coating to the promotion of ANP cell development from EpiSCs.** The complete culture medium contained Fgf2, activin and the culture substrate was coated with fibronectin. The expression levels of genes under various culture conditions, as determined by qRT-PCR, are shown relative to those in EpiSCs grown in complete culture medium. The effects of the removal of Fgf2 on the epiblast state were modest, in contrast to the strong downregulation of *Fgf5*, *Pou5f1*, *brachyury (T)*, *Sox17* and *Nanog* after the removal of activin. However, downregulation of *Pou5f1*, *T* and *Sox17* was significantly enhanced by the simultaneous removal of Fgf2 and activin. *Sox2* and *Sip1* were strongly activated by the removal of Fgf2 and activin together. The activation of *Otx2* and *Zic2*, which will be shown to be crucial for ANP development in a later section, was promoted by altering the substrate coating from fibronectin to gelatin.




**Fig. S3. Responses of the indicated genes to varying amounts of transfected Sox2 and Otx2 expression vectors.** Expression levels were determined by qRT-PCR and are shown relative to the endogenous levels in EpiSCs. Exogenous *Sox2* and *Otx2* increased linearly with the vector input over the range of 0-50 ng per well. **(A)** Sox2 transfection. Endogenous *Gbx2* was activated linearly with the exogenous Sox2 level. **(B)** Otx2 transfection. The repression of brachyury (*T*) and activation of *Hesx1* by Otx2 reached a plateau with ~10 ng of expression vector.






**Fig. S4. Responses of genes to transcription factor manipulations according to their activation and/or repression status.** The expected responses are shown schematically in the case of activation (A) and repression (B). The gene *X* level without knockdown or overexpression of factor Y (A) or factor Z (B) under a given condition is indicated as 1 for bar graphs. **(A)** In the case where gene *X* is activated by transcription factor Y, the effects of knockdown (K) and overexpression (O) of factor Y depend on the status of activation: (a) if gene *X* is not yet activated by Y, only the overexpression of Y increases gene *X* expression above basal levels; (b) when gene *X* has already been activated but not to the level of saturation, the knockdown of Y reduces, whereas the overexpression of Y augments, gene *X* expression; (c) if gene *X* is already activated by Y to saturation levels, only a knockdown of Y downregulates gene *X*. **(B)** Cases in which gene *X* is activated by transcription factor Y and repressed by Z: (d) if gene *X* is not yet activated, neither knockdown nor overexpression of Z affects its basal expression level; (e) under conditions where gene *X* is activated by a transcription factor (Y), but is not repressed by Z, then only the overexpression of Z will reduce gene *X* expression; (f) if gene *X* is in the middle of repression by Z, then knockdown of Z augments, and its overexpression reduces, gene *X* expression; (g) when gene *X* is completely repressed by Z, only knockdown of Z augments its expression.




**Fig. S5.** Continuation of the analysis shown in Fig. 3, representing cases with miscellaneous responses to exogenous transcription factor manipulations.




**Fig. S6.** The effects of overexpressing Otx2 or Sox2 and of Pou5f1 knockdown on the expression of five representative transcription factor genes in epiblastic state EpiSCs. Analysis was in parallel with those described in Fig. 4A. O2, Otx2; S, Sox2; KD-P5, Pou5f1 knockdown.

A



B



**Fig. S7. Activation of *Otx2* AN and *Hesx1* ANP enhancers.** (A) *Otx2* AN enhancer (Kurokawa et al., 2004) (165 bp EB fragment) containing a Pou binding site and a putative Zic binding site (top) was joined to the luciferase reporter, and its activation by Zic2 and Pou factors was measured in 10T1/2 fibroblasts. (B) The *Hesx1* ANP enhancer used in this study (top) contained three previously characterized Otx2 binding sites (Spieler et al., 2004). Its activation by Otx2 was measured by luciferase reporter assay.

A

N2 core CCCTGCCCGTTCGCCTTCATTTCCATAAAGGAGATTAGGAGAGGAGGGGAACCCACTCAAATGCAGATGCAGGAG

Probe A ACAGGTCCCCTGCCGTTCGCCTTC

Mut-ZIC ACAGGTCTAGGTACGTTCGCCTTC


Probe B GCCTTCATTTCCATAAAGGAGATTAGGAGAG

Mut-POU GCCTTCGCGTAACTAAGGAGATTAGGAGAG


Mut-INT GCCTTCATTTCCATCGATAGATTAGGAGAG

Mut-OTX GCCTTCATTTCCATAAGGCTCGGCGAGAG

B




C



**Fig. S8. EMSA analysis of the binding of Zic2 and Otx2 to the N2 enhancer core sequence.** (A) Probes and mutated competitor sequences used in the EMSA assay. (B) In vitro synthesized transcription factors used in the EMSA assay. The domains of these proteins, indicated by light-blue shading, were synthesized in vitro. HOM, homeodomain. (C) Competitive inhibition of factor binding to radiolabeled probes using various mutated competitor sequences. The *hTNC* (human tenascin C) competitor contains an OTX binding sequence (Briata et al., 1999). The band positions of Zic2 plus Probe A and Otx2 plus Probe B are indicated by open arrowheads. The formation of the former complex was inhibited by the probe sequence, but not by the Mut-ZIC sequence. The latter complex was inhibited by competitors carrying intact Otx2 binding sequences, but not by the Mut-OTX sequence. Zic3 gave identical results to Zic2 (data not shown).





**Fig. S9. Comparison of the activity of *Zic2* and *Zic3* in the activation of the N2 core enhancer.** Under the same conditions used for the transfection of 10T1/2 cells with N2[73bp]<sub>2</sub>-luciferase as described in Fig. 8, varying amounts of expression vectors for *Zic2* (solid circle and solid line) or *Zic3* (open circle and broken line) were co-transfected with (A) pCAGGS-*Pou5f1* (2.5 ng), (B) pCAGGS-*Pou3f1* (2.5 ng) and pCAGGS-*Otx2* (5 ng), or (C) alone. *Zic3* activated the N2 core enhancer in a manner analogous to that of *Zic2* under the conditions given in A and B.

**Table S1.** Primers used in qRT-PCR analyses

Target gene	Forward primer	Tm (C)	Reverse primer	Tm (C)	Reference sequence (the range of coding sequence)	Target position	of PCR product (bp)	Reference
<i>Gapdh</i>	CATGGCCTTCGGTGTTCCTA	60	GCGGCACGTCAGATCCA	59	NM_008084.2 (51-1052)	734-788	55	This study
<i>Fgf5</i>	AGAGTGGGCATCGGTTTCC	59	TGGGAGCCATTGACTTTGC	58	NM_010203.4 (229-1023)	502-560	59	This study
<i>Nanog</i>	AGGCCTGGACCGCTCAGT	59	AGTTATGGAGCGGAGCAGCAT	60	NM_028016.2 (216-1133)	907-996	60	This study
<i>Eomes</i>	GCCTTCCACCTTTGATGTATCC	58	AAAGCTTTGGCGCCTTCTCT	59	NM_010136.3 (472-2595)	2680-2740	61	This study
<i>Sox2</i>	CCATGGGCTCTGTGGTCAAG	60	CCCTGGAGTGGGAGGAAGAG	60	NM_011443 (412-1371)	1133-1204	72	This study
<i>Sox3</i>	CTGGGACCGTTGCCCTTGA	58	CCGACAGTTACGGCCAACT	59	NM_009237.2 (1-1353)	1439-1498	60	This study
<i>Sox1</i>	CGGAGGACAAAAGACAAAACC	59	AAGTTACAGAGCCGGCAGTCA	59	NM_009233.3 (843-2018)	2310-2377	80	This study
<i>Pou5f1</i>	TTCCCTCTGTTCCCGTCACT	58	TGGTGCTCAGTTTGAATGC	58	NM_013633.2 (62-1120)	1071-1127	57	This study
<i>Pou3f1</i>	AGACCACCATCTGCCGTTTC	59	TCCAGCCACTTGTGAGCAG	59	NM_011141 (62-1411)	924-1008	85	This study
<i>Pou3f2</i>	CGGATTTACTCAAGCAGACGTG	59	CAACAAAGGCTTCAGCTTGC	58	NM_008899.1 (1-1338)	858-990	133	This study
<i>Pou3f3</i>	CCCTTGACTTCTGCCCTCAA	59	CATCCGACGACCCATTC	60	NM_008900.2 (1-1494)	2639-2695	57	This study
<i>Pou3f4</i>	GCCACAGCTGCCTCGAAT	59	CATGGACAAGGGAGCTGGAA	60	NM_008901 (1-1086)	4-58	55	This study
<i>Otx2</i>	GAAATCAACTTGCAGAAATCCA	59	GCGGCACCTTAGCTCTTCGAT	58	NM_144841.3 (333-1202)	551-614	64	This study
<i>Zic2</i>	GCAGGGCCACCTTCTTTTC	57	GCCCATTGAGCAGCTTCTG	57	NM_009574.3 (293-1882)	751-824	74	This study
<i>Zic3</i>	TTTTAGGGTGCTGTTGGTTATTGA	59	ACAATTCCTTATCTCCACTTTTCTGTTA	58	NM_009575.2 (554-1954)	3729-3800	72	This study
<i>Sip1</i>	ACACTTTCCTTTTCGCTATTTCATGA	58	TGAGGCCTAAAAGTGTGGTTAC	58	NM_015753.3 (527-4174)	4650-4721	72	This study
<i>Zfp521</i>	TCCCCCGCCAAACTTCA	60	GTACCACCCATCCCTTCGAA	58	NM_145492.4 (235-4170)	762-820	59	This study
<i>Hesx1</i>	TCAGCTCCGGGAAAGCAA	59	CCAGTCTAAAATGCTCTCAATTG	58	MN_010420.2 (359-916)	385-446	62	This study
<i>Otx1</i>	GCGTCACCCCTTCAAGTCTTT	59	AACAGAGGGTCAGAGCGAAGAG	59	NM_011023.3 (282-1349)	1635-1712	78	This study
<i>Pax6</i>	ATGGGCATTGGTATGTTATAATGAAG	58	AACACAGATCCGCGATCCA	59	NM_013627.4 (526-1836)	2094-2158	65	This study
<i>Gbx2</i>	CAGCGACCACCTTCCCATAC	59	CGCAGTGTTTGTCTTGTGTCT	59	NM_010262.3 (422-1468)	1793-1853	61	This study
<i>Nkx1.2</i>	CCAGAGGGCGAGGAGAAGT	58	GACCCCTCAGTGGCTTGTGT	59	NM_009123.2 (112-1029)	1156-1212	57	This study
<i>T</i>	TTGAACTTTCCTCCATGTGCTGA	61	TCCCAAGAGCCTGCCACTTT	61	NM_009309.2 (109-1419)	1421-1502	82	Greber et al., 2010
<i>Sox17</i>	ATAAGCCCAGATGGGTCTTC	58	CCGTGGCTGTCTGAGAGGTT	59	NM_011441.4 (1083-2342)	2209-2275	67	This study

**Table S2.** Sequences inserted in pSilencerU6puro (Ambion) to produce shRNAs

Target genes	Sequences inserted (SENSE-loop-ANTISENSE)	References
Control	AGACAGCGAAACTGTTCTC-ttcaagaga-GAGAACAGTTTCGCTGTCT	Zeineddine et al., 2006
<i>Zic2/3*</i>	GAGAACCTCAAGATCCACA-ttcaagaga-TGTGGATCTTGAGGTTCTC	This study
<i>Otx2</i>	GGCTTCAGGTTATAGTCAAGG-ttcaagaga-CCTTGACTATAACCTGAAGCC	This study
<i>Sox2</i>	GGTTGATATCGTTGGTAAT-ttcaagaga-ATTACCAACGATATCAACC	Ivanova et al., 2006
<i>Pou5f1</i>	AGGTGTTTCAGCCAGACCAC-ttcaagaga-GTGGTCTGGCTGAACACCT	Zeineddine et al., 2006
<i>Pou3f1</i>	GCCCATGGACGACGTTTATGC-ttcaagaga-GCATAAACGTCGTCCATGGGC	Iwafuchi-Doi et al., 2011
<i>Nanog</i>	GAGACAGTGAGGTGCATAT-ttcaagaga-ATATGCACCTCACTGTCTC	Wang et al., 2007

\*The shRNA vector for *Zic2/3* was designed to target both *Zic* factor genes because of their functional similarities.

**Table S3. References for expression patterns in embryos from E6.5 to E8.5**

Genes	References
<i>Fgf5</i>	Hebert et al., 1991
<i>Nanog</i>	Hart et al., 2004
<i>Eomes</i>	Ciruna and Rossant, 1999; Hancock et al., 1999
<i>Sox2</i>	Uchikawa et al., 2011; Wood and Episkopou, 1999
<i>Sox3</i>	Uchikawa et al., 2011; Wood and Episkopou, 1999
<i>Sox1</i>	Uchikawa et al., 2011; Wood and Episkopou, 1999
<i>Pou5f1</i>	Perea-Gomez et al., 1999
<i>Pou3f1</i>	Zwart et al., 1996
<i>Pou3f2</i>	Bouchard et al., 2005
<i>Pou3f3</i>	Bouchard et al., 2005
<i>Pou3f4</i>	Bouchard et al., 2005
<i>Otx2</i>	Ang et al., 1994; Martinez-Barbera et al., 2001
<i>Zic2</i>	Elms et al., 2004; Inoue et al., 2007
<i>Zic3</i>	Elms et al., 2004; Inoue et al., 2007
<i>Sip1</i>	Miyoshi et al., 2006
<i>Zfp521</i>	Kamiya et al., 2011
<i>Hesx1</i>	Yang and Klingensmith, 2006
<i>Otx1</i>	Sakurai et al., 2010; Suda et al., 1999
<i>Pax6</i>	Inoue et al., 2000
<i>Gbx2</i>	Waters et al., 2003
<i>Nkx1.2</i>	Tamplin et al., 2008
<i>T</i>	Wilkinson et al., 1990
<i>Sox17</i>	Kanai-Azuma et al., 2002

**Link to Table 4.**

<http://dev.biologists.org/content/vol0/issue2012/images/data/dev.085936/DC1/DEV085936TableS4.xls>

**Table S5. Transfection efficiency and selection of transfected cells using puromycin (Pur), as estimated from CAGGS-EGFP expression**

Cells	EGFP-expressing cell fraction (%)	
	Without Pur selection	With Pur selection
Epiblastic EpiSCs	77	93
NP1 cells	56	95

**Table S6. Knockdown and overexpression efficiencies of transcription factors**

Transcription factor genes	Knockdown in epiblastic cells	Knockdown in NP1 cells	Overexpression in epiblastic cells
<i>Zic2</i>	0.59	0.73	256*
<i>Zic3</i>	0.19	0.34	—
<i>Otx2</i>	0.36	0.55	211
<i>Sox2</i>	0.33		257
<i>Pou5f1</i>	0.37	0.14	100
<i>Pou3f1</i>	0.63		146
<i>Nanog</i>	0.18		14

The values indicate the transcript expression levels determined by qRT-PCR after transfection of knockdown or overexpression vectors relative to untreated cells, which was taken as 1.

\*pCAGGS-*Zic2* was used to overexpress *Zic2/3* functions.

**Table S7. Endogenous expression of transcription factor genes in 10T1/2 fibroblasts used for enhancer transactivation analysis, as compared with EpiSCs**

Gene	10T1/2	EpiSC
<i>Sox2</i>	0.0445±0.0022	7.62±1.51
<i>Zic2</i>	0.0481±0.0149	1.07±0.09
<i>Zic3</i>	0.00872±0.00155	12.0±2.36
<i>Otx2</i>	0.0185±0.0039	7.82±0.45
<i>Pou5f1</i>	0.0161±0.0014	74.4±5.5
<i>Pou3f1</i>	Undetectable	1.78±1.03
<i>Pou3f2</i>	0.00761±0.00154	0.0101±0.0012
<i>Pou3f4</i>	Undetectable	0.0158±0.0043

Expression levels are relative to  $10^{-3}$  *Gapdh* and indicated with standard errors.