

RESEARCH ARTICLE

STEM CELLS AND REGENERATION

FGF ligands of the postnatal mammary stroma regulate distinct aspects of epithelial morphogenesis

Xiaohong Zhang, Denisse Martinez, Zuzana Koledova, Guijuan Qiao, Charles H. Streuli and Pengfei Lu*

ABSTRACT

FGF signaling is essential for mammary gland development, yet the mechanisms by which different members of the FGF family control stem cell function and epithelial morphogenesis in this tissue are not well understood. Here, we have examined the requirement of *Fgfr2* in mouse mammary gland morphogenesis using a postnatal organ regeneration model. We found that tissue regeneration from basal stem cells is a multistep event, including luminal differentiation and subsequent epithelial branching morphogenesis. Basal cells lacking *Fgfr2* did not generate an epithelial network owing to a failure in luminal differentiation. Moreover, *Fgfr2* null epithelium was unable to undergo ductal branch initiation and elongation due to a deficiency in directional migration. We identified FGF10 and FGF2 as stromal ligands that control distinct aspects of mammary ductal branching. FGF10 regulates branch initiation, which depends on directional epithelial migration. By contrast, FGF2 controls ductal elongation, requiring cell proliferation and epithelial expansion. Together, our data highlight a pleiotropic role of *Fgfr2* in stem cell differentiation and branch initiation, and reveal that different FGF ligands regulate distinct aspects of epithelial behavior.

KEY WORDS: Fibroblast growth factor, Branching morphogenesis, Collective epithelial migration, Stem cell, Mammary gland, Breast, FGFR2, FGF10

INTRODUCTION

An outstanding question in developmental biology is how different signaling pathways regulate diverse aspects of organ development. The presence of multiple members of a gene family, which are often functionally redundant, and the requirement of an essential gene at different stages of organ development pose a challenge to understanding the molecular basis of vertebrate organogenesis. The mouse mammary gland has become a powerful model for delineating the molecular basis of fundamental developmental processes, including epithelial differentiation and morphogenesis (Lu and Werb, 2008). Consistent with its relatively late emergence during metazoan evolution, the formation of the mammary gland depends on many of the same molecules; for example, components of the WNT and BMP signaling pathways and major transcription factors essential for the ontogeny of other body appendages (Mikkola and Millar, 2006).

Signaling via fibroblast growth factors (FGFs) is another major pathway essential for mammary gland biology (Dillon et al., 2004). The mammalian FGF family consists of 22 members, the majority of which function by binding with low affinity to heparan sulfate and

with high affinity to their cognate receptors FGFR1–4 (Itoh and Ornitz, 2008). Binding of paracrine FGFs to their receptors causes receptor dimerization, phosphorylation, and activation via one or more of several downstream signaling sub-branches, including signaling via Ras-Raf-MEK and PI3K (Turner and Grose, 2010). FGF has roles in both breast cancer and mammary gland development. Excessive FGF signaling due to overactive ligand or receptor causes breast tumors (Peters et al., 1983; Welm et al., 2002; Xian et al., 2005, 2007). FGFR2 upregulation as a result of allelic polymorphism is associated with human breast cancer, suggesting a causal role of excessive FGFR2 activities in the disease (Hunter et al., 2007; Meyer et al., 2008). By contrast, a reduction in FGF signaling due to removal of *Fgf10* or its receptor *Fgfr2* results in a failure to form mammary placodes during embryogenesis (Mailleux et al., 2002; Kim et al., 2013).

We have previously shown that during postnatal mammary gland development a reduction of FGF signaling by conditional removal of *Fgfr1* or *Fgfr2* severely compromises ductal branching (Lu et al., 2008; Parsa et al., 2008; Pond et al., 2012). Genetic mosaic analysis has indicated that *Fgfr2* null cells are outcompeted by wild-type cells when they co-exist during postnatal branching (Lu et al., 2008). Moreover, although *Fgfr2* null cells contribute to the branching network of mammary ducts in a mosaic environment, they could not regenerate the mammary gland when a pure population of mutant cells was transplanted into the fat pad. Although these studies have revealed that *Fgfr2* promotes cell proliferation at the invasion front, the precise function of *Fgfr2* in postnatal mammary development and homeostasis has remained largely elusive. In this study, we have used a combination of *in vivo* and *in vitro* models to examine *Fgfr2* functions in the developing mouse mammary gland.

RESULTS

Mammary epithelial cells lacking *Fgfr2* have reduced self-renewal ability and fail to regenerate the gland

Our previous study showing that FGFR2 is present in basal mammary epithelial cells (MECs) suggests that it plays a role in regulating mammary gland stem cell biology. We examined whether FGFR2 removal affects stem cell self-renewal using the ‘mammosphere’-forming assay, which measures the ability of mammary stem cells to form spheres in suspension (Moraes et al., 2007). We removed the *Fgfr2* gene by infecting MECs from either *Fgfr2*^{+/+} or *Fgfr2*^{fl/fl} mice with adenovirus-Cre-GFP to generate control cells (*Fgfr2*^{+/+}) or mutant (*Fgfr2*^{Δ/Δ}) cells (Fig. 1A) (Lu et al., 2008). About 80% of the MECs were transduced by adenovirus, as indicated by GFP expression. Immunoblotting revealed that FGFR2 protein levels were greatly reduced in samples that had been infected for 24 h (Fig. 1B). MECs lacking *Fgfr2* showed ~60% reduction in their ability to form primary mammospheres, indicating partially reduced self-renewal ability (Fig. 1C).

We also performed an aldefluor assay in which aldehyde dehydrogenase (ALDH), which is mainly expressed in the

Wellcome Trust Centre for Cell Matrix Research, Faculty of Life Sciences, University of Manchester, Manchester M13 9PT, UK.

*Author for correspondence (paul.pf.lu@gmail.com)




Fig. 1. Mammary epithelial cells lacking *Fgfr2* show reduced self-renewal activity and are unable to regenerate the mammary gland.

(A) The experimental procedure of sample preparation, adenoviral infection and FACS (for details see the Materials and Methods). (B) Western blotting analysis of FGFR2 expression by mammary gland organoids with (Ad⁺) or without (Ad⁻) 24 h infection by adenovirus-Cre-GFP. About 80% of mammary epithelial cells (MECs) were transduced and express GFP protein, whereas ~20% of MECs remained uninfected, leading to the residual expression of FGFR2 in experimental samples. (C) Quantitative analysis of mammosphere-forming efficiency by control or mutant MECs during primary mammosphere formation. Data were from three independent analyses. (D-G) ALDH⁺ stem/progenitor cells lacking *Fgfr2* function had reduced acini-forming efficiency in a 3D Matrigel assay. ALDH turns a colorless substrate into a green fluorescent product, a reaction that can be blocked by the inhibitor DEAB. Unlike ALDH⁻ cells, ALDH⁺ progenitor cells that were control (*Fgfr2^{+/+}*, E) or mutant (*Fgfr2^{Δ/Δ}*, F) had a relatively high efficiency in forming acini after 15 days culture. (G) Quantitative comparisons of acini-forming efficiency in MECs of different genotypes. Values shown are the mean \pm s.d. (H) The experimental procedure of basal cell (CD24^{med} CD49f^{hi}) enrichment and transplantation. About 10,000 basal cells were injected into the cleared fat pad. (I, J) Assays for β -Gal activity in the mammary glands 8 weeks after basal cell transplantation of either control (*Fgfr2^{+/+};R26R^{Δ/Δ}*, I) or mutant (*Fgfr2^{Δ/Δ};R26R^{Δ/Δ}*, n=32, J) cells. Scale bars: 100 μ m in E, F; 1 mm in I, J.

luminal epithelium, converts a colorless substrate into a green fluorescent product (Fig. 1D) (Ginestier et al., 2007; Eirew et al., 2012). ALDH⁺ progenitor cells readily formed hollow cysts after 15 days culture on Matrigel, whereas ALDH⁻ cells did not, indicating that the cyst-forming ability of MECs arises from ALDH⁺ progenitors (Fig. 1E-G). *Fgfr2* null MECs showed a minor (28.5%) reduction in cyst-forming efficiency compared with control MECs (Fig. 1G). Thus, mammary stem cells are compromised but not completely inhibited by FGFR2 removal.

To directly examine the role of *Fgfr2* in stem cell biology, we purified basal stem cells lacking *Fgfr2* and subjected them to the mammary gland regeneration assay (Fig. 1H). To eliminate the possibility of contamination by control cells, which have a competitive advantage over mutant cells in a mosaic environment, we crossed *Fgfr2^{fl/fl}* mice with the *R26R^{fl/fl}* reporter allele to mark cells transduced by β -Gal adenovirus (Soriano, 1999). MECs were isolated from *Fgfr2^{+/+};R26R^{fl/fl}* and *Fgfr2^{fl/fl};R26R^{fl/fl}* mice, and infected with

adenovirus-Cre-GFP to generate control (*Fgfr2^{+/+};R26R^{Δ/Δ}*) and mutant (*Fgfr2^{Δ/Δ};R26R^{Δ/Δ}*) cells, respectively. Infected MECs were enriched for basal cells (GFP⁺ CD24^{med} CD49f^{hi}) and transplanted into cleared fat pads of immune-deficient, nude mice (Fig. 1H). Control basal cells readily regenerated an epithelial network (Fig. 1I). However, *Fgfr2* mutant basal cells were unable to regenerate a mammary tree (n=32; Fig. 1J).

The complete inability of MECs lacking *Fgfr2* to regenerate the gland could not be explained by a reduction in the self-renewal ability of stem cells (Fig. 1A-G). Instead, the data suggest that *Fgfr2* plays a hitherto undefined role in mammary gland regeneration from basal cells.

***Fgfr2* function is required for luminal cell differentiation from basal stem cells**

The inability of *Fgfr2* null basal cells to regenerate mammary gland could result from a failure to produce luminal cells, a major

component of the epithelium; alternatively, *Fgfr2* null basal cells could generate luminal cells but their derivatives may not be able to undergo branching morphogenesis. To test these possibilities, we characterized the early morphological changes that basal *Fgfr2*-deleted cells undergo after transplantation *in vivo*. Transplants derived from 10,000 control basal cells initially grew in size and started forming small branches by 2 weeks, becoming further elaborated after 3 weeks (Fig. 2A,B). No epithelial outgrowths were observed from *Fgfr2* null basal cells or from control luminal cells (not shown).

Consistent with their inability to regenerate mammary gland, K8-expressing control (wild-type) luminal cells were unable to differentiate into K14-expressing basal cells 2 weeks after

transplantation (Fig. 2C-C'') (Shackleton et al., 2006; Stingl et al., 2006; Van Keymeulen et al., 2011). By contrast, K14-expressing control basal cells gave rise to K8-expressing luminal cells at this stage (Fig. 2D-D''). Accurate tissue architecture was confirmed by the bilayered organization of K14-expressing basal and K8-expressing luminal epithelium. However, in grafts derived from *Fgfr2* null basal cells, there were no luminal cells and the growths remained very small (Fig. 2E-E'').

These results reveal that *Fgfr2* is required for the generation of luminal cells. To determine whether this is due to FGFR2 activating genes involved in luminal differentiation, or cooperating with other pathways, we treated FACS-isolated CD24^{med} CD49f^{hi} basal cells




Fig. 2. *Fgfr2* function is required for luminal cell differentiation from basal stem cells. (A,B) Analysis of epithelial morphogenesis in recombinant glands derived from control basal cells at 2 (A) or 3 (B) weeks after transplantation. Epithelial branching was revealed by β -Gal assays. No epithelial outgrowth was observed in recombinant glands derived from mutant basal cells (not shown). Note that the outgrowths derived from control basal cells contained both branched and unbranched epithelia at 2 weeks and the branching network became progressively more elaborated 3 weeks after transplantation. (C-E'') Immunofluorescence using anti-K14 (green) or anti-K8 (red) antibodies on frozen sections of recombinant mammary glands derived from control luminal (*Fgfr2*^{+/+}; *R26R*^{Δ/Δ}, C-C'') and basal cells (D-D'') or mutant basal cells (*Fgfr2*^{Δ/Δ}; *R26R*^{Δ/Δ}, E-E'') 2 weeks after transplantation. Nuclei were counterstained by DAPI. K14 and K8 mark the basal and luminal epithelium, respectively. Note that control luminal cells did not give rise to K14⁺ basal cells after transplantation, whereas control basal cells had differentiated into luminal cells as marked by K8 expression, and that the overall architecture, in which basal epithelium encapsulates the luminal epithelium, had been established by this stage (D-D''). Luminal differentiation was not detected in transplants derived from mutant basal cells (E-E''). (F) Gene expression in basal stem cells in response to FGF2 (500 ng/ml) treatment after 24 h. Values shown are the mean \pm s.d. **P* < 0.05, unpaired two-tailed Student's *t*-tests. Scale bars: 1 mm in A,B; 100 μ m in C-E''.

with FGF2 and analyzed the expression of genes in the luminal and basal cell differentiation pathway (Fig. 2F). Expression of *Etv5*, which is a transcriptional target of FGF signaling, was upregulated by FGF2 (Zhang et al., 2009). Target genes of Notch signaling, including *Notch1*, which promotes luminal differentiation (Bouras et al., 2008; Yalcin-Ozuysal et al., 2010), were not significantly changed, although *Hes6* expression was upregulated (Fig. 2F; data not shown). However, whereas expression of basal markers including *K14* (also known as *Krt14*) and *p63* (*Trp63*) was increased by FGF2 treatment, that of luminal markers including *K8* (*Krt8*), *K18* (*Krt18*) and *Gata3* was reduced.

These results show that *Fgfr2* is required for mammary stem cells to differentiate into luminal cells. This provides one mechanism to explain why *Fgfr2* null basal cells are unable to regenerate a functional mammary gland.

Circumvention of *Fgfr2* requirement for luminal differentiation reveals its role in epithelial branching

Fgfr2 function might also be required for other aspects of mammary development, such as epithelial branching. To examine this possibility, we took advantage of the observation that basal and luminal lineages remain separate during both normal development and organ regeneration, and that basal-to-luminal differentiation only occurs if luminal cells are under-represented in the epithelium (Van Keymeulen et al., 2011). We tested whether the requirement of *Fgfr2* for luminal differentiation could be circumvented by co-transplanting *Fgfr2*^{ΔΔ} luminal cells together with basal cells.

We first determined whether adenovirus-Cre-GFP preferentially infected basal or luminal cells: MECs were infected with adenovirus-Cre-GFP before they were subjected to FACS based on CD24 and CD49f staining. We found that the basal-to-luminal ratio in the uninfected GFP⁻ MECs (0.76±0.34) was similar to that

in the infected GFP⁺ MECs (0.79±0.07; Fig. 3A-C). These data indicate that adenovirus-Cre-GFP does not preferentially infect basal or luminal cells and that *Fgfr2*^{ΔΔ} basal and luminal cells would be transplanted in the normal ratio observed under physiological conditions. Next, we transplanted sorted control or *Fgfr2* null MECs into the fat pads of nude mice. Based on the above results we concluded that *Fgfr2*^{ΔΔ} luminal cells are unlikely to be under-represented when compared with *Fgfr2*^{ΔΔ} basal cells, and thus the *Fgfr2* null cells would be able to form the mammary epithelial ductal network if luminal differentiation is the only process in mammary gland regeneration that requires *Fgfr2* function.

Recombinant glands derived from control MECs readily gave rise to an epithelial tree. However, those from mutant MECs were unable to form a new ductal network (*n*=24; Fig. 3D,E). The complete inability of *Fgfr2* null mammary epithelium, containing both basal and luminal cells, to build an epithelial network could not be explained by a reduction in cell proliferation, which is regulated by *Fgfr2* in the terminal end buds (TEBs) (Lu et al., 2008). Rather, the data suggest that *Fgfr2* plays additional roles in regulating epithelial branching of the mammary gland.

To further examine the role of *Fgfr2* in branching, we used a 3D culture model (Ewald et al., 2008). GFP-expressing control (*Fgfr2*^{+/+}) and mutant (*Fgfr2*^{ΔΔ}) MECs were aggregated, embedded in Matrigel, and cultured in basal medium containing FGF2. Control MEC aggregates remained unbranched without FGF2 (Fig. 4B), but formed nascent epithelial branches after 5 days in the presence of FGF2 and fully branched structures by 10 days (Fig. 3F-H). Surprisingly, the *Fgfr2* null MEC aggregates were also able to branch under the same conditions (Fig. 3I-K).

These results indicate that FGF2 induces epithelial branching *in vitro* via a receptor other than FGFR2. Thus, the lack of epithelial




Fig. 3. MECs lacking *Fgfr2* fail to branch *in vivo* but form branched structures when stimulated by FGF2 *in vitro*. (A-C) FACS analysis of MECs transduced by adenovirus-Cre-GFP. MECs were sorted based on their expression of integrin $\alpha 6$ (CD49f), CD24 and GFP. CD24^{med} CD49f^{hi} cells were basal, whereas CD24^{hi} CD49f^{ow} cells were luminal. The basal-to-luminal ratios were similar between untransduced GFP⁻ MECs (A,C) and transduced GFP⁺ MECs (B,C), at 0.76±0.34 and 0.79±0.07, respectively. Note that, unlike both luminal (lu) and basal (ba) populations, much of the stromal population (st) that existed in organoid preparations was not transduced well by adenovirus-Cre-GFP. Values shown are the mean±s.d. from at least three independent experiments. (D,E) Epithelial network as revealed by Carmine staining in recombinant mammary glands derived from control (*Fgfr2*^{+/+}) or mutant (*Fgfr2*^{ΔΔ}; *n*=24) MECs 8 weeks after transplantation. About 10,000 MECs were used for transplantation. (F-K) FGF2-based *in vitro* branching assay using control (F-H) or mutant (I-K) MEC aggregates. Scale bars: 1 mm in D,E; 100 μ m in F-K.




Fig. 4. FGF10 is a predominant stromal FGF ligand and directs mammary epithelium to undergo collective migration *in vitro*. (A) Relative expression of FGF genes in 5-week-old mammary glands. Analysis was based on microarray data retrieved from the NCBI Gene Expression Omnibus under accession numbers GSE2988 and GSE5602 (Kouros-Mehr and Werb, 2006). TEB/stroma (Cy5/Cy3) expression ratios and duct/stroma ratios are each shown for six independent experiments (lanes 1-6) and their respective means (M) using a color scale with black indicating no difference in expression, red indicating relative enrichment within the TEB or duct regions and green representing relative enrichment in the pure distal stroma. Intense red indicates a greater than fourfold enrichment in and around TEBs or ducts. (B-D) Differential responses of mammary organoids to basal medium alone (B) or to medium containing either FGF2 (C) or FGF10 (D). Note that FGF2 induced epithelial branching (C) whereas FGF10 did not (D). (E-T) Timecourse of mammary epithelial response to beads pre-soaked in BSA (E-H), FGF10 (I-L), FGF2 (M-P, $n=8$) or FGF7 (Q-T, $n=7$). Heparin acrylic beads of $\sim 100 \mu\text{m}$ in diameter were juxtaposed with mammary organoids at a distance of $\sim 100 \mu\text{m}$. Scale bars: $100 \mu\text{m}$.

branching by *Fgfr2* null MECs *in vivo* is not due to an intrinsic inability of mutant cells to form branched structures.

FGF10 is the predominant stromal FGF ligand during postnatal epithelial branching

The discrepancy between the requirement of FGFR2 for *in vivo* and *in vitro* branching could be explained if ligands other than FGF2 drive morphogenesis during normal development. To examine this possibility, we surveyed a microarray database that compares gene expression between ductal or TEB epithelia with distal stroma in 5-week-old mice (i.e. at a time of rapid branching *in vivo*) (Kouros-Mehr and Werb, 2006). *Fgfr2* is expressed in both the ductal and TEB epithelia (red, Fig. 4A) but not in the stroma. By contrast, FGF genes were detected in the stroma (green, Fig. 4A). Among them, *Fgf10* was the most abundantly expressed FGF gene at this developmental stage, although others were also present. qPCR confirmed that *Fgf10* was highly expressed during the stages of postnatal mammary ductal branching, and that *Fgf2*, 7, 9, 13 and 18 were also expressed (Table 1).

We therefore reasoned that in an *in vitro* assay FGF10 might influence epithelial branching more than FGF2. Surprisingly, when we replaced FGF2 with FGF10 in the above *in vitro* branching assay and re-examined the epithelial response, FGF10 did not induce epithelial branching, whereas FGF2 triggered robust epithelial branching from mammary organoids as expected (Fig. 4B-D). These data indicate that mammary epithelium responds differentially to FGF2 and FGF10.

FGF10 and FGF2 trigger distinct epithelial responses when delivered from a point source

In the branching systems of the trachea and air sacs in *Drosophila*, FGF/Branchless functions as a guidance cue and dictates the branching pattern (Affolter et al., 2003). We therefore reasoned that mammary organoids might only respond to FGF10 when delivered from a point source, rather than uniformly as in the above experiments. To test this, we examined the response of MECs to FGF-soaked beads. When mammary organoids were juxtaposed

Table 1. Expression of FGF genes in the mouse mammary gland at 3, 5 and 10 weeks as assayed by qPCR

Gene name	3 weeks	5 weeks	10 weeks
<i>Fgf1</i>	++	++	++
<i>Fgf2</i>	+	+	+
<i>Fgf3</i>	-	-	-
<i>Fgf4</i>	-	-	-
<i>Fgf5</i>	-	-	-
<i>Fgf6</i>	-	-	-
<i>Fgf7</i>	+	+	+
<i>Fgf8</i>	-	-	-
<i>Fgf9</i>	+	+	-
<i>Fgf10</i>	+++	+++	++
<i>Fgf11</i>	+	+	+
<i>Fgf12</i>	-	-	-
<i>Fgf13</i>	++	+	++
<i>Fgf14</i>	-	-	-
<i>Fgf15</i>	-	-	-
<i>Fgf16</i>	-	-	-
<i>Fgf17</i>	-	-	-
<i>Fgf18</i>	+	+	+
<i>Fgf20</i>	-	-	-
<i>Fgf21</i>	-	-	-
<i>Fgf22</i>	-	-	-
<i>Fgf23</i>	-	-	-

Levels of gene expression are based on the Ct number, with Ct ≥ 30 regarded as not expressed (-), $28 \leq \text{Ct} < 30$ as expressed (+), $27 \leq \text{Ct} < 28$ (++) and Ct < 27 (+++).

to beads pre-soaked in BSA, they remained at a relatively constant distance of $\sim 100 \mu\text{m}$ from the beads (Fig. 4E-H). By contrast, organoids moved towards FGF10-soaked beads, eventually making contact after 72 h (Fig. 4I-L). However, no epithelial migration was observed towards a point source of FGF2, as would be predicted by the current literature, although the organoids expanded in size and formed a large lumen (Fig. 4M-P; supplementary material Movie 1) (Ewald et al., 2008, 2012). Interestingly, FGF7 induced both cyst formation and movement (Fig. 4Q-T; supplementary material Movie 2). These data reveal that FGF10 promotes epithelial migration and suggest that it functions as a guidance cue for branching.

The differential epithelial responses to FGF2 and FGF10 could result from an insufficient amount of heparan sulfate, which is an essential low-affinity FGF receptor (Makarenkova et al., 2009). If so, then addition of heparan sulfate to the culture system should

abolish the differences observed. However, the addition of heparan sulfate had no effect on the response of mammary epithelium to FGF10 (Fig. 5A,B) or FGF2 (Fig. 5C,D). This confirms that the behaviors towards different FGFs are most likely an intrinsic property of the mammary epithelium.

As an initial step toward developing a complex system that mimics the *in vivo* stromal environment, in which multiple stromal FGF ligands co-exist, we examined how the presence of FGF2 and FGF10, either delivered together in the same position or separately in different positions, affected epithelial responsiveness. When delivered together in the same position, FGF2 and FGF10 triggered a similar response to that from FGF2 alone (Fig. 5E-H). However, FGF10 triggered directional migration when the ligands were delivered from separate positions (Fig. 5I-L).

Thus, FGF2 and FGF10 elicit distinct responses from mammary epithelium *in vitro*, suggesting that they regulate different aspects of epithelial branching *in vivo*. Our data also reveal that the spatial control of ligand delivery influences epithelial behavior. This shows that the spatiotemporal regulation of ligand expression in the postnatal mammary gland stroma is an important determinant of epithelial branching pattern.

Directional migration of stratified epithelium is not a result of localized cell proliferation or death and lacks an obvious front-rear polarity

The apparent movement of mammary organoids towards FGF10 beads could result from localized growth and/or apoptosis at the front or rear of the epithelium, or alternatively from collective epithelial migration. To examine these possibilities, we used phospho-histone H3 and LysoTracker staining to detect proliferating and dying cells, respectively (Lu et al., 2006). No localized cell proliferation occurred preferentially in the front region of the mammary organoids (Fig. 6A-J), and similarly there was no preferential death in the rear of mammary organoids juxtaposed with either BSA- or FGF10-soaked beads (supplementary material Fig. S1A-J).

Instead, time-lapse video recording revealed that mammary organoids underwent directional migration toward FGF10-soaked beads (supplementary material Movies 3 and 4). High-magnification imaging showed that the epithelium moved as a coherent entity during the course of its migration (supplementary material Movie 5). We saw no evidence of the existence of lamellipodia, filopodia and other cellular processes that are evident in other systems undergoing directional migration (Friedl and




Fig. 5. Mammary epithelium responds differentially to localized FGF2 and FGF10. Timecourse of mammary epithelial response toward beads pre-soaked in FGFs in the presence (A-D) or absence (E-L) of heparan sulfate (0.5 $\mu\text{g}/\text{ml}$). (E-L) FGF2 and FGF10 were delivered simultaneously to mammary epithelium either in the same bead (E-H) or in separate beads (I-L); the FGF10 beads are circled (dashed red line). Heparin acrylic beads of $\sim 100 \mu\text{m}$ in diameter were juxtaposed with mammary organoids at a distance of $\sim 100\text{-}150 \mu\text{m}$. Scale bars: 100 μm .




Fig. 6. Directional migration of stratified epithelium is not a result of localized cell proliferation. (A-H) Cell proliferation as detected by phospho-histone 3 (pH3) immunofluorescence (green) during the timecourse of mammary epithelial migration toward beads soaked in BSA (A-D) or FGF10 (E-H). Cell proliferation was quantified in one of the three evenly divided regions of an organoid – the front (f), middle (m) or rear (r) – depending on the distance between the region and the bead (asterisk). Scale bars: 100 μ m. (I, J) Quantification of cell proliferation in different regions of mammary organoids during epithelial migration. Only signals that overlap with nuclei were counted as proliferating cells, and background noise was discounted. No statistically significant differences were apparent among the different regions at the times indicated (unpaired two-tailed Student's *t*-test). Values shown are the mean \pm s.d. from at least three independent experiments.

Gilmour, 2009; Iliina and Friedl, 2009). Moreover, we did not detect a front-rear difference in the presence of basal cells, which enveloped luminal cells (supplementary material Fig. S2A-G''). Nor did we observe differential localization of the tissue polarity markers β 1-integrin, E-cadherin and ZO-1 (also known as TJP1) (supplementary material Fig. S2H-I'', L-M''). Expression of none of the marker genes examined was affected by whether the bead was pre-soaked in BSA or FGF10 (Fig. 2B-E''); data not shown). Interestingly, E-cadherin was found on all of the cell surfaces in the epithelium (supplementary material Fig. S2J-K''), rather than only on the basal-lateral surfaces, as would be expected for a simple epithelium. Thus, there is a partial loss of tissue polarity in the mammary epithelium during directional migration, presumably as a feature of stratified epithelium.

Collective epithelial migration depends on MEK and PI3K signaling and on MMP activities

To understand the molecular basis of the directional migration of the stratified epithelium, we examined the effect of several pharmacological inhibitors on the FGF10-induced migration. SU5402, an FGFR inhibitor, blocked the movement of mammary organoids toward FGF10 beads (Fig. 7A-D), confirming that collective epithelial migration requires this FGF. Similarly, the MEK inhibitors U0126 and PD98059 and the PI3K inhibitor LY294002 all inhibited MEC collective migration (Fig. 7E-J). To determine whether these two sub-branches of FGF signaling work independently or cooperatively, we serially diluted the MEK and PI3K inhibitors and found that concentrations that on their own did not affect epithelial migration worked in combination to either completely (1 of 4 samples) or partially (3 of 4 samples) block migration (supplementary material Table S2).

The pan-matrix metalloproteinase (MMP) inhibitor GM6001 partially inhibited epithelial migration (Fig. 7K-L), suggesting that MMP activities play a role in this process. To determine whether FGF signaling regulates MMP activities at the transcriptional level, we analyzed MMP gene expression (Mori et al., 2013; Wiseman et al., 2003) and found that *Mmp2*, *Mmp3* and *Mmp14* were upregulated in response to FGF10 treatment (Fig. 7M).

Together, these results show that directional migration of the mammary epithelium depends on MEK/ERK and PI3K/AKT signaling, which cooperate synergistically. Moreover, MMP activities promote epithelial migration and this is in part due to upregulation of MMP gene expression by FGF signaling.

Directional migration of mammary epithelium requires *Fgfr2* function

Finally, we hypothesized that the inability of *Fgfr2* null epithelium to form a branching ductal network *in vivo* is due to the inability of the cells to respond to the migratory cues induced by FGF10. *Fgfr2*^{+/+} and *Fgfr2* ^{Δ} MECs were aggregated before being subjected to the *in vitro* migration assay based on FGF10 stimulation. The control MEC aggregates underwent migration toward FGF10-soaked beads (Fig. 8A-D), whereas the *Fgfr2* null aggregates were unable to do so (Fig. 8E-H).

Thus, the *Fgfr2* gene is required for collective epithelial migration induced by FGF10, thereby providing a mechanism to explain the role of FGFR2 in the formation of mammary epithelial ducts *in vivo*.

DISCUSSION

In the current study, we circumvented the functional requirement of *Fgfr2* for embryonic development and studied its various roles during mammary gland formation. We found that mammary gland regeneration from basal cells is a multistep event, including luminal differentiation and subsequent epithelial branching, both of which depend on *Fgfr2* function (Fig. 8I). As a result of *Fgfr2* deletion, the cells fail to regenerate the mammary gland due to both an inability to generate luminal cells and a failure in directional migration, which is essential for branch initiation and epithelial patterning. We identified FGF10 and FGF2 as two stromal FGFs in the pubertal gland. FGF10 is a guidance cue to direct branch outgrowth, whereas FGF2 provides the driving force for epithelial expansion and ductal elongation. Together, our data highlight the pleiotropic roles of *Fgfr2* during mammary gland regeneration and underscore the novel phenomenon that FGF ligands regulate distinct aspects of mammary epithelial morphogenesis.




Fig. 7. FGF10-induced directional epithelial migration depends on MEK/ERK and PI3K/AKT signaling and on MMP activities. (A–L) Timecourse of mammary epithelial migration toward beads soaked in FGF10 in the absence (A,B) or presence (C–H) of various pharmacological inhibitors. Inhibition of signaling via FGFR, MEK or PI3K was achieved using media containing SU5402 (20 μ M, C,D), U0126 (2.0 μ M, E,F), PD98059 (20 μ M, G,H) and LY294002 (200 μ M, I,J). Note that collective epithelial migration was blocked in the presence of these inhibitors. GM6001 (100 μ M, K,L) was used as a general inhibitor against MMP activities. Collective epithelial migration was only partially blocked in the presence of GM6001; higher concentrations of GM6001 were without additional effect on migration but caused cell death (not shown). Scale bars: 100 μ m. (M) qPCR analysis of the expression of several MMP genes that play a role in mammary epithelial branching in response to FGF10 (500 ng/ml) treatment after 24 h. Values are the mean \pm s.d. * P < 0.05, unpaired two-tailed Student's t -tests.

FGFR2 regulates stem cell biology during mammary gland regeneration

Our data show that, whereas basal cells can give rise to luminal cells, the reciprocal process does not occur. These data are consistent with, and provide a functional explanation for, the well-documented observation that basal cells, but not luminal cells, can regenerate the mammary gland when transplanted into the cleared fat pad (Shackleton et al., 2006; Stingl et al., 2006). At present, it remains unclear whether luminal cell differentiation in the transplantation model shares a similar or distinct molecular basis with the *in vivo* event during normal development and whether what we learn from the former event could be applied to understanding the latter process.

We show that *Fgfr2* regulates at least two aspects of stem cell biology in the mammary gland. First, *Fgfr2* null cells have a reduced ability to form acini or mammospheres when compared with wild-type cells, suggesting that *Fgfr2* promotes self-renewal in adult stem cells of the mammary gland. Second, unlike wild-type basal cells, *Fgfr2* null basal cells are unable to differentiate into luminal cells. We conclude that failure in the generation of luminal cells is a cause of the inability to regenerate mammary gland from *Fgfr2* null basal cells (Fig. 8I).

FGF signaling has been shown to regulate cell differentiation in various other contexts, including the differentiation of embryonic stem cells and lens fiber (Zhao et al., 2008; Stavridis et al., 2010). Interestingly, however, FGF2 treatment of mammary basal cells does not upregulate the expression of luminal markers or target genes of Notch signaling, which promotes luminal differentiation (Bouras et al., 2008; Yalcin-Ozuyysal et al., 2010). These data suggest that *Fgfr2* does not directly promote luminal differentiation at the RNA level. Future studies aimed at addressing the mechanism by which *Fgfr2* regulates luminal differentiation could potentially be facilitated by the development of an appropriate *in vitro* model in which basal cell differentiation occurs.

Mammary epithelium responds differentially to FGF2 and FGF10 stimulation

We show that FGF2 and FGF10 elicit distinct responses from the mammary epithelium (Table 2). FGF2 triggers mostly epithelial

expansion and the formation of branched structures when delivered ubiquitously or large cysts when delivered locally. By contrast, FGF10 does not trigger an obvious epithelial expansion but instead elicits a conspicuous collective movement of the epithelium. Interestingly, similar differential responses from different FGF ligands have been reported in other developmental settings. For example, FGF4 attracts whereas FGF8 repels gastrulating cells in the early chick embryo (Yang et al., 2002). The basis for such differential responses, however, has remained unknown.

The distinct responsiveness of mammary epithelium to FGF2 and FGF10 was unaffected by heparan sulfate, suggesting that it results from intrinsic differences in the downstream events triggered by the ligands. Indeed, FGF2-mediated branch formation is driven by a high rate of cell proliferation, which when inhibited causes a failure in epithelial expansion and, presumably, ductal elongation (Fata et al., 2007; Ewald et al., 2008). This is in contrast to our data showing that cell proliferation is not preferentially localized to the front or rear of the epithelium and thus is unlikely to be a driving force for migration.

Furthermore, FGF2-based epithelial expansion does not depend on *Fgfr2* function, whereas FGF10-based directional migration does. Although it has remained unclear, the receptor that FGF2 activates is likely to be FGFR1, which like FGFR2 is expressed in the epithelium and plays a role in mammary branching (Lu et al., 2008; Pond et al., 2012). Interestingly, there is evidence to suggest that FGF2 and FGF10 preferentially activate FGFR1 and FGFR2, respectively (Ornitz et al., 1996; Zhang et al., 2006). Therefore, the distinct responses of mammary epithelium to FGF2 and FGF10 could, at least in part, be ascribed to differential binding to FGF receptors.

FGF ligands regulate distinct aspects of epithelial branching and coordinate organ development

Our data show that the FGF2-based *in vitro* system, which has been used to study various aspects of epithelial branching, is not suitable for the analysis of collective epithelial migration as has been suggested in the literature (Nguyen-Ngoc et al., 2012). This is because, despite the dynamic movement of individual cells, as a




Fig. 8. Directional migration of mammary epithelium requires *Fgfr2* function. (A-H) Timecourse of mammary epithelial migration in response to beads soaked in FGF10. Mammary epithelium of either control (*Fgfr2*^{+/+}, *n*=6, A-D) or mutant (*Fgfr2*^{Δ/Δ}, *n*=5, E-H) genotype was juxtaposed with FGF10 beads at a distance of ~100 μm. Note that although *Fgfr2*^{Δ/Δ} MEC aggregates did not migrate toward FGF10-soaked beads they did grow somewhat in size during culture, as did *Fgfr2*^{+/+} MEC aggregates, presumably due to cell proliferation. Scale bars: 100 μm. (I) Model of mammary gland regeneration from basal cell transplantation, which involves at least four stages. (1) FGFR2-dependent luminal cell differentiation from basal cells. (2) Expansion of mammary epithelium composed of both basal and luminal cells. (3) Epithelial budding and ductal elongation. FGF10-FGFR2 function via both MEK and PI3K signaling is essential for collective epithelial migration and branch initiation. By contrast, ductal elongation is primarily driven by FGF2 function in an FGFR2-independent manner. (4) Repeated branching morphogenesis until the mammary fat pad is filled and a functional mammary gland is generated. (1') In the absence of FGFR2 function, at least two of the above four steps fail to occur: *Fgfr2* null basal cells fail to differentiate into luminal cells and, consequently, a functional gland fails to regenerate; (3') when *Fgfr2*^{Δ/Δ} basal and luminal cells are transplanted, epithelial migration does not occur, leading to failure in epithelial invasion, budding and thus branch initiation. As a result, a functional gland fails to regenerate.

collective group cells do not move in a certain direction. Rather, the FGF2 model is most useful for understanding other aspects of branching morphogenesis, including cell rearrangements and shape changes during epithelial expansion and ductal elongation, an event that depends on cell proliferation (Table 2). By contrast, the FGF10-based system is best suited for understanding the basis of directional migration of stratified epithelium, which most likely employs a very

different strategy than that used by single cells, simple epithelial cells or loosely connected fibroblasts, where differences along the front-rear axis appear to be important for directional movement (Friedl and Gilmour, 2009; Iliina and Friedl, 2009). Interestingly, our analysis has detected no such differences in the presence of lamellipodia, filopodia and other cellular extensions or any difference in tissue polarity between the front and rear of the organoid epithelium.

Table 2. Comparison of mammary epithelial responses to FGF2 and FGF10

	FGF2	FGF10	References
Ubiquitous delivery	Epithelial expansion; branched structures	No branched structures	
Local delivery	Epithelial expansion; large cysts	Collective migration	
Directionality	No; random	Yes; toward FGF10	
Cell proliferation	Dependent	Independent	(Ewald et al., 2008)
Potential receptor	FGFR1	FGFR2	(Ornitz et al., 1996; Zhang et al., 2006)
<i>In vitro</i> model	Epithelial outgrowth	Directional migration	
<i>In vivo</i> process	Ductal elongation	Branch initiation and patterning	

Despite the clear differences between the directional migration of stratified mammary epithelium and that of individual cells in the *Drosophila* trachea and air sac systems, they share some interesting similarities. In the fly systems, for example, the direction of cell migration and branching pattern is controlled by FGF signaling (Affolter et al., 2003). In the absence of FGF signaling, owing to a lack of ligand (Branchless/FGF) or its receptor (Breathless/FGFR), cell migration does not occur, branches fail to initiate and, consequently, the tracheal and air sac systems fail to develop (Sutherland et al., 1996; Sato and Kornberg, 2002). In the mammary gland, our data show that epithelial cells lacking *Fgfr2* fail to undergo epithelial migration, which most likely leads to a failure of branch initiation, as observed in the fly system, and subsequently to the failure to form an epithelial branching tree.

Taken together, our working model is that FGF2 and FGF10 regulate distinct aspects of epithelial branching during postnatal mammary gland development. FGF10 regulates the branch initiation process, which depends on directional epithelial migration, and FGF2 controls the ductal elongation process, which depends on cell proliferation and epithelial expansion. Remarkably, similar situations have been observed in the fly air sac system. Although cell migration and branch initiation are controlled by FGFR signaling, cell proliferation and branch outgrowth are governed by a different receptor tyrosine kinase, EGFR (Cabernard and Affolter, 2005). In both cases, therefore, whether it is FGFR and EGFR in the fly or FGFR1 and FGFR2 in the mammary gland, members of the RTK family regulate distinct epithelial behaviors during branching morphogenesis (Xian et al., 2007).

MATERIALS AND METHODS

Mouse strains

Immunologically deficient female nude (*nu/nu*) mice were from Charles River Laboratories. Mice carrying the *R26R^{fl}* Cre-reporter allele were purchased from the Jackson Laboratory (Soriano, 1999). Mice carrying the *Fgfr2^{fl}* allele were provided by Dr David Ornitz (Yu et al., 2003). Offspring from crosses of the various lines were genotyped according to methods given in the publications describing the mouse lines. Mice were housed and maintained according to the University of Manchester and UK Home Office guidelines for animal research.

Preparation of mammary gland epithelial cells, adenovirus infection and transplantation

Preparation of MECs was as described (Zhang et al., 2014). Mammary gland fat pads of nude mice at 3 weeks of age were cleared based on an established protocol (Lu et al., 2008). About 10,000 MECs were injected into the cleared fat pad and harvested after 8 weeks.

Fluorescence activated cell sorting (FACS) and quantitative real-time PCR

Single cells were dissociated and labeled by infection using adenovirus-Cre-GFP or by antibody staining. Antibodies against CD24 (FITC; eBioscience, 48-0242), CD49f (PE; eBioscience, 12-0495) and CD45 (PeCy7; BD Biosciences, 552848) were used together with 7-AAD (BD Biosciences) to sort for luminal and basal populations using Aria and FACSCalibur systems (BD Biosciences). Data were processed using FACSDiva software (BD Biosciences). qPCR was performed using the 7500 Fast Real-Time PCR system (Applied Biosystems) and data were normalized to the expression of at least two reference genes, which included *Actb*, *Rn18s*, *Eef1g* and *Gapdh* (supplementary material Table S1).

In vitro epithelial branching and migration assays

Branching and epithelial invasion assays and live imaging were as described (Ewald et al., 2008; Zhang et al., 2014). FGF2 and FGF10 (2.5 nM; Sigma) were used in the 7- to 10-day culture. Primary antibodies were against smooth muscle actin (Sigma, 1A4, C6198), K8 (DSHB, Troma-1), K14 (Covance,

PRB-155P), phospho-histone H3 (Upstate, 06-570), β 1-integrin (Millipore, MAB1997), E-cadherin (BD Biosciences, 610181) and ZO-1 (Millipore, MABT11). Samples were counterstained with To-Pro-3 (Invitrogen, T3605) or 4',6-diamidino-2-phenylindole (DAPI; Sigma, 32670). Assays for cell death using LysoTracker (Molecular Probes, L-7528) staining were as previously reported (Lu et al., 2006).

Statistical data were analyzed using two-factorial analysis of variance (ANOVA), having time and section of organoid as the two factors. Student's *t*-test was used to compare the number of phospho-histone H3-positive nuclei between control (BSA-soaked beads) and experimental (FGF10-soaked beads) groups.

Data mining of expression microarray

Microarray data were from the NCBI Gene Expression Omnibus under the accession numbers GSE2988 and GSE5602 (Kouros-Mehr and Werb, 2006). Expression of FGF genes was determined by calculating $M = \log_2(Cy5/Cy3)$, where Cy5 values of TEB/ductal epithelium were compared with Cy3 values of the distal stroma from female mice at 5 weeks of age. TEB/stroma (Cy5/Cy3) expression ratios and duct/stroma ratios are each shown for six independent experiments.

Acknowledgements

We thank Dr Yaoyong Li for bioinformatics analyses on FGF gene expression, and Drs Zena Werb, Rob Clark and Karel Dorey for critical reading of the manuscript.

Competing interests

The authors declare no competing financial interests.

Author contributions

X.Z. and P.L. designed the research. X.Z., D.M., Z.K., G.Q. and P.L. performed the experiments. All authors analyzed the data. P.L. wrote the paper.

Funding

This work was supported by grants from the Wellcome Trust [no. 081203/Z/06/Z] to C.H.S.; and the National Institutes of Health [R03 HD060807] and Breakthrough Breast Cancer to P.L. The Wellcome Trust Centre for Cell-Matrix Research, University of Manchester, is supported by core funding from the Wellcome Trust [088785/Z/09/Z]. Deposited in PMC for release after 6 months.

Supplementary material

Supplementary material available online at <http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.106732/-/DC1>

References

- Affolter, M., Bellusci, S., Itoh, N., Shilo, B., Thiery, J.-P. and Werb, Z. (2003). Tube or not tube: remodeling epithelial tissues by branching morphogenesis. *Dev. Cell* **4**, 11-18.
- Bouras, T., Pal, B., Vaillant, F., Harburg, G., Asselin-Labat, M.-L., Oakes, S. R., Lindeman, G. J. and Visvader, J. E. (2008). Notch signaling regulates mammary stem cell function and luminal cell-fate commitment. *Cell Stem Cell* **3**, 429-441.
- Cabernard, C. and Affolter, M. (2005). Distinct roles for two receptor tyrosine kinases in epithelial branching morphogenesis in *Drosophila*. *Dev. Cell* **9**, 831-842.
- Dillon, C., Spencer-Dene, B. and Dickson, C. (2004). A crucial role for fibroblast growth factor signaling in embryonic mammary gland development. *J. Mammary Gland Biol. Neoplasia* **9**, 207-215.
- Eirew, P., Kannan, N., Knapp, D. J. H. F., Vaillant, F., Eberman, J. T., Lindeman, G. J., Visvader, J. E. and Eaves, C. J. (2012). Aldehyde dehydrogenase activity is a biomarker of primitive normal human mammary luminal cells. *Stem Cells* **30**, 344-348.
- Ewald, A. J., Brenot, A., Duong, M., Chan, B. S. and Werb, Z. (2008). Collective epithelial migration and cell rearrangements drive mammary branching morphogenesis. *Dev. Cell* **14**, 570-581.
- Ewald, A. J., Huebner, R. J., Palsdottir, H., Lee, J. K., Perez, M. J., Jorgens, D. M., Tauscher, A. N., Cheung, K. J., Werb, Z. and Auer, M. (2012). Mammary collective cell migration involves transient loss of epithelial features and individual cell migration within the epithelium. *J. Cell Sci.* **125**, 2638-2654.
- Fata, J. E., Mori, H., Ewald, A. J., Zhang, H., Yao, E., Werb, Z. and Bissell, M. J. (2007). The MAPK(ERK-1,2) pathway integrates distinct and antagonistic signals from TGF α and FGF7 in morphogenesis of mouse mammary epithelium. *Dev. Biol.* **306**, 193-207.
- Friedl, P. and Gilmour, D. (2009). Collective cell migration in morphogenesis, regeneration and cancer. *Nat. Rev. Mol. Cell Biol.* **10**, 445-457.

- Ginestier, C., Hur, M. H., Charafe-Jauffret, E., Monville, F., Dutcher, J., Brown, M., Jacquemier, J., Viens, P., Kleer, C. G., Liu, S. et al. (2007). ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell* **1**, 555-567.
- Hunter, D. J., Kraft, P., Jacobs, K. B., Cox, D. G., Yeager, M., Hankinson, S. E., Wacholder, S., Wang, Z., Welch, R., Hutchinson, A. et al. (2007). A genome-wide association study identifies alleles in FGFR2 associated with risk of sporadic postmenopausal breast cancer. *Nat. Genet.* **39**, 870-874.
- Iliina, O. and Friedl, P. (2009). Mechanisms of collective cell migration at a glance. *J. Cell Sci.* **122**, 3203-3208.
- Itoh, N. and Ornitz, D. M. (2008). Functional evolutionary history of the mouse Fgf gene family. *Dev. Dyn.* **237**, 18-27.
- Kim, E.-J., Jung, H.-S. and Lu, P. (2013). Pleiotropic functions of fibroblast growth factor signaling in embryonic mammary gland development. *J. Mammary Gland Biol. Neoplasia* **18**, 139-142.
- Kouros-Mehr, H. and Werb, Z. (2006). Candidate regulators of mammary branching morphogenesis identified by genome-wide transcript analysis. *Dev. Dyn.* **235**, 3404-3412.
- Lu, P. and Werb, Z. (2008). Patterning mechanisms of branched organs. *Science* **322**, 1506-1509.
- Lu, P., Minowada, G. and Martin, G. R. (2006). Increasing Fgf4 expression in the mouse limb bud causes polysyndactyly and rescues the skeletal defects that result from loss of Fgf8 function. *Development* **133**, 33-42.
- Lu, P., Ewald, A. J., Martin, G. R. and Werb, Z. (2008). Genetic mosaic analysis reveals FGF receptor 2 function in terminal end buds during mammary gland branching morphogenesis. *Dev. Biol.* **321**, 77-87.
- Mailleux, A. A., Spencer-Dene, B., Dillon, C., Ndiaye, D., Savona-Baron, C., Itoh, N., Kato, S., Dickson, C., Thiery, J. P. and Bellusci, S. (2002). Role of FGF10/FGFR2b signaling during mammary gland development in the mouse embryo. *Development* **129**, 53-60.
- Makarenkova, H. P., Hoffman, M. P., Beenken, A., Eliseenkova, A. V., Meech, R., Tsau, C., Patel, V. N., Lang, R. A. and Mohammadi, M. (2009). Differential interactions of FGFs with heparan sulfate control gradient formation and branching morphogenesis. *Sci. Signal.* **2**, ra55.
- Meyer, K. B., Maia, A.-T., O'Reilly, M., Teschendorff, A. E., Chin, S.-F., Caldas, C. and Ponder, B. A. J. (2008). Allele-specific up-regulation of FGFR2 increases susceptibility to breast cancer. *PLoS Biol.* **6**, e108.
- Mikkola, M. L. and Millar, S. E. (2006). The mammary bud as a skin appendage: unique and shared aspects of development. *J. Mammary Gland Biol. Neoplasia* **11**, 187-203.
- Moraes, R. C., Zhang, X., Harrington, N., Fung, J. Y., Wu, M.-F., Hilsenbeck, S. G., Allred, D. C. and Lewis, M. T. (2007). Constitutive activation of smoothened (SMO) in mammary glands of transgenic mice leads to increased proliferation, altered differentiation and ductal dysplasia. *Development* **134**, 1231-1242.
- Mori, H., Lo, A. T., Inman, J. L., Alcaraz, J., Ghajar, C. M., Mott, J. D., Nelson, C. M., Chen, C. S., Zhang, H., Bascom, J. L. et al. (2013). Transmembrane/cytoplasmic, rather than catalytic, domains of Mmp14 signal to MAPK activation and mammary branching morphogenesis via binding to integrin beta1. *Development* **140**, 343-352.
- Nguyen-Ngoc, K.-V., Cheung, K. J., Brenot, A., Shamir, E. R., Gray, R. S., Hines, W. C., Yaswen, P., Werb, Z. and Ewald, A. J. (2012). ECM microenvironment regulates collective migration and local dissemination in normal and malignant mammary epithelium. *Proc. Natl. Acad. Sci. USA* **109**, E2595-E2604.
- Ornitz, D. M., Xu, J., Colvin, J. S., McEwen, D. G., MacArthur, C. A., Coulier, F., Gao, G. and Goldfarb, M. (1996). Receptor specificity of the fibroblast growth factor family. *J. Biol. Chem.* **271**, 15292-15297.
- Parsa, S., Ramasamy, S. K., De Langhe, S., Gupte, V. V., Haigh, J. J., Medina, D. and Bellusci, S. (2008). Terminal end bud maintenance in mammary gland is dependent upon FGFR2b signaling. *Dev. Biol.* **317**, 121-131.
- Peters, G., Brookes, S., Smith, R. and Dickson, C. (1983). Tumorigenesis by mouse mammary tumor virus: evidence for a common region for provirus integration in mammary tumors. *Cell* **33**, 369-377.
- Pond, A. C., Bin, X., Batts, T., Roarty, K., Hilsenbeck, S. and Rosen, J. M. (2012). Fibroblast growth factor receptor signaling is essential for normal mammary gland development and stem cell function. *Stem Cells* **31**, 178-189.
- Sato, M. and Kornberg, T. B. (2002). FGF is an essential mitogen and chemoattractant for the air sacs of the drosophila tracheal system. *Dev. Cell* **3**, 195-207.
- Shackleton, M., Vaillant, F., Simpson, K. J., Stingl, J., Smyth, G. K., Asselin-Labat, M.-L., Wu, L., Lindeman, G. J. and Visvader, J. E. (2006). Generation of a functional mammary gland from a single stem cell. *Nature* **439**, 84-88.
- Soriano, P. (1999). Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat. Genet.* **21**, 70-71.
- Stavridis, M. P., Collins, B. J. and Storey, K. G. (2010). Retinoic acid orchestrates fibroblast growth factor signalling to drive embryonic stem cell differentiation. *Development* **137**, 881-890.
- Stingl, J., Eirew, P., Ricketson, I., Shackleton, M., Vaillant, F., Choi, D., Li, H. I. and Eaves, C. J. (2006). Purification and unique properties of mammary epithelial stem cells. *Nature* **439**, 993-997.
- Sutherland, D., Samakovlis, C. and Krasnow, M. A. (1996). branchless encodes a Drosophila FGF homolog that controls tracheal cell migration and the pattern of branching. *Cell* **87**, 1091-1101.
- Turner, N. and Grose, R. (2010). Fibroblast growth factor signalling: from development to cancer. *Nat. Rev. Cancer* **10**, 116-129.
- Van Keymeulen, A., Rocha, A. S., Ousset, M., Beck, B., Bouvencourt, G., Rock, J., Sharma, N., Dekoninck, S. and Blanpain, C. (2011). Distinct stem cells contribute to mammary gland development and maintenance. *Nature* **479**, 189-193.
- Welm, B. E., Freeman, K. W., Chen, M., Contreras, A., Spencer, D. M. and Rosen, J. M. (2002). Inducible dimerization of FGFR1: development of a mouse model to analyze progressive transformation of the mammary gland. *J. Cell Biol.* **157**, 703-714.
- Wiseman, B. S., Sternlicht, M. D., Lund, L. R., Alexander, C. M., Mott, J., Bissell, M. J., Soloway, P., Itohara, S. and Werb, Z. (2003). Site-specific inductive and inhibitory activities of MMP-2 and MMP-3 orchestrate mammary gland branching morphogenesis. *J. Cell Biol.* **162**, 1123-1133.
- Xian, W., Schwertfeger, K. L., Vargo-Gogola, T. and Rosen, J. M. (2005). Pleiotropic effects of FGFR1 on cell proliferation, survival, and migration in a 3D mammary epithelial cell model. *J. Cell Biol.* **171**, 663-673.
- Xian, W., Schwertfeger, K. L. and Rosen, J. M. (2007). Distinct roles of fibroblast growth factor receptor 1 and 2 in regulating cell survival and epithelial-mesenchymal transition. *Mol. Endocrinol.* **21**, 987-1000.
- Yalcin-Ozuysal, O., Fiche, M., Guitierrez, M., Wagner, K.-U., Raffoul, W. and Briskin, C. (2010). Antagonistic roles of Notch and p63 in controlling mammary epithelial cell fates. *Cell Death Differ.* **17**, 1600-1612.
- Yang, X., Dormann, D., Münsterberg, A. E. and Weijer, C. J. (2002). Cell movement patterns during gastrulation in the chick are controlled by positive and negative chemotaxis mediated by FGF4 and FGF8. *Dev. Cell* **3**, 425-437.
- Yu, K., Xu, J., Liu, Z., Sosic, D., Shao, J., Olson, E. N., Towler, D. A. and Ornitz, D. M. (2003). Conditional inactivation of FGF receptor 2 reveals an essential role for FGF signaling in the regulation of osteoblast function and bone growth. *Development* **130**, 3063-3074.
- Zhang, X., Ibrahim, O. A., Olsen, S. K., Umemori, H., Mohammadi, M. and Ornitz, D. M. (2006). Receptor specificity of the fibroblast growth factor family: the complete mammalian FGF family. *J. Biol. Chem.* **281**, 15694-15700.
- Zhang, Z., Verheyden, J. M., Hassell, J. A. and Sun, X. (2009). FGF-regulated ETV genes are essential for repressing Shh expression in mouse limb buds. *Dev. Cell* **16**, 607-613.
- Zhang, X., Qiao, G. and Lu, P. (2014). Modulation of fibroblast growth factor signaling is essential for mammary epithelial morphogenesis. *PLoS ONE* **9**, e92735.
- Zhao, H., Yang, T., Madakashira, B. P., Thiels, C. A., Bechtler, C. A., Garcia, C. M., Zhang, H., Yu, K., Ornitz, D. M., Beebe, D. C. et al. (2008). Fibroblast growth factor receptor signaling is essential for lens fiber cell differentiation. *Dev. Biol.* **318**, 276-288.




Figure S1: Directional migration of stratified epithelium is not a result of localized cell death. (A-J) Cell death as detected by Lysotracker assay (green) during the time course of mammary epithelial migration toward beads soaked in BSA (A-D) or FGF10 (E-H). Cell death was quantified in one of the three evenly divided regions of an organoid, the front (f), middle (m), or rear (r) regions, depending on their distances from the bead (asterisk). Scale bars: 100 μ m. (I, J) Quantification of cell death in different regions of mammary organoids during epithelial migration. Only signals that overlap with cell bodies were counted as dying cells whereas background noise was discounted. Statistical data were analyzed using two-factorial Analysis of Variance (ANOVA), having time and section of organoid as the two factors.





Figure S2: Directional migration of stratified epithelium lacks obvious front-rear polarity. (A) Schematic diagram depicting optical sectioning protocol of the mammary organoid in relation to bead location. Samples were sectioned optically along the Z-axis and images from the bottom (Bot) and central (Cen) are shown in (B-M). The front (Fro) and rear (Rea) areas of the central optical section were further shown in close-up views (C'-M''). (B-M) Immunofluorescence on mammary organoids to detect basal cells using antibodies against SMA (B-C'') and K14 (C-E'') and tissue polarity using antibodies against $\beta 1$ integrin (H-I''), E-Cadherin (J-K''), and Zo-1 (L-M''). Scale bars: 25 μm .




Movie 1: Time-lapse movie of mammary organoids stimulated by FGF2-soaked beads




Movie 2: Time-lapse movie of mammary organoids stimulated by FGF7-soaked beads



Movie 3: Time-lapse movie of mammary organoids stimulated by BSA-soaked beads



Movie 4: Time-lapse movie of mammary organoids stimulated by FGF10-soaked beads



Movie 4': Time-lapse movie of close-up view of mammary collective epithelial migration when stimulated by FGF10-soaked beads



Movie 5: High-resolution time-lapse movie of mammary organoids stimulated by FGF10-soaked beads

SUPPLEMENTARY TABLE S1: Primers used in qPCR.

Gene name	Forward sequence (5' → 3')	Reverse sequence (5' → 3')
<i>Fgf1</i>	ggacaccgaagggtttat	gcatgcttctggagggttaa
<i>Fgf2</i>	cggctctactgcaagaacg	tgcttgagttgtagttgacg
<i>Fgf3</i>	tgagaacagcgcctatagca	gtaccgccagaaaagagc
<i>Fgf4</i>	gcaagctcttcggtgtgc	cgtaggattcgtaggcgtg
<i>Fgf5</i>	cgaggagtttcagcaacaaa	tccgtaaattggcacttgc
<i>Fgf6</i>	tcagtggaacacacgaggag	cccgttctaccgtggagat
<i>Fgf7</i>	aagggacccaggagatgaag	actgccacggctctgattt
<i>Fgf8</i>	caggtcctggccaacaag	ggtctccacaatgagcttcg
<i>Fgf9</i>	actgcaggactggatttcattag	ccaggcccactgctatactg
<i>Fgf10</i>	cgggaccaagaatgaagact	aacaactccgatttccactga
<i>Fgf11</i>	ctttgccagaaacagctcct	gcctttgagctgaggctct
<i>Fgf12</i>	gacgaaaacagcgactacacc	tctccattcatggccacata
<i>Fgf13</i>	caggcagatggaaccattg	cccacagggatgagggtaaa
<i>Fgf14</i>	tgctgtacaggcaacaggag	ttctcggtacatggcaacttc
<i>Fgf15</i>	ggcaagatatacgggctgatt	tccatttctccctgaaggt
<i>Fgf16</i>	agtggactctggcctgtaccta	cattcacgtgtgagtttctcg
<i>Fgf17</i>	tatgaacaagaggggcaagc	ctcgggtgaacacgcagtct
<i>Fgf18</i>	aggacggggacaagtatgc	ggacttgactcccgaaggtat
<i>Fgf20</i>	cggcaggatcacagtctctt	aagggtacaggccactgtcca
<i>Fgf21</i>	agatggagctctctatggatcg	gggcttcagactggtacacatt
<i>Fgf22</i>	ctatgtggccatgaatcgc	cggaacctacagtccacagag
<i>Fgf23</i>	atctccacggcaacattttt	gtccactggcggaaacttg
<i>Etv5</i>	aggaccccaggctgtacttt	tggccgattcttctggatac
<i>K8</i>	atcgagatcaccacctaccg	tgaagccagggtctagtgagt
<i>K18</i>	agatgacaccaacatcacaagg	tccagaccttgacttctc
<i>Notch1</i>	acaacaacgagtgtgagtcc	acacgtggctcctgtatatg
<i>Hes6</i>	gctgctctcgtttgtaacaca	cgatgggatggcaaccaa
<i>Gata3</i>	agccacatctctcccttcag	agggctctgcctcttaacc
<i>K14</i>	atcgaggacctgaagagcaa	ggctctcaatctgcatctcc
<i>p63</i>	ggatgatttgcaagtctga	acttggggtctcaggagat
<i>Mmp2</i>	taacctggatgccgtcgt	ttcaggttaataagcacctttaa
<i>Mmp3</i>	ttgttctttgatgcagtcagc	gatttgcgcaaaaagtgc
<i>Mmp14</i>	gagaacttcgtgttgcctga	ctttgtgggtgaccctgact

<i>Actb</i>	ggctgtattccctccatg	ccagttggaacaatgcatgt
<i>18S</i>	gtaaccggtgaacccatt	gtaaccggtgaacccatt
<i>EEF1g</i>	ggattctgtgtttgagagca	cagcaaagetgaccactg
<i>GAPDH</i>	ttcaccacatggagaaggc	ccctttggctccacct

SUPPLEMENTARY TABLE S2. Effects of MEK inhibitor (U0126) and PI3K inhibitor (LY294002), alone or in combination, on migration of mammary epithelium when stimulated by beads soaked in FGF10.

U0126 [μ M]	LY294002 [μ M]	Complete migration	Partial migration	No migration	Numbers tested
2.0	0	0	2	3	5
1.3	0	4	0	0	4
1.0	0	3	0	1	4
0	200.0	0	0	9	9
0	20.0	0	0	6	6
0	13.3	0	0	4	4
0	10.0	1	1	2	4
0	5.0	2	0	2	4
0	3.3	4	0	0	4
2.0	3.0	0	3	0	3
2.0	5.0	0	3	1	4
1.3	13.3	0	0	3	3
1.3	10.0	0	0	7	7
1.3	5.0	0	2	2	4
1.3	3.3	0	3	1	4