

RESEARCH REPORT

TECHNIQUES AND RESOURCES

Translational profiling through biotinylation of tagged ribosomes in zebrafish

Michael P. Housley^{1,*,\$}, Sven Reischauer^{1,*}, Marc Dieu², Martine Raes², Didier Y. R. Stainier^{1,*,\$} and Benoit Vanhollebeke^{1,*,\$,‡}

ABSTRACT

Heterogeneity within a population of cells of the same type is a common theme in metazoan biology. Dissecting complex developmental and physiological processes crucially relies on our ability to probe the expression profile of these cell subpopulations. Current strategies rely on cell enrichment based on sequential or simultaneous use of multiple intersecting markers starting from a heterogeneous cell suspension. The extensive tissue manipulations required to generate single-cell suspensions, as well as the complexity of the required equipment, inherently complicate these approaches. Here, we propose an alternative methodology based on a genetically encoded system in the model organism *Danio rerio* (zebrafish). In transgenic fish, we take advantage of the combinatorial biotin transfer system, where polysome-associated mRNAs are selectively recovered from cells expressing both a tagged ribosomal subunit, Rpl10a, and the bacterial biotin ligase BirA. We have applied this technique to skeletal muscle development and identified new genes with interesting temporal expression patterns. Through this work we have thus developed additional tools for highly specific gene expression profiling.

KEY WORDS: TRAP, Gene expression profiling, Muscle development, BirA

INTRODUCTION

Cellular diversity is a defining feature of multicellular organisms. Growth and development requires the production of various cell types from undifferentiated progenitors that differ in fates and behaviors. Differential gene expression patterns maintain and refine this heterogeneity. Genome-wide assessment of cell type-specific expression patterns is thus central to our understanding of development, physiology and disease. The constraints associated with the currently available methods for cell type-specific RNA isolation calls for the implementation of novel strategies.

Most commonly used methods rely on cell-based purification strategies (Heiman et al., 2008). The target cell population is enriched from a heterogeneous cell suspension typically through fluorescence-activated cell sorting (FACS), although other methods of cell enrichment based on affinity purification (Wysocki and Sato, 1978) have also been employed. RNA is isolated from the purified cell population in a second step. A key asset of this methodology is

that multiple markers can be evaluated, refining the cellular subtypes that can be isolated. However, because of the time required to isolate cell populations and the harsh dissociation methods often required to generate a single cell suspension, the cellular stress response must be considered.

To overcome these limitations, alternative strategies aimed at recovering the target RNA directly from intact complex tissues have been implemented. Genetically encoded phosphoribosyltransferase (UPRT) biosynthetically labels nascent RNA in the target cell type when provided with 4-thiouracil. Chemically modified RNA molecules can then be affinity purified, a technique called TU-tagging (Miller et al., 2009; Gay et al., 2013). Alternatively, mRNA-binding proteins can be affinity tagged in specific cells and used as anchors to co-purify associated mRNAs. Both poly(A)-binding proteins (Roy et al., 2002; Kunitomo et al., 2005) and ribosomal proteins (Heiman et al., 2008) have been successfully employed to measure cellular mRNAs. This latter technique, translating ribosome affinity purification (TRAP), the purification of labeled polysomes for subsequent analysis of actively translating mRNAs, has the additional advantage of assaying translating mRNAs, and not total cellular pools, which should more accurately represent protein expression. A common limitation of those methods compared with FACS is that they currently rely on a single genetically encoded transgene. For example, recent applications of TRAP in zebrafish use the *cmc2* promoter (Fang et al., 2013) or the *ftyfp* promoter (Tryon et al., 2013) to drive expression of EGFP-Rpl10a.

Here, we describe a combinatorial version of the TRAP method where polysome-associated RNA is selectively recovered from cells with intersecting expression of two independently encoded transgenes. The first transgene controls the expression of an Avi (biotin ligase recognition peptide)-tagged Rpl10a ribosomal protein that serves as the substrate for the biotin transfer reaction controlled by a second transgene: the bacterial biotin ligase BirA. We have implemented this technology in zebrafish embryos, which for the study of vertebrate developmental genetics possess several advantages, including external fertilization, optical transparency, rapid organ development and facile transgenesis. Additionally, we have generated vectors for Tol2-based transgenesis of specific BirA and TRAP zebrafish lines and tested this system on skeletal muscle development.

RESULTS AND DISCUSSION

BirA activity in zebrafish embryos

We first sought to determine whether the biotin ligase system would be functional in zebrafish. We transiently expressed Avi-tagged EGFP and BirA in zebrafish embryos exposed to exogenous biotin and assessed the presence of biotinylated proteins with streptavidin-HRP (Fig. 1A) and streptavidin-Alexa Fluor 647 (supplementary material Fig. S1). We found that in addition to at least three endogenously biotinylated proteins (with approximate molecular

¹Department of Biochemistry & Biophysics, UCSF, San Francisco, CA 94158, USA.

²URBC-Narilis, University of Namur, Namur B-5000, Belgium.

*Present address: Department of Developmental Genetics, Max Planck Institute for Heart and Lung Research, Ludwigstrasse 43, Bad Nauheim 61231, Germany.

‡Present address: IBMM, Université Libre de Bruxelles, 12 rue des Prof. Jeener et Brachet, Gosselies B-6041, Belgium.

\$Authors for correspondence (michael.housley@mpi-bn.mpg.de; didier.stainier@mpi-bn.mpg.de; benoit.vanhollebeke@ulb.ac.be)




Fig. 1. Specific biotinylation of Avi-tagged EGFP in zebrafish embryos. (A) Avi-tagged EGFP (arrowhead) is selectively biotinylated when co-expressed with BirA, as shown by SDS-PAGE and streptavidin-HRP blotting (upper panel) of lysates from zebrafish embryos transiently expressing Avi-EGFP and BirA. The middle panel is an anti-GFP antibody blot confirming expression of Avi-EGFP. The bottom panel is an anti- β -tubulin (Tubb) antibody blot shown as a loading control. BirA and Avi-EGFP were transiently expressed by co-injection of 500 pg of *in vitro* transcribed mRNA into one-cell stage zebrafish embryos. Lysates for blotting analysis were generated from 24 hpf embryos. (B) Streptavidin gel shift assay (upper panel, anti-GFP antibody blot) performed on zebrafish embryos treated with indicated amounts of exogenous biotin following transient expression of Avi-EGFP and BirA. The lower panel is a streptavidin-HRP blot of the endogenously biotinylated 130 kDa protein shown as a loading control. Transient expression was performed as in A. (C) SDS-PAGE and Coomassie Blue G250 staining of streptavidin-agarose purified Avi-EGFP from lysates generated from 50 zebrafish embryos transiently expressing Avi-EGFP and BirA. Lane 2 contains 50 ng of BSA for comparison. Transient expression was performed by mRNA injection as in A. Thin lines in B,C mark positions where lanes from the same gel have been spliced together.

weights of 75, 130 and 260 kDa), a single streptavidin-reactive band consistent with the anticipated molecular weight of biotinylated EGFP could be readily detected only when Avi-tagged EGFP and the BirA ligase were co-expressed (arrowhead, Fig. 1A).

To confirm the identity of the streptavidin-reactive band, and to investigate the efficiency of the biotinylation process and determine if exogenous biotin is required, we performed a streptavidin shift assay on samples from embryos exposed to biotin ranging from 2 to 200 μM (Fig. 1B). In this assay, after co-expression of Avi-EGFP and BirA, zebrafish lysates were incubated with an excess of unlabeled streptavidin resulting in a ~ 53 kDa shift in apparent molecular weight when biotinylated. GFP immunodetection in such lysates revealed that in the presence of 50 μM of exogenous biotin, Avi-tagged EGFP appeared to be completely biotinylated (Fig. 1B, lane 6). A streptavidin-HRP blot of the endogenously biotinylated 130 kDa protein is shown as a loading control.

To determine whether we could recover biotinylated proteins from zebrafish lysates, we transiently biotinylated Avi-EGFP *in vivo* and performed a pull down using streptavidin agarose from a lysate of 50 zebrafish embryos (Fig. 1C). The purified product was eluted, separated by SDS-PAGE, stained with Coomassie Blue G250 (lane 3) and compared with 50 ng of BSA (lane 2), indicating that ~ 1 ng of Avi-EGFP protein could be purified per embryo. We therefore conclude that the BirA ligase is functional, at least transiently, in early zebrafish embryos, that ectopic expression of the bacterial enzyme results in no discernible developmental defects and that the Avi-tag may be used to purify fusion proteins.

Biotinylation of tagged ribosomes in zebrafish

Previous studies have shown that EGFP fused to the N terminus of the large subunit ribosomal protein L10a is compatible with its incorporation into functional polysomes in mammalian cells and in zebrafish (Heiman et al., 2008; Fang et al., 2013; Tryon et al., 2013).

We evaluated whether an N-terminal Avi-tagged version of EGFP-Rpl10a would be similarly incorporated into functional ribosomes and would act as a suitable substrate for the biotin ligase BirA *in vivo* through transient expression assays similar to those described above.

Probing zebrafish lysates with streptavidin-HRP revealed a reactive band compatible with biotinylated Avi-tagged EGFP-Rpl10a when BirA and Avi-EGFP-Rpl10a were co-expressed (arrowhead, Fig. 2A) suggesting that the Avi-tag is accessible to the biotin ligase in the ribosomal fusion protein. To evaluate whether the biotinylated ribosomal subunit was incorporated into functional ribosomes, we performed streptavidin affinity purification of biotinylated proteins under native conditions. Under conditions where the TRAP is expected to function, i.e. when both the ligase and the tagged ribosomal protein are co-expressed, we could co-purify proteins with molecular weights corresponding to ribosomal proteins (Fig. 2B; see also supplementary material Fig. S3 and Table S1 for further validation) as well as ribosomal RNA (Fig. 2C), indicating that the biotinylated ribosomes can be incorporated into functional polysomes.

A binary genetically encoded ribosome biotinylation system in zebrafish

After demonstrating that transiently expressed Avi-EGFP-Rpl10a was biotinylated in zebrafish, we tested whether this system could be used with stable transgenic lines. We generated a line stably expressing BirA-EGFP driven by the *ubiquitin b* promoter, *Tg(ubb:birA-EGFP)^{s972}* (Fig. 3A, supplementary material Fig. S2), as well as a skeletal muscle-specific tagged ribosome line, *Tg(actc1b:Avi-EGFP-rpl10a)^{s973}* (Fig. 3A). Confocal imaging revealed that ribosomes were distributed throughout the multinucleated skeletal muscle cells, which appeared to be unaffected by transgene expression, and that localized fluorescence patterns in the




Fig. 2. Biotin ligase-acceptor peptide combinatorial TRAP in zebrafish. (A) Streptavidin-HRP blot of lysates generated from *Tg(ubb:birA-EGFP)⁹⁷²* embryos transiently expressing Avi-EGFP-Rpl10a (arrowhead). Transient expression was performed by injecting 300 pg of *in vitro* transcribed mRNA into one-cell stage embryos. Blots represent protein from 10 de-yolked embryos harvested at 24 hpf. (B) Sypro Ruby protein stain of eluates separated by SDS-PAGE following TRAP performed on *Tg(ubb:birA-EGFP)⁹⁷²* embryos transiently expressing Avi-EGFP-Rpl10a. Transient expression was performed as in A, and represents protein from 100 embryos. (C) Ethidium bromide stain of eluates separated electrophoretically following TRAP and Trizol extraction performed on *Tg(ubb:birA-EGFP)⁹⁷²* embryos transiently expressing Avi-EGFP-Rpl10a. Transient expression was performed as in A and represents RNA from 100 embryos.

nucleus appeared to be consistent with import to the nucleolus, indicating that Avi-tagged Rpl10a was being incorporated into ribosomes (Krüger et al., 2007). To determine the efficiency of biotinylation of skeletal muscle Avi-EGFP-Rpl10a when exposed to BirA expressed under control of the *ubb* promoter, we compared *Tg(ubb:birA-EGFP)⁹⁷²* or *Tg(actc1b:Avi-EGFP-rpl10a)⁹⁷³* crossed with wild-type fish with *Tg(ubb:birA-EGFP)⁹⁷²* crossed with *Tg(actc1b:Avi-EGFP-rpl10a)⁹⁷³* by streptavidin shift assay (Fig. 3B). Efficient biotinylation is indicated by the complete shift of Avi-tagged Rpl10a (arrowhead, Fig. 3B, lane 3). *Tg(ubb:birA-EGFP)^{972/+}* and *Tg(actc1b:Avi-EGFP-rpl10a)^{973/+}* were then crossed, and the offspring were harvested at 1.5 dpf and sorted into two pools based on the presence of skeletal muscle fluorescence. TRAP was then performed on the offspring of this cross (using the protocol described in supplementary material Fig. S5), and RNAs and proteins were extracted and subjected to electrophoretic separation (Fig. 3C). Protein gels were stained with Sypro Ruby and revealed that many proteins, generally smaller than 50 kDa, were specifically isolated by the TRAP (Fig. 3C, compare lane 1 with lane 2). Ethidium bromide stained agarose gels revealed bands characteristic of ribosomal RNA (Fig. 3C, compare lane 3 with lane 4). To confirm that the bands in Fig. 3C were ribosomal proteins, we performed immuno-blotting for Rpl7a (supplementary material Fig. S3) and mass spectrometric analysis and recovered at least one peptide from 27 out of 32 proteins that make up the small ribosomal subunit and 26 out of 47 proteins that make up the large subunit (Fig. 3D, supplementary material Table S1) (Nakao et al., 2004). Conspicuously absent from this list was Rpl10a itself. The lack of the tagged ribosomal protein is not surprising, however; the conditions under which the ribosomes were eluted (high salt) are not harsh enough to elute biotinylated proteins from the streptavidin column.

To test whether we could detect mRNAs using the combinatorial BirA and Avi-tagged Rpl10a system, we crossed *Tg(ubb:birA-EGFP)⁹⁷²* to *Tg(actc1b:Avi-EGFP-rpl10a)⁹⁷³* and performed TRAP on 48 hpf embryos. We then reverse transcribed RNA purified from samples that were eluted from the streptavidin column, as well as input samples. We then assayed by qPCR five non-skeletal muscle genes (*tie1*, *cmic2*, *pax7a*, *pax7b* and *efnb3b*) and five genes known to be expressed in skeletal muscle (*myod1*, *myog*, *myf5*, *myf6* and *myl1*), and found that the relative enrichment of mRNA bound to the column was greater for the skeletal muscle-

specific genes (Fig. 3E), indicating that this method is useful for separating mRNAs from cells that may be tightly associated with skeletal muscle, such as endothelial cells (*tie1*) and satellite cells (*pax7a* and *pax7b*).

Upon confirmation that we could biotinylate and purify translating ribosomes in skeletal muscle using the combinatorial TRAP, we set out to apply this approach to determine which genes are being actively translated in skeletal muscle during myoblast fusion. We crossed *Tg(ubb:birA-EGFP)⁹⁷²* with *Tg(actc1b:Avi-EGFP-rpl10a)⁹⁷³* and harvested embryos at four time points (21, 24, 27 and 34 hpf) surrounding the beginning of myoblast fusion in zebrafish (24 hpf) (Moore et al., 2007). Ribosome-associated mRNAs were purified and analyzed by microarray (Fig. 4). Fig. 4A depicts the relative ribosome association of individual mRNAs at 21 hpf on the *x*-axis plotted versus 34 hpf on the *y*-axis. As expected, most genes fall along the 45° line, i.e. their ribosome-association is not differentially regulated during this developmental window. However, a subset of genes was found to be either up- or downregulated between 21 and 34 hpf. Well-known skeletal muscle genes, such as *mylpfb*, *myoz1b*, *atp2a1*, *myh2*, *ckmb*, *acta1b*, *tmt3a*, *tmi2a.4* and *pvalb3* were upregulated (Fig. 4A, red circles; supplementary material Table S2) indicating that the TRAP was indeed assaying expression in developing skeletal muscle. Downregulated genes (Fig. 4A, blue circles; supplementary material Table S2) include *mespaa*, *msgn1*, *tbx16*, *rippy2*, *lfi1*, *her1*, *her5*, *cdx4*, *wnt8a*, *tbx6* and *ved*, which are associated with mesoderm-related tissue specification and non-skeletal muscle cells (Sokol et al., 1991; Chapman et al., 1996; Muller et al., 1996; Griffin et al., 1998; Bisgrove et al., 1999; Bally-Cuif et al., 2000; Sawada et al., 2000; Yoon and Wold, 2000; Henry et al., 2002; Shimizu et al., 2002; Wang et al., 2005; Davidson and Zon, 2006; Morimoto et al., 2007), and are, as expected, expressed to a lesser extent in later skeletal muscle development.

To examine the changes over time in more detail, we plotted the relative ribosome association at 21, 24, 27 and 34 hpf for several interesting genes from the skeletal muscle TRAP (Fig. 4B, supplementary material Fig. S4). *tie1* encodes an endothelial cell-specific receptor tyrosine kinase (Lyons et al., 1998), and, as expected, is minimally expressed. *murca*, *murcb*, *rtn2a*, *rtn2b* and *tgm2a* were found to be upregulated throughout the time course, indicating a potential role in muscle development. To confirm that these mRNAs with novel regulation were skeletal muscle specific,




Fig. 3. Skeletal muscle-specific translating ribosome purification in zebrafish embryos. (A) Fluorescence image of *Tg(ubb:birA-EGFP)* expression (upper panel) at 48 hpf. Merged fluorescence and bright-field images of *Tg(actc1b:Avi-EGFP-rpl10a)* expression at 48 hpf in the middle two panels show lateral and dorsal views. The bottom panel is a confocal image of *Tg(actc1b:Avi-EGFP-rpl10a)* expression in skeletal muscle. Anterior is towards the left; dorsal towards the top. (B) Streptavidin shift assay with lysates from 15 embryos from *Tg(ubb:birA-EGFP)^{S972}* crossed with wild type (lane 1), from *Tg(actc1b:Avi-EGFP-rpl10a)^{S973}* crossed with wild type (lane 2), or from *Tg(ubb:birA-EGFP)^{S972}* crossed with *Tg(actc1b:Avi-EGFP-rpl10a)^{S973}* (lane 3). Arrowhead indicates Avi-EGFP-Rpl10a bound with free streptavidin. Protein from four larvae at 4 dpf was applied to each lane. (C) Eluates from the skeletal muscle TRAP performed on 200 embryos (36 hpf) from *Tg(ubb:birA-EGFP)^{S972}* crossed with *Tg(actc1b:Avi-EGFP-rpl10a)^{S973}*. Eluates were separated by SDS-PAGE and stained with Sypro Ruby protein stain (lane 1 is from sibling controls, whereas lane 2 is from embryos expressing both transgenes). Lanes 3 (sibling controls) and 4 (TRAP embryos) depict an ethidium bromide stained agarose gel following Trizol extraction. (D) Illustration of ribosomal proteins with at least one peptide recovered by mass spectrometry analysis of skeletal muscle TRAP eluates. The full list is provided in supplementary material Table S1. (E) qPCR of skeletal muscle and non-skeletal muscle genes from reverse transcribed mRNA purified by TRAP or from whole lysate (input) from 48 hpf embryos generated by crossing *Tg(ubb:birA-EGFP)^{S972}* to *Tg(actc1b:Avi-EGFP-rpl10a)^{S973}*. The data are shown as relative abundance of purified mRNA divided by total input mRNA (i.e. TRAP/input) normalized to *act1b*. Error bars represent s.e.m. from three independent experiments.

we performed *in situ* hybridizations and found that *murca*, *murch*, *rtn2a*, *rtn2b* and *tgm2a* mRNAs were localized to the somites in 28 hpf zebrafish embryos (Fig. 4C).

Conclusions

We have applied a combinatorial biotin ligase-acceptor peptide TRAP to muscle development where biotin ligase expression was driven by the *ubiquitin* promoter and the *Avi-EGFP-rpl10a* gene was under the control of the skeletal muscle-specific *actc1b* promoter. Although the TRAP methodology is limited by how well promoters have been defined as well as the time required to generate the transgenic organisms, the development of the Tol2 transposase system has improved the efficiency of generating transgenic fish

(Urasaki et al., 2006), mitigating some of the limitations associated with TRAP. The skeletal muscle TRAP described here provides a proof-of-principle test of the combinatorial biotin ligase and acceptor peptide-based TRAP during vertebrate embryogenesis. The combinatorial TRAP allows one to define a target cell population for analysis using two markers, as is possible with FACS, but retains the advantages associated with TRAP, such as minimal tissue manipulations, rapid cell lysis and assessment of ribosome-associated mRNAs and not total mRNA pools. Currently, we are generating additional stable TRAP and BirA lines to answer more detailed developmental questions. The increased specificity in gene expression analysis will aid in the discovery of novel biomarkers and targets for drug screening, suggest ways to




Fig. 4. Translational profiling of zebrafish skeletal muscle. (A) Expression profile of 21 hpf versus 34 hpf skeletal muscle-specific TRAP mRNA assayed by microarray hybridization. Dots represent normalized hybridization signals of individual genes plotted on a log scale. Blue circles indicate downregulation over time whereas red circles indicate upregulation. The yellow line represents no difference in expression between samples whereas the black lines depict a fivefold change.

(B) Timecourse of relative expression (log scale, y-axis) of selected genes from A. *tie1* is an endothelial-specific gene and is minimally expressed. *mesogenin* (*msgn1*) is an example of a gene whose expression is downregulated during skeletal muscle development. (C) Whole-mount *in situ* hybridization at 28 hpf of selected upregulated mRNAs. Dorsal (left) and lateral (right) views.

challenge mutants generated by reverse genetic approaches, and further our understanding of cell-fate decisions, as well as cell behavior and function.

MATERIALS AND METHODS

DNA constructs, RNA synthesis and zebrafish lines

TRAP plasmids are based on the mini Tol2 transposase vectors (Urasaki et al., 2006) and are available at Addgene. See supplementary methods for details.

Antibodies, staining, blotting, gel shift assays and protein purification

Streptavidin shift assays were performed by lysing 10 embryos in Laemmli buffer, boiling for 5 minutes, then incubating with 30 μ g streptavidin (Sigma) prior to SDS-PAGE and anti-GFP blotting. See supplementary methods for details.

TRAP

Embryos were collected in egg water supplemented with 50 μ M biotin. Prior to harvesting at indicated time points, embryos were washed for 30 min in egg water to remove unincorporated biotin. Following

dechoriation, embryos were washed in cold Ringer's solution, pelleted in 1.5 ml tubes and snap frozen on dry ice. Samples were lysed in 0.5 ml ice-cold polysome buffer (supplementary material Fig. S5). Lysates were cleared by centrifugation and applied to 100 μ l of streptavidin-agarose (Pierce) followed by rotation at 4°C for 1 h. Beads were then pipetted into microspin columns (Bio-Rad) and washed twice with 1 ml wash buffer. RNAs and ribosomal proteins were eluted with 0.2 ml elution buffer and immediately added to 1 ml Trizol reagent (Invitrogen).

Mass spectrometry and protein identification

Peptides were analyzed by using nano-LC-ESI-MS/MS maXis Impact UHR-TOF (Bruker, Bremen, Germany) coupled with a 2D-LC Dionex UltiMate 3000 (Thermo). See supplementary methods for details.

RT-PCR and microarray profiling

Two-color microarray hybridization was performed using Zebrafish V3 44K gene expression arrays (Agilent Technologies) and the data have been deposited at NCBI (GEO GSE59355). See supplementary methods for details.

Whole-mount *in situ* hybridization

Antisense probes were generated from cDNA using PCR primers listed in supplementary material Table S3. PCR products were cloned into pGEM-T Easy (Promega) and whole-mount *in situ* hybridizations were performed according to established protocols (Thisse and Thisse, 2008).

Acknowledgements

We thank A. Ayala, M. Alva, P. Lopez Pazmino and M. Sklar for zebrafish husbandry. We also thank O. Stone for reading the manuscript.

Competing interests

The authors declare no competing financial interests.

Author contributions

B.V. developed the concept. M.P.H., S.R. and B.V. designed the experimental set up. M.P.H., M.D., M.R. and B.V. performed experiments and data analysis. M.P.H., D.Y.R.S. and B.V. prepared and edited the manuscript. D.Y.R.S. supervised the work.

Funding

The research was made possible by a grant from the California Institute for Regenerative Medicine [TG2-01153] to M.P.H., as well as by funds from the National Institutes of Health, Packard Foundation and Max Planck Society to D.Y.R.S. The MS equipment of the MS facility of the URBC-Narilis was funded by a Fonds de la Recherche Fondamentale Collective (FRFC) grant of the Fonds de la Recherche Scientifique (FRS-FNRS). B.V. was supported by postdoctoral fellowships from European Molecular Biology Organization (EMBO) and the Human Frontier Science Program (HFSP). Deposited in PMC for release after 12 months.

Supplementary material

Supplementary material available online at <http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.111849/-/DC1>

References

- Bally-Cuif, L., Goutel, C., Wassef, M., Wurst, W. and Rosa, F. (2000). Coregulation of anterior and posterior mesendodermal development by a hairy-related transcriptional repressor. *Genes Dev.* **14**, 1664-1677.
- Bisgrove, B. W., Essner, J. J. and Yost, H. J. (1999). Regulation of midline development by antagonism of lefty and nodal signaling. *Development* **126**, 3253-3262.
- Chapman, D. L., Agulnik, I., Hancock, S., Silver, L. M. and Papaioannou, V. E. (1996). Tbx6, a mouse T-box gene implicated in paraxial mesoderm formation at gastrulation. *Dev. Biol.* **180**, 534-542.
- Davidson, A. J. and Zon, L. I. (2006). The caudal-related homeobox genes *cdx1a* and *cdx4* act redundantly to regulate *hox* gene expression and the formation of putative hematopoietic stem cells during zebrafish embryogenesis. *Dev. Biol.* **292**, 506-518.
- Fang, Y., Gupta, V., Karra, R., Holdway, J. E., Kikuchi, K. and Poss, K. D. (2013). Translational profiling of cardiomyocytes identifies an early Jak1/Stat3 injury response required for zebrafish heart regeneration. *Proc. Natl. Acad. Sci. USA* **110**, 13416-13421.
- Gay, L., Miller, M. R., Ventura, P. B., Devasthali, V., Vue, Z., Thompson, H. L., Temple, S., Zong, H., Cleary, M. D., Stankunas, K. et al. (2013). Mouse TU tagging: a chemical/genetic intersectional method for purifying cell type-specific nascent RNA. *Genes Dev.* **27**, 98-115.
- Griffin, K. J., Amacher, S. L., Kimmel, C. B. and Kimmel, D. (1998). Molecular identification of spadetail: regulation of zebrafish trunk and tail mesoderm formation by T-box genes. *Development* **125**, 3379-3388.
- Heiman, M., Schaefer, A., Gong, S., Peterson, J. D., Day, M., Ramsey, K. E., Suárez-Fariñas, M., Schwarz, C., Stephan, D. A., Surmeier, D. J. et al. (2008). A translational profiling approach for the molecular characterization of CNS cell types. *Cell* **135**, 738-748.
- Henry, C. A., Urban, M. K., Dill, K. K., Merlie, J. P., Page, M. F., Kimmel, C. B. and Amacher, S. L. (2002). Two linked hairy/Enhancer of split-related zebrafish genes, *her1* and *her7*, function together to refine alternating somite boundaries. *Development* **129**, 3693-3704.
- Krüger, T., Zentgraf, H. and Scheer, U. (2007). Intranucleolar sites of ribosome biogenesis defined by the localization of early binding ribosomal proteins. *J. Cell Biol.* **177**, 573-578.
- Kunitomo, H., Uesugi, H., Kohara, Y. and Iino, Y. (2005). Identification of ciliated sensory neuron-expressed genes in *Caenorhabditis elegans* using targeted pull-down of poly(A) tails. *Genome Biol.* **6**, R17.
- Lyons, M. S., Bell, B., Stainier, D. and Peters, K. G. (1998). Isolation of the zebrafish homologues for the *tie-1* and *tie-2* endothelium-specific receptor tyrosine kinases. *Dev. Dyn.* **212**, 133-140.
- Miller, M. R., Robinson, K. J., Cleary, M. D. and Doe, C. Q. (2009). TU-tagging: cell type specific RNA isolation from intact complex tissues. *Nat. Methods* **6**, 439-441.
- Moore, C. A., Parkin, C. A., Bidet, Y. and Ingham, P. W. (2007). A role for the Myoblast city homologues *Dock1* and *Dock5* and the adaptor proteins *Crk* and *Crk-like* in zebrafish myoblast fusion. *Development* **134**, 3145-3153.
- Morimoto, M., Sasaki, N., Oginuma, M., Kiso, M., Igarashi, K., Aizaki, K.-I., Kanno, J. and Saga, Y. (2007). The negative regulation of *Mesp2* by mouse *Ripply2* is required to establish the rostro-caudal patterning within a somite. *Development* **134**, 1561-1569.
- Muller, M., Weizsacker, E. and Campos-Ortega, J. A. (1996). Expression domains of a zebrafish homologue of the *Drosophila* pair-rule gene *hairy* correspond to primordia of alternating somites. *Development* **122**, 2071-2078.
- Nakao, A., Yoshihama, M. and Kenmochi, N. (2004). RPG: the ribosomal protein gene database. *Nucleic Acids Res.* **32**, D168-D170.
- Roy, P. J., Stuart, J. M., Lund, J. and Kim, S. K. (2002). Chromosomal clustering of muscle-expressed genes in *Caenorhabditis elegans*. *Nature* **418**, 975-979.
- Sawada, A., Fritz, A., Jiang, Y., Yamamoto, A., Yamasu, K., Kuroiwa, A., Saga, Y. and Takeda, H. (2000). Zebrafish *Mesp* family genes, *mesp-a* and *mesp-b* are segmentally expressed in the presomitic mesoderm, and *Mesp-b* confers the anterior identity to the developing somites. *Development* **127**, 1691-1702.
- Shimizu, T., Yamanaka, Y., Nojima, H., Yabe, T., Hibi, M. and Hirano, T. (2002). A novel repressor-type homeobox gene, *ved*, is involved in dharma/bozozok-mediated dorsal organizer formation in zebrafish. *Mech. Dev.* **118**, 125-138.
- Sokol, S., Christian, J. L., Moon, R. T. and Melton, D. A. (1991). Injected Wnt RNA induces a complete body axis in *Xenopus* embryos. *Cell* **67**, 741-752.
- Thisse, C. and Thisse, B. (2008). High-resolution *in situ* hybridization to whole-mount zebrafish embryos. *Nat. Protoc.* **3**, 59-69.
- Tryon, R. C., Pisat, N., Johnson, S. L. and Dougherty, J. D. (2013). Development of translating ribosome affinity purification for zebrafish. *Genesis* **51**, 187-192.
- Urasaki, A., Morvan, G. and Kawakami, K. (2006). Functional dissection of the *Tol2* transposable element identified the minimal cis-sequence and a highly repetitive sequence in the subterminal region essential for transposition. *Genetics* **174**, 639-649.
- Wang, Y., Yates, F., Naveiras, O., Ernst, P. and Daley, G. Q. (2005). Embryonic stem cell-derived hematopoietic stem cells. *Proc. Natl. Acad. Sci. USA* **102**, 19081-19086.
- Wysocki, L. J. and Sato, V. L. (1978). "Panning" for lymphocytes: a method for cell selection. *Proc. Natl. Acad. Sci. USA* **75**, 2844-2848.
- Yoon, J. K. and Wold, B. (2000). The bHLH regulator *pMesogenin1* is required for maturation and segmentation of paraxial mesoderm. *Genes Dev.* **14**, 3204-3214.

Supplementary Methods

DNA Constructs, RNA Synthesis, and Zebrafish Lines. An N-terminal Avi-EGFP fusion construct was generated by PCR primer extension and cloned into pcGlobin (Ro et al., 2004) using BamHI and XhoI restriction sites. *rpl10a* (Fang et al., 2013) was cloned into the pcGlobin-N-Avi-EGFP vector. mRNA was generated using the mMessage mMachine kit (Invitrogen) and injected into one-cell stage zebrafish embryos. *Avi-EGFP-rpl10a* was sub-cloned into pT2KXIG in (Urasaki et al., 2006) to generate pT2KXIG in-MCS-Avi-EGFP-rpl10a (addgene.org/58380, Fig. S5). A 4 kb fragment of the *actc1b* promoter (Higashijima et al., 1997) was cloned by PCR into the pT2KXIG in-MCS-Avi-EGFP-rpl10a construct and used to generate the line *Tg(actc1b:Avi-EGFP-rpl10a)^{s973}*. *birA* (addgene.org/20857) was fused by PCR to EGFP and sub-cloned into pT2KXIG in to generate pT2KXIG in-MCS-BirA-EGFP (addgene.org/58377, Fig. S5). tagRFP was subcloned in place of EGFP to generate pT2KXIG in-MCS-BirA-tagRFP (addgene.org/58378, Fig. S5) The *ubb* promoter (addgene.org/27320) was sub-cloned into pT2KXIG in-MCS-BirA-EGFP to generate the line *Tg(ubb:birA-EGFP)^{s972}*.

Mass Spectrometry and Protein Identification. Peptides were analyzed by using nano-LC-ESI-MS/MS maXis Impact UHR-TOF (Bruker, Bremen, Germany) coupled with a 2D-LC Dionex UltiMate 3000 (Thermo). Spots were excised manually and proteins were digested with trypsin by in-gel digestion. The gel pieces were washed twice with distilled water and then shrunk with 100 % acetonitrile. The proteolytic digestion was performed by the addition of 8 μ l of modified trypsin (Promega) suspended in 50 mM NH_4HCO_3 cold buffer. Proteolysis was performed overnight at 37°C. The supernatant was collected and the eluates were kept at – 20°C prior to analysis.

The digests were separated by reverse-phase liquid chromatography using a 75 μ m X 150 mm reverse phase Thermo column (Acclaim PepMap 100 C18) in an Ultimate 3000 liquid chromatography system. Mobile phase A was 95 % of 0.1 % formic acid in water and 5 % acetonitrile. Mobile phase B was 0.1 % formic acid in acetonitrile. The digest (15 μ l) was injected, and the organic content of the mobile phase was increased linearly from 5 % B to 40 % B in 25 min and from 40 % B to 100 % B in 5min. The column effluent was connected to a Captive Spray (Bruker). In survey scan, MS spectra were acquired for 0.5 s in the m/z range between 50 and 2200. The 10 most intense peptides ions 2+ or 3+ were sequenced. The collision-induced dissociation (CID) energy was automatically set according to mass to charge (m/z) ratio and charge state of the precursor ion. MaXis and Thermo systems were piloted by Compass HyStar 3.2 (Bruker).

Peak lists were created using DataAnalysis 4.0 (Bruker) and saved as an XML file for use with ProteinScape 3.1 (Bruker) with Mascot 2.4 as the search engine (Matrix Science). Enzyme specificity was set to trypsin, and the maximum number of missed cleavages per peptide was set at one. Carbamidomethylation was allowed as fixed modification, oxidation of methionine and Gln – pyro-Glu were allowed as variable modification. Mass tolerance for monoisotopic peptide window was 5 ppm and MS/MS tolerance window was set to 0.05 Da. The peak lists were searched against the NCBI nr database.

Antibodies, Staining, Blotting, Gel Shift Assays, and Protein Purification. Anti-GFP antibodies (Aves Lab), anti-Tubb (Abcam), anti-Rpl7a (Genetex), and streptavidin-HRP (Pierce) were used in blotting experiments. Streptavidin gel shift assays were performed

by lysing 10 embryos in Laemmli buffer, boiling for 5 minutes, then incubating with 30 µg streptavidin (Sigma) prior to SDS-PAGE and anti-GFP blotting. Protein Staining was done with Sypro Ruby and visualized on a blue light trans-illuminator (Invitrogen) or using Coomassie Blue G-250 (Bio-Rad) as indicated. Avi-EGFP was purified by harvesting dechorionated and deyolked embryos, boiling in 1% SDS, and diluting samples with PBS. Streptavidin agarose was then incubated with samples for 30 minutes at room temperature after which the resin was washed twice and boiled with Laemmli buffer containing 1mM biotin for 10 minutes prior to SDS-PAGE. Whole-mount streptavidin staining was performed on PFA-fixed (4%, overnight at 4°C) embryos at 24 hpf. Embryos were permeabilized with proteinase K (10 µg/ml for 5 minutes), and fixed again with 4% PFA for 15 min in PBS-Tween 20 (0.1%). After PBS washes, embryos were blocked in PBS with 0.3% Triton X-100 and 4% BSA for 2 hours at room temperature. Staining was performed in blocking buffer with streptavidin-Alexa Fluor 647 (2 µg/ml, Life Technologies) overnight at 4°C followed by washes with PBS 0.3% Triton X-100.

RT-PCR and Microarray Profiling. For RT-PCR, cDNA was generated from equal amounts of total RNA using Maxima reagents (Thermo Scientific) and qPCR analysis was performed on an Illumina Eco system using SYBR Green (Thermo Scientific). Primer pairs are listed in Table S3. For qPCR, 3 technical replicates were performed for each TRAP experiment and averaged. To determine relative enrichment, fold differences between bound mRNA and input mRNA were calculated by subtracting threshold values and adjusting for amounts added to the reverse transcription reaction. Fold enrichment was then normalized to *act1b*. Two-color labeling and microarray hybridization was performed by Mogene using Zebrafish V3 44K gene expression arrays (Agilent Technologies) with 50 ng of RNA purified by TRAP. Amplification was performed with the NuGEN Pico Amp kit. Data were acquired using Agilent Feature Extraction with LOWESS dye normalization. Normalization across arrays was done using GeneSpring 11.5 default values. The data have been deposited at NCBI (GEO GSE59355).

References

- Fang, Y., Gupta, V., Karra, R., Holdway, J. E., Kikuchi, K. and Poss, K. D.** (2013). Translational profiling of cardiomyocytes identifies an early Jak1/Stat3 injury response required for zebrafish heart regeneration. *Proc. Natl. Acad. Sci.* 201309810.
- Higashijima, S., Okamoto, H., Ueno, N., Hotta, Y. and Eguchi, G.** (1997). High-Frequency Generation of Transgenic Zebrafish Which Reliably Express GFP in Whole Muscles or the Whole Body by Using Promoters of Zebrafish Origin. *Dev. Biol.* **192**, 289–299.
- Ro, H., Soun, K., Kim, E.-J. and Rhee, M.** (2004). Novel vector systems optimized for injecting in vitro-synthesized mRNA into zebrafish embryos. *Mol. Cells* **17**, 373–376.
- Urasaki, A., Morvan, G. and Kawakami, K.** (2006). Functional Dissection of the Tol2 Transposable Element Identified the Minimal cis-Sequence and a Highly Repetitive Sequence in the Subterminal Region Essential for Transposition. *Genetics* **174**, 639–649.

Fig. S1




Fig. S1. Streptavidin-conjugated Alexa Fluor 647 staining of embryos expressing Avi-tagged EGFP. 24 hpf embryos were injected with either *BirA*, *Avi-EGFP*, or both *birA* and *Avi-EGFP* mRNA and stained with streptavidin-Alexa Fluor 647.

Fig. S2




Fig. S2. Fluorescence images of *Tg(ubb:birA-EGFP)* expression at the indicated developmental stages.

Fig. S3




Fig. S3. Rpl7a immunoblotting of TRAP samples. Embryos injected with RNA for either *Avi-EGFP-rpl10a* or both *BirA* and *Avi-EGFP-rpl10a* were harvested at 24 hpf. Following TRAP, samples were subjected to SDS-PAGE, and analyzed by immunoblotting with anti-Rpl7a antibodies. Input samples for the TRAP were also blotted with anti-Rpl7a antibodies and streptavidin-HRP to confirm biotinylation.

Fig. S4




Fig. S4. Microarray expression data from two skeletal muscle TRAP time courses. Relative mRNA expression data from selected genes (log scale, y-axis) from skeletal muscle-specific TRAP (*Tg(ubb:birA-EGFP)^{s972}* crossed to *Tg(actc1b:Avi-EGFP-rpl10a)^{s973}*) assayed by microarray hybridization. Array 1 is reproduced from Fig. 4B. Array 2 represents expression from a second time course, TRAP, and microarray experiment. *tie1* is an endothelial-specific gene and is minimally expressed. *mesogenin (msgn1)* is an example of a gene whose expression is down-regulated during skeletal muscle development.

Fig. S5. Combinatorial Biotin Ligase - Avi-tag TRAP Protocol

Clone promoters of interest into TRAP vectors and generate transgenic lines

Image transgenic lines to confirm expression of BirA and Avi-EGFP-Rpl10a

Cross Tg(promoter a:birA-EGFP) to Tg(promoter b:Avi-EGFP-rpl10a) fish

Collect embryos and add 50 μ M biotin to eggwater

Incubate embryos to desired developmental stage, then wash out biotin

Collect and lyse embryos in polysome buffer

Bind labeled polysomes to streptavidin agarose


Transfer beads to column and wash 3 times

Elute RNA and ribosomal proteins

Extract RNA with Trizol

Purify and concentrate RNA on a micro spin column

Elute RNA and measure concentration



Dissolve biotin at 50 mM in DMSO then add to eggwater at a final concentration of 50 μ M. Change biotin-containing eggwater every 24 hours. Always make a fresh biotin solution and we recommend storing powdered biotin under inert gas.

Incubate embryos until desired developmental stage is reached. Wash away unincorporated biotin which would block streptavidin binding sites on the column by first anesthetizing fish, collecting them in a strainer, rinsing with egg water, and transferring them to a new dish containing fresh egg water. Incubate fish for 30 minutes. Prepare polysome buffer and streptavidin agarose columns or dry ice to snap freezing samples. Prepare cycloheximide fresh each time by dissolving 10 mg/ml in DMSO. Store under inert gas.

Label a clean, RNase-free, 1.5ml tube for each sample and add 100 μ l streptavidin agarose slurry. Add 1 ml polysome buffer to equilibrate the column. Incubate at room temperature for 10 minutes, pellet beads, remove supernatant, and place tubes on ice.

Collect embryos into a 1.5 ml tube, place on ice, wash once in cold Ringer's solution, and pellet embryos. Tail bud and younger embryos should be de-yolked by gentle pipetting with a p200, followed by pelleting of cells, and one wash in Ringer's. Snap freeze on dry ice or continue with the TRAP.

Lyse fish in 0.5 ml ice-cold polysome buffer with the aid of a clean pestle. Incubate on ice for 10 minutes then centrifuge at 4 $^{\circ}$ C for 5 min at 8000 g.

Remove 20 μ l of the supernatant and add to 1 ml Trizol for the input sample, if desired. Add remaining supernatant to polysome buffer equilibrated beads. Gently rotate at 4 $^{\circ}$ C for one hour. Prepare Wash and Elution Buffers.

Apply beads to an empty column and drain by gravity flow. Wash 3 times with 1 ml ice cold wash buffer.

Place spin column in a clean 1.5 ml tube. Elute RNA with 100 μ l elution buffer. Elute a second time with 100 μ l. Immediately add 1 ml Trizol to eluate.

Extract RNA by adding 200 μ l chloroform. Shake tubes and centrifuge at 12,000 g for 10 min at 4 $^{\circ}$ C. Prepare binding buffer for Zymo Clean & Concentrate RNA columns. Remove aqueous phase to a fresh 2 ml tube*.

Add two volumes of Binding Buffer per volume of aqueous phase. Mix. Add to the RNA column in multiple, successive loads. Wash column 3 times with 0.7 ml 80% ethanol 10 mM Tris pH 7.5*. With each wash, cap and invert column several times. Centrifuge 30 seconds.

Dry column by centrifugation. Transfer the column to a fresh tube and dry again.

Transfer the column to a fresh tube. Elute twice with 6 μ l water.

Notes:

Use clean, RNase-free tubes and barrier pipette tips and solutions.
*Deviates from manufacturer's recommendations
Catalog Numbers:
Pestles: Axygen PES-15-B-SI
Empty columns: BioRad 732-6204
RNA columns: Zymo R1016
Cycloheximide: Cell Signaling 2112s
Biotin: Thermo 29129
Streptavidin Agarose: Pierce 20359
RNasin: Promega N2511

Calcium free Ringer's:
116 mM NaCl
2.9 mM KCl
5.0 mM Hepes 7.2
Plasmids available from addgene.org
58377
58378
58380

Polysome Buffer:
10 mM Hepes pH 7.4
150 mM KCl
5 mM MgCl₂
1% Triton X-100
add fresh each time:
0.5 mM DTT
0.5 mM PMSF
10 units/ml RNasin
100 μ g/ml cycloheximide

Wash Buffer:
10 mM Hepes
300 mM KCl
5 mM MgCl₂
1% Triton X-100
add fresh each time:
0.5 mM DTT
0.5 mM PMSF
10 units/ml RNasin
100 μ g/ml cycloheximide

Elution Buffer:
1 M NaCl
10 mM Tris pH 7.5

Zymo RNA Column Binding Buffer:
1 volume binding buffer concentrate
1 volume 95% ethanol
(0.5 ml each per column is generally sufficient)

Table S1.**Mass spectrometric identification of ribosomal proteins from the skeletal-muscle specific TRAP.**

Small Subunit Protein	Gene Name	Accession Number	Description	Mascot Score	Peptides
SA	<i>rpsa</i>	gi 41054972	ribosomal protein SA [Danio rerio]	376.65	51
S2	<i>rps2</i>	gi 47086117	ribosomal protein S2 [Danio rerio]	113.52	5
S3	<i>rps3</i>	gi 37595356	ribosomal protein S3 [Danio rerio]	846.98	138
S3A	<i>rps3a</i>	gi 41152457	ribosomal protein S3A [Danio rerio]	832.99	61
S4	<i>rps4x</i>	gi 53933236	ribosomal protein S4, X-linked [Danio rerio]	597.65	57
S5	<i>rps5</i>				
S6	<i>rps6</i>	gi 28630207	ribosomal protein S6 [Scyliorhinus canicula]	84.22	10
S7	<i>rps7</i>	gi 41152175	ribosomal protein S7 [Danio rerio]	208.17	10
S8	<i>rps8</i>	gi 47550881	ribosomal protein S8 [Danio rerio]	380.39	41
S9	<i>rps9</i>	gi 198285475	ribosomal protein S9 [Salmo salar]	474.78	37
S10	<i>rps10</i>	gi 148235024	40S ribosomal protein S10 [Xenopus laevis]	204.41	13
S11	<i>rps11</i>	gi 47087057	40S ribosomal protein S11 [Danio rerio]	186.2	9
S12	<i>rps12</i>	gi 119568395	ribosomal protein S12, isoform CRA_c [Homo sapiens]	37.94	1
S13	<i>rps13</i>	gi 50344812	40S ribosomal protein S13 [Danio rerio]	314.67	15
S14	<i>rps14</i>	gi 41152464	ribosomal protein S14 [Danio rerio]	365.26	31
S15	<i>rps15</i>	gi 12733945	40S ribosomal protein S15a [Platichthys flesus]	24.85	1
S15A	<i>rps15a</i>	gi 12733945	40S ribosomal protein S15a [Platichthys flesus]	24.85	1
S16	<i>rps16</i>	gi 115529347	ribosomal protein S16 [Danio rerio]	429.71	21
S17	<i>rps17</i>	gi 41053565	40S ribosomal protein S17 [Danio rerio]	196.17	13
S18	<i>rps18</i>				
S19	<i>rps19</i>	gi 41152179	ribosomal protein S19 [Danio rerio]	159.94	4
S20	<i>rps20</i>	gi 47086001	40S ribosomal protein S20 [Danio rerio]	123.79	7
S21	<i>rps21</i>	gi 41055255	40S ribosomal protein S21 [Danio rerio]	115.05	3
S23	<i>rps23</i>				
S24	<i>rps24</i>	gi 220678148	ribosomal protein S24 [Danio rerio]	243.04	7
S25	<i>rps25</i>				
S26	<i>rps26</i>	gi 12641796	40S ribosomal protein S26 [Platichthys flesus]	126.68	17
S27	<i>rps27</i>	gi 226958509	40S ribosomal protein S27-like [Danio rerio]	139.58	5
S27A	<i>rps27a</i>	gi 226358625	ribosomal protein S27a [Hypophthalmichthys nobilis]	18.76	1
S28	<i>rps28</i>	gi 183212473	ribosomal protein S28 [Xenopus borealis]	56.36	3
S29	<i>rps29</i>				
S30	<i>rps30</i>	gi 149518006	40S ribosomal protein S30-like, partial [Ornithorhynchus anatinus]	48.17	1
Large Subunit Protein	Gene Name	Accession Number	Description	Mascot Score	Peptides
L3	<i>rpl3</i>	gi 60688481	Ribosomal protein L3 [Danio rerio]	428.42	17

L4	<i>rpl4</i>	gi 54261775	60S ribosomal protein L4 [Danio rerio]	500.09	18
L5	<i>rpl5</i>	gi 50344868	ribosomal protein L5 [Danio rerio]	301.57	29
L6	<i>rpl6</i>	gi 62202562	Ribosomal protein L6 [Danio rerio]	187.71	24
L7	<i>rpl7</i>				
L7A	<i>rpl7a</i>	gi 41152461	60S ribosomal protein L7a [Danio rerio]	425.05	24
L8	<i>rpl8</i>	gi 116488054	60S ribosomal protein L8 [Scophthalmus maximus]	197.92	14
L9	<i>rpl9</i>	gi 3088347	ribosomal protein L9 [Homo sapiens]	30.72	1
L10	<i>rpl10</i>				
L10A	<i>rpl10a</i>				
L11	<i>rpl11</i>	gi 209737976	60S ribosomal protein L11 [Salmo salar]	79.61	7
L12	<i>rpl12</i>	gi 60551102	Ribosomal protein L12 [Danio rerio]	260.25	15
L13	<i>rpl13</i>	gi 47086477	60S ribosomal protein L13a [Danio rerio]	149.33	8
L13A	<i>rpl13a</i>	gi 47086477	60S ribosomal protein L13a [Danio rerio]	149.33	8
L14	<i>rpl14</i>				
L15	<i>rpl15</i>	gi 31322590	ribosomal protein L15 [Coturnix japonica]	97.11	2
L17	<i>rpl17</i>	gi 47086529	60S ribosomal protein L17 [Danio rerio]	102.23	4
L18	<i>rpl18</i>	gi 51010947	60S ribosomal protein L18 [Danio rerio]	253.34	18
L18A	<i>rpl18a</i>	gi 41055022	60S ribosomal protein L18a [Danio rerio]	178.4	9
L19	<i>rpl19</i>				
L21	<i>rpl21</i>	gi 50344966	ribosomal protein L21 [Danio rerio]	119.57	6
L22	<i>rpl22</i>	gi 83415112	60S ribosomal protein L22 [Danio rerio]	337.05	23
L23	<i>rpl23</i>	gi 41282078	60S ribosomal protein L23 [Danio rerio]	58.7	3
L23A	<i>rpl23a</i>				
L24	<i>rpl24</i>	gi 318101831	60S ribosomal protein L24 [Ictalurus punctatus]	179.63	14
L26	<i>rpl26</i>	gi 47085861	60S ribosomal protein L26 [Danio rerio]	159.06	9
L27	<i>rpl27</i>	gi 41054351	60S ribosomal protein L27 [Danio rerio]		
L27A	<i>rpl27a</i>	gi 41053327	60S ribosomal protein L27a [Danio rerio]	89.77	8
L28	<i>rpl28</i>	gi 41055030	ribosomal protein L28-like [Danio rerio]	79.55	3
L29	<i>rpl29</i>				
L30	<i>rpl30</i>	gi 119612175	ribosomal protein L30, isoform CRA_a [Homo sapiens]	163.07	12
L31	<i>rpl31</i>				
L32	<i>rpl32</i>				
L34	<i>rpl34</i>				
L35	<i>rpl35</i>				
L35A	<i>rpl35a</i>				
L36	<i>rpl36</i>	gi 47086131	60S ribosomal protein L36 [Danio rerio]	101.43	8
L36A	<i>rpl36a</i>				
L37	<i>rpl37</i>				
L37A	<i>rpl37a</i>				
L38	<i>rpl38</i>				
L39	<i>rpl39</i>				
L40	<i>rpl40</i>				
L41	<i>rpl41</i>				
LP0	<i>rplp0</i>				
LP1	<i>rplp1</i>				
LP2	<i>rplp2</i>	gi 154426308	60S acidic ribosomal protein P2 [Danio rerio]	283.67	36

Table S2.**Detailed information on genes from skeletal muscle TRAP highlighted in Fig. 4.**

Gene Symbol	Gene Name	21 hpf	34 hpf	Fold Change	zfin link
<i>mespaa</i>	<i>mesoderm posterior aa</i>	203.1	1.2	166.9	http://zfin.org/ZDB-GENE-000406-8
<i>her7</i>	<i>hairy and enhancer of split related-7</i>	110	1.5	74.7	http://zfin.org/ZDB-GENE-000427-6
<i>cdx1a</i>	<i>caudal type homeo box transcription factor 1 a</i>	114.6	1.7	68.8	http://zfin.org/ZDB-GENE-050510-1
<i>msgn1</i>	<i>mesogenin 1</i>	5631.2	98.4	57.2	http://zfin.org/ZDB-GENE-030722-1
<i>tbx16</i>	<i>T-box gene 16</i>	692.4	15.3	45.4	http://zfin.org/ZDB-GENE-990615-5
<i>rippy2</i>	<i>rippy2</i>	203	4.9	41.6	http://zfin.org/ZDB-GENE-060113-2
<i>lft1</i>	<i>lefty1</i>	668.3	24.6	27.2	http://zfin.org/ZDB-GENE-990630-10
<i>her1</i>	<i>hairy-related 1</i>	207.1	8.3	25.1	http://zfin.org/ZDB-GENE-980526-125
<i>her5</i>	<i>hairy-related 5</i>	136.7	7.3	18.7	http://zfin.org/ZDB-GENE-990415-90
<i>cdx4</i>	<i>caudal type homeo box transcription factor 4</i>	2127.1	124.3	17.1	http://zfin.org/ZDB-GENE-980526-330
<i>wnt8a</i>	<i>wingless-type MMTV integration site family, member 8a</i>	146.4	10.2	14.3	http://zfin.org/ZDB-GENE-980526-332
<i>tbx6</i>	<i>T-box gene 6</i>	965.4	70.6	13.7	http://zfin.org/ZDB-GENE-020416-5
<i>ved</i>	<i>ventrally expressed dharma/bozozok antagonist</i>	3285.3	280.5	11.7	http://zfin.org/ZDB-GENE-030813-1
<i>rtn2a</i>	<i>reticulon 2a</i>	83.3	625.3	7.5	http://zfin.org/ZDB-GENE-060420-1
<i>rtn2b</i>	<i>reticulon 2b</i>	146.2	1615.7	11.1	http://zfin.org/ZDB-GENE-060331-95
<i>acta1b</i>	<i>actin, alpha 1b, skeletal muscle</i>	2872.6	47773.2	16.6	http://zfin.org/ZDB-GENE-030131-55
<i>murc</i>	<i>muscle-related coiled-coil protein b</i>	67.8	1134.1	16.7	http://zfin.org/ZDB-GENE-041212-87
<i>ckmb</i>	<i>creatine kinase, muscle b</i>	1368.8	26470.4	19.3	http://zfin.org/ZDB-GENE-040426-2128
<i>myhz2</i>	<i>myosin, heavy polypeptide 2, fast muscle specific</i>	609	12955.6	21.3	http://zfin.org/ZDB-GENE-020604-1
<i>myoz1b</i>	<i>myozenin 1b</i>	176.6	3830.3	21.7	http://zfin.org/ZDB-GENE-040718-146
<i>atp2a1</i>	<i>ATPase, Ca⁺⁺ transporting, cardiac muscle, fast twitch 1</i>	2180.5	48618.4	22.3	http://zfin.org/ZDB-GENE-020905-1
<i>tgm2a</i>	<i>transglutaminase 2, C polypeptide A</i>	58.9	1545.5	26.2	http://zfin.org/ZDB-GENE-040912-78
<i>mylpfb</i>	<i>myosin light chain, phosphorylatable, fast skeletal muscle fiber</i>	1147.7	36526.4	31.8	http://zfin.org/ZDB-GENE-040912-115
<i>pvalb3</i>	<i>parvalbumin 3</i>	29.5	2165.4	73.5	http://zfin.org/ZDB-GENE-040426-945
<i>tnnt3a</i>	<i>troponin T3a, skeletal, fast</i>	42.2	3865.7	91.7	http://zfin.org/ZDB-GENE-000322-3
<i>tnni2a.4</i>	<i>troponin I, skeletal, fast 2a, tandem duplicate 4</i>	48.5	8637.8	178	http://zfin.org/ZDB-GENE-040625-119

Table S3.
Oligonucleotides used to generate in situ hybridization probes and as qPCR primers.

Gene Symbol	Gene Name	Sequence
<i>in situ</i> probe primers		
<i>tmg2a</i>	<i>transglutaminase 2, C polypeptide A</i>	5'-AGTGACCCCAACACCAGCTGAAGACG 5'-TAGAAGCGCTCCATTATTAGGTCGCC
<i>rtn2a</i>	<i>reticulon 2a</i>	5'-ATGTGGAGAAATGTGGAGCTCACTG 5'-GAAACAAGTCACTTTAAAACAAGTCAT
<i>rtn2b</i>	<i>reticulon 2b</i>	5'-GTTTCAGAGATCAGGGACACATAG 5'-TTTTACAGATGAAATGTTAAAACAG
<i>murca</i>	<i>muscle-related coiled-coil protein a</i>	5'-TCTCAAAGTGGCTCGTGACCACGG 5'-CATATTGAGTTTTTTTTAATACAT
<i>murcb</i>	<i>muscle-related coiled-coil protein b</i>	5'-GCTTAAGCAGGCCTGACAGCCAC 5'-GCGGCGCGATTACGCAAATGAAG
qPCR primers		
<i>actb1</i>	<i>actin, beta 1</i>	5'-CGAGCAGGAGATGGGAACC 5'-CAACGGAAACGCTCATTGC
<i>tie1</i>	<i>endothelium-specific receptor tyrosine kinase 1</i>	5'-AATTTAAATGCTCAAGGAGTTCGCC 5'-AGGCTGGATCTGTCTCTAGGACTCG
<i>cmlc2 (myl7)</i>	<i>myosin, light polypeptide 7, regulatory</i>	5'-GCCCATAACTTCACTGTCTTCC 5'-CTGGTCAACCTCTTCTGCTG
<i>pax7a</i>	<i>paired box gene 7a</i>	5'-GGCTACTTTACCAGGAACAG 5'-AGGTGTTGAGACTTCTAATGGG
<i>pax7b</i>	<i>paired box gene 7b</i>	5'-AGTTCCTCAGTAAGTTCTATCAG 5'-CCTTTGTCTCCAGAATGCC
<i>efnb3b</i>	<i>ephrin B3b</i>	5'-CTATAAACTGTACCTGGTTTCGTC 5'-GACCATATGGACTCTGTCCC
<i>myod1</i>	<i>myogenic differentiation 1</i>	5'-AAACTACCAATGCTGACCGT 5'-AAATCCATCATGCCATCAGAG
<i>myog</i>	<i>myogenin</i>	5'-TCAGTTCCTCAACCAGCAG 5'-ATCACTAGAGGACGACACCC
<i>myf5</i>	<i>myogenic factor 5</i>	5'-GAACTACTACAGCCTGCCGA 5'-CCGCAGGATCTACAGAGGAC
<i>myf6</i>	<i>myogenic factor 6</i>	5'-CTGACCCTTACACATACAACCTG 5'-ACACGGCTCCTTCTCTATGAC
<i>myl1</i>	<i>myosin, light chain 1, alkali; skeletal, fast</i>	5'-TCCGGTGTGAAGCTCGATTCACCC 5'-CATAGGCAGGAAACCCTCAAAGTCC