

RESEARCH ARTICLE

REVOLUTA and WRKY53 connect early and late leaf development in *Arabidopsis*

Yakun Xie¹, Kerstin Huhn¹, Ronny Brandt¹, Maren Potschin¹, Stefan Bieker¹, Daniel Straub^{1,2}, Jasmin Doll¹, Thomas Drechsler¹, Ulrike Zentgraf^{1,*} and Stephan Wenkel^{1,2,*}

ABSTRACT

As sessile organisms, plants have to continuously adjust growth and development to ever-changing environmental conditions. At the end of the growing season, annual plants induce leaf senescence to reallocate nutrients and energy-rich substances from the leaves to the maturing seeds. Thus, leaf senescence is a means with which to increase reproductive success and is therefore tightly coupled to the developmental age of the plant. However, senescence can also be induced in response to sub-optimal growth conditions as an exit strategy, which is accompanied by severely reduced yield. Here, we show that class III homeodomain leucine zipper (HD-ZIPIII) transcription factors, which are known to be involved in basic pattern formation, have an additional role in controlling the onset of leaf senescence in *Arabidopsis*. Several potential direct downstream genes of the HD-ZIPIII protein REVOLUTA (REV) have known roles in environment-controlled physiological processes. We report that REV acts as a redox-sensitive transcription factor, and directly and positively regulates the expression of *WRKY53*, a master regulator of age-induced leaf senescence. HD-ZIPIII proteins are required for the full induction of *WRKY53* in response to oxidative stress, and mutations in *HD-ZIPIII* genes strongly delay the onset of senescence. Thus, a crosstalk between early and late stages of leaf development appears to contribute to reproductive success.

KEY WORDS: REVOLUTA, HD-ZIPIII, WRKY53, Leaf senescence, Hydrogen peroxide signaling

INTRODUCTION

Senescence is the final stage of leaf development and involves the concerted reallocation of nutrients from the leaves to developing parts of the plant, especially fruits and seeds. Thus, leaf senescence has a major impact on yield quantity and quality, e.g. salvaged nitrogen (N) from wheat leaves accounts for up to 90% of the total grain N content (Kichey et al., 2007). In order to minimize loss of nutrients, plants induce leaf senescence in response to endogenous cues such as plant age and altered hormone homeostasis. However, external factors, such as the availability of water or light quality can also induce senescence, referred to as premature senescence (Ballaré, 1999). Although age-induced senescence tends to

maximize seed production, premature senescence describes an exit strategy that is induced in response to sub-optimal growth conditions and is often correlated with severely decreased yields.

The onset and progression of leaf senescence is accompanied by immense changes in the leaf transcriptome. It is estimated that about 20% of all genes are altered in expression upon induction of senescence, implying an important role for transcriptional regulators (Balazadeh et al., 2008; Breeze et al., 2011; Buchanan-Wollaston et al., 2005; Zentgraf et al., 2004). NAC and WRKY transcription factors are over-represented in the senescence transcriptome (Guo et al., 2004) and some members of these two transcription factor families have been shown to play central roles in regulating senescence (Balazadeh et al., 2010, 2011; Besseau et al., 2012; Breeze et al., 2011; Miao et al., 2004; Uauy et al., 2006; Ülker et al., 2007; Yang et al., 2011). WRKY proteins are plant-specific transcriptional regulators that contain a DNA-binding domain of ~60 amino acids. This domain contains a WRKYGQK motif at the N terminus and a zinc-finger structure at the C terminus, and is called the WRKY domain. Diverse processes, such as the response to pathogens or wounding but also leaf senescence, are controlled by WRKY transcription factors (Rushton et al., 2010). WRKY53, a key player in age-induced leaf senescence, regulates a complex network of downstream targets that promote vast physiological changes associated with the reallocation of nutrients and the induction of cell death (Lin and Wu, 2004; Miao et al., 2004). Owing to its important function, *WRKY53* expression, activity and protein stability are tightly controlled (Zentgraf et al., 2010). When leaf senescence is induced, the *WRKY53* locus is activated by histone modifications H3K4me2 and H3K4me3 (Ay et al., 2009; Brusslan et al., 2012), whereas DNA methylation remains low and unchanged (Zentgraf et al., 2010). Several promoter-binding proteins have already been characterized for *WRKY53* regulation, including WRKY53 itself, other WRKYs and the activation domain protein (AD protein), which has some similarity to HPT kinases and works as an activator of *WRKY53* expression (Miao et al., 2008; Potschin et al., 2014). In addition, a mitogen-activated protein kinase kinase kinase (MEKK1) was characterized to bind directly to the DNA of the *WRKY53* promoter. The binding region of MEKK1 appears to be involved in the switch from leaf age-dependent to plant age-dependent expression of *WRKY53* (Hinderhofer and Zentgraf, 2001; Miao and Zentgraf, 2007). MEKK1 can directly phosphorylate the WRKY53 protein, thereby increasing its DNA-binding activity (Miao and Zentgraf, 2007). As almost all WRKY factors contain WRKY factor-binding sites (W-boxes) in their proximal promoter regions, a complex regulatory WRKY network exists. Besides the transcriptional regulation, WRKY53 protein stability is strongly controlled by a HECT E3-ubiquitin ligase (Miao and Zentgraf, 2010). Moreover, gene expression changes are accompanied by hormonal changes. Although the plant hormones cytokinin and auxin act to delay senescence (Kim et al., 2011;

¹Center for Plant Molecular Biology, University of Tuebingen, Auf der Morgenstelle 32, 72076 Tuebingen, Germany. ²Copenhagen Plant Science Centre, University of Copenhagen, Thorvaldsensvej 40, Frederiksberg C 1871, Denmark.

*Authors for correspondence (ulrike.zentgraf@zmbp.uni-tuebingen.de; wenkel@plen.ku.dk)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Li et al., 2012), ethylene, abscisic acid (ABA), salicylic acid (SA) and jasmonic acid (JA) strongly promote leaf senescence (Li et al., 2012). Besides hormone homeostasis, elevated hydrogen peroxide levels also trigger senescence (Bieker et al., 2012; Smykowski et al., 2010).

Here, we identify REVOLUTA (REV), a transcription factor known to regulate polarity-associated growth processes in embryos, leaves, stems, vasculature and roots (Carlsbecker et al., 2010; McConnell et al., 2001; Smith and Long, 2010), as a direct regulator of *WRKY53* expression. During early leaf development, REV is involved in establishing the dorsoventral axis of leaves by specifying the domain that will later develop into the upper side of the leaf (Byrne, 2006). REV, also known as INTERFASCICULAR FIBERLESS (IFL), has been shown to play multiple roles in meristem organization, leaf polarity set-up and vascular development (Otsuga et al., 2001; Talbert et al., 1995; Zhong and Ye, 1999). Using a ChIP-Seq approach, we identified REV-binding sites in the *WRKY53* promoter and by qRT-PCR demonstrate that REV promotes *WRKY53* expression. Conversely, plants that carry loss-of-function mutations in *REV* and other *HD-ZIP* genes show lower levels of *WRKY53* expression, confirming that *HD-ZIP* genes are also required for *WRKY53* expression. By performing a detailed expression analysis using both *REV* and *WRKY53* GUS-reporter lines, we reveal that both genes have partially overlapping patterns of expression. In wild-type plants, *WRKY53* expression is strongly induced in response to hydrogen peroxide. However, in *rev* mutant plants and in transgenic plants with reduced *HD-ZIP* activity, this response is significantly dampened. Furthermore, the ability of REV to bind to the *WRKY53* promoter is also dependent on the redox environment and, under oxidative conditions, less binding is observed. In line with the lower *WRKY53* expression levels, *rev* mutant plants are considerably delayed in age-induced leaf senescence, suggesting a role for *HD-ZIP* genes in this physiological process. Taken together, we conclude that REV is a positive regulator of *WRKY53* expression, which influences the onset of leaf senescence in response to changes in the cellular redox state. Obviously, early and late leaf development are tightly linked by transcriptional networks between *HD-ZIP* and *WRKY* factors, in which disturbed early development is coupled to extended life span of leaves and delayed senescence.

RESULTS

REVOLUTA is a positive regulator of *WRKY53* expression, a major factor controlling age-induced leaf senescence

REVOLUTA is a member of the class III homeodomain leucine zipper (*HD-ZIP*) transcription factor family that regulates various polarity-associated growth processes during development (Carlsbecker et al., 2010; McConnell et al., 2001; Smith and Long, 2010), but plays an additional role in shade-induced growth promotion (Bou-Torrent et al., 2012; Brandt et al., 2012). REVOLUTA expression is controlled by the microRNAs *miR165* and *miR166* at the post-transcriptional level (Rhoades et al., 2002), and by the association with small leucine-zipper-type microProteins at the post-translational level (Kim et al., 2008; Staudt and Wenkel, 2011; Wenkel et al., 2007). Using a genome-wide chromatin-immunoprecipitation sequencing approach (ChIP-Seq), we recently identified binding regions for REV across the *Arabidopsis* genome (Brandt et al., 2012). This analysis revealed binding of REV to the promoter of the *WRKY53* transcription factor (Fig. 1A). Transient promoter-GUS experiments in *Arabidopsis* protoplasts revealed an induction of *WRKY53* expression after co-transformation of *35S::REVd*, a dominant microRNA-resistant version of REV (Fig. 1B). Quantitative ChIP-PCRs confirmed the

binding of REV to the ChIP-Seq identified binding motifs (Fig. 1C). For better control of REV activity, we constructed transgenic plants expressing REVd fused to the rat glucocorticoid receptor carrying an N terminal FLAG epitope. In response to dexamethasone (DEX) induction, the chimeric FLAG-GR-REVd fusion protein translocates to the nucleus, where it can associate with DNA and alter the expression of target genes. In response to DEX induction, REV can significantly upregulate *WRKY53* expression (Fig. 1D), while seedlings carrying mutations in *REV* and plants with globally reduced *HD-ZIP* activity show reduced levels of *WRKY53* mRNA (Fig. 1E), thus supporting a new role for REV as a direct and positive regulator of *WRKY53*.

REVOLUTA and *WRKY53* have overlapping patterns of expression

REVOLUTA, as well as the other class III *HD-ZIP* transcription factors of *Arabidopsis*, have a distinct expression pattern, confining their expression to the adaxial domain of developing leaves, the xylem part of the vasculature, the pro-vasculature and the shoot apical meristem. Both *WRKY53* and *REV* are expressed in young seedlings (Fig. 2A,B). Even though REV function was initially described for polarity-associated growth processes during early leaf development, REV is still expressed at later stages of development (supplementary material Fig. S1) and an additional function in shade avoidance has recently been assigned to REV (Brandt et al., 2012). In comparison with the vascular expression pattern of *REV*, *WRKY53* shows a broader less-specific pattern of expression and is most highly expressed in old leaves (Miao and Zentgraf, 2007). In genetic backgrounds with reduced *REV* mRNA [*rev-5* (Fig. 2C), *35S::miR165a* (Fig. 2D)] or with reduced REV protein activity (*35S::ZPR3*; Fig. 2E), the spatial expression of *WRKY53* is more restricted to hydrotodes and overall expression levels appear to be much lower in leaf tissue. In older seedlings, expression of both genes is found in vascular strands (Fig. 2F–M). Surprisingly, high co-expression is observed in the root vasculature at all investigated stages of development. It is not known whether *WRKY53* has an additional function in root development but it might be important to note that the expression in the root vascular appears to be independent of *HD-ZIP* function (Fig. 2B–E).

Using publicly available microarray data (<http://bar.utoronto.ca>), we also analyzed at which stages of development and in response to which treatments *REV* and *WRKY53* are co-expressed (supplementary material Fig. S2). We find evidence for co-expression during early developmental stages but not during the later stages of leaf development. This discrepancy suggests that *REV* mRNA is not upregulated at late stages of leaf development but residual protein could respond to a cellular signal and induce the expression of REV-regulated senescence targets. However, our GUS expression analyses using *REV::GUS* plants indicate that REV is still expressed to certain extends in older leaves (supplementary material Fig. S1).

In order to identify other direct REV targets that show an expression pattern resembling *WRKY53*, we surveyed recently published timecourse microarray datasets (Reinhart et al., 2013) that revealed 119 genes to be upregulated in response to REVOLUTA induction. Our ChIP-Seq datasets resulted in the identification of 286 high confidence REV-binding sites (corresponding to 552 potentially regulated genes) across the entire *Arabidopsis* genome (Brandt et al., 2012). By comparing both datasets, we could identify 18 of the 119 REV-regulated genes (15% of the REV upregulated set) to have REV-binding sites in their respective promoters (Table 1). *WRKY53* is among these 18 genes and we investigated whether other senescence-related genes could be identified in this dataset. A




Fig. 1. REVOLUTA binds to the *WRKY53* promoter and is a direct and positive regulator of *WRKY53* expression. (A) ChIP-Seq results for the binding of REV to the *WRKY53* promoter. Two binding sites (BS) were identified, located -1.3 kb and -2.1 kb upstream of the transcriptional start site. Traces in gray are sequence reads derived from sequencing ChIP DNA from Col-0 wild-type plants; red plots ChIP DNA from dexamethasone-induced *35S::FLAG-GR-REVd* transgenic plants. (B) Transient expression assay in *Arabidopsis* protoplasts. A plasmid with a 2.8 kb *WRKY53* promoter fragment fused to the *GUS* gene was transformed along with a second plasmid containing a *CaMV35S*-promoter (control) or the *CaMV35S*-promoter driving expression of *REVd*. *GUS* activity was determined ~ 15 h after transformation. Data are mean \pm s.d. * $P < 0.05$. (C) Chromatin-immunoprecipitation qPCR experiments with two biological replicates for *35S::FLAG-GR-REVd* without DEX (gray bars) and *35S::FLAG-GR-REVd* with DEX (red bars) plants testing four positions in the *WRKY53* promoter. Y-axis shows the fold enrichment normalized to the non-induced IPs. Gene map above the chart shows the localization of the REV-binding site identified by ChIP-Seq and the regions that were tested. Distance between two marks along the chromosomes represents 1.0 kb. (D) Real-time quantitative PCR experiments showing expression changes of *WRKY53* in Col-0 (black) and *35S::GR-REVd* (orange) in response to 60 min DEX induction in the presence of the protein biosynthesis inhibitor cycloheximide (CHX). Data are mean \pm s.d. * $P < 0.05$. (E) Expression of *WRKY53* was analyzed in different *rev* mutant plants (*rev-5*, *rev-6*, *phb phv rev/+* and *phb phv rev*) and in plants with reduced activity of HD-ZIPIII proteins (*35S::ZPR3*). The bars indicate expression levels relative to wild type, including standard errors of the mean of three individual biological experiments. * $P < 0.05$.

genome-wide survey with a high temporal resolution classified thousands of genes as differentially expressed senescence genes (DESGs) (Breeze et al., 2011). Interestingly, REV was also classified as a DESG, showing a dip of expression at the onset of leaf senescence. Furthermore, nine out of the 18 potential direct REV targets (Table 1) were also classified as DESGs, implying that REV might have an additional function in late developmental stages.

***WRKY53* expression is modulated in response to oxidative stress in a REVOLUTA-dependent manner**

WRKY53 expression is strongly upregulated in response to hydrogen peroxide as part of the age-induced senescence-promotion pathway (Miao et al., 2004). Because REV is a novel upstream regulator of *WRKY53* expression and possesses a domain that is suggestive of sensing changes in the redox state of the cell, we investigated whether REV is required for the induction of *WRKY53* expression in response to oxidative stress. Therefore, we grew Col-0 wild-type plants and mutant plants with reduced HD-ZIPIII activity (*rev5*, *35S::miR165a* and *35S::ZPR3*) on soil for 3 weeks in long-day conditions. In order

to elicit oxidative stress, plants were sprayed with hydrogen peroxide solutions of different concentrations (0.01%, 0.1% and 1%) and plant material was harvested before and after spraying. Subsequent RNA isolation, cDNA synthesis and quantitative PCR analysis revealed a strong induction of *WRKY53* in response to H_2O_2 application in Col-0 wild-type plants. These changes of *WRKY53* mRNA levels were significantly dampened in *rev* mutant plants (*rev-5*) and *35S::miR165a*, and in plants with reduced HD-ZIPIII activity (*35S::ZPR3*), indicating that REV activity is required for high-level *WRKY53* induction in response to oxidative stress signaling (Fig. 3). To assess which externally applied hydrogen peroxide concentration is able to elicit redox changes that would occur under natural conditions, we measured intracellular hydrogen peroxide levels after applying heat stress and compared them with the intracellular levels reached after external application of H_2O_2 by spraying. To be sure that only intracellular H_2O_2 is measured, we used non-fluorescent H_2DCFDA (2',7'-dichlorodihydrofluorescein diacetate), which is converted to the highly fluorescent 2',7'-dichlorofluorescein upon cleavage of the acetate groups by intracellular esterases and




Fig. 2. Expression analysis of *REV* and *WRKY53*. (A–I) Spatial patterns of expression of *REV* (A,F,G) and *WRKY53* (B–E,H,I) in 8-day-old *Arabidopsis* seedlings. GUS staining of *REV::GUS* (A), *WRKY53::GUS* (B) in the Col-0 ecotype and *WRKY53::GUS, rev5* (C), *WRKY53::GUS, 35S::miR165* (D), *WRKY53::GUS, 35S::ZPR3* (E) seedlings. Scale bars: 1 mm. (F–I) Hypocotyls (F,H) and roots (G,I). (J,K) The pattern of GFP accumulation in the hypocotyl (J) and root (K) vascular tissue of 8-day-old plants carrying the *REV::REV-GFP* transgene. Scale bars: 50 μ m. (L,M) Cross-sections of roots of 10-day-old seedlings reveal *REV* (L) and *WRKY53* (M) expression in the vascular cylinder.

subsequent oxidation. The increase in intracellular H₂O₂ was similar 1 h after heat treatment and 1 h after spraying 0.1% H₂O₂ but dropped more rapidly in the H₂O₂-treated samples. This indicates that external application of 0.1% H₂O₂ leads to intracellular changes in the range of an oxidative burst in stress response (supplementary material Fig. S3).

REVOLUTA is a redox-sensitive transcription factor

REV is a positive regulator of *WRKY53* expression and is required for high level of *WRKY53* induction in response to oxidative stress.

This could be either due to an upregulation of *REV* mRNA in response to oxidative stress or to a response of the REV protein to altered redox conditions. To test whether *REV* mRNA is upregulated in response to hydrogen peroxide treatment, we treated Col-0 wild-type plants with H₂O₂ and performed quantitative RT-PCRs. We detected no induction of *REV* mRNA but a slight decrease in response to high levels of hydrogen peroxide (supplementary material Fig. S4), excluding the idea that *REV* is transcriptionally upregulated in response to oxidative stress.

It has been shown that proteins of the class II homeodomain leucine-zipper (HD-ZIPII) family from sunflower interact with DNA in a redox-sensitive manner (Tron et al., 2002). To test whether REV shows also redox-dependent DNA binding, we performed redox-sensitive DPI-ELISA experiments. Therefore, crude lysate of *E. coli* cells expressing HIS-tagged REV protein were prepared and incubated with streptavidin plates pre-loaded with biotinylated oligonucleotides containing the REV-binding site 1 of the *WRKY53* promoter (W53-BS1). ELISA plates were then washed and subsequently incubated with HRP-tagged anti-HIS antibodies. Enhanced signal was detected in the control binding reaction (HIS-REV lysate versus a lysate from BL21 cells expressing the empty vector control), indicating that HIS-REV binds to the W53-BS1 element (Fig. 4A). As observed for the sunflower HD-ZIPII proteins (Tron et al., 2002), REV also showed enhanced binding in response to reducing conditions (10 mM DTT), whereas in response to oxidative conditions (10 mM H₂O₂) DNA-binding was reduced (Fig. 4A). This negative effect is reversible as the subsequent addition of 10 mM DTT was able to restore REV DNA binding.

We examined the possibility of whether the C-terminal PAS-domain of REV might act as a redox sensor domain. Redox-DPI-ELISA experiments with HIS-REV lacking the PAS-domain (HIS-REVΔPAS) showed the same redox-sensitive behavior as observed for HIS-REV (Fig. 4B). However, without the PAS-domain, REV-DNA binding was strongly enhanced, supporting the idea that the PAS-domain regulates REV activity via a steric masking mechanism, as proposed by Magnani and Barton (2011). It is conceivable that the observed redox effects in the ELISA system

Table 1. Identification of potentially direct REV target genes by comparing ChIP-Seq and microarray experiments with an inducible version of REV

AGI	Name	Microarray	ChIP-Seq				
		Fold change	q_rank	Enrichment	Distance	Location	DESG*
AT2G41940	ZFP8	2.0	469	7.5	1691	Down	Yes
AT5G47370	HAT2	3.1	253	8.0	1548	Up	No
AT2G39705	DVL11/RTFL8	2.8	1626	7.6	2509	Down	No
AT5G06710	HAT14	2.7	272	9.3	5364	Up	No
AT5G47180	Plant VAMP protein	1.7	35	15.2	168	Up	Yes
AT5G19590	DUF538 protein	1.3	465	9.4	2810	Up	Yes
AT4G18700	CIPK12	3.0	169	13.2	282	Down	No
AT4G27730	OPT6	2.7	30	16.1	1305	Up	No
AT4G03510	RMA1	7.0	33	14.9	1989	Up	No
AT1G17970	RING/U-Box protein	5.1	1173	9.7	8	Up	Yes
AT5G14730	DUF1645	5.8	726	6.6	2299	Up	No
AT2G45450	ZPR1	13.1	400	9.2		5'UTR	No
AT5G05690	CPD	1.9	202	8.0	4847	Up	No
AT1G74940	DUF581	2.2	106	17.5	81	Up	Yes
AT3G60390	HAT3	2.9	115	8.7	5597	Up	Yes
AT4G23810	WRKY53	3.9	450	8.7	2132	Up	Yes [†]
AT5G16030	Unknown protein	2.8	789	9.6	2193	Up	Yes
AT1G49200	RING/U-Box protein	14.4	18	14.2	187	Up	No
AT2G02080	IDD4	0.5	528	8.2	8334	Down	No
AT3G13810	IDD11	0.4	1643	6.7	1422	Down	Yes

*Differentially expressed genes during senescence (Breeze et al., 2011).
[†]Senescence-associated gene not included in the Breeze et al. (2011) analysis.




Fig. 3 HD-ZIPIII activity is required for H₂O₂-mediated upregulation of *WRKY53*. Real-time qPCR experiment showing *WRKY53* induction in response to hydrogen peroxide treatment in wild-type and *rev* mutant plants. Three-week-old plants were treated with different concentrations of H₂O₂ [0% (mock; gray bars), 0.01% (yellow bars), 0.1% (orange bars) and 1% (red bars)] for 40 min. Data are representative relative expression changes (fold change) of the mean of four technical replicates \pm s.d. Similar expression changes have been observed in at least two independent biological experiments.

are due to an influence of *E. coli* proteins on the activity of REV. To exclude such effects, we purified GST-REV protein from *E. coli* and performed *in vitro* gel retardation assays in the presence of reducing agents (DTT) and oxidizing agents (H₂O₂) (Fig. 4C). These gel-shift experiments largely confirm the results obtained by redox-DPI-ELISA and confirm that REV activity can be modulated by the intracellular redox state.

To validate redox-sensitive DNA binding *in planta*, we treated 35S::FLAG-GR-REVd transgenic plants with either a mock

substrate (0.1% ethanol), dexamethasone (DEX) or DEX+0.1% H₂O₂. In 12-day-old seedlings, we detected REV binding to binding site 2 (fragment II) and no binding was observed to binding site 1 (fragment III). When treated with hydrogen peroxide prior DEX induction, binding to binding site 2 was significantly affected (Fig. 4D), indicating that REV DNA binding is indeed redox sensitive. The same experiment with 7-week-old plants revealed that, at later developmental stages, both binding sites are occupied by REV and the binding seems to be enhanced but exhibits the same redox sensitivity (Fig. 4E). Taken together, we demonstrate that REV shows a stage-specific redox-dependent DNA-binding behavior and that oxidizing conditions decrease the ability to bind DNA *in vitro* and *in vivo*.

Mutations in the *REVOLUTA* gene or the overall reduction of HD-ZIPIII activity delay the onset of leaf senescence

One function of the WRKY53 protein is the regulation of the onset of senescence, documented by the phenotype of the *wrky53* mutant showing delayed senescence. As REV is an activator of *WRKY53* expression, we expected *rev* mutant plants to also display a delayed senescence phenotype. Our analysis revealed that plants carrying mutations in REV or plants with greatly reduced HD-ZIPIII activity are significantly delayed in senescence, while overall development is not retarded, which clearly confirms a role of HD-ZIPIII proteins in this process (Fig. 5; supplementary material Figs S5, S6). Furthermore, the phenotype of *rev5* was even stronger than that of *wrky53*, indicating that *WRKY53* might not be the only senescence-associated gene regulated by REV.

Overexpression of the small leucine-zipper-type microProtein ZPR3, which largely reduces the activity of HD-ZIPIII, led to a further enhancement of the senescence phenotype, which was ameliorated in the *wrky53* mutant background (supplementary material Fig. S3). This confirms that the senescence phenotype is




Fig. 4. Redox-mediated regulation of REVOLUTA-DNA-binding capability and influence of the PAS domain. (A,B) Redox-DPI-ELISAs. The DNA-protein interaction assays were performed by using 5' biotinylated complementary annealed oligonucleotides coupled to a streptavidin-coated ELISA plate. Crude *E. coli* extracts (25 μ g) expressing recombinant REV or REVΔPAS were pre-incubated with different concentrations of DTT and H₂O₂ to examine a redox state-dependent binding of REV. In order to test the reversibility of the redox effect, high concentrations of H₂O₂ were added first and then oxidizing conditions were reversed by addition of DTT. After binding, biotinylated DNA-protein complexes were detected using anti His-HRP conjugated antibodies. Results for REV binding site 1 of the *WRKY53* promoter are shown. *E. coli* BL21 cells transformed with the empty vector were used as background control. (C) Non-radioactive electrophoretic mobility shift assays. Purified GST-REV protein was incubated with a biotinylated oligonucleotide containing the HB9-binding motif (Wenkel et al., 2007) in the presence of different redox conditions. After gel electrophoresis and subsequent blotting, the biotinylated DNA probe was detected with a HRP-streptavidin substrate. (D,E) Chromatin-immunoprecipitation qPCR assays of 35S::FLAG-GR-REVd plants. Twelve-day-old seedlings (D) and 7-week-old transgenic plants (E) were treated with mock substrate (0.1% ethanol), DEX or 0.1% H₂O₂ and DEX. H₂O₂ was given 15 min prior to 45 min of DEX induction. Fold enrichment for the same primer sets as in Fig. 1 is shown.

mediated by deregulation of *WRKY53* expression through HD-ZIP IIIs but also suggests that additional HD-ZIP IIIs are involved, as the senescence phenotype of *35S::ZPR3* plants is much stronger compared with *rev5* mutants (Fig. 5; supplementary material Figs S5,S6). Consistent with the phenotype, two typical senescence-related physiological parameters, the decrease in chlorophyll content and the increase in lipid peroxidation, were also delayed in *wrky53*, *rev5* and *rev5 wrky53* mutants (Fig. 6A,B). Furthermore, the mRNA expression levels of *SENESCENCE ASSOCIATED GENE 12* (*SAG12*) and *SAG13*, which are commonly used as senescence marker genes, were significantly reduced at the late developmental stages in *wrky53*, *rev5* and *rev5 wrky53* mutants compared with Col-0 wild-type plants (Fig. 6C,D). Taken together, these results confirm that REV acts upstream of *WRKY53* in the control of age-induced senescence.

Depletion of *REV* delays the onset of leaf senescence more efficiently than depletion of *WRKY53*. To further investigate the possibility that REV acts upstream of several senescence-associated genes, we focused our attention on the potential direct REV targets classified as DESGs (Table 1). Here, we decided to investigate three groups of genes: (1) genes whose expression decreases with age (*HAT3* and *AT1G49200*); (2) genes whose expression increases with age (*AT1G74940* and *IDD11*); and (3) genes whose expression decreases with age but rises during senescence (*AT5G47180* and *ZFP8*). In the first group of genes, we found that expression in *wrky53*, *rev5* and *rev5 wrky53* mutants is maintained at a higher level towards the onset of senescence (weeks 5 and 6), whereas expression levels are dropping rapidly in wild-type plants (Fig. 7A,B). For the second group of genes whose expression increases with age in wild-type plants, we detected elevated levels in *wrky53*, *rev5* and *rev5*




Fig. 5. Genetic interaction of REV with WRKY53. (A) Rosette leaves of 6- and 7-week-old representative plants were sorted according to their age; whole rosettes were also photographed upside down to visualize the older leaves. (B) For a quantitative evaluation of leaf senescence, plants were harvested in a weekly rhythm and leaves of at least six plants were categorized into four groups according to their leaf color: (1) 'green'; (2) leaves starting to become yellow from the tip as 'yellow-green'; (3) completely yellow leaves as 'yellow'; and (4) dry and/or brown leaves as 'brown/dry'. The percentages of each group with respect to total leaf numbers are presented. Error bars indicate s.d. Student's *t*-test was performed comparing leaf counts of *wrky53*, *rev5* and *rev5wrky53* with Col-0 numbers, **P*<0.05, ***P*<0.005, ****P*<0.0005. *n*=7-15.




Fig. 6. Molecular senescence parameters. (A) Chlorophyll contents of number 5 leaves from *Arabidopsis* Col-0, *wrky53*, *rev5* and *rev5wrky53* plants. Left axis indicates atLeaf+ values. Plant age is indicated in days after seeding (DAS). (B) Lipid peroxidation in Col-0, *wrky53*, *rev5* and *rev5wrky53* plants. Values represent mean of at least three biological replicates \pm s.d. Comparison of means and the determination of statistical differences was carried out using Student's *t*-test (* $P < 0.05$, ** $P < 0.005$ and *** $P < 0.0005$). (C,D) qRT-PCR expression analysis of the senescence marker genes *SAG12* and *SAG13*. All values were normalized to *GAPDH* expression. Error bars indicate s.d. of four technical replicates.

wrky53 mutants at early developmental stages (weeks 4 and 5) and decreased levels at the late stages (Fig. 7C,D). Expression of the third group of genes is also altered at various time points in *wrky53*, *rev5* and *rev5 wrky53* mutants compared with Col-0, but in all lines the transcriptional increase during senescence is diminished (Fig. 7E,F), further corroborating the idea that loss of *REV* function profoundly alters the senescence transcriptome, which might be causative for the strong senescence phenotype of *rev* mutant plants.

Loss-of-function *wrky53* mutant plants do not show obvious developmental defects during early leaf development, indicating that *WRKY53* is not required for *REV* function at these stages of development. However, the severe *35S::ZRP3*-induced leaf phenotype is ameliorated in the *wrky53* mutant background, suggesting that the action of other HD-ZIP IIIs involves *WRKY53* also at early stages (supplementary material Fig. S7). Nonetheless, *WRKY53* protein levels are most likely very low during these early




Fig. 7. qRT-PCR of other *REV* target genes differentially expressed during senescence. Quantitative real-time PCR profiling of putative *REV* target genes at late developmental stages in wild-type and mutant plants (4-, 5-, 6-, 7- and 8-week-old plants). (A–F) Expression changes over time of *HAT3*, *AT1G49200*, *AT1G74940*, *IDD11*, *AT5G47180* and *ZFP8*. The Y-axis represents the relative expression level normalized to *GAPDH*. Error bars indicate s.d. of four technical replicates.

stages of development due to the degradation of WRKY53 by the HECT domain ubiquitin ligase UPL5, which is highly expressed in young leaves (Miao and Zentgraf, 2010). Taken together, we discovered that HD-ZIPIII factors interact with *WRKY53* genetically to promote age-induced leaf senescence, and disruption of early leaf development correlates with delayed senescence and extended life span of leaves.

Functional analyses of root-specific co-expression patterns of *REV* and *WRKY53*

It is unknown which tissues are involved in the perception of senescence signals and conversion of these into the senescence triggers. We find co-expression of *REV* and *WRKY53* during the early stages of leaf development. Later in development, co-expression was very obvious in the vasculature of the leaves and in the root vascular cylinder (Fig. 2L,M), although both *REV* and *WRKY53* are expressed throughout development (supplementary material Fig. S1). This is in agreement with the finding that *REV* is involved in the induction of *WRKY53* expression by hydrogen peroxide and that very high levels of hydrogen peroxide were observed in vascular tissue indicated by DAB staining of leaf sections (Zimmermann et al., 2006). Moreover, it remains tempting to speculate that the root might also act as a senescence sensor; however, whether roots play a role during onset and progression of senescence has not yet been determined and whether and to what extent hydrogen peroxide is transported through the vasculature over long distances is also not known so far. Auto-propagating waves of reactive oxygen species (ROS) that rapidly spread from the initial site of exposure to abiotic stress to the entire plant are involved in conferring systemic acquired acclimation, also allowing a much faster transcriptome and metabolome reprogramming of systemic tissues in response to abiotic stress (Mittler et al., 2011; Suzuki et al., 2013).

To further investigate the spatial aspects of *REV* and *WRKY53* expression, we decided to perform grafting experiments with Col-0 wild-type, *rev5* and *wrky53* mutant plants. When the aerial parts of Col-0 were grafted onto either *wrky53* or *rev5* rootstocks, no significant delays in the onset of senescence were observed. However, the converse grafting of the aerial parts of either *wrky53* or *rev5* to Col-0 rootstocks significantly delayed the onset of senescence where the latter again showed a much stronger effect (Fig. 8A,B). The grafting experiments revealed that the root seems not to be involved in the *REV*/*WRKY53*-mediated senescence pathway and that depletion of *REV* and *WRKY53* in only aerial tissue strongly affects senescence.

DISCUSSION

Plants induce leaf senescence to provide carbon, nitrogen and mineral resources to the developing fruits or seeds. Senescence is induced in response to plant age but environmental signals such as light, the availability of water and temperature strongly influence this process. A high-resolution temporal transcript profiling of senescing *Arabidopsis* leaves gives insight into the temporal order of gene activation and repression (Breeze et al., 2011). Approximately 6500 genes are up- or downregulated during the course of leaf senescence, implying an important role for transcription factors in this process. Transcription factors themselves are transcriptionally upregulated in senescing leaves the largest groups being NAC, WRKY, C2H2-type zinc-finger, AP2/EREBP and MYB proteins (Guo and Gan, 2005). Here, we now show that HD-ZIPIII factors, which are known to be involved in basic patterning processes, have an additional role in the latest




Fig. 8. Grafting experiments and senescence phenotype. (A) Nine combinations of grafted plants were generated between the wild-type and mutant plants (*rev6* and *wrky53*), including three self-grafted controls, e.g. wild type to wild type (Col-0/Col-0; scion/root). Error bars indicate s.d. ($n=4-6$ independent grafted plants with the exception of Col-0/*wrky53*, where we achieved only two successful grafts). The quantitative evaluation of leaf senescence of the non-grafted plants is shown in Fig. 5. Asterisks represent significant differences from the Col-0/Col-0 graft, as determined using Student's *t*-test (* $P<0.05$, ** $P<0.005$, *** $P<0.0005$). (B) The leaf-senescence phenotypes of grafts. Photographs were taken 7 weeks after grafting.

step of leaf development, the regulation of senescence. *REV* is a direct and positive regulator of *WRKY53* expression and mutations in *REV* and other HD-ZIPIII genes delay the onset of leaf senescence. Interestingly, the delay of the onset of leaf senescence in plants lacking *REV* is stronger compared with plants lacking only *WRKY53*, implying that *REV* acts also upstream of other senescence-associated genes. In plant lines with even more reduced HD-ZIPIII activity, achieved by overexpression of

miRNA165a (35S::miR165a), rosette leaves were so strongly downward curled that it was impossible to determine the onset of senescence. The loss of several *HD-ZIPIII* genes, as in the case of the *phb phv rev* triple mutant, causes severe developmental defects, including consumption of the apical stem cells (Emery et al., 2003; Prigge et al., 2005). The severity of these developmental defects largely precludes a thorough analysis of the general role of *HD-ZIPIII* proteins at later stages of development. Nevertheless, our findings clearly suggest that the role of *HD-ZIPIII* proteins in promoting senescence is more complex and involves regulation of several senescence-associated target genes. In the *rev5/wrky53* double mutant, leaf yellowing and chlorophyll loss were less severe at later stages than in the *rev* single mutant, whereas senescence-associated gene expression was more severely affected for some senescence-related genes. This clearly points towards a complex network that is altered in different aspects if one or more components are depleted from the system. It was already shown that *WRKY53* acts as an upstream regulator, downstream target and protein-interaction partner of *WRKY18*, which is a negative regulator of leaf senescence, illustrating the complexity of the network and possibly explaining the partially intermediate phenotype of the double mutant (Potschin et al., 2014).

The mechanism by which *REV* promotes senescence appears to involve transcriptional regulation of direct target genes. Here, we have identified nine genes that are potential direct *REV* targets that are also differentially expressed during senescence. One of these target genes is *HAT3*, which has been shown to play an important role downstream of *REV* in the process of setting up polarity in the young leaf primordium (Bou-Torrent et al., 2012; Brandt et al., 2012; Turchi et al., 2013). In young seedlings, *HAT3* expression depends partly on the presence of *REV*, which is supported by lower levels of *HAT3* mRNA in *rev* mutant seedlings (Brandt et al., 2012). During senescence, *HAT3* mRNA levels decrease with plant age (Fig. 7A). In *rev* mutant seedlings, however, *HAT3* mRNA is more abundant compared with wild type (Fig. 7A). Moreover, the expression levels of several other senescence-related target *REV* genes changed in a complex way (Fig. 7B–F). These findings suggest that the transcriptome of *rev* mutant plants is profoundly altered, resulting in stage-dependent mis-expression of many differentially expressed senescence-associated genes.

It still remains unclear to which endogenous or exogenous signals *HD-ZIPIII*s respond in order to promote senescence. The finding that *WRKY53* expression is strongly upregulated in response to hydrogen peroxide treatment and that this induction is dampened in *hd-zipIII* mutant plants implies that *HD-ZIPIII*s might be involved in signal transduction processes in response to changes in the intracellular redox state. Many senescence-associated genes, especially transcription factors of the *WRKY* and the *NAC* family, transcriptionally respond to elevated levels of hydrogen peroxide but the mechanism by which the hydrogen peroxide signal is perceived and transmitted is still unclear. Remarkably, the subcellular compartment of hydrogen peroxide production appears to play a role in senescence signaling in which the cytoplasmic H_2O_2 is more effective in senescence induction than peroxisomal or mitochondrial H_2O_2 (Bieker et al., 2012; Zentgraf et al., 2012). Thus, sensors and mediators of hydrogen peroxide-induced senescence are most likely cytoplasmic and/or nuclear proteins or molecules. During bolting, intracellular hydrogen peroxide levels increase in leaf tissue. This increase is thought to be mediated by a complex regulation of the hydrogen peroxide scavenging enzymes and promotes the onset of senescence (Bieker et al., 2012; Smykowski et al., 2010).

Analysis of the redox sensitivity of the *REV* protein revealed a reduced DNA-binding ability of *REV* in response to oxidative conditions, which appears to be a direct effect on the *REV* protein and does not involve accessory proteins. These results contradict the finding that upregulation of *WRKY53* partially requires *HD-ZIPIII*s and indicate a more complex regulatory mechanism. Most likely, DNA-binding of *REV* is affected by redox changes and also the transactivation activity or protein-protein interfaces, which will be further dissected in the future. However, two of the direct *REV* target genes encode *EAR*-domain proteins that are part of transcriptional repressor complexes (Causier et al., 2012). Among these transcriptional repressors are *HAT3* and *ZFP8*, the mRNA levels of which are altered in the senescence process. Therefore, it seems plausible to conclude that *REV* is a redox-sensitive transcription factor, which among other targets, regulates genes encoding transcriptional repressors. Decreasing *REV* DNA-binding activity will result in lower expression levels of these transcriptional repressors, alleviating the repressive activity on their targets. Thus, modulation of *REV* activity in response to alterations of the intracellular redox state will profoundly affect the *REV*-regulated transcriptome. It is tempting to speculate that also within the shoot apical meristem, domains with different cellular redox states might exist that could serve as positional signals affecting *HD-ZIPIII* activity.

Developmental age is a major determinant for the induction of leaf senescence in an optimal growth environment. However, when plants are exposed to situations that strongly permit normal growth, senescence is accelerated in order to bypass these adverse conditions and produce seeds that can withstand these adverse conditions. We have tried to depict the complex interplay between *REV* and *WRKY* during early and late development in a model (Fig. 9) in which the regulatory cues of *REV* involving miRNA-dependent regulation through *miR165*, *miR166* and the *LITTLE ZIPPER* microProteins *ZRP1-4* is connected to the MAP kinase-triggered *WRKY* transcriptional network. Several intersections can be detected between the formerly independently described players in early and late leaf development in which hydrogen peroxide might play a central role.

Shade causes profound developmental changes in shade-sensitive plants aimed at outgrowing competitor plants. We have previously shown that the leaf regulatory module consisting of *HD-ZIPIII* and *KANADI* transcription factors is involved in modulating




Fig. 9. Model *HD-ZIPIII*/senescence. A model summarizing our findings and showing the relationship between early leaf development processes and senescence. Both *REV* and *WRKY53* intersect to regulate the late stages of leaf development.

growth in response to shade (Brandt et al., 2012). Consistent with this, shade can also trigger leaf senescence (Brouwer et al., 2012), suggesting that leaf patterning, shade avoidance and leaf senescence are interconnected by differential activity of HD-ZIPIII proteins, thus linking early and late leaf development, and adjusting plant growth and development to changing external conditions.

Perspectives

It was recently shown that embryonic growth and patterning of mammals largely depends on cellular senescence as a developmental mechanism to shape organ growth (Muñoz-Espín et al., 2013; Storer et al., 2013). This mechanism partly relies on macrophages, which are mobile cells that invade the tissue to remove senescent cells. In this context, senescent cells also produce secreted compounds that can act as positional signals triggering pattern formation and proliferation in adjacent tissue (Storer et al., 2013). The immune system of plants is substantially different from animals and does not involve macrophage-mediated cell clearing. However, it is conceivable that local cellular senescence could provide positional information to direct growth responses. Our finding that HD-ZIPIII, which are known basic patterning factors, can influence senescence processes, suggest not only that early and late leaf development are coupled and processes that influence patterning in the early organ control the concerted degradation of tissue during the late phase of development, but also that physiological processes related to senescence, such as nutrient mobilization or lipid peroxidation, might be part of early leaf patterning processes. Furthermore, the puzzling reduction of DNA-binding activity under oxidizing conditions that contradicts the finding that upregulation of *WRKY53* expression by hydrogen peroxide partially requires *REVOLUTA* prompts us to decipher the redox-dependent changes in the *REVOLUTA* protein outside the DNA-binding domain in more detail. This, however, will be the subject of further investigations.

MATERIALS AND METHODS

Plant material and growth conditions

The following *rev/hd-zipIII* mutant lines were used in this study: *rev-5* (A260V) and *rev-6* (R346STOP), two strong ethyl-methylsulfonate (EMS) alleles (Otsuga et al., 2001), *phb phv rev* triple mutants introgressed in Col-0 (Prigge et al., 2005), *35S::ZPR3* (Wenkel et al., 2007) and *35S::miR165* (Kim et al., 2010). For senescence phenotyping, *Arabidopsis thaliana* plants were grown in a climatic chamber at 20°C under long-day conditions (16 h of light) with only moderate light intensity (60–100 $\mu\text{mol s}^{-1} \text{m}^{-2}$) to slow down development for better analyses. Under these conditions, the plants developed bolts and flowers within 5–6 weeks. During growth and development of the leaves, the respective positions within the rosette were color coded with different colored threads, so that even at very late stages of development, individual leaves could be analyzed according to their age. Plants were harvested in a weekly rhythm and samples were always taken at the same time in the morning to avoid circadian effects. For the evaluation of leaf senescence phenotypes, leaves of at least six plants were categorized in four groups according to their leaf color: (1) ‘green’; (2) leaves starting to get yellow from the tip as ‘yellow-green’; (3) completely yellow leaves as ‘yellow’; and (4) dry and/or brown leaves as ‘brown/dry’. Exogenous hydrogen peroxide treatment was conducted by spraying 1%, 0.1% or 0.01% hydrogen peroxide solution including 0.1% Tween20. Grafting experiments were carried out according to Marsch-Martínez et al. (2013).

Intracellular hydrogen peroxide measurements

After stress treatment, leaf 7 (0.1% H_2O_2 treatment) and leaf 8 (heat stress, 2 h at 39°C) were harvested and incubated for exactly 45 min in DCFDA working-solution (2',7'-dichlorodihydrofluorescein diacetate, 200 μg in 40 ml MS-Medium, pH 5.7–5.8). Leaves were then rinsed with water and

frozen in liquid nitrogen. After homogenization on ice, 500 μl 40 mM Tris (pH 7.0) were added and the samples were centrifuged at 4°C for 30 min. Fluorescence (480 nm excitation, 525 nm emission) of the supernatant was measured in a Berthold TriStar LB941 plate reader.

Chromatin-immunoprecipitation and quantitative PCRs

ChIP and ChIP-qPCRs were carried out as described by Brandt et al. (2012). To quantify gene expression changes, RNA was isolated from seedlings using the roboklon GeneMATRIX universal RNA purification kit following manufacturer's recommendations. One microgram of total RNA was reverse transcribed using the Fermentas RevertAid Premium Reverse transcriptase with oligo-dT primers. cDNAs were diluted 10-fold and 3.5 μl were used for RT-PCR reactions. Quantitative measurements were performed on a Bio-Rad CFX384 using the Fermentas SYBR Green qPCR master mix. Relative quantities were calculated using the delta Ct method and normalized relative to a standard curve. Oligonucleotide sequences are listed in supplementary material Table S1. Further descriptions of the methods can be found in the supplementary material. The ChIP-Seq dataset has been published in the Gene Expression Omnibus database (accession number GSE26722).

Redox-DPI-ELISA

Recombinant 6xHis-tagged REV protein with and without the PAS domain was expressed in *E. coli* and DNA-protein interaction ELISA was basically performed as described previously (Brand et al., 2010). Crude extracts were pre-incubated with different concentrations of DTT and H_2O_2 to examine a redox state-dependent binding of REV (for a detailed description, see methods in the supplementary material).

Transformation of *Arabidopsis* protoplasts and transient promoter-GUS expression

Protoplasts were derived from a cell culture of *Arabidopsis thaliana* var. Columbia 0 and were transformed with effector and reporter plasmids following roughly the protocol of Negrutiu et al. (1987). The GUS activity assays were carried out as described by Jefferson et al. (1987). A detailed description is presented in the methods in the supplementary material.

Chlorophyll measurements and phenotypic analysis

For assessment of the leaf senescence state, chlorophyll content of leaf 5 was measured using an atLeaf+ chlorophyll meter (<http://www.atleaf.com>), lipid peroxidation of leaf 6 was measured using the improved thiobarbituric acid/ reactive substances assay, as described previously (Hodges and Fomey, 2000), and expression of the senescence-associated marker genes *SAG12* (At5g45890) and *SAG13* (At2g29350) was analyzed by qRT-PCR. A detailed description is presented in the methods in the supplementary material.

Acknowledgements

We thank Gesine Seibold and Ingrid Blumberg for excellent technical support.

Competing interests

The authors declare no competing financial interests.

Author contributions

K.H., M.P. and J.D. performed senescence phenotyping experiments and redox-ELISA; R.B., T.D. and Y.X. carried out the molecular analysis; D.S. did the gel shift experiment; S.B. measured hydrogen peroxide levels; U.Z. and S.W. designed research, analyzed the data and wrote the article.

Funding

The authors' research is supported by grants from the Deutsche Forschungsgemeinschaft [WE4281/6-1, ZE 313/8-2, ZE 313/9-1 and the Sonderforschungsbereich SFB1101], from the European Union [FP7, ITN MERIT] and from the European Research Council (to S.W.). Deposited in PMC for immediate release.

Supplementary material

Supplementary material available online at <http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.117689/-DC1>

References

- Ay, N., Irmiler, K., Fischer, A., Uhlemann, R., Reuter, G. and Humbeck, K. (2009). Epigenetic programming via histone methylation at WRKY53 controls leaf senescence in *Arabidopsis thaliana*. *Plant J.* **58**, 333-346.
- Balazadeh, S., Riaño-Pachón, D. M. and Mueller-Roeber, B. (2008). Transcription factors regulating leaf senescence in *Arabidopsis thaliana*. *Plant Biol.* **10** Suppl. 1, 63-75.
- Balazadeh, S., Siddiqui, H., Allu, A. D., Matallana-Ramirez, L. P., Caldana, C., Mehrnia, M., Zanon, M. I., Kohler, B. and Mueller-Roeber, B. (2010). A gene regulatory network controlled by the NAC transcription factor ANAC092/AtNAC2/ORE1 during salt-promoted senescence. *Plant J.* **62**, 250-264.
- Balazadeh, S., Kwasniewski, M., Caldana, C., Mehrnia, M., Zanon, M. I., Xue, G.-P. and Mueller-Roeber, B. (2011). ORS1, an H₂O₂-responsive NAC transcription factor, controls senescence in *Arabidopsis thaliana*. *Mol. Plant* **4**, 346-360.
- Ballaré, C. L. (1999). Keeping up with the neighbours: phytochrome sensing and other signalling mechanisms. *Trends Plant Sci.* **4**, 97-102.
- Besseau, S., Li, J. and Palva, E. T. (2012). WRKY54 and WRKY70 co-operate as negative regulators of leaf senescence in *Arabidopsis thaliana*. *J. Exp. Bot.* **63**, 2667-2679.
- Bieker, S., Riester, L., Stahl, M., Franzaring, J. and Zentgraf, U. (2012). Senescence-specific alteration of hydrogen peroxide levels in *Arabidopsis thaliana* and oilseed rape spring variety *Brassica napus* L. cv. Mozart. *J. Integr. Plant Biol.* **54**, 540-554.
- Bou-Torrent, J., Salla-Martret, M., Brandt, R., Musielak, T., Palauqui, J.-C., Martínez-García, J. F. and Wenkel, S. (2012). ATHB4 and HAT3, two class II HD-ZIP transcription factors, control leaf development in *Arabidopsis*. *Plant Signal. Behav.* **7**, 1382-1387.
- Brand, L. H., Kirchler, T., Hummel, S., Chaban, C. and Wanke, D. (2010). DPI-ELISA: a fast and versatile method to specify the binding of plant transcription factors to DNA in vitro. *Plant Methods* **6**, 25.
- Brandt, R., Salla-Martret, M., Bou-Torrent, J., Musielak, T., Stahl, M., Lanz, C., Ott, F., Schmid, M., Greb, T., Schwarz, M. et al. (2012). Genome-wide binding-site analysis of REVOLUTA reveals a link between leaf patterning and light-mediated growth responses. *Plant J.* **72**, 31-42.
- Breeze, E., Harrison, E., McHattie, S., Hughes, L., Hickman, R., Hill, C., Kiddle, S., Kim, Y.-s., Penfold, C. A., Jenkins, D. et al. (2011). High-resolution temporal profiling of transcripts during *Arabidopsis* leaf senescence reveals a distinct chronology of processes and regulation. *Plant Cell* **23**, 873-894.
- Brouwer, B., Ziolkowska, A., Bagard, M., Keech, O. and Gardestrom, P. (2012). The impact of light intensity on shade-induced leaf senescence. *Plant Cell Environ.* **35**, 1084-1098.
- Brusslan, J. A., Rus Alvarez-Canterbury, A. M., Nair, N. U., Rice, J. C., Hitchler, M. J. and Pellegrini, M. (2012). Genome-wide evaluation of histone methylation changes associated with leaf senescence in *Arabidopsis*. *PLoS ONE* **7**, e33151.
- Buchanan-Wollaston, V., Page, T., Harrison, E., Breeze, E., Lim, P. O., Nam, H. G., Lin, J.-F., Wu, S.-H., Swidzinski, J., Ishizaki, K. et al. (2005). Comparative transcriptome analysis reveals significant differences in gene expression and signalling pathways between developmental and dark/starvation-induced senescence in *Arabidopsis*. *Plant J.* **42**, 567-585.
- Byrne, M. E. (2006). Shoot meristem function and leaf polarity: the role of class III HD-ZIP genes. *PLoS Genet.* **2**, e89.
- Carlsbecker, A., Lee, J.-Y., Roberts, C. J., Dettmer, J., Lehesranta, S., Zhou, J., Lindgren, O., Moreno-Risueno, M. A., Váten, A., Thitamadee, S. et al. (2010). Cell signalling by microRNA165/6 directs gene dose-dependent root cell fate. *Nature* **465**, 316-321.
- Causier, B., Ashworth, M., Guo, W. and Davies, B. (2012). The TOPLESS interactome: a framework for gene repression in *Arabidopsis*. *Plant Biol.* **158**, 423-438.
- Emery, J. F., Floyd, S. K., Alvarez, J., Eshed, Y., Hawker, N. P., Izhaki, A., Baum, S. F. and Bowman, J. L. (2003). Radial patterning of *Arabidopsis* shoots by class III HD-ZIP and KANADI genes. *Curr. Biol.* **13**, 1768-1774.
- Guo, Y. and Gan, S. (2005). Leaf senescence: signals, execution, and regulation. *Curr. Top. Dev. Biol.* **71**, 83-112.
- Guo, Y., Cai, Z. and Gan, S. (2004). Transcriptome of *Arabidopsis* leaf senescence. *Plant Cell Environ.* **27**, 521-549.
- Hinderhofer, K. and Zentgraf, U. (2001). Identification of a transcription factor specifically expressed at the onset of leaf senescence. *Planta* **213**, 469-473.
- Hodges, D. M., DeLong, J. M., Forney, C. F. and Prange, R. K. (1999). Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* **270**, 604-611.
- Jefferson, R. A., Kavanagh, T. A. and Bevan, M. W. (1987). GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* **6**, 3901-3907.
- Karimi, M., Inzé, D. and Depicker, A. (2002). GATEWAY™ vectors for Agrobacterium-mediated plant transformation. *Trends Plant Sci.* **7**, 193-195.
- Kichey, T., Hirel, B., Heumez, E., Dubois, F. and Le Gouis, J. (2007). In winter wheat (*Triticum aestivum* L.), post-anthesis nitrogen uptake and remobilisation to the grain correlates with agronomic traits and nitrogen physiological markers. *Field Crop Res.* **102**, 22-32.
- Kim, Y.-S., Kim, S.-G., Lee, M., Lee, I., Park, H.-Y., Seo, P. J., Jung, J.-H., Kwon, E.-J., Suh, S. W., Paek, K.-H. et al. (2008). HD-ZIP III activity is modulated by competitive inhibitors via a feedback loop in *Arabidopsis* shoot apical meristem development. *Plant Cell* **20**, 920-933.
- Kim, H.-S., Kim, S. J., Abbasi, N., Bressan, R. A., Yun, D.-J., Yoo, S.-D., Kwon, S.-Y. and Choi, S.-B. (2010). The DOF transcription factor Dof5.1 influences leaf axial patterning by promoting *Revoluta* transcription in *Arabidopsis*. *Plant J.* **64**, 524-535.
- Kim, J. I., Murphy, A. S., Baek, D., Lee, S.-W., Yun, D.-J., Bressan, R. A. and Narasimhan, M. L. (2011). YUCCA6 over-expression demonstrates auxin function in delaying leaf senescence in *Arabidopsis thaliana*. *J. Exp. Bot.* **62**, 3981-3992.
- Li, Z., Peng, J., Wen, X. and Guo, H. (2012). Gene network analysis and functional studies of senescence-associated genes reveal novel regulators of *Arabidopsis* leaf senescence. *J. Integr. Plant Biol.* **54**, 526-539.
- Lin, J.-F. and Wu, S.-H. (2004). Molecular events in senescing *Arabidopsis* leaves. *Plant J.* **39**, 612-628.
- Magnani, E. and Barton, M. K. (2011). A per-ARNT-sim-like sensor domain uniquely regulates the activity of the homeodomain leucine zipper transcription factor REVOLUTA in *Arabidopsis*. *Plant Cell* **23**, 567-582.
- Marsch-Martínez, N., Franken, J., Gonzalez-Aguilera, K. L., de Folter, S., Angenent, G. and Alvarez-Buylla, E. R. (2013). An efficient flat-surface collar-free grafting method for *Arabidopsis thaliana* seedlings. *Plant Methods* **9**, 14.
- McConnell, J. R., Emery, J., Eshed, Y., Bao, N., Bowman, J. and Barton, M. K. (2001). Role of PHABULOSA and PHAVOLUTA in determining radial patterning in shoots. *Nature* **411**, 709-713.
- Miao, Y. and Zentgraf, U. (2007). The antagonist function of *Arabidopsis* WRKY53 and ESR/ESP in leaf senescence is modulated by the jasmonic and salicylic acid equilibrium. *Plant Cell* **19**, 819-830.
- Miao, Y. and Zentgraf, U. (2010). A HECT E3 ubiquitin ligase negatively regulates *Arabidopsis* leaf senescence through degradation of the transcription factor WRKY53. *Plant J.* **63**, 179-188.
- Miao, Y., Laun, T., Zimmermann, P. and Zentgraf, U. (2004). Targets of the WRKY53 transcription factor and its role during leaf senescence in *Arabidopsis*. *Plant Mol. Biol.* **55**, 853-867.
- Miao, Y., Smykowski, A. and Zentgraf, U. (2008). A novel upstream regulator of WRKY53 transcription during leaf senescence in *Arabidopsis thaliana*. *Plant Biol.* **10** Suppl. 1, 110-120.
- Mittler, R., Vanderauwera, S., Suzuki, N., Miller, G., Tognetti, V. B., Vandepeole, K., Gollery, M., Shulaev, V. and Van Breusegem, F. (2011). ROS signaling: the new wave? *Trends Plant Sci.* **16**, 300-309.
- Muñoz-Espín, D., Cañamero, M., Maraver, A., Gómez-López, G., Contreras, J., Murillo-Cuesta, S., Rodríguez-Baeza, A., Varela-Nieto, I., Ruberte, J., Collado, M. et al. (2013). Programmed cell senescence during mammalian embryonic development. *Cell* **155**, 1104-1118.
- Negrutiu, I., Shillito, R., Potrykus, I., Biasini, G. and Sala, F. (1987). Hybrid genes in the analysis of transformation conditions: I. Setting up a simple method for direct gene transfer in plant protoplasts. *Plant Mol. Biol.* **8**, 363-373.
- Otsuga, D., DeGuzman, B., Prigge, M. J., Drews, G. N. and Clark, S. E. (2001). REVOLUTA regulates meristem initiation at lateral positions. *Plant J.* **25**, 223-236.
- Potschin, M., Schlienger, S., Bieker, S. and Zentgraf, U. (2014). Senescence networking: WRKY18 is an upstream regulator, a downstream target gene, and a protein interaction partner of WRKY53. *J. Plant Growth Regul.* **33**, 106-118.
- Prigge, M. J., Otsuga, D., Alonso, J. M., Ecker, J. R., Drews, G. N. and Clark, S. E. (2005). Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in *Arabidopsis* development. *Plant Cell* **17**, 61-76.
- Reinhart, B. J., Liu, T., Newell, N. R., Magnani, E., Huang, T., Kerstetter, R., Michaels, S. and Barton, M. K. (2013). Establishing a framework for the Ad/Abaxial regulatory network of *Arabidopsis*: ascertaining targets of class III HOMEODOMAIN LEUCINE ZIPPER and KANADI regulation. *Plant Cell* **25**, 3228-3249.
- Rhoades, M. W., Reinhart, B. J., Lim, L. P., Burge, C. B., Bartel, B. and Bartel, D. P. (2002). Prediction of plant microRNA targets. *Cell* **110**, 513-520.
- Rushton, P. J., Somssich, I. E., Ringler, P. and Shen, Q. J. (2010). WRKY transcription factors. *Trends Plant Sci.* **15**, 247-258.
- Smith, Z. R. and Long, J. A. (2010). Control of *Arabidopsis* apical-basal embryo polarity by antagonistic transcription factors. *Nature* **464**, 423-426.
- Smykowski, A., Zimmermann, P. and Zentgraf, U. (2010). G-Box binding factor1 reduces CATALASE2 expression and regulates the onset of leaf senescence in *Arabidopsis*. *Plant Physiol.* **153**, 1321-1331.
- Staudt, A.-C. and Wenkel, S. (2011). Regulation of protein function by 'microProteins'. *EMBO Rep.* **12**, 35-42.
- Storer, M., Mas, A., Robert-Moreno, A., Pecoraro, M., Ortells, M. C., Di Giacomo, V., Yosef, R., Pilpel, N., Krizhanovsky, V., Sharpe, J. et al. (2013). Senescence is a developmental mechanism that contributes to embryonic growth and patterning. *Cell* **155**, 1119-1130.

- Suzuki, N., Miller, G., Salazar, C., Mondal, H. A., Shulaev, E., Cortes, D. F., Shuman, J. L., Luo, X., Shah, J., Schlauch, K. et al. (2013). Temporal-spatial interaction between reactive oxygen species and abscisic acid regulates rapid systemic acclimation in plants. *Plant Cell* **25**, 3553-3569.
- Talbert, P. B., Adler, H. T., Parks, D. W. and Comai, L. (1995). The REVOLUTA gene is necessary for apical meristem development and for limiting cell divisions in the leaves and stems of *Arabidopsis thaliana*. *Development* **121**, 2723-2735.
- Tron, A. E., Bertocini, C. W., Chan, R. L. and Gonzalez, D. H. (2002). Redox regulation of plant homeodomain transcription factors. *J. Biol. Chem.* **277**, 34800-34807.
- Turchi, L., Carabelli, M., Ruzza, V., Possenti, M., Sassi, M., Penalosa, A., Sessa, G., Salvi, S., Forte, V., Morelli, G. et al. (2013). *Arabidopsis* HD-Zip II transcription factors control apical embryo development and meristem function. *Development* **140**, 2118-2129.
- Uauy, C., Distelfeld, A., Fahima, T., Blechl, A. and Dubcovsky, J. (2006). A NAC Gene regulating senescence improves grain protein, zinc, and iron content in wheat. *Science* **314**, 1298-1301.
- Ülker, B., Shahid Mukhtar, M. and Somssich, I. E. (2007). The WRKY70 transcription factor of *Arabidopsis* influences both the plant senescence and defense signaling pathways. *Planta* **226**, 125-137.
- Weigel, M., Varotto, C., Pesaresi, P., Finazzi, G., Rappaport, F., Salamini, F. and Leister, D. (2003). Plastocyanin is indispensable for photosynthetic electron flow in *Arabidopsis thaliana*. *J. Biol. Chem.* **278**, 31286-31289.
- Wenkel, S., Emery, J., Hou, B.-H., Evans, M. M. S. and Barton, M. K. (2007). A feedback regulatory module formed by LITTLE ZIPPER and HD-ZIP III genes. *Plant Cell* **19**, 3379-3390.
- Yang, S.-D., Seo, P. J., Yoon, H.-K. and Park, C.-M. (2011). The *Arabidopsis* NAC transcription factor VNI2 integrates abscisic acid signals into leaf senescence via the COR/RD genes. *Plant Cell* **23**, 2155-2168.
- Zentgraf, U., Jobst, J., Kolb, D. and Rentsch, D. (2004). Senescence-related gene expression profiles of rosette leaves of *Arabidopsis thaliana*: leaf age versus plant age. *Plant Biol.* **6**, 178-183.
- Zentgraf, U., Laun, T. and Miao, Y. (2010). The complex regulation of WRKY53 during leaf senescence of *Arabidopsis thaliana*. *Eur. J. Cell Biol.* **89**, 133-137.
- Zentgraf, U., Zimmermann, P. and Smykowski, A. (2012). Role of intracellular hydrogen peroxide as signalling molecule for plant senescence. In *Senescence* (T. Nagata, ed.). <http://www.intechopen.com/books/senescence/role-of-intracellular-hydrogen-peroxide-as-signalling-molecule-for-plant-senescence>
- Zhong, R. Q. and Ye, Z. H. (1999). IFL1, a gene regulating interfascicular fiber differentiation in *Arabidopsis*, encodes a homeodomain-leucine zipper protein. *Plant Cell* **11**, 2139-2152.
- Zimmermann, P., Heinlein, C., Orendi, G. and Zentgraf, U. (2006). Senescence-specific regulation of catalases in *Arabidopsis thaliana* (L.) Heynh. *Plant Cell Environ.* **29**, 1049-1060.

Supplementary Material Xie et al.

Supplementary methods

GUS staining and histology. For histochemical GUS staining, *REV::GUS* (*rev-9*) and *WRKY53::GUS* plants were incubated overnight in GUS staining solution (100mM NaPO₄ pH7.0, 10mM EDTA, 0.5mM K₄Fe(CN)₆, 0.5mM K₃Fe₆, 0.01% Triton-X and 1mM X-Gluc). Samples were cleared with an ethanol series. For histological sections, roots of ten day-old seedlings were stained overnight in GUS staining solution following transfer into fixative (50mM NaPh pH=7.2; 1% Glutaraldehyde; 4% Formaldehyde) for two days. Afterwards, the roots were dehydrated in an ethanol series (30%/ 50%/ 70% each for two hours) and finally stored in 100% ethanol prior embedding in Technovit (Heraeus). Two-micron sections were cut using a Leica microtome.

Redox-DPI-ELISA. The coding sequence of *REV* with and without the PAS domain was cloned into the pET-32b vector for expression of the proteins fused to a 6xHis-tag. The *E. coli* strain BL21-SI was used for protein expression. The cells were grown in 10ml of selective medium overnight and subsequently diluted 1:20 in a final volume of 100ml in medium without antibiotics. Protein expression was induced after 1h by the addition of 1mM IPTG and the cells were grown a further hour at 37°C. After centrifugation (2500g, 20min, 4°C) and washing (10mM Tris-HCl pH 7.5, 100mM NaCl), the bacterial pellet was resuspended in protein extraction buffer (4mM HEPES pH 7.5, 100mM KCl, 8% (v/v) glycerol, 1x complete proteinase inhibitor without EDTA (Roche) and protein extraction was performed by sonication under native conditions. The protein concentration of the crude extract was measured by

Bradford assay (Bio-Rad). Expression of the recombinant proteins was confirmed by Western blot and immune detection using anti-His-antibodies.

The DNA-protein interaction assay was basically performed as described before (Brand et al., 2010). 5' biotinylated complementary oligonucleotides were annealed (final concentration 2M) to get double-stranded DNA fragments. The double-stranded binding sites of REV in the *WRKY53* promoter (BS1 and BS2) were added to a streptavidin-coated ELISA plate (Nunc Immobilizer) for binding for 1h at 37°C. After blocking using blocking reagent (Roche) for 30min, blocking reagent was removed and 25µg crude extracts was added and incubated for 1h at room temperature. Crude extracts were pre-incubated with different concentrations of DTT and H₂O₂ to examine a redox-state dependant binding of REV. In order to test reversibility of the redox-effect, first high concentrations of H₂O₂ were added and subsequently oxidizing conditions were reversed by addition of DTT. After binding, biotinylated DNA-protein complexes were washed two times for 10min at room temperature (blocking solution, Qiagen) and incubated with Anti His-HRP conjugate antibodies (Qiagen) 1:1500 diluted in blocking solution for 1h at room temperature. After washing, interaction was detected by a peroxidase reaction with ortho-phenylenediamine (OPD-tablets, Agilent technologies (Dako)) for 15min in darkness. After stopping the reaction with 0.5M H₂SO₄ solution, positive interactions which resulted in a yellow color could be measured with an ELISA-reader (TECAN Safire XFluor4).

Transformation of *Arabidopsis* Protoplasts and transient Promoter-GUS Expression.


The protoplasts used for the transient expression assay were derived from a cell culture of *Arabidopsis thaliana* var. *Columbia 0*. Cells were transformed with 5 µg of

effector and reporter plasmid DNA each roughly following the protocol of Negrutiu et al. (1987). For details see the following protocol: <http://www.zmbp.uni-tuebingen.de/CentralFacilities/transf/index.html>. As an internal control 0.1 µg of a Luciferase construct was co-transfected. The protoplasts were incubated overnight in the dark and then used for GUS-assays. For the *WRKY53*-reporter, a 2.8kb promoter sequence upstream of the *WRKY53* start codon was cloned into the binary vector pBGWFS7.0 (Karimi et al., 2002) and served as the reporter construct. The *REVOLUTA* CDS in the pJAN33 vector (Weigel et al., 2003) served as an effector. The GUS activity assays were carried out as described by Jefferson (Jefferson et al., 1987). To normalize differences in protoplast transfection a luciferase-assay was performed using the Dual-Luciferase[®] Reporter Assay System from Promega following the user's manual (Promega, <http://www.promega.de/~media/Files/Resources/Protocols/Technical%20Bulletins/0/Luciferase%20Assay%20System%20Protocol.pdf>). In total 6 biological replicates with 3 technical replicates each were performed for the experiment.


Chlorophyll measurements and phenotypic analysis. For assessment of the leaf senescence state, color-coded leaves were used. Chlorophyll content of leaf No. 5 was measured using an atLeaf+ chlorophyll meter (<http://www.atleaf.com>). Total chlorophyll content of leaves can be obtained by converting the atLEAF+ values in SPAD and considering the relation among chlorophyll content and SPAD units (<http://www.atleaf.com>). Each leaf was measured in triplicate at three different positions. Chlorophyll was determined for three independent plants and values were averaged. In addition, expression of the senescence associated marker genes *SAG12* (At5g45890) encoding a cysteine protease and *SAG13* (At2g29350) encoding a short-

chain alcohol dehydrogenase was analyzed by qRT-PCR and normalized to the expression of the GAPDH gene (At1g13440). Lipid peroxidation was measured using the improved thiobarbituric acid-reactive-substances assay as described previously (Hodges et al., 1999). Leaf No. 6 of three different plants was homogenized in liquid nitrogen and 25 mg of the fine leaf tissue powder was used for analyzes.


Supplementary figures




Suppl Fig. S1. Expression analysis of *REV* and *WRKY53*. Spatial expression patterns of *REV* and *WRKY53* using histochemical staining of 2-week to 6-week-old rosettes of p*WRKY53*::GUS and p*REV*::GUS plant lines.




Suppl Fig. S2. Publicly available microarray-based expression analysis using Arabidopsis eFP browser showing co-expression of *REV* and *WRKY53* in different tissues. Plotted are images showing relative expression values where red color indicates high expression and blue color indicates low expression. Intermediated expression levels are displayed in yellow and variations between yellow and red and yellow and blue.



Suppl. Fig. S3. Measurement of intracellular H₂O₂ contents using DCFDA after 0.1% H₂O₂ application or heat treatment. Error bars indicate standard deviation of at least three biological replicates, *** p<0.0005 (T-Test).




Suppl. Fig. S4. Analysis of REV mRNA changes in response to hydrogen peroxide. Real-time qPCR experiment showing *REV* expression in response to hydrogen peroxide treatment in wild type. 3-week old plants were treated with different concentrations of H₂O₂ (0%, 0.01%, 0.1% and 1%) for 40 min. Plotted are representative relative expression changes (fold change) of the mean of four technical replicates including standard deviations. Similar expression changes have been observed in at least two independent biological experiments.




Suppl. Fig. S5. Quantitative evaluation of senescence in additional lines with altered HD-ZIPIII expression.

For a quantitative evaluation of leaf senescence plants were harvested in a weekly rhythm and leaves of at least ten plants were categorized into four groups according to their leaf color: 1) “green”, 2) leaves starting to get yellow from the tip as “yellow-green”, 3) completely yellow leaves as “yellow” and 4) dry and/or brown leaves as “brown/dry”. The percentages of each group with respect to total leaf numbers are presented. Error bars indicate standard deviations. Col-0 (wild type); *rev6* (different

rev mutant line); *35S::ZPR3* (plant line overexpressing the small leucine-zipper-type microProtein *ZPR3*, a micro protein inhibiting HD-ZIPIII function by protein-protein interaction); *wrky53/35S::ZPR3* (plant line overexpressing *ZPR3* in the *wrky53* mutant background).



Suppl. Fig. S6. Comparison of plant development of WT, *wrky53*, *rev5*, *wrky53/rev5*, *35S::ZRP3* and *35S::ZRP3/wrky53* and *rev6* plants from 5-week to 7-week-old plants



Suppl. Fig S7. Phenotype of leaves of 6-week-old plants of different plant lines.

Table 1. Oligonucleotides

Primers	Sequence	Use
W53CpAf	tgctcaagtttctgggaaaaa	ChIP-qPCR (Primer I)
W53CpAr	aagacgtgctgtcctctgaa	ChIP-qPCR (Primer I)
W53CpBf	caatgaatgaatgacgcaaaa	ChIP-qPCR (Primer II)
W53CpBr	ttcaaaaggacaggattgga	ChIP-qPCR (Primer II)
W53CpCf	cgtatcacacagtactggtttt	ChIP-qPCR (Primer III)
W53CpCr	ctgaccaagtcacatggaa	ChIP-qPCR (Primer III)
qW53f	aaactgttgggcaacgaaac	ChIP-qPCR (Primer IV)
qW53r	aatggctggttgactctgg	ChIP-qPCR (Primer IV)
REVe56F	ttctttgcctaattgcttgg	rev5 genotyping (with rev5R and BaeI digest)
rev5R	tcataaaggggtcgaagcac	rev5 genotyping (with REVe56F and BaeI digest)
vam-F_rev6	tgctgcttcagcttctcagt	rev6 genotyping (with REV-3R and TaqI digest)
REV-3R	gaagcacctccaaccgtaga	rev6 genotyping (with vam-F_rev6 and TaqI digest)
W53tDNAr	gttcaagtcctgttgaaattcc	wrky53 genotyping
W53tDNAf	ggcagtggtccagaatctcc	wrky53 genotyping
LB-Salk	tggttcacgtagtgggccatcg	wrky53 genotyping
AtREV-topo_F	caccaacatacatgacatgtgaaat	Cloning of the pREV::REV::GFP construct
AtREV-topo_R	cacaaaagaccagtttacaaggag	Cloning of the pREV::REV::GFP construct
qGAPDH_F	aaagtgttgccatccctcaa	AT5G47180 qRT-qPCR
qGAPDH_R	tcggtagacacaacatcatcct	AT5G47180 qRT-qPCR
qWRKY53_F	ggcagtggtccagaatctcc	WRKY53 qRT-PCR
qWRKY53_R	gcctctctctgggcttattc	WRKY53 qRT-PCR
qSAG12_F	ggaggaaaacaatcgtctacg	SAG12 RT-qPCR
qSAG12_R	acggcgacatttttagtttg	SAG12 RT-qPCR
qSAG13_F	gtgccagagacgaaactc	SAG13 RT-qPCR
qSAG13_R	gctgtaaactctgtggtc	SAG13T-qPCR
qHAT3_2f	ggagtttcgtctccgaacag	HAT3 RT-qPCR
qHAT3_2r	atcggagtttcttctctgaa	HAT3 RT-qPCR
qZFP8_F	ccgccattattcgtctcttc	ZFP8 RT-qPCR
qZFP8_R	gatgtccacctagggttga	ZFP8 RT-qPCR
qIDD11_F	tggcaacaacagattcgtg	IDD11 RT-qPCR
qIDD11_R	gatggatcatggtggacaca	IDD11 RT-qPCR
qAT5G47180_F	caagcgcaacgagagtatcc	ZFP8 RT-qPCR
qAT5G47180_R	ttgggtgttagacggggtaa	ZFP8 RT-qPCR
qAT1G74940_F	ggtattgttgctgcgttgga	AT1G74940 qRT-qPCR
qAT1G74940_R	acaattcaaacccgtcgtcg	AT1G74940 qRT-qPCR
qAT1G49200_F	tctgttgctcgggttacat	AT1G49200 qRT-qPCR
qAT1G49200_R	cggcaagttcatctcaggtg	AT1G49200 qRT-qPCR

Supplementary references

Karimi, M., Inze, D. and Depicker, A. (2002). GATEWAY vectors for

Agrobacterium-mediated plant transformation. Trends Plant Sci 7, 193-195.

Weigel, M., Varotto, C., Pesaresi, P., Finazzi, G., Rappaport, F., Salamini, F. and

Leister, D. (2003). Plastocyanin Is Indispensable for Photosynthetic Electron Flow in

Arabidopsis thaliana. Journal of Biological Chemistry 278, 31286-31289.