

REVIEW

Heparan sulfate proteoglycans: a sugar code for vertebrate development?

Fabienne E. Poulin¹ and H. Joseph Yost^{2,*}

ABSTRACT

Heparan sulfate proteoglycans (HSPGs) have long been implicated in a wide range of cell-cell signaling and cell-matrix interactions, both *in vitro* and *in vivo* in invertebrate models. Although many of the genes that encode HSPG core proteins and the biosynthetic enzymes that generate and modify HSPG sugar chains have not yet been analyzed by genetics in vertebrates, recent studies have shown that HSPGs do indeed mediate a wide range of functions in early vertebrate development, for example during left-right patterning and in cardiovascular and neural development. Here, we provide a comprehensive overview of the various roles of HSPGs in these systems and explore the concept of an instructive heparan sulfate sugar code for modulating vertebrate development.

KEY WORDS: Patterning, Left/right asymmetry, Nervous system, Heart, Glycan, Sugars

Introduction

Vertebrate development – from a fertilized egg to a whole organism – is a complex process orchestrated by many signaling pathways. The generation of various cell types and organs from a single cell is achieved through successive developmental programs that require the spatial and temporal coordination of developmental signaling molecules, including morphogens and growth factors. Evidence accumulated over the past twenty years has shown that most, if not all, of these developmental factors are regulated by heparan sulfate proteoglycans (HSPGs).

HSPGs are composed of a core protein to which long linear glycosaminoglycan heparan sulfate (HS) chains are covalently linked. Syndecans (Sdc) and glypcans (Gpc), which are among the main classes of core protein, are anchored to the cell surface by a transmembrane domain or a glycosylphosphatidylinositol anchor, respectively, and can be released into the extracellular space following cleavage by various enzymes (Fig. 1). Other HSPGs, including agrin, collagen XVIII and perlecan, are directly secreted into the extracellular matrix (ECM). Core proteins have specific numbers of HS chain addition sites, and some also harbor chondroitin sulfate (CS) as another type of glycosaminoglycan (Table S1). HS chains are synthesized in the Golgi apparatus in a multi-step process that involves several enzymes (Fig. 2; Table S2). Exostosin (Ext) enzymes elongate HS chains by adding alternating glucuronic acid and *N*-acetylglucosamine residues, whereas *N*-deacetylase/*N*-sulfotransferase (Ndst) enzymes, C5 epimerase and

2-O-, 3-O-, and 6-O-sulfotransferases modify them by catalyzing deacetylation, epimerization and sulfations at different positions (for nomenclature, see Lawrence et al., 2008). These modifications do not occur uniformly along the HS chains but instead are concentrated in different sub-regions, creating domains with variably modified disaccharides that can interact with other proteins in distinct manners.

Given their various HS chains and localization (i.e. at the cell surface or in the ECM), HSPGs have been proposed to regulate signaling pathways in many different ways (Fig. 3). They can act cell-autonomously as receptors or co-receptors [for instance, during Wnt or fibroblast growth factor (FGF) signaling], as ‘recruiters’ to lipid rafts (thereby increasing ligand or receptor concentration at the cell surface), by regulating receptor membrane trafficking (during endocytosis) or by controlling ligand secretion. They can also act non-cell-autonomously as direct cues, or by controlling the distribution of signaling gradients as well as the composition of the ECM. Adding to the complexity of their signaling functions, HSPGs can interact with proteins through various mechanisms (for reviews, see Lindahl and Li, 2009; Xu and Esko, 2014). Binding to proteins is generally mediated by HS chains, but some factors such as Hedgehog (Hh) have been shown to directly interact with core proteins (Capurro et al., 2008). Whereas some interactions require specific structural motifs along the HS chains (Thunberg et al., 1982; de Paz et al., 2007), others are non-selective in terms of motifs but require a certain type of sulfation (for instance, 2-O-sulfation is essential for Fgf2 binding; Ashikari-Hada et al., 2004) or a certain negative charge density. Interestingly, it has been demonstrated that distinct organs or tissues produce unique compositions of HS, and that HS epitope expression changes during embryonic development in a tissue-specific manner (Allen and Rapraeger, 2003; Ledin et al., 2004). This observation, together with the large variability of HS chains that can be generated by the HS-modifying enzymes, has led to the hypothesis of a complex ‘sugar code’ (Holt and Dickson, 2005), in which specific HS modifications present in a region-specific manner in the embryo would orchestrate developmental programs by interacting with and modulating defined signaling pathways. This sugar code would thus depend on the spatial and temporal regulation of HS-modifying enzymes, and perhaps on different members of the same family of HS-modifying enzymes having slightly different activities or specificities of actions. For example, some family members of the 3-O-sulfotransferase family might have a greater or lesser propensity to make a modification adjacent to other sulfation marks, changing both the configuration of sulfation marks and the charge density in particular regions of an HS chain. The sugar code would also have to be dynamic, as HS chains can be degraded or modified at the cell surface by heparanase and sulfatase enzymes, respectively.

Although studies in *Caenorhabditis elegans* have strongly supported the sugar code hypothesis (Bülow et al., 2008; Tecle et al., 2013; Díaz-Balzac et al., 2014), evidence for specific roles of HS modifications and core proteins in vertebrates is only beginning to

¹Department of Biological Sciences, University of South Carolina, Columbia, SC 29208, USA. ²University of Utah, Department of Neurobiology and Anatomy, Department of Pediatrics, Salt Lake City, UT 84132, USA.

*Author for correspondence (jyost@genetics.utah.edu)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.




Fig. 1. Examples of key cell surface and extracellular HSPGs. Glycans (Gpc) are attached to the cell surface via a glycosylphosphatidylinositol (GPI) anchor. They possess a large globular domain stabilized by conserved di-sulfide (S-S) bonds, and HS chains (represented by chains of pink, blue and purple circles) in their C-terminal part. They can be released into the extracellular matrix following cleavage of their GPI anchor by the lipase Notum. A furin-like convertase can also cleave Gpc at the C-terminal end of their globular domain, leading to the formation of two subunits that remain attached to each other by disulfide bonds. Syndecans (Sdcs) are single-pass transmembrane proteins with HS chains attached to their N-terminal part. Their intracellular region interacts with many different partners through two conserved domains, constant 1 (C1) and constant 2 (C2), that are separated by a more variable region (V). Like Gpc, Sdcs can be shed into the extracellular environment after cleavage by proteases such as matrix metalloproteases and a disintegrin and metalloproteinase (ADAM) disintegrins. Agrin and perlecan are large multidomain proteoglycans that carry several HS chains and are secreted as different isoforms generated by alternative splicing. The cleavage of the C-terminal region of perlecan by metalloproteases releases endorepellin, an angiogenesis inhibitor. C-ter, C-terminus; N-ter, N-terminus.

emerge. Nonetheless, accumulative data clearly indicate that HSPGs have distinct instructive roles during vertebrate development. For example, studies have shown that many organs, including the hematopoietic and musculoskeletal systems, and the liver, lungs and kidneys, do not form properly when HSPGs are absent or modified (Tables S1, S2; for reviews, see Thompson et al., 2010; Huegel et al., 2013; Nigam and Bush, 2014). These studies have utilized various model organisms, including mice, zebrafish and *Xenopus*, although it should be noted that some disparities between results have arisen, perhaps owing to the technical challenges of assessing HSPG functions *in vivo* (see Box 1). In this Review, we discuss the specific functions of HSPGs during vertebrate development, focusing on the early stages of development and the formation of the cardiovascular and nervous systems. We then evaluate the concept of the sugar code hypothesis, discussing evidence for and against it, and highlighting future challenges for the field.

Roles for HSPGs in early development

The first stages of development following egg fertilization (including cell cleavage, gastrulation and patterning) establish the different body axes and overall architecture of the embryo. Studies of mutants that are defective for genes encoding HSPG biosynthetic enzymes promote the view that HS sugar chains regulate multiple steps throughout these early developmental processes (Fig. 4A). HS chain polymerization has been shown to be crucial for patterning and early developmental function. Homozygous knockout (KO) mutants of *Ext1* or *Ext2* in mice are embryonic lethal, exhibiting defects in gastrulation (Lin et al., 2000; Mitchell et al., 2001; Stickens et al., 2005; Shimokawa et al., 2011). Zebrafish *ext2* (*dackel*) or *extl3* (*boxer*) mutants undergo normal gastrulation owing to a maternal contribution of both enzymes, but later develop cartilage and limb defects and display impaired FGF, Wnt and Shh

signaling (Grandel et al., 2000; Norton et al., 2005; Clément et al., 2008; Holmborn et al., 2012). HS chain modifications also appear to be important, as the absence of both *Ndst1* and *Ndst2*, which encode enzymes that catalyze HS deacetylation and N-sulfation, is embryonic lethal in mice (Grobe et al., 2002), and epiboly is disrupted in zebrafish embryos injected with morpholinos (MOs) against *hs2st*, which encodes a sulfotransferase that adds sulfate to the C2 position of hexuronic acid residues along the HS chain (Cadwalader et al., 2012). Interestingly, the other reported mutants in HS-modifying enzymes only show later developmental defects (Table S2), suggesting functional redundancy between multiple gene family members that can contribute to specific biochemical steps in HSPG synthesis. Analyzing combinations of mutants will be required to uncover the early developmental functions of HSPGs.

Studies of mutants that are defective for individual core proteins also provide evidence for early developmental roles for HSPGs. In zebrafish, *gpc4* (*knyapek*) mutants exhibit defects in Wnt11-dependent convergent-extension cell movements during gastrulation, suggesting that the potentiation of Wnt11 signaling is dependent on HSPG function (Topczewski et al., 2001). Gastrulation and posterior elongation defects are also observed in *Xenopus* embryos lacking or overexpressing *gpc4* (Ohkawara et al., 2003). By contrast, *Gpc4* KO mutants in mice develop normally and only show later defects in neural synapse formation (Allen et al., 2012). Other reported core proteins mutants have later onset developmental defects (Table S1), including a wide range of specific neurodevelopmental defects (discussed below), that often result in perinatal death.

Roles for HSPGs in left-right patterning

Kupffer's vesicle (KV) is a ciliated organ that controls left-right (LR) patterning in the zebrafish embryo. During normal




Fig. 2. The synthesis of heparan sulfate chains. Heparan sulfate (HS) chains are linked by a xylose and a tetrasaccharide linkage to a specific serine residue of a core protein. This linkage region is identical in HS and chondroitin (CS) chains. The addition of the first N-acetylglucosamine is then catalyzed by the enzyme EXT1. Elongation of the chains is achieved through the alternative addition of glucuronic acid and N-acetylglucosamine by the transferases EXT1 and EXT2. During polymerization, chains undergo several modifications including epimerization of glucuronic acid and sulfations at different positions. These modifications occur in different clusters and generate N-acetylated, N-sulfated and mixed domains differently involved in interactions with partners. Ser, serine; NS, N-sulfate group on glucosamine residues; 2S, 2-O-sulfate group on glucuronic or iduronic acid residues; 3S, 3-O-sulfate group on glucosamine residues; 6S, 6-O-sulfate group on glucosamine residues. Adapted from Esko et al., 2009.

development, KV cells migrate and undergo a mesoderm-to-epithelial transition to form a vesicle. As the vesicle is inflated, each KV cell grows a specialized apical cilium that is motile and moves fluid asymmetrically in the KV, thereby establishing LR patterning throughout the embryo. A number of studies have implicated a role for HSPGs during this process of LR patterning (Fig. 4B).

The lineage-specific targeting of *sdc2* in zebrafish KV cells with MOs results in disrupted KV morphogenesis and shortened cilia, giving rise to altered LR patterning throughout the embryo (Arrington et al., 2013). Whole-embryo *sdc2* knockdown also causes later defects in vasculogenesis (Chen et al., 2004). Roles for HSPGs in LR patterning have also been assessed in *Xenopus* embryos, which offer the ability to identify and target cell lineages more precisely by microinjection. For example, the lineage-specific expression of dominant-negative mutant Sdc2 isoforms by targeted injection of mRNAs into individual *Xenopus* cells results in LR patterning defects (Kramer and Yost, 2002). Strikingly, the dominant-negative Sdc2 isoforms, including phosphomimetic and phosphodeficient forms that have altered phosphorylation sites in the cytoplasmic domain of the core protein, indicate that normal LR patterning requires both non-phosphorylated Sdc2 on the left and protein kinase C gamma (PKC γ)-dependent phosphorylated Sdc2 on the right side of the embryo (Kramer et al., 2002). This reveals a developmental role for asymmetric Sdc2 phosphorylation in LR patterning. However, it is possible that both the dominant-negative constructs and the MOs affect more than just Sdc2; dominant-negative Sdc2 might, for example, disrupt the complex formed by Sdc2 and Sdc4. It is

therefore important to confirm these observations using combinations of mutants in the Sdc family.

HS chain modifications catalyzed by 3-O-sulfotransferases (encoded by *Hs3st* genes) have also been implicated in LR patterning. The MO-based lineage-specific targeting of *hs3st1l2* (also known as *hs3st5*) and *hs3st3l* (also known as *hs3st6*) in the zebrafish KV results in normal KV morphogenesis, but causes LR patterning defects that appear to be due to distinct perturbations in KV cilium length and cilia motility (Neugebauer et al., 2013). This study also implicated 3-O-sulfotransferase modifications in the regulation of Fgf8 and other cell signaling pathways. Few other mutants have been reported for the *Hs3st* family, although it should be noted that knockouts of *Hs3st1* in mice have subtle intrauterine growth retardation that is background dependent (Shworak et al., 2002; HajMohammadi et al., 2003).

HSPG functions in cardiovascular development

Roles for core proteins as well as HS modifications in the development of the cardiovascular system have been identified. During early organogenesis, the mesoderm lineages that give rise to the heart and the mesoderm of the gut are present at the lateral edges of the vertebrate embryo and migrate towards the embryonic midline, fusing to form the cardiac and gut tube, respectively. The MO-mediated lineage-specific knockdown of zebrafish *sdc2* reveals a role for this HSPG core protein in the non-cell-autonomous regulation of ECM formation, which is required for cardiac and gut mesoderm migration. In particular, the disruption of *sdc2* function in embryonic yolk cells leads to altered ECM formation throughout




Fig. 3. The regulation of cell signaling pathways by HSPGs. HSPGs regulate signaling pathways in many different ways, both at the intracellular level and in the extracellular matrix (ECM). HSPGs have been shown to: (1) mediate signal transduction by acting as receptors or co-receptors; (2) regulate receptor trafficking to and from the plasma membrane; (3) control the secretion of ligands; (4) signal to other cells by acting themselves as cues or presenting ligands to their receptors; and (5) regulate the structure of the ECM and the establishment of signaling gradients.

the embryo and impedes the migration of both cardiac and gut precursor cells, resulting in cardia bifida (two separate heart tubes) and bifid gut tubes (Arrington and Yost, 2009). Among the other core proteins, only Gpc3 and Perlecan (Hspg2 – Zebrafish Information Network) have so far been implicated in heart development. Congenital cardiac malformation, including ventricular septal defects or double outlet right ventricle, have been reported in mice lacking *Gpc3* and seem to result from impaired Shh signaling in the heart (Ng et al., 2009). Perlecan-deficient mice show a different phenotype, characterized by malformations of the cardiac outflow tract (Costell et al., 2002).

An important insight into the role of HS chains in cardiovascular development came from the analysis of *Ndst1* KO mice, which show several heart defects reminiscent of those observed in the absence of *Gpc3* (Pan et al., 2014). FGF signaling is also strongly reduced in *Ndst1* KO mice, and impaired Fgf8 function appears to contribute to their phenotype. The conditional deletion of *Ndst1* in neural crest cells in a systemic *Ndst2*-deficient background further indicates an important function of HS in the neural crest cell lineage that migrates into the developing cardiac tube during normal cardiac development, and suggests at least partially compensatory roles among *Ndst* family members. The functions of the other HS chain modifications in cardiovascular development remain difficult to address. Mice lacking C5-epimerase or Hs2st die at birth and show severe developmental defects (Table S2). Furthermore, various phenotypes have been reported for mice lacking *Hs6st1* (Pratt et al., 2006; Habuchi et al., 2007; Izvolsky et al., 2008), whereas double *Hs6st1* and *Hs6st2* mutants have enhanced phenotypes and are embryonic lethal (Conway et al., 2011b), indicating some level of functional redundancy between the members of the 6-O-sulfotransferase family. Zebrafish embryos injected with MOs against the 3-O-sulfotransferase *hs3st7* (*hs3stIII*) exhibit normal LR patterning and early development but show defects in contraction of the cardiac ventricle. This myocardium contraction defect can be

rescued by the transgenic expression of *hs3st7* in the endocardium tissue that lines the cardiac tube, but not in the myocardium, indicating that the regulation of myocardium contraction by *hs3st7* is non-cell-autonomous, implicating cell-cell signaling. Strikingly, genetic reduction of bone morphogenetic protein (BMP) signaling in the heart also rescues contraction in *hs3st7* morphants (Samson et al., 2013). Together, these results suggest that different members of the *Hs3st* family control distinct steps, and perhaps distinct cell signaling pathways, during cardiac development.

Roles for HSPGs during nervous system development

Nervous system development proceeds through several partially concomitant steps that are regulated by a large diversity of environmental factors, such as morphogens, guidance cues, growth factors, adhesion molecules and components of the ECM, distribution and/or function of which are in large part controlled by HSPGs (Fig. 3). Although the crucial role of HS synthesis in nervous system formation was demonstrated over ten years ago, novel functions for HS modifications as well as core proteins are being discovered, suggesting that the diversity of these molecules might provide a 'code' regulating each aspect of nervous system development in an instructive manner (Fig. 5).

Nervous system patterning

During its early development, the neural tube becomes regionalized into distinct domains that give rise to the major structures of the central nervous system. The crucial role of HSPGs in brain patterning was first demonstrated in *Nestin-cre; Ext1* conditional knockout (cKO) mice (*Nestin-cre; Ext1^{fl/fl}*), in which HS synthesis is abolished in neural stem cells (NSCs) and, consequently, in the three lineages originating from them: neurons, astrocytes and oligodendrocytes (Inatani et al., 2003). These mice die at birth and show several patterning defects, including a lack of discrete cerebellum and the absence of olfactory bulbs. Interestingly, a

Box 1. Analyzing HSPG functions in development: approaches and challenges

The functions of HSPGs in development have been assessed using various approaches, including targeted mutants in mice, genetic mutants in zebrafish and morpholino (MO) knockdowns in zebrafish and *Xenopus*. However, a growing challenge in the field is to reconcile the disparity among the phenotypes generated by these different methods (Tables S1, S2). In the absence of genetic mutants, the roles of HSPGs in zebrafish have mostly been assessed based on MO phenotypes, and these studies have provided some interesting insights into the specific roles of HSPG core proteins and biosynthetic enzymes in embryonic morphogenesis. There are, however, increasing concerns about potential MO artifacts and the possibility that mutants can be functionally compensated by other family members or other regulatory events (Kok et al., 2015; Stainier et al., 2015). As such, conclusions from MO-based studies await confirmation by engineered mutants in zebrafish. It is also somewhat surprising that mutants for some of the core protein genes, such as *sdc2* or *gpc5*, have not yet been reported in mice, and that mutations in the same gene (for instance *gpc4*) cause drastically different phenotypes in mice versus zebrafish or *Xenopus*. It is thus likely that a fuller understanding of HSPG functions will require the combination of multiple mutants, transgenic and other techniques to test lineage-specific roles of HSPG core proteins and biosynthetic pathways in vertebrate development.

similar phenotype is observed in hypomorphic *Fgf8* mutants (Meyers et al., 1998) and in the *swaying* mice, which exhibit a recessive mutation in *Wnt1* (Thomas et al., 1991). The analysis of *Fgf8* immunoreactivity and of the expression domains of genes downstream of FGF in *Nestin-cre; Ext1^{fl/fl}* mice revealed an expanded distribution and activity of *Fgf8* in the absence of HS, suggesting that HSPGs regulate midbrain/hindbrain patterning by controlling *Fgf8* levels in the environment. Supporting this hypothesis, a more recent study showed that the formation of an FGF posterior-to-anterior gradient along the midbrain is impaired after treatment with heparitinase or heparin (Chen et al., 2009).


Further investigations revealed that HS modifications appear to be essential for brain patterning. Mice lacking the N-deacetylase/N-sulfotransferase *Ndst1* display forebrain defects, including cerebral hypoplasia and a lack of olfactory bulbs (Grobe et al., 2005). The regulated removal of 6-O moieties from HS chains at the cell surface also appears important, as mice lacking the endosulfatase *Sulf2* show brain abnormalities, including neural tube closure defects, an enlargement of the ventricular system and hydrocephalus (although with partial penetrance) (Kalus et al., 2009).

More recently, specific core proteins involved in nervous system patterning have been identified. For example, mice lacking *Gpc1* do not form the most anterior lobe of the cerebellum, a defect possibly related to the cerebellar agenesis seen in the *Nestin-cre; Ext1* cKO mice (Jen et al., 2009). The knockdown of *gpc4* in *Xenopus* leads to a different phenotype characterized by dorsal forebrain defects and an absence of anterior neural tube closure due to impaired FGF signaling (Galli et al., 2003). Defective neural tube closure is also observed in *Xenopus sdc4* morphants, and genetic interaction studies suggest that *sdc4* and *vangl2* act in the Wnt/PCP pathway to regulate this process (Muñoz and Larraín, 2006; Escobedo et al., 2013). *Sdc4* has also been shown to activate the FGF/ERK and PKC signaling pathways to trigger neural induction (Kuriyama and Mayor, 2009).

Neurogenesis

Neurogenesis – the generation and differentiation of neurons from NSCs – occurs predominantly during development, but also continues in the adult in specific regions of the brain, such as the

A Convergent extension



B Left-right patterning




Fig. 4. Functions of HSPGs in early development. (A) Convergent extension allows body axis elongation along the anteroposterior (AP) axis during gastrulation. The collective migration of lateral mesendoderm progenitors (presumptive somites, green) towards the midline (red arrows) and the cell intercalation (white arrows) of notochord progenitors (pink) lead to the extension of the notochord along the AP axis (green arrows). In zebrafish and *Xenopus*, *Gpc4* regulates these convergent extension (CE) movements by modulating the *Wnt11* pathway. (B) Normal left-right (LR) patterning in *Xenopus* requires both non-phosphorylated *Sdc2* on the left and *PKC γ* -dependent phosphorylated *Sdc2* on the right side of the embryo. *Sdc2* signals non-cell-autonomously from the ectoderm, possibly by activating or mediating *Vg1* signaling on the left side, which leads to asymmetric patterning of the mesoderm characterized by the expression of *Nodal*, *Lefty* and *Pitx2* on the left side.

olfactory bulb, the rostral migratory stream, the subgranular zone in the dentate gyrus of the hippocampus, or the subventricular zone (SVZ) lining the ventricles. Neurogenesis is regulated by several morphogens and growth factors, including *Fgf2*, which plays a dominant role (Guillemot and Zimmer, 2011) and requires HS for its activity. Accordingly, the proliferation of cultured cortical progenitors from *Nestin-cre; Ext1* cKO mice in response to *Fgf2* is dramatically reduced compared with that of progenitors from wild-type mice (Inatani et al., 2003). Consequently, these mutants show impaired neurogenesis leading to a smaller forebrain and a strongly reduced cortical thickness. Interestingly, a similar phenotype is observed in the absence of *Hs2st*, suggesting an important role for 2-O-sulfation in NSC proliferation (McLaughlin et al., 2003; Pratt et al., 2006).

Specific core proteins also play major roles in neurogenesis. The expression of perlecan is increased in response to *Fgf2* (Mashayekhi et al., 2011), and mice lacking perlecan (*Hspg2* – Mouse Genome Informatics) show reduced cell proliferation leading to microcephaly with thinner cerebral walls (Girós et al., 2007). Perlecan is also required for the maintenance of NSCs, as both the size of the NSC population and the extent of neurogenesis are strongly reduced in the SVZ of adult mice lacking perlecan (Kerever et al., 2014). Among membrane-associated HSPGs, *Gpc1* appears






A Neurogenesis**B Neuronal migration****C Axon elongation and growth cone guidance****D Synapse formation****E Neuromuscular junction****Fig. 5.** See next page for legend.

Fig. 5. Functions of HSPGs in nervous system development. (A) During neurogenesis, neuroepithelial cells (NECs) act as neural stem cells (NSCs) that proliferate and differentiate in the ventricular and subventricular zones (VZ and SVZ) of the cortex, giving rise to neuronal progenitors (NPs), neurons and radial glia. Several factors required for neurogenesis are regulated by HSPGs: the Fgf2 pathway is controlled by Gpc1 and Gpc4; Sdc1 modulates the canonical Wnt pathway. Perlecan also acts in the extracellular matrix (ECM) to regulate NEC proliferation. (B) During neuronal migration in the cortex, Sdc3 regulates cell migration along radial glia (blue arrows), from the intermediate zone (IZ) to the marginal zone (MZ), by mediating pleiotrophin and EGFR signaling. Sdc3 also acts as a receptor for GDNF and mediates the tangential migration of inhibitory neurons along a GDNF gradient present from the medial ganglionic eminence (MGE) towards the cortex (pink arrow). CP, cortical plate; LGE, lateral ganglionic eminence; SP, subplate. (C) During axon elongation, guidance cues present at the midline, such as Netrin, Slits or Shh, require HSPGs for signaling through their respective receptors DCC, Robo and Patched, both *in vitro* and *in vivo* in *Drosophila* and *C. elegans*. *In vivo* in vertebrates, Gpc1 mediates the response of axons to Shh: in pre-crossing axons, Gpc1 interacts with Shh bound to Patched to promote the transcription of specific genes, including that of the Shh receptor Hhip. In post-crossing axons, Gpc1 binds to Shh, triggering a repulsive response through Hhip signaling. (D) During synaptogenesis, pre-synaptic Gpc's, in particular Gpc4, interact with the pre-synaptic receptor PTP σ and the post-synaptic protein Lrrtm4 to promote excitatory synapse development. Gpc4 and Gpc6 are also released from astrocytes and promote synapse formation by clustering post-synaptic AMPA glutamate receptors (AMPARs). On the post-synaptic membrane, EphB2 phosphorylates Sdc2 and induces its clustering. The interaction of Sdc2 with adaptor proteins, such as Cask or neurofibromin, promotes the formation of dendritic spines. (E) At the neuromuscular junction, agrin is released from the nerve terminal and becomes stabilized in the basal lamina. It binds and activates the receptor Musk, leading to AChR clustering via the cytoplasmic adaptor protein Rapsn.

to contribute significantly to the regulation of neurogenesis. Mice lacking *Gpc1* exhibit a reduced brain size as a result of decreased proliferation (by 20%) from embryonic day (E) 8.5 to E9.5 (Jen et al., 2009). This modest but significant effect seems to be mediated by a reduction in Fgf17 signaling, and might be partly compensated by *Gpc4*. The analysis of compound mutants (*Gpc1* and a gene-trap allele of *Gpc4*) does indeed suggest that *Gpc1* and *Gpc4* might have overlapping functions in regulating brain size (Jen et al., 2009). *Gpc4* is strongly expressed in the ventricular zone of the telencephalon as well as in neurogenic regions in adult mice. Its ability to bind Fgf2 further suggests a possible role for *Gpc4* as a modulator of FGF signaling during neurogenesis (Hagihara et al., 2000). Additional studies will be required to test the contribution of *Gpc4*, as the gene-trap allele of *Gpc4* is thought not to be completely null. Finally, a recent study has highlighted a novel function for *Sdc1* in neurogenesis (Wang et al., 2012). *Sdc1* is highly expressed in progenitors in the ventricular zone and SVZ during development. Inhibiting its expression by electroporating a short hairpin RNA *in utero* reduces the proliferation of NPCs in response to Wnt and leads to premature differentiation. Interestingly, *Sdc1* modulates the canonical Wnt pathway but does not appear to regulate the FGF effector Erk1/2 (Mapk3/Mapk1) (Wang et al., 2012). This is in stark contrast to the mode of action employed by *Gpc1* (Jen et al., 2009), supporting the hypothesis that the diversity of HSPGs might fine-tune the regulation of the different pathways orchestrating nervous system development.

Neuronal migration

After neurogenesis and the establishment of neuronal identity, individual post-mitotic neurons migrate to their final functional destination. Several types of migration occurring in different waves have been described. Some involve the movement of neurons along

radial glial fibers, such as the radial migration of newborn neurons from the ventricular zone to form the different layers of the cortex (Nadarajah and Parnavelas, 2002). In others (for instance, the migration of neurons into the olfactory bulb), neurons do not use glia as a scaffold and instead migrate ‘in chain’ (Lois et al., 1996). HSPGs appear to be involved in regulating both of these types of migration. *Sdc3* in particular appears to play an essential role in neuronal migration in mammals (Hienola et al., 2006; Bespalov et al., 2011). Mice lacking *Sdc3* have a disorganized laminar structure of their cortex owing to impaired radial migration along glia. ‘In chain’ migration along the rostral migratory stream towards the olfactory bulb is also abnormal. Both defects seem to arise from a defective response of neural cells to the migration-promoting factors pleiotrophin and epidermal growth factor (EGF). Accordingly, it was shown that the *Sdc3* ectodomain binds to pleiotrophin and the epidermal growth factor receptor (EGFR), and that forebrain cells isolated from *Sdc3* KO mice show a reduced migration response to EGF and pleiotrophin *in vitro* (Hienola et al., 2006). Whether *Sdc3* requires its HS chains to mediate pleiotrophin and EGF signaling remains to be determined. More recently, *Sdc3* has been identified as a novel receptor for glial cell-derived neurotrophic factor (GDNF) (Bespalov et al., 2011). GDNF binds to *Sdc3* in a HS-dependent manner and requires *Sdc3* to promote the migration of cortical neurons *in vitro*. This observation translates *in vivo*: cortical GABAergic neurons known to migrate along a GDNF gradient from the medial ganglionic eminence towards the cortex (Pozas and Ibáñez, 2005) do not migrate properly in mice lacking *Sdc3* (Bespalov et al., 2011).

Surprisingly, although the inhibition of global HS synthesis in *C. elegans* by RNAi against *rib-1* or *rib-2* (homologs of the mammalian Ext genes) disrupts neuronal migration (Pedersen et al., 2013), no obvious defects in cortical lamination patterns or localization of specific neuronal populations [calbindin 2 (also known as calretinin)-positive and Pou3f1 (also known as Tst-1 and SCIP)-positive neurons] could be observed in *Nestin-cre; Ext1* cKO mice (Inatani et al., 2003). This suggests that the effects of HSPGs on neuronal migration may be species specific, or that compensatory mechanisms may exist in mice.

Axon growth and guidance

Concomitantly or subsequent to their migration, neurons extend axonal processes towards their future synaptic partners. During its navigation, the axon is guided by a plethora of attractive and repulsive guidance cues that act via direct contact or over a distance in the environment. A crucial role for HS during axon pathfinding has been known for a long time. Major guidance cues, such as Netrins, Slits, and Ephrins, have been shown to bind HS with a high affinity (Serafini et al., 1994; Bennett et al., 1997; Hussain et al., 2006; Irie et al., 2008). Furthermore, adding HS to the developing *Xenopus* retinotectal pathway or removing HS with heparitinase prevents retinal axons from entering their brain target, the tectum (Walz et al., 1997; Irie et al., 2002). Not surprisingly, *Nestin-cre; Ext1* cKO mice lacking HS in the nervous system show major axon guidance defects, including an absence of commissural tracts and errors from retinal axons (Inatani et al., 2003). Similarly, retinal axons erroneously project into the forebrain, the hindbrain and the opposite eye in *ext2; ext3* zebrafish mutants, which also exhibit decreased HS levels (Lee et al., 2004). Mis-sorting of retinal axons along the optic tract is also observed in these mutants and has revealed an important function of HS in developmental axon degeneration (Poulain and Chien, 2013). Indeed, the topographic organization of retinal axons along the tract is achieved through the

selective degeneration of mis-sorted axons in wild-type animals. This correction mechanism requires the presence of HS in the environment and is impaired in *ext2* mutants.

The first observation that different HS-modifying enzymes regulate distinct aspects of axon navigation came from elegant work in *C. elegans* that built the foundation for the hypothesis of a ‘sugar code’ (Bülow and Hobert, 2004; Bülow et al., 2008). *C. elegans* has single orthologs of the various HS-modifying enzymes, making genetic manipulation and analysis of the resulting phenotype easier: deacetylation and N-sulfation are catalyzed by HST-1, epimerization is mediated by HSE-5, and 2-O, 3-O and 6-O sulfations are performed by HST-2, HST-3.1 and HST-3.2, and HST-6. Interestingly, different classes of neurons require the activity of HSE-5, HST-2 and HST-6 in distinct combinations for their axons to be correctly guided. HST-3.1 and HST-3.2 regulate more refined steps of morphogenesis, controlling branching in a context-dependent manner (Tecle et al., 2013; Gysi et al., 2013). In mammals, however, the contribution of specific HS motifs to axon guidance is only beginning to emerge. Like *Nestin-cre; Ext1 cKO* mice, mouse mutants lacking *Ndst1* show absent or hypoplastic anterior and hippocampal commissures (Grobe et al., 2005). Supporting the sugar code hypothesis, mouse retinal axons lacking *Hs2st* or *Hs6st1* make distinct errors at the chiasm resulting from impaired responses to Slit1 and Slit2 (Pratt et al., 2006; Conway et al., 2011a; Conway et al., 2011b). Similarly, callosal axon navigation at the midline is differently impaired in the absence of *Hs2st* or *Hs6st1* (Conway et al., 2011a; Conway et al., 2011b; Clegg et al., 2014). Interestingly, *Fgf8* expression is increased and leads to a strong activation of ERK in the telencephalon of *Hs6st1* KO mice, but not in *Hs2st* mutants. In addition, suppressing components of the *Fgf8/Erk* pathway improves the phenotype of mice lacking *Hs6st1*, suggesting that *Hs2st* and *Hs6st1* might regulate callosal axon guidance in a molecularly distinct manner.

Although the phenotypes observed in the absence of HS strongly suggest a role for HSPGs in axon pathfinding, very little is known about the roles of the HSPG core proteins. Most studies describing functions for specific core proteins in axon guidance have been performed in *Drosophila* and *C. elegans*, in which one *Sdc* and two *Gpc* genes have been identified. For example, it was shown in both organisms that mutations in *Sdc* induce defects in midline axon guidance through impaired Slit/Robo signaling (Johnson et al., 2004; Steigemann et al., 2004; Rhiner et al., 2005; Smart et al., 2011). Both *Sdc* and the *Gpc* Dally-like are required for proper axon pathfinding and visual system function in *Drosophila* (Rawson et al., 2005). Finally, the *Gpc lon-2* regulates motor axon guidance in *C. elegans* (Bülow et al., 2008). In vertebrates, only one recent study has demonstrated a role for *Gpc1* in mediating the repulsive response of postcrossing axons to Shh at the floorplate in chick (Wilson and Stoeckli, 2013).

Dendritogenesis

As axon outgrowth proceeds, neurons develop branched dendritic arbors that are responsible for receiving, integrating and propagating signals received from other neurons. The resulting dendritic morphology is unique to each specific neuronal class and is thought to be determined by both environmental and cell-intrinsic factors. *Sdc2* has been shown to regulate dendrite formation in hippocampal neurons in culture by interacting with the multidomain adaptor molecule *Sarm1* and activating the *JNK/MKK4* pathway (Chen et al., 2011). It would be interesting to generate *Sdc2* KO mice and test whether *Sdc2* has a similar role *in vivo*. Very little is

known about the role of other HSPG core proteins or HS-synthesizing enzymes in dendritogenesis, and further studies are required to address their functions.

Synaptogenesis

Synapses are specialized adhesive junctions that enable the transmission of information between neurons (in the central nervous system) or between neurons and muscles (at neuromuscular junctions, NMJs). Neurotransmitters are released from presynaptic axonal termini into the synaptic cleft and bind to their receptors on the postsynaptic membranes of dendrites or muscle fibers. In the central nervous system, post-synaptic membranes are organized on dendrites into small protrusions of various shapes called spines. These spines evolve from actin-rich thin filopodial protrusions and develop into mushroom-like or stubby shapes during their formation and maturation.

The role of HSPGs in synaptogenesis was first demonstrated at the NMJ nearly 20 years ago. This early demonstration showed that the HSPG agrin, named after its ability to induce the aggregation of nicotinic acetylcholine receptors (AChRs) on developing skeletal muscle fibers in culture (Godfrey et al., 1984), is secreted from the terminal end of motor axons and initiates the formation of neuromuscular synapses by binding to and activating the receptor tyrosine kinase Musk on the muscle surface (Glass et al., 1996). Mice lacking agrin do not form NMJs and die *in utero* (Gautam et al., 1996). Similarly, a homozygous missense mutation in the agrin gene has been identified in a case of congenital myasthenic syndrome in humans, in which abnormal neuromuscular transmission leads to muscle weakness accentuated by exertion (Huzé et al., 2009). In addition to its high expression levels in motor neurons, agrin is detected in the central nervous system, and it was shown that blocking agrin function in cultured hippocampal neurons leads to the abnormal formation of synapses (Ferreira, 1999; Bose et al., 2000). Synapse loss has also recently been described in the cortex of transgenic adult mice lacking agrin expression everywhere but in motor neurons, confirming a role for agrin in the formation and/or maintenance of synapses in the brain (Ksiazek et al., 2007).

Studies of cultured hippocampal neurons have also revealed a role for *Sdc2* in the formation of dendritic spines. Such a role had been suggested by the high expression levels of *Sdc2* during synaptogenesis in the nervous system, and the identification of the synaptic protein Cask (calcium/CaM-dependent serine protein kinase) as an *Sdc2*-interacting partner (Cohen et al., 1998; Hsueh et al., 1998). Overexpressing *Sdc2* in rat hippocampal neurons accelerates spine formation, whereas inhibiting its expression prevents the development of dendritic protrusions (Ethell and Yamaguchi, 1999; Lin et al., 2007). The ability of *Sdc2* to promote filopodia formation and synaptogenesis requires different domains in its intracellular region (Fig. 1). The C1 domain interacts with neurofibromin, activates PKA and promotes ENA/VASP activity underlying filopodia formation (Lin et al., 2007). Phosphorylation of the C1 domain by the EphB2 receptor tyrosine kinase has also been shown to be required for *Sdc2* clustering on dendrites and the induction of dendritic spines (Ethell et al., 2001). The C2 region, which contains a type II PDZ-binding motif (residues EFYA) that mediates interactions with several adaptor proteins such as syntenin (*Sdcbp*), Cask, synbindin (*Trappc4*) and syndecan (*Gipc1*), is also required for *Sdc2* function in spines. However, the contribution of HS chains to *Sdc2* function, as well as the role of *Sdc2* *in vivo*, remains unknown.

In addition to *Sdc2*, some *Gpc*s have recently been implicated in the regulation of synaptogenesis. Two independent studies in

rodents and *in vitro* identified Gpc genes (most notably Gpc4) as pre-synaptic partners of the leucine-rich repeat transmembrane neuronal protein Lrrtm4, which is located at the post-synaptic membrane (de Wit et al., 2013; Siddiqui et al., 2013). This trans-synaptic interaction involves the HS-dependent binding of Gpc4 to the pre-synaptic receptor protein tyrosine phosphatase PTP σ (Ko et al., 2015) and is required for Lrrtm4 to promote excitatory synapse development. Additionally, Gpc4 and Gpc6 have been shown to be secreted from astrocytes and to promote excitatory synapse formation in retinal ganglion cells through clustering of glutamate receptors (Allen et al., 2012). Generating and analyzing mutants for the different Gpc genes will be of high interest to validate these exciting results *in vivo*.

Although the functions of the various core proteins in synaptogenesis are emerging, at least *in vitro*, the roles of HS chain modifications remain poorly characterized. HS is enriched in synapses of adult neurons, and cKO mice lacking *Ext1* in post-mitotic neurons exhibit altered glutamatergic synaptic transmission and develop numerous autism-like behavioral deficits (Irie et al., 2012). Mental impairment and autism have similarly been described in patients with deletion mutations in *EXT1* (Li et al., 2002), demonstrating a role for HS in synaptic and cognitive function. Interestingly, a recent screen performed in *Drosophila* showed different effects of *Sulf1* and *Hs6st* on synapse formation and function at the NMJ (Dani et al., 2012): mutants lacking *Sulf1* or *Hs6st* both have an increased number of synaptic boutons but show striking opposite differences in their neurotransmission strength. *Sulf1* and *Hs6st* thus appear to share a similar function in synapse morphogenesis, but play opposing roles in synaptic functional differentiation. In addition, the cell surface expression of the Gpc Dlp is reduced in synapses lacking *Hs6st*, whereas both Sdc and Dlp are increased in the absence of *Sulf1*. These changes correlate with differences in Wnt anterograde and BMP retrograde trans-synaptic signaling, both of which are responsible for the pre- and postsynaptic molecular differentiation defects observed in mutants. The absence of *Sulf1* in mammals also appears to disturb synaptogenesis (Kalus et al., 2009), as hippocampal neurons lacking *Sulf1* display a decreased density of dendritic spines and an altered synapse structure with a reduced number of synaptic vesicles. These defects lead to an impaired synaptic plasticity in adult *Sulf1* KO mice.

Future directions and implications for the sugar code hypothesis

HSPGs have been shown in cell culture studies to regulate a wide range of cell signaling pathways, cell-cell interactions, extracellular matrix formation, and cellular behaviors implicated in developmental mechanisms. Most recent studies have confirmed that HSPGs do indeed play major roles, many of them developmental, *in vivo*. However, there remain a number of major challenges in the field.

One key aim is to use genetics, transgenics and other functional assays to discern the roles of HSPG core proteins and the biosynthetic pathways that generate modified sugar chains. Although mutant analyses of some of the genes have yielded exciting results, most genes (including most of the multigene Hs3st family) have not yet been tested (Tables S1, S2). Genetic functional analysis is likely to be complicated by partially overlapping functional redundancies, for example within the same biosynthetic enzyme family, between some members of the same core protein family, and perhaps between different core families. Although it is clear that some core proteins have distinct

functions (Table S1), those functions could be independent of HS attachments to that core protein. Conversely, one could speculate that an Sdc core protein and Gpc core protein might be able to perform some of the same functions as long as they can acquire the same modified sugar chains (sugar code) and deliver that information to the cell surface in a particular cell lineage during development. Similarly, it might be possible for multiple members of the same biosynthetic gene family, for example the seven or eight members of the Hs3st family in vertebrates, to participate in generating the same sugar code in a particular cell lineage. Thus, it might be necessary to generate double mutants and beyond, as has been done for a few biosynthetic enzyme genes (Table S2). Despite these challenges, genetic analysis will be essential for testing the functions of HSPGs in development.

It should also be noted that the sugar code hypothesis is exciting but controversial. A number of key findings are in support of the hypothesis: distinct cells can generate unique compositions of HS; distinct HS structural motifs are associated with defined developmental events, especially in invertebrates; and some HS-protein interactions show specificity and selectivity. Of particular interest, it has been shown that 3-O-sulfation enzymes contribute to a very small level of HS modification and yet are represented by the largest gene family in the pathway (Thacker et al., 2014), suggesting that individual family members have branched out during evolution to perform slightly different functions. The notion of a family of enzymes having distinct functions in the generation of a sugar code is supported by biochemical evidence for differences in substrate preference within the same family of enzymes (Moon et al., 2012). However, a number of arguments against the sugar code have also been put forward. For example, the genetic absence of certain individual enzymes/core proteins does not trigger major defects, perhaps owing to functional redundancy or compensatory mechanisms, as seen in the comparison of single versus double mutants, or perhaps because the loss of one enzyme can also induce global changes in HS chain contents, which might provide some functional compensation. It has also been noted, *in vitro*, that some HS-protein interactions are relatively non-selective, as long as a specific type of sulfation (Ashikari-Hada et al., 2004) or an overall density of negative charges is provided by HS.

When considering genetic functional analyses of individual enzyme family members, it should also be noted that a phenotype could arise not because other enzymes in the same family cannot compensate for function, but because those enzymes are not expressed in the affected tissue. One of the great challenges is that, in contrast to RNA synthesis, HS synthesis is not template driven. Thus, if there is a sugar code, building it is likely to be dependent on the innate properties of the biosynthetic enzymes and their complexes, which are perhaps dependent on subtle substrate specificities as they decide where to add modifications to HS chains in the context of other modifications on that chain.

Finally, although powerful methods have been developed to study HS-protein interactions *in vitro* (Powell et al., 2004; Shipp and Hsieh-Wilson, 2007), with new technologies rapidly evolving, determining HS structure in biochemical and functional analyses in the context of *in vivo* development remains a real challenge. In analogy to advances in RNA expression profiling, one would ideally like to be able to analyze the ‘sequence’ of HS modifications, from one end of individual chains to the other, and to do so from a very heterogeneous mix of HS chains derived from single cells obtained from developing organisms. With these many challenges, there is clearly much work to be done to understand the roles of heparan sulfate proteoglycans in development.

Acknowledgements

We apologize to authors whose work was omitted from this article owing to space limitations.

Competing interests

The authors declare no competing or financial interests.

Funding

Work in the F.E.P. laboratory is supported in part by the National Institute of Neurological Disorders and Stroke at the National Institutes of Health [R00NS083714 to F.E.P.]. Work in the H.J.Y. laboratory is supported by the National Heart, Lung, and Blood Institute at the National Institutes of Health. Deposited in PMC for release after 12 months.

Supplementary information

Supplementary information available online at
<http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.098178/-/DC1>

References

- Allen, B. L. and Rapraeger, A. C.** (2003). Spatial and temporal expression of heparan sulfate in mouse development regulates FGF and FGF receptor assembly. *J. Cell Biol.* **163**, 637-648.
- Allen, N. J., Bennett, M. L., Foo, L. C., Wang, G. X., Chakraborty, C., Smith, S. J. and Barres, B. A.** (2012). Astrocyte glycans 4 and 6 promote formation of excitatory synapses via GluA1 AMPA receptors. *Nature* **486**, 410-414.
- Arrington, C. B. and Yost, H. J.** (2009). Extra-embryonic syndecan 2 regulates organ primordia migration and fibrillogenesis throughout the zebrafish embryo. *Development* **136**, 3143-3152.
- Arrington, C. B., Peterson, A. G. and Yost, H. J.** (2013). Sdc2 and Tbx16 regulate Fgf2-dependent epithelial cell morphogenesis in the ciliated organ of asymmetry. *Development* **140**, 4102-4109.
- Ashikari-Hada, S., Habuchi, H., Kariya, Y., Itoh, N., Reddi, A. H. and Kimata, K.** (2004). Characterization of growth factor-binding structures in heparin/heparan sulfate using an octasaccharide library. *J. Biol. Chem.* **279**, 12346-12354.
- Bennett, K. L., Bradshaw, J., Youngman, T., Rodgers, J., Greenfield, B., Aruffo, A. and Linsley, P. S.** (1997). Deleted in colorectal carcinoma (DCC) binds heparin via its fifth fibronectin type III domain. *J. Biol. Chem.* **272**, 26940-26946.
- Bespalov, M. M., Sidorova, Y. A., Tumova, S., Ahonen-Bishopp, A., Magalhães, A. C., Kulesskiy, E., Pavlevic, M., Rivera, C., Rauvala, H. and Saarma, M.** (2011). Heparan sulfate proteoglycan syndecan-3 is a novel receptor for GDNF, neurturin, and artemin. *J. Cell Biol.* **192**, 153-169.
- Bose, C. M., Qiu, D., Bergamaschi, A., Gravante, B., Bossi, M., Villa, A., Rupp, F. and Malgaroli, A.** (2000). Agrin controls synaptic differentiation in hippocampal neurons. *J. Neurosci.* **20**, 9086-9095.
- Bülow, H. E. and Hobert, O.** (2004). Differential sulfations and epimerization define heparan sulfate specificity in nervous system development. *Neuron* **41**, 723-736.
- Bülow, H. E., Tjoe, N., Townley, R. A., Didiano, D., van Kuppevelt, T. H. and Hobert, O.** (2008). Extracellular sugar modifications provide instructive and cell-specific information for axon-guidance choices. *Curr. Biol.* **18**, 1978-1985.
- Cadwalader, E. L., Condic, M. L. and Yost, H. J.** (2012). 2-O-sulfotransferase regulates Wnt signaling, cell adhesion and cell cycle during zebrafish epiboly. *Development* **139**, 1296-1305.
- Capurro, M. I., Xu, P., Shi, W., Li, F., Jia, A. and Filmus, J.** (2008). Glycan-3 inhibits Hedgehog signaling during development by competing with patched for Hedgehog binding. *Dev. Cell* **14**, 700-711.
- Chen, E., Hermanson, S. and Ekker, S. C.** (2004). Syndecan-2 is essential for angiogenic sprouting during zebrafish development. *Blood* **103**, 1710-1719.
- Chen, Y., Mohammadi, M. and Flanagan, J. G.** (2009). Graded levels of FGF protein span the midbrain and can instruct graded induction and repression of neural mapping labels. *Neuron* **62**, 773-780.
- Chen, C.-Y., Lin, C.-W., Chang, C.-Y., Jiang, S.-T. and Hsueh, Y.-P.** (2011). Sarm1, a negative regulator of innate immunity, interacts with syndecan-2 and regulates neuronal morphology. *J. Cell Biol.* **193**, 769-784.
- Clegg, J. M., Conway, C. D., Howe, K. M., Price, D. J., Mason, J. O., Turnbull, J. E., Basson, M. A. and Pratt, T.** (2014). Heparan sulfotransferases Hs6st1 and Hs2st keep Erk in check for mouse corpus callosum development. *J. Neurosci.* **34**, 2389-2401.
- Clément, A., Wiweger, M., von der Hardt, S., Rusch, M. A., Selleck, S. B., Chien, C.-B. and Roehl, H. H.** (2008). Regulation of zebrafish skeletogenesis by ext2/dackel and papst1/pinscher. *PLoS Genet.* **4**, e1000136.
- Cohen, A. R., Wood, D. F., Marfatia, S. M., Walther, Z., Chishti, A. H. and Anderson, J. M.** (1998). Human CASK/LIN-2 binds syndecan-2 and protein 4.1 and localizes to the basolateral membrane of epithelial cells. *J. Cell Biol.* **142**, 129-138.
- Conway, C. D., Howe, K. M., Nettleton, N. K., Price, D. J., Mason, J. O., Pratt, T.** (2011a). Heparan sulfate sugar modifications mediate the functions of slits and other factors needed for mouse forebrain commissure development. *J. Neurosci.* **31**, 1955-1970.
- Conway, C. D., Price, D. J., Pratt, T. and Mason, J. O.** (2011b). Analysis of axon guidance defects at the optic chiasm in heparan sulphate sulphotransferase compound mutant mice. *J. Anat.* **219**, 734-742.
- Costell, M., Carmona, R., Gustafsson, E., González-Iriarte, M., Fässler, R. and Muñoz-Chápuli, R.** (2002). Hyperplastic conotruncal endocardial cushions and transposition of great arteries in perlecan-null mice. *Circ. Res.* **91**, 158-164.
- Dani, N., Nahm, M., Lee, S. and Broadie, K.** (2012). A targeted glycan-related gene screen reveals heparan sulfate proteoglycan sulfation regulates WNT and BMP trans-synaptic signaling. *PLoS Genet.* **8**, e1003031.
- de Paz, J. L., Moseman, E. A., Noti, C., Polito, L., von Andrian, U. H. and Seeberger, P. H.** (2007). Profiling heparin-chemokine interactions using synthetic tools. *ACS Chem. Biol.* **2**, 735-744.
- de Wit, J., O'Sullivan, M. L., Savas, J. N., Condormitti, G., Caccese, M. C., Vennekens, K. M., Yates, J. R., III and Ghosh, A.** (2013). Unbiased discovery of glycan as a receptor for LRRTM4 in regulating excitatory synapse development. *Neuron* **79**, 696-711.
- Díaz-Balzac, C. A., Lázaro-Peña, M. I., Tecle, E., Gomez, N. and Bülow, H. E.** (2014). Complex cooperative functions of heparan sulfate proteoglycans shape nervous system development in *Caenorhabditis elegans*. *G3* **4**, 1859-1870.
- Escobedo, N., Contreras, O., Muñoz, R., Farías, M., Carrasco, H., Hill, C., Tran, U., Pryor, S. E., Wessely, O., Copp, A. J. et al.** (2013). Syndecan 4 interacts genetically with Vangl2 to regulate neural tube closure and planar cell polarity. *Development* **140**, 3008-3017.
- Ethell, I. M. and Yamaguchi, Y.** (1999). Cell surface heparan sulfate proteoglycan syndecan-2 induces the maturation of dendritic spines in rat hippocampal neurons. *J. Cell Biol.* **144**, 575-586.
- Ethell, I. M., Irie, F., Kalo, M. S., Couchman, J. R., Pasquale, E. B. and Yamaguchi, Y.** (2001). EphB/syndecan-2 signaling in dendritic spine morphogenesis. *Neuron* **31**, 1001-1013.
- Esko, J. D., Kimata, K. and Lindahl, U.** (2009). Proteoglycans and sulfated glycosaminoglycans. In *Essentials of Glycobiology*, 2nd edn. Cold Spring Harbor (New York, USA): Cold Spring Harbor Laboratory Press.
- Ferreira, A.** (1999). Abnormal synapse formation in agrin-depleted hippocampal neurons. *J. Cell Sci.* **112**, 4729-4738.
- Galli, A., Roure, A., Zeller, R. and Dono, R.** (2003). Glycan 4 modulates FGF signalling and regulates dorsoventral forebrain patterning in *Xenopus* embryos. *Development* **130**, 4919-4929.
- Gautam, M., Noakes, P. G., Moscoso, L., Rupp, F., Scheller, R. H., Merlie, J. P. and Sanes, J. R.** (1996). Defective neuromuscular synaptogenesis in agrin-deficient mutant mice. *Cell* **85**, 525-535.
- Girós, A., Morante, J., Gil-Sanz, C., Fairén, A. and Costell, M.** (2007). Perlecan controls neurogenesis in the developing telencephalon. *BMC Dev. Biol.* **7**, 29.
- Glass, D. J., Bowen, D. C., Stitt, T. N., Radziejewski, C., Bruno, J., Ryan, T. E., Gies, D. R., Shah, S., Mattsson, K., Burden, S. J. et al.** (1996). Agrin acts via a MuSK receptor complex. *Cell* **85**, 513-523.
- Godfrey, E. W., Nitkin, R. M., Wallace, B. G., Rubin, L. L. and McMahan, U. J.** (1984). Components of Torpedo electric organ and muscle that cause aggregation of acetylcholine receptors on cultured muscle cells. *J. Cell Biol.* **99**, 615-627.
- Grandel, H., Draper, B. W. and Schulte-Merker, S.** (2000). dackel acts in the ectoderm of the zebrafish pectoral fin bud to maintain AER signaling. *Development* **127**, 4169-4178.
- Grobe, K., Ledin, J., Ringvall, M., Holmborn, K., Forsberg, E., Esko, J. D. and Kjellén, L.** (2002). Heparan sulfate and development: differential roles of the N-acetylgalactosamine N-deacetylase/N-sulfotransferase isozymes. *Biochim. Biophys. Acta* **1573**, 209-215.
- Grobe, K., Inatani, M., Pallerla, S. R., Castagnola, J., Yamaguchi, Y. and Esko, J. D.** (2005). Cerebral hypoplasia and craniofacial defects in mice lacking heparan sulfate Ndst1 gene function. *Development* **132**, 3777-3786.
- Guillemot, F. and Zimmer, C.** (2011). From cradle to grave: the multiple roles of fibroblast growth factors in neural development. *Neuron* **71**, 574-588.
- Gysi, S., Rhiner, C., Flibotte, S., Moerman, D. G. and Hengartner, M. O.** (2013). A network of HSPG core proteins and HS modifying enzymes regulates netrin-dependent guidance of D-type motor neurons in *Caenorhabditis elegans*. *PLoS ONE* **8**, e74908.
- Habuchi, H., Nagai, N., Sugaya, N., Atsumi, F., Stevens, R. L. and Kimata, K.** (2007). Mice deficient in heparan sulfate 6-O-sulfotransferase-1 exhibit defective heparan sulfate biosynthesis, abnormal placentation, and late embryonic lethality. *J. Biol. Chem.* **282**, 15578-15588.
- Hagihara, K., Watanabe, K., Chun, J. and Yamaguchi, Y.** (2000). Glycan-4 is an FGF2-binding heparan sulfate proteoglycan expressed in neural precursor cells. *Dev. Dyn.* **219**, 353-367.
- HajMohammadi, S., Enyoyi, K., Princivalle, M., Christi, P., Lech, M., Beeler, D., Rayburn, H., Schwartz, J. J., Barzegar, S., de Agostini, A. I. et al.** (2003). Normal levels of anticoagulant heparan sulfate are not essential for normal hemostasis. *J. Clin. Invest.* **111**, 989-999.
- Hienola, A., Tumova, S., Kulesskiy, E. and Rauvala, H.** (2006). N-syndecan deficiency impairs neural migration in brain. *J. Cell Biol.* **174**, 569-580.
- Holmborn, K., Habicher, J., Kasza, Z., Eriksson, A. S., Filipek-Gorniak, B., Gopal, S., Couchman, J. R., Ahlberg, P. E., Wiweger, M., Spillmann, D.**

- et al.** (2012). On the roles and regulation of chondroitin sulfate and heparan sulfate in zebrafish pharyngeal cartilage morphogenesis. *J. Biol. Chem.* **287**, 33905-33916.
- Holt, C. E. and Dickson, B. J. (2005). Sugar codes for axons? *Neuron* **46**, 169-172.
- Hsueh, Y.-P., Yang, F.-C., Kharazia, V., Naisbitt, S., Cohen, A. R., Weinberg, R. J. and Sheng, M. (1998). Direct interaction of CASK/LIN-2 and syndecan heparan sulfate proteoglycan and their overlapping distribution in neuronal synapses. *J. Cell Biol.* **142**, 139-151.
- Huegel, J., Sgariglia, F., Enomoto-Iwamoto, M., Koyama, E., Dormans, J. P. and Pacifici, M. (2013). Heparan sulfate in skeletal development, growth, and pathology: the case of hereditary multiple exostoses. *Dev. Dyn.* **242**, 1021-1032.
- Hussain, S.-A., Piper, M., Fukuhara, N., Strochlic, L., Cho, G., Howitt, J. A., Ahmed, Y., Powell, A. K., Turnbull, J. E., Holt, C. E. et al. (2006). A molecular mechanism for the heparan sulfate dependence of slit-robo signaling. *J. Biol. Chem.* **281**, 39693-39698.
- Huzé, C., Bauché, S., Richard, P., Chevessier, F., Goillot, E., Gaudon, K., Ben Ammar, A., Chaboud, A., Grosjean, I., Lecuyer, H.-A. et al. (2009). Identification of an agrin mutation that causes congenital myasthenia and affects synapse function. *Am. J. Hum. Genet.* **85**, 155-167.
- Inatani, M., Irie, F., Plump, A. S., Tessier-Lavigne, M. and Yamaguchi, Y. (2003). Mammalian brain morphogenesis and midline axon guidance require heparan sulfate. *Science* **302**, 1044-1046.
- Irie, A., Yates, E. A., Turnbull, J. E. and Holt, C. E. (2002). Specific heparan sulfate structures involved in retinal axon targeting. *Development* **129**, 61-70.
- Irie, F., Okuno, M., Matsumoto, K., Pasquale, E. B. and Yamaguchi, Y. (2008). Heparan sulfate regulates ephrin-A3/EphA receptor signaling. *Proc. Natl. Acad. Sci. USA* **105**, 12307-12312.
- Irie, F., Badie-Mahdavi, H. and Yamaguchi, Y. (2012). Autism-like socio-communicative deficits and stereotypies in mice lacking heparan sulfate. *Proc. Natl. Acad. Sci. USA* **109**, 5052-5056.
- Izvolsky, K. I., Lu, J., Martin, G., Albrecht, K. H. and Cardoso, W. V. (2008). Systemic inactivation of Hs6st1 in mice is associated with late postnatal mortality without major defects in organogenesis. *Genesis* **46**, 8-18.
- Jen, Y.-H., Musacchio, M. and Lander, A. D. (2009). Glycan-1 controls brain size through regulation of fibroblast growth factor signaling in early neurogenesis. *Neural Dev.* **4**, 33.
- Johnson, K. G., Ghose, A., Epstein, E., Lincecum, J., O'Connor, M. B. and Van Vactor, D. (2004). Axonal heparan sulfate proteoglycans regulate the distribution and efficiency of the repellent slit during midline axon guidance. *Curr. Biol.* **14**, 499-504.
- Kalus, I., Salmen, B., Viebahn, C., von Figura, K., Schmitz, D., D'Hooge, R. and Dierks, T. (2009). Differential involvement of the extracellular 6-O-endosulfatases Sulf1 and Sulf2 in brain development and neuronal and behavioural plasticity. *J. Cell. Mol. Med.* **13**, 4505-4521.
- Kerever, A., Mercier, F., Nonaka, R., de Vega, S., Oda, Y., Zalc, B., Okada, Y., Hattori, N., Yamada, Y. and Arikawa-Hirasawa, E. (2014). Perlecan is required for FGF-2 signaling in the neural stem cell niche. *Stem Cell Res.* **12**, 492-505.
- Ko, J. S., Pramanik, G., Um, J. W., Shim, J. S., Lee, D., Kim, K. H., Chung, G.-Y., Condorimitti, G., Kim, H. M., Kim, H. et al. (2015). PTP γ functions as a presynaptic receptor for the glycan-4/LRRTM4 complex and is essential for excitatory synaptic transmission. *Proc. Natl. Acad. Sci. USA* **112**, 1874-1879.
- Kok, F. O., Shin, M., Ni, C.-W., Gupta, A., Grossé, A. S., van Impel, A., Kirchmaier, B. C., Peterson-Maduro, J., Kourkoulis, G., Male, I. et al. (2015). Reverse genetic screening reveals poor correlation between morpholino-induced and mutant phenotypes in zebrafish. *Dev. Cell* **32**, 97-108.
- Kramer, K. L. and Yost, H. J. (2002). Ectodermal syndecan-2 mediates left-right axis formation in migrating mesoderm as a cell-nonautonomous Vg1 cofactor. *Dev. Cell* **2**, 115-124.
- Kramer, K. L., Barnette, J. E. and Yost, H. J. (2002). PKC γ regulates syndecan-2 inside-out signaling during *Xenopus* left-right development. *Cell* **111**, 981-990.
- Ksiazek, I., Burkhardt, C., Lin, S., Seddik, R., Maj, M., Bezakova, G., Jucker, M., Arber, S., Caroni, P., Sanes, J. R. et al. (2007). Synapse loss in cortex of agrin-deficient mice after genetic rescue of perinatal death. *J. Neurosci.* **27**, 7183-7195.
- Kuriyama, S. and Mayor, R. (2009). A role for Syndecan-4 in neural induction involving ERK- and PKC-dependent pathways. *Development* **136**, 575-584.
- Lawrence, R., Lu, H., Rosenberg, R. D., Esko, J. D., and Zhang, L. (2008). Disaccharide structure code for the easy representation of constituent oligosaccharides from glycosaminoglycans. *Nat. Methods* **5**, 291-292.
- Ledin, J., Staatz, W., Li, J.-P., Gotte, M., Selleck, S., Kjellen, L. and Spillmann, D. (2004). Heparan sulfate structure in mice with genetically modified heparan sulfate production. *J. Biol. Chem.* **279**, 42732-42741.
- Lee, J.-S., von der Hardt, S., Rusch, M. A., Stringer, S. E., Stickney, H. L., Talbot, W. S., Geisler, R., Nüsslein-Volhard, C., Selleck, S. B., Chien, C.-B. et al. (2004). Axon sorting in the optic tract requires HSPG synthesis by ext2 (dackel) and ext3 (boxer). *Neuron* **44**, 947-960.
- Li, H., Yamagata, T., Mori, M. and Momoi, M. Y. (2002). Association of autism in two patients with hereditary multiple exostoses caused by novel deletion mutations of EXT1. *J. Hum. Genet.* **47**, 262-265.
- Lin, X., Wei, G., Shi, Z., Dryer, L., Esko, J. D., Wells, D. E. and Matzuk, M. M. (2000). Disruption of gastrulation and heparan sulfate biosynthesis in EXT1-deficient mice. *Dev. Biol.* **224**, 299-311.
- Lin, Y.-L., Lei, Y.-T., Hong, C.-J. and Hsueh, Y.-P. (2007). Syndecan-2 induces filopodia and dendritic spine formation via the neurofibromin-PKA-Ena/VASP pathway. *J. Cell Biol.* **177**, 829-841.
- Lindahl, U. and Li, J.-P. (2009). Interactions between heparan sulfate and proteins – design and functional implications. *Int. Rev. Cell Mol. Biol.* **276**, 105-159.
- Lois, C., García-Verdugo, J. M. and Alvarez-Buylla, A. (1996). Chain migration of neuronal precursors. *Science* **271**, 978-981.
- Mashayekhi, F., Sadeghi, M. and Rajaei, F. (2011). Induction of perlecan expression and neural cell proliferation by FGF-2 in the developing cerebral cortex: an *in vivo* study. *J. Mol. Neurosci.* **45**, 87-93.
- McLaughlin, D., Karlsson, F., Tian, N., Pratt, T., Bullock, S. L., Wilson, V. A., Price, D. J. and Mason, J. O. (2003). Specific modification of heparan sulphate is required for normal cerebral cortical development. *Mech. Dev.* **120**, 1481-1488.
- Meyers, E. N., Lewandoski, M. and Martin, G. R. (1998). An Fgf8 mutant allelic series generated by Cre- and Flp-mediated recombination. *Nat. Genet.* **18**, 136-141.
- Mitchell, K. J., Pinson, K. I., Kelly, O. G., Brennan, J., Zupich, J., Scherz, P., Leighton, P. A., Goodrich, L. V., Lu, X., Avery, B. J. et al. (2001). Functional analysis of secreted and transmembrane proteins critical to mouse development. *Nat. Genet.* **28**, 241-249.
- Moon, A. F., Xu, Y., Woody, S. M., Krahm, J. M., Lindardt, R. J., Liu, J. and Pedersen, L. C. (2012). Dissecting the substrate recognition of 3-O-sulfotransferase for the biosynthesis of anticoagulant heparin. *Proc. Natl. Acad. Sci.* **109**, 5265-5270.
- Muñoz, R. and Larraín, J. (2006). xSyndecan-4 regulates gastrulation and neural tube closure in *Xenopus* embryos. *ScientificWorldJournal* **6**, 1298-1301.
- Nadarajah, B. and Parnavelas, J. G. (2002). Modes of neuronal migration in the developing cerebral cortex. *Nat. Rev. Neurosci.* **3**, 423-432.
- Neugebauer, J. M., Cadwallader, A. B., Amack, J. D., Bisgrove, B. W. and Yost, H. J. (2013). Differential roles for 3-O-STS in the regulation of cilia length and motility. *Development* **140**, 3892-3902.
- Ng, A., Wong, M., Viviano, B., Erlich, J. M., Alba, G., Pflederer, C., Jay, P. Y. and Saunders, S. (2009). Loss of glycan-3 function causes growth factor-dependent defects in cardiac and coronary vascular development. *Dev. Biol.* **335**, 208-215.
- Nigam, S. K. and Bush, K. T. (2014). Growth factor-heparan sulfate "switches" regulating stages of branching morphogenesis. *Pediatr. Nephrol.* **29**, 727-735.
- Norton, W. H. J., Ledin, J., Grandel, H. and Neumann, C. J. (2005). HSPG synthesis by zebrafish Ext2 and Ext3 is required for Fgf10 signalling during limb development. *Development* **132**, 4963-4973.
- Ohkawara, B., Yamamoto, T. S., Tada, M. and Ueno, N. (2003). Role of glycan-4 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* **130**, 2129-2138.
- Pan, Y., Carbe, C., Kupich, S., Pickhine, U., Ohlig, S., Frye, M., Seeliger, R., Pallerla, S. R., Moon, A. M., Lawrence, R. et al. (2014). Heparan sulfate expression in the neural crest is essential for mouse cardiogenesis. *Matrix Biol.* **35**, 253-265.
- Pedersen, M. E., Snieckute, G., Kagias, K., Nehammer, C., Multhaupt, H. A. B., Couchman, J. R. and Pocock, R. (2013). An epidermal microRNA regulates neuronal migration through control of the cellular glycosylation state. *Science* **341**, 1404-1408.
- Poulain, F. E. and Chien, C.-B. (2013). Proteoglycan-mediated axon degeneration corrects pretarget topographic sorting errors. *Neuron* **78**, 49-56.
- Powell, A. K., Yates, E. A., Fernig, D. G. and Turnbull, J. E. (2004). Interactions of heparan/heparan sulfate with proteins: appraisal of structural factors and experimental approaches. *Glycobiology* **14**, 17R-30R.
- Pozas, E. and Ibáñez, C. F. (2005). GDNF and GFRalpha1 promote differentiation and tangential migration of cortical GABAergic neurons. *Neuron* **45**, 701-713.
- Pratt, T., Conway, C. D., Tian, N. M. M.-L., Price, D. J. and Mason, J. O. (2006). Heparan sulphation patterns generated by specific heparan sulfotransferase enzymes direct distinct aspects of retinal axon guidance at the optic chiasm. *J. Neurosci.* **26**, 6911-6923.
- Rawson, J. M., Dimitroff, B., Johnson, K. G., Ge, X., Van Vactor, D. and Selleck, S. B. (2005). The heparan sulfate proteoglycans Dally-like and Syndecan have distinct functions in axon guidance and visual-system assembly in *Drosophila*. *Curr. Biol.* **15**, 833-838.
- Rhiner, C., Gysi, S., Frohli, E., Hengartner, M. O. and Hajnal, A. (2005). Syndecan regulates cell migration and axon guidance in *C. elegans*. *Development* **132**, 4621-4633.
- Samson, S. C., Ferrer, T., Jou, C. J., Sachse, F. B., Shankaran, S. S., Shaw, R. M., Chi, N. C., Tristani-Firouzi, M. and Yost, H. J. (2013). 3-OST-7 regulates BMP-dependent cardiac contraction. *PLoS Biol.* **11**, e1001727.
- Serafini, T., Kennedy, T. E., Gaiko, M. J., Mirzayan, C., Jessell, T. M. and Tessier-Lavigne, M. (1994). The netrins define a family of axon outgrowth-promoting proteins homologous to *C. elegans* UNC-6. *Cell* **78**, 409-424.
- Shimokawa, K., Kimura-Yoshida, C., Nagai, N., Mukai, K., Matsubara, K., Watanabe, H., Matsuda, Y., Mochida, K. and Matsuo, I. (2011). Cell surface

- heparan sulfate chains regulate local reception of FGF signaling in the mouse embryo. *Dev. Cell* **21**, 257-272.
- Shipp, E. L. and Hsieh-Wilson, L. C.** (2007). Profiling the sulfation specificities of glycosaminoglycan interactions with growth factors and chemotactic proteins using microarrays. *Chem. Biol.* **14**, 195-208.
- Shworak, N. W., HajMohammadi, S., de Agostini, A. I. and Rosenberg, R. D.** (2002). Mice deficient in heparan sulfate 3-O-sulfotransferase-1: normal hemostasis with unexpected perinatal phenotypes. *Glycoconj. J.* **19**, 355-361.
- Siddiqui, T. J., Tari, P. K., Connor, S. A., Zhang, P., Dobie, F. A., She, K., Kawabe, H., Wang, Y. T., Brose, N. and Craig, A. M.** (2013). An LRRTM4-HSPG complex mediates excitatory synapse development on dentate gyrus granule cells. *Neuron* **79**, 680-695.
- Smart, A. D., Course, M. M., Rawson, J., Selleck, S., Van Vactor, D. and Johnson, K. G.** (2011). Heparan sulfate proteoglycan specificity during axon pathway formation in the Drosophila embryo. *Dev. Neurobiol.* **71**, 608-618.
- Stainier, D. Y. R., Kontarakis, Z. and Rossi, A.** (2015). Making sense of anti-sense data. *Dev. Cell* **32**, 7-8.
- Steigemann, P., Molitor, A., Fellert, S., Jäckle, H. and Vorbrüggen, G.** (2004). Heparan sulfate proteoglycan syndecan promotes axonal and myotube guidance by slit/robo signaling. *Curr. Biol.* **14**, 225-230.
- Stickens, D., Zak, B. M., Rougier, N., Esko, J. D. and Werb, Z.** (2005). Mice deficient in Ext2 lack heparan sulfate and develop exostoses. *Development* **132**, 5055-5068.
- Tecle, E., Diaz-Balzac, C. A. and Bülow, H. E.** (2013). Distinct 3-O-sulfated heparan sulfate modification patterns are required for kal-1-dependent neurite branching in a context-dependent manner in *Caenorhabditis elegans*. *G3* **3**, 541-552.
- Thacker, B. E., Xu, D., Lawrence, R. and Esko, J. D.** (2014). Heparan sulfate 3-O-sulfation: a rare modification in search of a function. *Matrix Biol.* **35**, 60-72.
- Thomas, K. R., Musci, T. S., Neumann, P. E. and Capecchi, M. R.** (1991). Swaying is a mutant allele of the proto-oncogene *Wnt-1*. *Cell* **67**, 969-976.
- Thompson, S. M., Jesudason, E. C., Turnbull, J. E. and Fernig, D. G.** (2010). Heparan sulfate in lung morphogenesis: the elephant in the room. *Birth Defects Res. C Embryo. Today* **90**, 32-44.
- Thunberg, L., Bäckström, G. and Lindahl, U.** (1982). Further characterization of the antithrombin-binding sequence in heparin. *Carbohydr. Res.* **100**, 393-410.
- Topczewski, J., Sepich, D. S., Myers, D. C., Walker, C., Amores, A., Lele, Z., Hammerschmidt, M., Postlethwait, J. and Solnica-Krezel, L.** (2001). The zebrafish glycan knypek controls cell polarity during gastrulation movements of convergent extension. *Dev. Cell* **1**, 251-264.
- Walz, A., McFarlane, S., Brickman, Y. G., Nurcombe, V., Bartlett, P. F. and Holt, C. E.** (1997). Essential role of heparan sulfates in axon navigation and targeting in the developing visual system. *Development* **124**, 2421-2430.
- Wang, Q., Yang, L., Alexander, C. and Temple, S.** (2012). The niche factor syndecan-1 regulates the maintenance and proliferation of neural progenitor cells during mammalian cortical development. *PLoS ONE* **7**, e42883.
- Wilson, N. H. and Stoeckli, E. T.** (2013). Sonic hedgehog regulates its own receptor on postcrossing commissural axons in a glycan1-dependent manner. *Neuron* **79**, 478-491.
- Xu, D. and Esko, J. D.** (2014). Demystifying heparan sulfate–protein interactions. *Annu. Rev. Biochem.* **83**, 129-157.

Tables S1 and S2

[Click here to Download Table S1 and S2](#)

References of tables S1 and S2

- Abramsson, A., Kurup, S., Busse, M., Yamada, S., Lindblom, P., Schallmeiner, E., Stenzel, D., Sauvaget, D., Ledin, J., Ringvall, M. et al.** (2007). Defective N-sulfation of heparan sulfate proteoglycans limits PDGF-BB binding and pericyte recruitment in vascular development. *Genes Dev.* **21**, 316-331.
- Adhikari, N., Basi, D.L., Townsend, D., Rusch, M., Mariash, A., Mullegama, S., Watson, A., Larson, J., Tan, S., Lerman, B. et al.** (2010). Heparan sulfate Ndst1 regulates vascular smooth muscle cell proliferation, vessel size and vascular remodeling. *J. Mol. Cell. Cardiol.* **49**, 287-293.
- Adhikari, N., Billaud, M., Carlson, M., Lake, S.P., Montaniel, K.R., Staggs, R., Guan, W., Walek, D., Desir, S. et al.** (2014). Vascular biomechanical properties in mice with smooth muscle specific deletion of Ndst1. *Mol. Cell. Biochem.* **385**, 225-238.
- Ahn, J., Lüdecke, H.J., Lindow, S., Horton, W.A., Lee, B., Wagner, M.J., Horsthemke, and B., Wells, D.E.** (1995). Cloning of the putative tumour suppressor gene for hereditary multiple exostoses (EXT1). *Nat. Genet.* **11**, 137-143.
- Ai, X., Kitazawa, T., Do, A.T., Kusche-Gullberg, M., Labosky, P.A., and Emerson, C.P. Jr.** (2007). SULF1 and SULF2 regulate heparan sulfate-mediated GDNF signaling for esophageal innervation. *Development* **134**, 3327-3338.
- Aikawa, T., Whipple, C.A., Lopez, M.E., Gunn, J., Young, A., Lander, A.D., and Korc, M.** (2008). Glypican-1 modulates the angiogenic and metastatic potential of human and mouse cancer cells. *J. Clin. Invest.* **118**, 89-99.

- Aikio, M., Hurskainen, M., Brideau, G., Hägg, P., Sormunen, R., Heljasvaara, R., Gould, D.B., and Pihlajaniemi, T.** (2013). Collagen XVIII short isoform is critical for retinal vascularization, and overexpression of the Tsp-1 domain affects eye growth and cataract formation. *Invest. Ophthalmol. Vis. Sci.* **54**, 7450-7462.
- Aikio, M., Elamaa, H., Vicente, D., Izzi, V., Kaur, I., Seppinen, L., Speedy, H.E., Kaminska, D., Kuusisto, S., Sormunen, R. et al.** (2014). Specific collagen XVIII isoforms promote adipose tissue accrual via mechanisms determining adipocyte number and affect fat deposition. *Proc. Natl. Acad. Sci. USA* **111**, E3043-3052.
- Alexander, C.M., Reichsman, F., Hinkes, M.T., Lincecum, J., Becker, K.A., Cumberledge, S., and Bernfield, M.** (2000). Syndecan-1 is required for Wnt-1-induced mammary tumorigenesis in mice. *Nat. Genet.* **25**, 329-332.
- Allen, N.J., Bennett, M.L., Foo, L.C., Wang, G.X., Chakraborty, C., Smith, S.J., and Barres, B.A.** (2012). Astrocyte glycans 4 and 6 promote formation of excitatory synapses via GluA1 AMPA receptors. *Nature* **486**, 410-414.
- Arikawa-Hirasawa, E., Watanabe, H., Takami, H., Hassell, J.R., and Yamada, Y.** (1999). Perlecan is essential for cartilage and cephalic development. *Nat. Genet.* **23**, 354-358.
- Arikawa-Hirasawa, E., Wilcox, W.R., Le, A.H., Silverman, N., Govindraj, P., Hassell, J.R., and Yamada, Y.** (2001). Dyssegmental dysplasia, Silverman-Handmaker type, is caused by functional null mutations of the perlecan gene. *Nat. Genet.* **27**, 431-434.
- Arikawa-Hirasawa, E., Le, A.H., Nishino, I., Nonaka, I., Ho, N.C., Francomano, C.A., Govindraj, P., Hassell, J.R., Devaney, J.M., Spranger, J. et al.** (2002a). Structural and functional mutations of the perlecan gene cause Schwartz-Jampel syndrome, with myotonic myopathy and chondrodysplasia. *Am. J. Hum. Genet.* **70**, 1368-1375.
- Arikawa-Hirasawa, E., Rossi, S.G., Rotundo, R.L., and Yamada, Y.** (2002b). Absence of acetylcholinesterase at the neuromuscular junctions of perlecan-null mice. *Nat. Neurosci.* **5**, 119-123.
- Arking, D.E., Reinier, K., Post, W., Jui, J., Hilton, G., O'Connor, A., Prineas, R.J., Boerwinkle, E., Psaty, B.M., Tomaselli, G.F. et al.** (2010). Genome-wide association study identifies GPC5 as a novel genetic locus protective against sudden cardiac arrest. *PLoS One* **5**, e9879.

- Arrington, C.B. and Yost, H.J.** (2009). Extra-embryonic syndecan 2 regulates organ primordia migration and fibrillogenesis throughout the zebrafish embryo. *Development* **136**, 3143-3152.
- Arrington, C.B., Peterson, A.G., and Yost, H.J.** (2013). Sdc2 and Tbx16 regulate Fgf2-dependent epithelial cell morphogenesis in the ciliated organ of asymmetry. *Development* **140**, 4102-4109.
- Astudillo, P., Carrasco, H., and Larraín, J.** (2014). Syndecan-4 inhibits Wnt/β-catenin signaling through regulation of low-density-lipoprotein receptor-related protein (LRP6) and R-spondin 3. *Int. J. Biochem. Cell Biol.* **46**, 103-112.
- Axelsson, J., Xu, D., Kang, B.N., Nussbacher, J.K., Handel, T.M., Ley, K., Sriramaraao, P., and Esko, J.D.** (2012). Inactivation of heparan sulfate 2-O-sulfotransferase accentuates neutrophil infiltration during acute inflammation in mice. *Blood* **120**, 1742-1751.
- Baker, A.B., Groothuis, A., Jonas, M., Ettenson, D.S., Shazly, T., Zcharia, E., Vlodavsky, I., Seifert, P., and Edelman, E.R.** (2009). Heparanase alters arterial structure, mechanics, and repair following endovascular stenting in mice. *Circ. Res.* **104**, 380-387.
- Banerjee, S., Isaacman-Beck, J., Schneider, V.A., and Granato, M.** (2013). A novel role for Lh3 dependent ECM modifications during neural crest cell migration in zebrafish. *PLoS One* **8**, e54609.
- Baranzini, S.E., Wang, J., Gibson, R.A., Galwey, N., Naegelin, Y., Barkhof, F., Radue, E.W., Lindberg, R.L., Uitdehaag, B.M., Johnson, M.R. et al.** (2009). Genome-wide association analysis of susceptibility and clinical phenotype in multiple sclerosis. *Hum. Mol. Genet.* **18**, 767-778.
- Bertolesi, G.E., Michaiel, G., and McFarlane, S.** (2008). Two heparanase splicing variants with distinct properties are necessary in early Xenopus development. *J. Biol. Chem.* **283**, 16004-16016.
- Bespakov, M.M., Sidorova, Y.A., Tumova, S., Ahonen-Bishopp, A., Magalhães, A.C., Kulesskiy, E., Paveliev, M., Rivera, C., Rauvala, H., and Saarma, M.** (2011). Heparan sulfate proteoglycan syndecan-3 is a novel receptor for GDNF, neurturin, and artemin. *J. Cell Biol.* **192**, 153-169.
- Bink, R.J., Habuchi, H., Lele, Z., Dolk, E., Joore, J., Rauch, G.J., Geisler, R., Wilson, S.W., den Hertog, J., Kimata, K., and Zivkovic, D.** (2003). Heparan sulfate 6-o-sulfotransferase is essential for muscle development in zebrafish. *J. Biol. Chem.* **278**, 31118-31127.

- Bishop, J.R., Passos-Bueno, M.R., Fong, L., Stanford, K.I., Gonzales, J.C., Yeh, E., Young, S.G., Bensadoun, A., Witztum, J.L., Esko, J.D., and Moulton, K.S.** (2010). Deletion of the basement membrane heparan sulfate proteoglycan type XVIII collagen causes hypertriglyceridemia in mice and humans. *PLoS One* **5**, e13919.
- Bode, L., Salvestrini, C., Park, P.W., Li, J.P., Esko, J.D., Yamaguchi, Y., Murch, S., and Freeze, H.H.** (2008). Heparan sulfate and syndecan-1 are essential in maintaining murine and human intestinal epithelial barrier function. *J. Clin. Invest.* **118**, 229-238.
- Bogdanik, L.P. and Burgess, R.W.** (2011). A valid mouse model of AGRIN-associated congenital myasthenic syndrome. *Hum. Mol. Genet.* **20**, 4617-4633.
- Bullock, S.L., Fletcher, J.M., Beddington, R.S., and Wilson, V.A.** (1998). Renal agenesis in mice homozygous for a gene trap mutation in the gene encoding heparan sulfate 2-sulfotransferase. *Genes Dev.* **12**, 1894-1906.
- Burgess, R.W., Nguyen, Q.T., Son, Y.J., Lichtman, J.W., and Sanes, J.R.** (1999). Alternatively spliced isoforms of nerve- and muscle-derived agrin: their roles at the neuromuscular junction. *Neuron* **23**, 33-44.
- Bush, K.T., Crawford, B.E., Garner, O.B., Nigam, K.B., Esko, J.D., and Nigam, S.K.** (2012). N-sulfation of heparan sulfate regulates early branching events in the developing mammary gland. *J. Biol. Chem.* **287**, 42064-42070.
- Cadwalader, E.L., Condic, M.L., and Yost, H.J.** (2012). 2-O-sulfotransferase regulates Wnt signaling, cell adhesion and cell cycle during zebrafish epiboly. *Development* **139**, 1296-1305.
- Cai, Z., Grobe, K., and Zhang, X.** (2014). Role of heparan sulfate proteoglycans in optic disc and stalk morphogenesis. *Dev. Dyn.* **243**, 1310-1316.
- Campos-Xavier, A.B., Martinet, D., Bateman, J., Belluocchio, D., Rowley, L., Tan, T.Y., Baxová, A., Gustavson, K.H., Borochowitz, Z.U., Innes, A.M. et al.** (2009). Mutations in the heparan-sulfate proteoglycan glypcan 6 (GPC6) impair endochondral ossification and cause recessive omodysplasia. *Am. J. Hum. Genet.* **84**, 760-770.
- Cano-Gauci, D.F., Song, H.H., Yang, H., McKerlie, C., Choo, B., Shi, W., Pullano, R., Piscione, T.D., Grisaru, S., Soon, S. et al.** (1999). Glypican-3-deficient mice exhibit developmental

overgrowth and some of the abnormalities typical of Simpson-Golabi-Behmel syndrome. *J. Cell. Biol.* **146**, 255-264.

Capurro, M.I., Xu, P., Shi, W., Li, F., Jia, A., and Filmus, J. (2008). Glypican-3 inhibits Hedgehog signaling during development by competing with patched for Hedgehog binding. *Dev. Cell* **14**, 700-711.

Capurro, M.I., Li, F., and Filmus, J. (2009). Overgrowth of a mouse model of Simpson-Golabi-Behmel syndrome is partly mediated by Indian hedgehog. *EMBO Rep.* **10**, 901-907.

Cénit, M.D., Blanco-Kelly, F., de las Heras, V., Bartolomé, M., de la Concha, E.G., Urcelay, E., Arroyo, R., and Martínez, A. (2009). Glypican 5 is an interferon-beta response gene: a replication study. *Mult. Scler.* **15**, 913-917.

Chand, A.L., Robertson, D.M., Shelling, A.N., and Harrison, C.A. (2007). Mutational analysis of betaglycan/TGF-betaRIII in premature ovarian failure. *Fertil. Steril.* **87**, 210-212.

Chen, E., Hermanson, S., and Ekker, S.C. (2004). Syndecan-2 is essential for angiogenic sprouting during zebrafish development. *Blood* **103**, 1710-1719.

Chen, E., Stringer, S.E., Rusch, M.A., Selleck, S.B., and Ekker, S.C. (2005). A unique role for 6-O sulfation modification in zebrafish vascular development. *Dev. Biol.* **284**, 364-376.

Chen, J., Repunte-Canonigo, V., Kawamura, T., Lefebvre, C., Shin, W., Howell, L.L., Hemby, S.E., Harvey, B.K., Califano, A., Morales, M. et al. (2013). Hypothalamic proteoglycan syndecan-3 is a novel cocaine addiction resilience factor. *Nat. Commun.* **4**, 1955.

Chen, S., Wassenhove-McCarthy, D.J., Yamaguchi, Y., Holzman, L.B., van Kuppevelt, T.H., Jenniskens, G.J., Wijnhoven, T.J., Woods, A.C., and McCarthy, K.J. (2008). Loss of heparan sulfate glycosaminoglycan assembly in podocytes does not lead to proteinuria. *Kidney Int.* **74**, 289-299.

Clegg, J.M., Conway, C.D., Howe, K.M., Price, D.J., Mason, J.O., Turnbull, J.E., Basson, M.A., and Pratt, T. (2014) Heparan sulfotransferases Hs6st1 and Hs2st keep Erk in check for mouse corpus callosum development. *J. Neurosci.* **34**, 2389-2401.

- Clément, A., Wiweger, M., von der Hardt, S., Rusch, M.A., Selleck, S.B., Chien, C.B., and Roehl, H.H.** (2008). Regulation of zebrafish skeletogenesis by ext2/dackel and papst1/pinscher. *PLoS Genet.* **4**, e1000136.
- Compton, L.A., Potash, D.A., Brown, C.B., and Barnett, J.V.** (2007). Coronary vessel development is dependent on the type III transforming growth factor beta receptor. *Circ Res.* **101**, 784-791.
- Conway, C.D., Howe, K.M., Nettleton, N.K., Price, D.J., Mason, J.O., and Pratt, T.** (2001). Heparan sulfate sugar modifications mediate the functions of slits and other factors needed for mouse forebrain commissure development. *J. Neurosci.* **31**, 1955-1970.
- Conway, C.D., Price, D.J., Pratt, T., and Mason, J.O.** (2011). Analysis of axon guidance defects at the optic chiasm in heparan sulphate sulphotransferase compound mutant mice. *J. Anat.* **219**, 734-742.
- Cornelison, D.D., Wilcox-Adelman, S.A., Goetinck, P.F., Rauvala, H., Rapraeger, A.C., and Olwin, B.B.** (2004). Essential and separable roles for Syndecan-3 and Syndecan-4 in skeletal muscle development and regeneration. *Genes Dev.* **18**, 2231-2236.
- Costell, M., Carmona, R., Gustafsson, E., González-Iriarte, M., Fässler, R., and Muñoz-Chápuli, R.** (2002). Hyperplastic conotruncal endocardial cushions and transposition of great arteries in perlecan-null mice. *Circ. Res.* **91**, 158-164.
- Cottereau, E., Mortemousque, I., Moizard, M.P., Bürglen, L., Lacombe, D., Gilbert-Dussardier, B., Sigaudy, S., Boute, O., David, A., Faivre, L. et al.** (2013). Phenotypic spectrum of Simpson-Golabi-Behmel syndrome in a series of 42 cases with a mutation in GPC3 and review of the literature. *Am. J. Med. Genet. C. Semin. Med. Genet.* **163C**, 92-105.
- Coulson-Thomas, V.J., Gesteira, T.F., Esko, J., and Kao, W.** (2014). Heparan sulfate regulates hair follicle and sebaceous gland morphogenesis and homeostasis. *J. Biol. Chem.* **289**, 25211-25226.
- Crawford, B.E., Garner, O.B., Bishop, J.R., Zhang, D.Y., Bush, K.T., Nigam, S.K., and Esko, J.D.** (2010). Loss of the heparan sulfate sulfotransferase, Ndst1, in mammary epithelial cells selectively blocks lobuloalveolar development in mice. *PLoS One* **5**, e10691.

- Cui, S., Leyva-Vega, M., Tsai, E.A., EauClaire, S.F., Glessner, J.T., Hakonarson, H., Devoto, M., Haber, B.A., Spinner, N.B., and Matthews, R.P.** (2013). Evidence from human and zebrafish that GPC1 is a biliary atresia susceptibility gene. *Gastroenterology* **144**, 1107-1115.
- Daly, S.B., Urquhart, J.E., Hilton, E., McKenzie, E.A., Kammerer, R.A., Lewis, M., Kerr, B., Stuart, H., Donnai, D., Long, D.A. et al.** (2010). Mutations in HPSE2 cause urofacial syndrome. *Am. J. Hum. Genet.* **86**, 963-969.
- De Cat, B., Muyldeermans, S.Y., Coomans, C., Degeest, G., Vanderschueren, B., Creemers, J., Biemar, F., Peers, B., and David, G.** (2003). Processing by proprotein convertases is required for glypcan-3 modulation of cell survival, Wnt signaling, and gastrulation movements. *J. Cell Biol.* **163**, 625-635.
- Dedkov, E.I., Thomas, M.T., Sonka, M., Yang, F., Chittenden, T.W., Rhodes, J.M., Simons, M., Ritman, E.L., and Tomanek, R.J.** (2007). Synectin/syndecan-4 regulate coronary arteriolar growth during development. *Dev. Dyn.* **236**, 2004-2010.
- Dhamija, R., Graham, J.M. Jr., Smaoui, N., Thorland, E., and Kirmani, S.** (2014). Novel de novo SPOCK1 mutation in a proband with developmental delay, microcephaly and agenesis of corpus callosum. *Eur. J. Med. Genet.* **57**, 181-184.
- Dixit, H., Rao, K.L., Padmalatha, V.V., Kanakavalli, M., Deenadayal, M., Gupta, N., Chakrabarty, B.N., and Singh, L.** (2006). Mutational analysis of the betaglycan gene-coding region in susceptibility for ovarian failure. *Hum. Reprod.* **21**, 2041-2046.
- Duchez, S., Pascal, V., Cogné, N., Jayat-Vignoles, C., Julien, R., and Cogné, M.** (2011). Glycotranscriptome study reveals an enzymatic switch modulating glycosaminoglycan synthesis during B-cell development and activation. *Eur. J. Immunol.* **41**, 3632-3644.
- Echaniz-Laguna, A., Rene, F., Marcel, C., Bangratz, M., Fontaine, B., Loeffler, J.P., and Nicole, S.** (2009). Electrophysiological studies in a mouse model of Schwartz-Jampel syndrome demonstrate muscle fiber hyperactivity of peripheral nerve origin. *Muscle Nerve* **40**, 55-61.
- Echtermeyer, F., Streit, M., Wilcox-Adelman, S., Saoncella, S., Denhez, F., Detmar, M., and Goetinck, P.** (2001). Delayed wound repair and impaired angiogenesis in mice lacking syndecan-4. *J. Clin. Invest.* **107**, R9-R14.

- Elamaa, H., Sormunen, R., Rehn, M., Soininen, R., and Pihlajaniemi, T.** (2005). Endostatin overexpression specifically in the lens and skin leads to cataract and ultrastructural alterations in basement membranes. *Am J Pathol.* **166**, 221-229.
- Elliott, L., Ashley-Koch, A.E., De Castro, L., Jonassaint, J., Price, J., Ataga, K.I., Levesque, M.C., Brice Weinberg, J., Eckman, J.R., Orringer, E.P. et al.** (2007). Genetic polymorphisms associated with priapism in sickle cell disease. *Br. J. Haematol.* **137**, 262-267.
- Escobedo, N., Contreras, O., Muñoz, R., Farías, M., Carrasco, H., Hill, C., Tran, U., Pryor, S.E., Wessely, O., Copp, A.J., and Larraín, J.** (2013). Syndecan 4 interacts genetically with Vangl2 to regulate neural tube closure and planar cell polarity. *Development* **140**, 3008-3017.
- Fan, G., Xiao, L., Cheng, L., Wang, X., Sun, B., and Hu, G.** (2000). Targeted disruption of NDST-1 gene leads to pulmonary hypoplasia and neonatal respiratory distress in mice. *FEBS Lett.* **467**, 7-11.
- Fellgett, S.W., Maguire, R.J., and Pownall, M.E.** (2015). Sulf1 has ligand dependent effects on canonical and non-canonical Wnt signalling. *J. Cell Sci.* pii: jcs.164467.
- Filipek-Górniok, B., Carlsson, P., Haitina, T., Habicher, J., Ledin, J., and Kjellén, L.** (2015). The ndst gene family in zebrafish: role of ndst1b in pharyngeal arch formation. *PLoS One* **10**, e0119040.
- Fischer, S., Filipek-Gorniok, B., and Ledin J.** (2011). Zebrafish Ext2 is necessary for Fgf and Wnt signaling, but not for Hh signaling. *BMC Dev. Biol.* 2011 Sep 5, 11:53.
- Floer, M., Götte, M., Wild, M.K., Heidemann, J., Gassar, E.S., Domschke, W., Kiesel, L., Luegering, A., and Kucharzik, T.** (2010). Enoxaparin improves the course of dextran sodium sulfate-induced colitis in syndecan-1-deficient mice. *Am. J. Pathol.* **176**, 146-157.
- Forsberg, E., Pejler, G., Ringvall, M., Lunderius, C., Tomasini-Johansson, B., Kusche-Gullberg, M., Eriksson, I., Ledin, J., Hellman, L., and Kjellén, L.** (1999). Abnormal mast cells in mice deficient in a heparin-synthesizing enzyme. *Nature* **400**, 773-776.
- Freeman, S.D., Moore, W.M., Guiral, E.C., Holme, A.D., Turnbull, J.E., and Pownall, M.E.** (2008). Extracellular regulation of developmental cell signaling by XtSulf1. *Dev. Biol.* **320**, 436-445.

- Fuerst, P.G., Rauch, S.M., and Burgess, R.W.** (2007). Defects in eye development in transgenic mice overexpressing the heparan sulfate proteoglycan agrin. *Dev. Biol.* **303**, 165-180.
- Fukai, N., Eklund, L., Marneros, A.G., Oh, S.P., Keene, D.R., Tamarkin, L., Niemelä, M., Ilves, M., Li, E., Pihlajaniemi, T., and Olsen, B.R.** (2002). Lack of collagen XVIII/endostatin results in eye abnormalities. *EMBO J.* **21**, 1535-1544.
- Fuster, M.M., Wang, L., Castagnola, J., Sikora, L., Reddi, K., Lee, P.H., Radek, K.A., Schuksz, M., Bishop, J.R., Gallo, R.L. et al.** (2007). Genetic alteration of endothelial heparan sulfate selectively inhibits tumor angiogenesis. *J. Cell. Biol.* **177**, 539-549.
- Galli, A., Roure, A., Zeller, R., and Dono, R.** (2003). Glypican 4 modulates FGF signalling and regulates dorsoventral forebrain patterning in Xenopus embryos. *Development* **130**, 4919-4929.
- Garner, O.B., Yamaguchi, Y., Esko, J.D., and Videm, V.** (2008). Small changes in lymphocyte development and activation in mice through tissue-specific alteration of heparan sulphate. *Immunology* **125**, 420-429.
- Garner, O.B., Bush, K.T., Nigam, K.B., Yamaguchi, Y., Xu, D., Esko, J.D., and Nigam, S.K.** (2011). Stage-dependent regulation of mammary ductal branching by heparan sulfate and HGF-cMet signaling. *Dev. Biol.* **355**, 394-403.
- Gautam, M., Noakes, P.G., Moscoso, L., Rupp, F., Scheller, R.H., Merlie, J.P., and Sanes, J.R.** (1996). Defective neuromuscular synaptogenesis in agrin-deficient mutant mice. *Cell* **85**, 525-535.
- Gautam, M., DeChiara, T.M., Glass, D.J., Yancopoulos, G.D., and Sanes, J.R.** (1999). Distinct phenotypes of mutant mice lacking agrin, MuSK, or rapsyn. *Brain Res. Dev. Brain Res.* **114**, 171-178.
- Ge, X.N., Ha, S.G., Rao, A., Greenberg, Y.G., Rushdi, M.N., Esko, J.D., Rao, S.P., and Sriramaraao, P.** (2014). Endothelial and leukocyte heparan sulfates regulate the development of allergen-induced airway remodeling in a mouse model. *Glycobiology* **24**, 715-727.
- Ghiselli, G. and Farber, S.A.** (2005). D-glucuronyl C5-epimerase acts in dorso-ventral axis formation in zebrafish. *BMC Dev. Biol.* **12**, 5-19.

- Gil, N., Goldberg, R., Neuman, T., Garsen, M., Zcharia, E., Rubinstein, A.M., van Kuppevelt, T., Meirovitz, A., Pisano, C., Li, J.P. et al.** (2012). Heparanase is essential for the development of diabetic nephropathy in mice. *Diabetes* **61**, 208-216.
- Girós, A., Morante, J., Gil-Sanz, C., Fairén, A., and Costell, M.** (2007). Perlecan controls neurogenesis in the developing telencephalon. *BMC Dev. Biol.* **5**, 7-29.
- Godfrey, E.W., Roe, J., and Heathcote, R.D.** (1999). Overexpression of agrin isoforms in Xenopus embryos alters the distribution of synaptic acetylcholine receptors during development of the neuromuscular junction. *Dev. Biol.* **205**, 22-32.
- Gorsi, B., Liu, F., Ma, X., Chico, T.J., Shrinivasan, A., Kramer, K.L., Bridges, E., Monteiro, R., Harris, A.L., Patient, R., and Stringer, S.E.** (2014). The heparan sulfate editing enzyme Sulf1 plays a novel role in zebrafish VegfA mediated arterial venous identity. *Angiogenesis* **17**, 77-91.
- Gotha, L., Lim, S.Y., Osherov, A.B., Wolff, R., Qiang, B., Erlich, I., Nili, N., Pillarisetti, S., Chang, Y.T., Tran, P.K. et al.** (2014). Heparan sulfate side chains have a critical role in the inhibitory effects of perlecan on vascular smooth muscle cell response to arterial injury. *Am. J. Physiol. Heart. Circ. Physiol.* **307**, H337-345.
- Götte, M., Joussen, A.M., Klein, C., Andre, P., Wagner, D.D., Hinkes, M.T., Kirchhof, B., Adamis, A.P., and Bernfield, M.** (2002). Role of syndecan-1 in leukocyte-endothelial interactions in the ocular vasculature. *Invest. Ophthalmol. Vis. Sci.* **43**, 1135-1141.
- Grandel, H., Draper, B.W., and Schulte-Merker, S.** (2000). dackel acts in the ectoderm of the zebrafish pectoral fin bud to maintain AER signaling. *Development* **127**, 4169-4178.
- Grisaru, S., Cano-Gauci, D., Tee, J., Filmus, J., and Rosenblum, N.D.** (2001). Glypican-3 modulates BMP- and FGF-mediated effects during renal branching morphogenesis. *Dev. Biol.* **231**, 31-46.
- Grobe, K. and Esko, J.D.** (2002). Regulated translation of heparan sulfate N-acetylglicosamine N-deacetylase/n-sulfotransferase isozymes by structured 5'-untranslated regions and internal ribosome entry sites. *J. Biol. Chem.* **277**, 30699-30706.
- Grobe, K., Inatani, M., Pallerla, S.R., Castagnola, J., Yamaguchi, Y., and Esko, J.D.** (2005). Cerebral hypoplasia and craniofacial defects in mice lacking heparan sulfate Ndst1 gene function. *Development* **132**, 3777-3786.

- Ha, E., Kim, M.J., Choi, B.K., Rho, J.J., Oh, D.J., Rho, T.H., Kim, K.H., Lee, H.J., Shin, D.H., and Yim, S.V.** (2006). Positive association of obesity with single nucleotide polymorphisms of syndecan 3 in the Korean population. *J. Clin. Endocrinol. Metab.* **91**, 5095-5099.
- Habuchi, H., Nagai, N., Sugaya, N., Atsumi, F., Stevens, R.L., and Kimata, K.** (2007). Mice deficient in heparan sulfate 6-O-sulfotransferase-1 exhibit defective heparan sulfate biosynthesis, abnormal placentation, and late embryonic lethality. *J. Biol. Chem.* **282**, 15578-15588.
- HajMohammadi, S., Enjyoji, K., Princivalle, M., Christi, P., Lech, M., Beeler, D., Rayburn, H., Schwartz, J.J., Barzegar, S., de Agostini, A.I. et al.** (2003). Normal levels of anticoagulant heparan sulfate are not essential for normal hemostasis. *J. Clin. Invest.* **111**, 989-999.
- Hanai, J., Gloy, J., Karumanchi, S.A., Kale, S., Tang, J., Hu, G., Chan, B., Ramchandran, R., Jha, V., Sukhatme, V.P., and Sokol, S.** (2002). Endostatin is a potential inhibitor of Wnt signaling. *J. Cell Biol.* **158**, 529-539.
- Harfouche, R., Hentschel, D.M., Piecewicz, S., Basu, S., Print, C., Eavarone, D., Kiziltepe, T., Sasisekharan, R., and Sengupta, S.** (2009). Glycome and transcriptome regulation of vasculogenesis. *Circulation* **120**, 1883-1892.
- Hart, M., Li, L., Tokunaga, T., Lindsey, J.R., Hassell, J.R., Snow, A.D., and Fukuchi, K.** (2001). Overproduction of perlecan core protein in cultured cells and transgenic mice. *J. Pathol.* **194**, 262-269.
- Hartmann, U., Hülsmann, H., Seul, J., Röll, S., Midani, H., Breloy, I., Hechler, D., Müller, R., and Paulsson, M.** (2013). Testican-3: a brain-specific proteoglycan member of the BM-40/SPARC/osteonectin family. *J. Neurochem.* **125**, 399-409.
- Hartwig, S., Hu, M.C., Cella, C., Piscione, T., Filmus, J., and Rosenblum, N.D.** (2005). Glypican-3 modulates inhibitory Bmp2-Smad signaling to control renal development in vivo. *Mech. Dev.* **122**, 928-938.
- Hausser, H.J., Ruegg, M.A., Brenner, R.E., and Ksiazek, I.** (2007). Agrin is highly expressed by chondrocytes and is required for normal growth. *Histochem. Cell. Biol.* **127**, 363-374.
- Hayano, S., Kurosaka, H., Yanagita, T., Kalus, I., Milz, F., Ishihara, Y., Islam, M.N., Kawanabe, N., Saito, M., Kamioka, H. et al.** (2012). Roles of heparan sulfate sulfation in dentinogenesis. *J. Biol. Chem.* **287**, 12217-12229.

- Hayashida, K., Chen, Y., Bartlett, A.H., and Park, P.W.** (2008). Syndecan-1 is an in vivo suppressor of Gram-positive toxic shock. *J. Biol. Chem.* **283**, 19895-19903.
- Hienola, A., Tumova, S., Kulesskiy, E., and Rauvala, H.** (2006). N-syndecan deficiency impairs neural migration in brain. *J. Cell. Biol.* **174**, 569-580.
- Hilgenberg, L.G., Ho, K.D., Lee, D., O'Dowd, D.K., and Smith, M.A.** (2002). Agrin regulates neuronal responses to excitatory neurotransmitters in vitro and in vivo. *Mol. Cell. Neurosci.* **19**, 97-110.
- Hilton, M.J., Gutiérrez, L., Martinez, D.A., and Wells, D.E.** (2005). EXT1 regulates chondrocyte proliferation and differentiation during endochondral bone development. *Bone* **36**, 379-386.
- Holmborn, K., Habicher, J., Kasza, Z., Eriksson, A.S., Filipek-Gorniok, B., Gopal, S., Couchman, J.R., Ahlberg, P.E., Wiweger, M., Spillmann, D. et al.** (2012). On the roles and regulation of chondroitin sulfate and heparan sulfate in zebrafish pharyngeal cartilage morphogenesis. *J. Biol. Chem.* **287**, 33905-33916.
- Hu, Z., Yu, M., and Hu, G.** (2007). NDST-1 modulates BMPR and PTHrP signaling during endochondral bone formation in a gene knockout model. *Bone* **40**, 1462-1474.
- Hu, Z., Wang, C., Xiao, Y., Sheng, N., Chen, Y., Xu, Y., Zhang, L., Mo, W., Jing, N., and Hu, G.** (2009). NDST1-dependent heparan sulfate regulates BMP signaling and internalization in lung development. *J. Cell Sci.* **122**, 1145-1154.
- Huegel, J., Mundy, C., Sgariglia, F., Nygren, P., Billings, P.C., Yamaguchi, Y., Koyama, E., and Pacifici, M.** (2013). Perichondrium phenotype and border function are regulated by Ext1 and heparan sulfate in developing long bones: a mechanism likely deranged in Hereditary Multiple Exostoses. *Dev. Biol.* **377**, 100-112.
- Huzé, C., Bauché, S., Richard, P., Chevessier, F., Goillot, E., Gaudon, K., Ben Ammar, A., Chaboud, A., Grosjean, I., Lecuyer, H.A. et al.** (2009). Identification of an agrin mutation that causes congenital myasthenia and affects synapse function. *Am. J. Hum. Genet.* **85**, 155-167.
- Ida-Yonemochi, H., Satokata, I., Ohshima, H., Sato, T., Yokoyama, M., Yamada, Y., and Saku, T.** (2011). Morphogenetic roles of perlecan in the tooth enamel organ: an analysis of overexpression using transgenic mice. *Matrix Biol.* **30**, 379-388.

- Inatani, M., Irie, F., Plump, A.S., Tessier-Lavigne, M., and Yamaguchi, Y.** (2003). Mammalian brain morphogenesis and midline axon guidance require heparan sulfate. *Science* **302**, 1044-1046.
- Inomata, T., Ebihara, N., Funaki, T., Matsuda, A., Watanabe, Y., Ning, L., Xu, Z., Murakami, A., and Arikawa-Hirasawa, E.** (2012). Perlecan-deficient mutation impairs corneal epithelial structure. *Invest. Ophthalmol. Vis. Sci.* **53**, 1277-1284.
- Irie, F., Badie-Mahdavi, H., and Yamaguchi, Y.** (2012). Autism-like socio-communicative deficits and stereotypies in mice lacking heparan sulfate. *Proc. Natl. Acad. Sci. USA* **109**, 5052-5056.
- Ishiguro, K., Kadomatsu, K., Kojima, T., Muramatsu, H., Tsuzuki, S., Nakamura, E., Kusugami, K., Saito, H., and Muramatsu, T.** (2000a). Syndecan-4 deficiency impairs focal adhesion formation only under restricted conditions. *J. Biol. Chem.* **275**, 5249-5252.
- Ishiguro, K., Kadomatsu, K., Kojima, T., Muramatsu, H., Nakamura, E., Ito, M., Nagasaka, T., Kobayashi, H., Kusugami, K., Saito, H., and Muramatsu, T.** (2000b). Syndecan-4 deficiency impairs the fetal vessels in the placental labyrinth. *Dev. Dyn.* **219**, 539-544.
- Ishiguro, K., Kadomatsu, K., Kojima, T., Muramatsu, H., Matsuo, S., Kusugami, K., Saito, H., and Muramatsu, T.** (2001a). Syndecan-4 deficiency increases susceptibility to kappa-carrageenan-induced renal damage. *Lab. Invest.* **81**, 509-516.
- Ishiguro, K., Kadomatsu, K., Kojima, T., Muramatsu, H., Iwase, M., Yoshikai, Y., Yanada, M., Yamamoto, K., Matsushita, T., Nishimura, M. et al.** (2001b). Syndecan-4 deficiency leads to high mortality of lipopolysaccharide-injected mice. *J. Biol. Chem.* **276**, 47483-47488.
- Ishijima, M., Suzuki, N., Hozumi, K., Matsunobu, T., Kosaki, K., Kaneko, H., Hassell, J.R., Arikawa-Hirasawa, E., and Yamada, Y.** (2012). Perlecan modulates VEGF signaling and is essential for vascularization in endochondral bone formation. *Matrix Biol.* **31**, 234-245.
- Iwabuchi, T. and Goetinck, P.F.** (2006). Syndecan-4 dependent FGF stimulation of mouse vibrissae growth. *Mech. Dev.* **123**, 831-841.
- Iwao, K., Inatani, M., Matsumoto, Y., Ogata-Iwao, M., Takihara, Y., Irie, F., Yamaguchi, Y., Okinami, S., and Tanihara, H.** (2009). Heparan sulfate deficiency leads to Peters anomaly in mice by disturbing neural crest TGF-beta2 signaling. *J. Clin. Invest.* **119**, 1997-2008.

- Iwao, K., Inatani, M., Ogata-Iwao, M., Yamaguchi, Y., Okinami, S., and Tanihara, H.** (2010). Heparan sulfate deficiency in periocular mesenchyme causes microphthalmia and ciliary body dysgenesis. *Exp. Eye Res.* **90**, 81-88.
- Izvolsky, K.I., Lu, J., Martin, G., Albrecht, K.H., and Cardoso, W.V.** (2008). Systemic inactivation of Hs6st1 in mice is associated with late postnatal mortality without major defects in organogenesis. *Genesis* **46**, 8-18.
- Jen, Y.H., Musacchio, M., and Lander, A.D.** (2009). Glypican-1 controls brain size through regulation of fibroblast growth factor signaling in early neurogenesis. *Neural Dev.* **4**, 4:33.
- Jia, J., Maccarana, M., Zhang, X., Bespalov, M., Lindahl, U., and Li, J.P.** (2009). Lack of L-iduronic acid in heparan sulfate affects interaction with growth factors and cell signaling. *J. Biol. Chem.* **284**, 15942-15950.
- Jiang, D., Liang, J., Campanella, G.S., Guo, R., Yu, S., Xie, T., Liu, N., Jung, Y., Homer, R., Meltzer, E.B. et al.** (2010). Inhibition of pulmonary fibrosis in mice by CXCL10 requires glycosaminoglycan binding and syndecan-4. *J. Clin. Invest.* **120**, 2049-2057.
- Jones, K.B., Piombo, V., Searby, C., Kurriger, G., Yang, B., Grabelius, F., Roughley, P.J., Morcuende, J.A., Buckwalter, J.A., Capecchi, M.R. et al.** (2010). A mouse model of osteochondromagenesis from clonal inactivation of Ext1 in chondrocytes. *Proc. Natl. Acad. Sci. USA* **107**, 2054-2059.
- Kaksonen, M., Pavlov, I., Võikar, V., Lauri, S.E., Hienola, A., Riekki, R., Lakso, M., Taira, T., and Rauvala, H.** (2002). Syndecan-3-deficient mice exhibit enhanced LTP and impaired hippocampus-dependent memory. *Mol. Cell. Neurosci.* **21**, 158-172.
- Kalus, I., Salmen, B., Viebahn, C., von Figura, K., Schmitz, D., D'Hooge, R., and Dierks, T.** (2009). Differential involvement of the extracellular 6-O-endosulfatases Sulf1 and Sulf2 in brain development and neuronal and behavioural plasticity. *J. Cell. Mol. Med.* **13**, 4505-4521.
- Kaneko, H., Ishijima, M., Futami, I., Tomikawa-Ichikawa, N., Kosaki, K., Sadatsuki, R., Yamada, Y., Kurosawa, H., Kaneko, K., and Arikawa-Hirasawa, E.** (2013). Synovial perlecan is required for osteophyte formation in knee osteoarthritis. *Matrix Biol.* **32**, 178-187.

- Karlsson-Lindahl, L., Schmidt, L., Haage, D., Hansson, C., Taube, M., Egecioglu, E., Tan, Y.X., Admyre, T., Jansson, J.O., Vlodavsky, I, et al.** (2012). Heparanase affects food intake and regulates energy balance in mice. *PLoS One* **7**, e34313.
- Karlstrom, R.O., Trowe, T., Klostermann, S., Baier, H., Brand, M., Crawford, A.D., Grunewald, B., Haffter, P., Hoffmann, H., Meyer, S.U. et al.** (1996). Zebrafish mutations affecting retinotectal axon pathfinding. *Development*. **123**, 427-438.
- Kasza, I., Suh, Y., Wollny, D., Clark, R.J., Roopra, A., Colman, R.J., MacDougald, O.A., Shedd, T.A., Nelson, D.W., Yen, M.I. et al.** (2014). Syndecan-1 is required to maintain intradermal fat and prevent cold stress. *PLoS Genet.* **10**, e1004514.
- Kehoe, O., Kalia, N., King, S., Eustace, A., Boyes, C., Reizes, O., Williams, A., Patterson, A., and Middleton, J.** (2014). Syndecan-3 is selectively pro-inflammatory in the joint and contributes to antigen-induced arthritis in mice. *Arthritis Res. Ther.* **16**, R148.
- Keren, B., Suzuki, O.T., Gérard-Blanluet, M., Brémont-Gignac, D., Elmaleh, M., Titomanlio, L., Delezoide, A.L., Passos-Bueno, M.R., and Verloes, A.** (2007). CNS malformations in Knobloch syndrome with splice mutation in COL18A1 gene. *Am. J. Med. Genet. A*. **143A**, 1514-1518.
- Kerever, A., Mercier, F., Nonaka, R., de Vega, S., Oda, Y., Zalc, B., Okada, Y., Hattori, N., Yamada, Y., and Arikawa-Hirasawa, E.** (2014). Perlecan is required for FGF-2 signaling in the neural stem cell niche. *Stem Cell Res.* **12**, 492-505.
- Kim, M.J., Liu, I.H., Song, Y., Lee, J.A., Halfter, W., Balice-Gordon, R.J., Linney, E., and Cole, G.J.** (2007). Agrin is required for posterior development and motor axon outgrowth and branching in embryonic zebrafish. *Glycobiology* **17**, 231-247.
- Kinnunen, A.I., Sormunen, R., Elamaa, H., Seppinen, L., Miller, R.T., Ninomiya, Y., Janmey, P.A., and Pihlajaniemi, T.** (2011). Lack of collagen XVIII long isoforms affects kidney podocytes, whereas the short form is needed in the proximal tubular basement membrane. *J. Biol. Chem.* **286**, 7755-7764.
- Kobayashi, T., Habuchi, H., Tamura, K., Ide, H., and Kimata, K.** (2007). Essential role of heparan sulfate 2-O-sulfotransferase in chick limb bud patterning and development. *J. Biol. Chem.* **282**, 19589-19597.

- Kobayashi, T., Habuchi, H., Nogami, K., Ashikari-Hada, S., Tamura, K., Ide, H., and Kimata, K.** (2010). Functional analysis of chick heparan sulfate 6-O-sulfotransferases in limb bud development. *Dev. Growth Differ.* **52**, 146-156.
- Kon, S., Ikesue, M., Kimura, C., Aoki, M., Nakayama, Y., Saito, Y., Kurotaki, D., Diao, H., Matsui, Y., Segawa, T. et al.** (2008). Syndecan-4 protects against osteopontin-mediated acute hepatic injury by masking functional domains of osteopontin. *J Exp. Med.* **205**, 25-33.
- Kramer, K.L. and Yost, H.J.** (2002a). Ectodermal syndecan-2 mediates left-right axis formation in migrating mesoderm as a cell-nonautonomous Vg1 cofactor. *Dev. Cell* **2**, 115-124.
- Kramer, K.L., Barnette, J.E., and Yost, H.J.** (2002b). PKCgamma regulates syndecan-2 inside-out signaling during xenopus left-right development. *Cell* **111**, 981-990.
- Ksiazek, I., Burkhardt, C., Lin, S., Seddik, R., Maj, M., Bezakova, G., Jucker, M., Arber, S., Caroni, P., Sanes, J.R. et al.** (2007). Synapse loss in cortex of agrin-deficient mice after genetic rescue of perinatal death. *J. Neurosci.* **27**, 7183-7195.
- Kuriyama, S. and Mayor, R.** (2009). A role for Syndecan-4 in neural induction involving ERK- and PKC-dependent pathways. *Development* **136**, 575-584.
- Lamanna, W.C., Baldwin, R.J., Padva, M., Kalus, I., Ten Dam, G., van Kuppevelt, T.H., Gallagher, J.T., von Figura, K., Dierks, T., and Merry, C.L.** (2006). Heparan sulfate 6-O-endosulfatases: discrete in vivo activities and functional co-operativity. *Biochem. J.* **400**, 63-73.
- LeClair, E.E., Mui, S.R., Huang, A., Topczewska, J.M., and Topczewski, J.** (2009). Craniofacial skeletal defects of adult zebrafish Glypican 4 (knypek) mutants. *Dev. Dyn.* **238**, 2550-2563.
- Lee, J.S., von der Hardt, S., Rusch, M.A., Stringer, S.E., Stickney, H.L., Talbot, W.S., Geisler, R., Nüsslein-Volhard, C., Selleck, S.B., Chien, C.B., and Roehl, H.** (2004). Axon sorting in the optic tract requires HSPG synthesis by ext2 (dackel) and extl3 (boxer). *Neuron* **44**, 947-960.
- Lencz, T., Guha, S., Liu, C., Rosenfeld, J., Mukherjee, S., DeRosse, P., John, M., Cheng, L., Zhang, C., Badner, J.A. et al.** (2013). Genome-wide association study implicates NDST3 in schizophrenia and bipolar disorder. *Nat. Commun.* **4**, 2739.

- Li, H., Yamagata, T., Mori, M., and Momoi, M.Y.** (2002). Association of autism in two patients with hereditary multiple exostoses caused by novel deletion mutations of EXT1. *J. Hum. Genet.* **47**, 262-265.
- Li, J., Partovian, C., Li, J., Hampton, T.G., Metais, C., Tkachenko, E., Sellke, F.W., and Simons, M.** (2002). Modulation of microvascular signaling by heparan sulfate matrix: studies in syndecan-4 transgenic mice. *Microvasc. Res.* **64**, 38-46.
- Li, J.P., Gong, F., Hagner-McWhirter, A., Forsberg, E., Abrink, M., Kisilevsky, R., Zhang, X., and Lindahl, U.** (2003). Targeted disruption of a murine glucuronyl C5-epimerase gene results in heparan sulfate lacking L-iduronic acid and in neonatal lethality. *J. Biol. Chem.* **278**, 28363-28366.
- Li, L., Wang, B., Gao, T., Zhang, X., Hao, J.X., Vlodavsky, I., Wiesenfeld-Hallin, Z., Xu, X.J., and Li, J.P.** (2012). Heparanase overexpression reduces carrageenan-induced mechanical and cold hypersensitivity in mice. *Neurosci. Lett.* **511**, 4-7.
- Lin, X., Wei, G., Shi, Z., Dryer, L., Esko, J.D., Wells, D.E., and Matzuk, M.M.** (2000). Disruption of gastrulation and heparan sulfate biosynthesis in EXT1-deficient mice. *Dev. Biol.* **224**, 299-311.
- Liu, B., Bell, A.W., Paranjpe, S., Bowen, W.C., Khillan, J.S., Luo, J.H., Mars, W.M., and Michalopoulos, G.K.** (2010). Suppression of liver regeneration and hepatocyte proliferation in hepatocyte-targeted glycan 3 transgenic mice. *Hepatology* **52**, 1060-1067.
- Liu, I.H., Zhang, C., Kim, M.J., and Cole, G.J.** (2008). Retina development in zebrafish requires the heparan sulfate proteoglycan agrin. *Dev. Neurobiol.* **68**, 877-898.
- Liu, L., Yang, X., Wang, H., Cui, G., Xu, Y., Wang, P., Yuan, G., Wang, X., Ding, H., and Wang, D.W.** (2013). Association between variants of EXT2 and type 2 diabetes: a replication and meta-analysis. *Hum. Genet.* **132**, 139-145.
- Liu, Y., Echtermeyer, F., Thilo, F., Theilmeier, G., Schmidt, A., Schülein, R., Jensen, B.L., Loddenkemper, C., Jankowski, V., Marcussen, N. et al.** (2012). The proteoglycan syndecan 4 regulates transient receptor potential canonical 6 channels via RhoA/Rho-associated protein kinase signaling. *Arterioscler. Thromb. Vasc. Biol.* **32**, 378-385.
- Lowe, D.A., Lepori-Bui, N., Fomin, P.V., Sloofman, L.G., Zhou, X., Farach-Carson, M.C., Wang, L., and Kirn-Safran, C.B.** (2014). Deficiency in perlecan/HSPG2 during bone

- development enhances osteogenesis and decreases quality of adult bone in mice. *Calcif. Tissue Int.* **95**, 29-38.
- Lum, D.H., Tan, J., Rosen, S.D., and Werb, Z.** (2007). Gene trap disruption of the mouse heparan sulfate 6-O-endosulfatase gene, Sulf2. *Mol. Cell. Biol.* **27**, 678-688.
- Maltseva, I., Chan, M., Kalus, I., Dierks, T., and , S.D.** (2013). The SULFs, extracellular sulfatases for heparan sulfate, promote the migration of corneal epithelial cells during wound repair. *PLoS One* **8**, e69642.
- Marneros, A.G. and Olsen, B.R.** (2003). Age-dependent iris abnormalities in collagen XVIII/endostatin deficient mice with similarities to human pigment dispersion syndrome. *Invest. Ophthalmol. Vis. Sci.* **44**, 2367-2372.
- Marneros, A.G., Keene, D.R., Hansen, U., Fukai, N., Moulton, K., Goletz, P.L., Moiseyev, G., Pawlyk, B.S., Halfter, W., and Dong, S.** (2004). Collagen XVIII/endostatin is essential for vision and retinal pigment epithelial function. *EMBO J.* **23**, 89-99.
- Matsui, Y., Ikesue, M., Danzaki, K., Morimoto, J., Sato, M., Tanaka, S., Kojima, T., Tsutsui, H., and Uede, T.** (2011). Syndecan-4 prevents cardiac rupture and dysfunction after myocardial infarction. *Circ. Res.* **108**, 1328-1339.
- Matsumoto, Y., Irie, F., Inatani, M., Tessier-Lavigne, M., and Yamaguchi, Y.** (2007). Netrin-1/DCC signaling in commissural axon guidance requires cell-autonomous expression of heparan sulfate. *J. Neurosci.* **27**, 4342-4350.
- Matsumoto, K., Irie, F., Mackem, S., and Yamaguchi, Y.** (2010). A mouse model of chondrocyte-specific somatic mutation reveals a role for Ext1 loss of heterozygosity in multiple hereditary exostoses. *Proc. Natl. Acad. Sci. USA* **107**, 10932-10937.
- Matthews, H.K., Marchant, L., Carmona-Fontaine, C., Kuriyama, S., Larraín, J., Holt, M.R., Parsons, M., and Mayor, R.** (2008). Directional migration of neural crest cells in vivo is regulated by Syndecan-4/Rac1 and non-canonical Wnt signaling/RhoA. *Development* **135**, 1771-1780.
- McDermott, S.P., Ranheim, E.A., Leatherberry, V.S., Khwaja, S.S., Klos, K.S., and Alexander, C.M.** (2007). Juvenile syndecan-1 null mice are protected from carcinogen-induced tumor development. *Oncogene* **26**, 1407-1416.

McLaughlin, D., Karlsson, F., Tian, N., Pratt, T., Bullock, S.L., Wilson, V.A., Price, D.J., and Mason, J.O. (2003). Specific modification of heparan sulphate is required for normal cerebral cortical development. *Mech. Dev.* **120**, 1481-1488.

Menzel, O., Bekkeheien, R.C., Reymond, A., Fukai, N., Boye, E., Kosztolanyi, G., Aftimos, S., Deutsch, S., Scott, H.S., Olsen, B.R. et al. (2004). Knobloch syndrome: novel mutations in COL18A1, evidence for genetic heterogeneity, and a functionally impaired polymorphism in endostatin. *Hum. Mutat.* **23**, 77-84.

Meyers, J.R., Planamento, J., Ebrom, P., Kruelewitz, N., Wade, E., and Pownall, M.E. (2013). Sulf1 modulates BMP signaling and is required for somite morphogenesis and development of the horizontal myoseptum. *Dev. Biol.* **378**, 107-121.

Mitchell, K.J., Pinson, K.I., Kelly, O.G., Brennan, J., Zupicich, J., Scherz, P., Leighton, P.A., Goodrich, L.V., Lu, X., Avery, B.J. et al. (2001). Functional analysis of secreted and transmembrane proteins critical to mouse development. *Nat. Genet.* **28**, 241-249.

Montaniel, K.R., Billaud, M., Graham, C., Kim, S.K., Carlson, M., Zeng, W., Zeng, O., Pan, W., Isakson, B.E., Hall, J.L., and Adhikari, N. (2012). Smooth muscle specific deletion of Ndst1 leads to decreased vessel luminal area and no change in blood pressure in conscious mice. *J. Cardiovasc. Transl. Res.* **5**, 274-279.

Morimoto, K., Shimizu, T., Furukawa, K., Morio, H., Kurosawa, H., and Shirasawa, T. (2002). Transgenic expression of the EXT2 gene in developing chondrocytes enhances the synthesis of heparan sulfate and bone formation in mice. *Biochem. Biophys. Res. Commun.* **292**, 999-1009.

Morita, H., Yoshimura, A., Inui, K., Ideura, T., Watanabe, H., Wang, L., Soininen, R., and Tryggvason, K. (2005). Heparan sulfate of perlecan is involved in glomerular filtration. *J. Am. Soc. Nephrol.* **16**, 1703-1710.

Mundy, C., Yasuda, T., Kinumatsu, T., Yamaguchi, Y., Iwamoto, M., Enomoto-Iwamoto, M., Koyama, E., and Pacifici, M. (2011). Synovial joint formation requires local Ext1 expression and heparan sulfate production in developing mouse embryo limbs and spine. *Dev. Biol.* **351**, 70-81.

Muñoz, R., Moreno, M., Oliva, C., Orbenes, C., and Larraín, J. (2006). Syndecan-4 regulates non-canonical Wnt signalling and is essential for convergent and extension movements in Xenopus embryos. *Nat. Cell. Biol.* **8**, 492-500.

- Nadanaka, S., Kagiyama, S., and Kitagawa, H.** (2013a). Roles of EXTL2, a member of the EXT family of tumour suppressors, in liver injury and regeneration processes. *Biochem. J.* **454**, 133-145.
- Nadanaka, S., Zhou, S., Kagiyama, S., Shoji, N., Sugahara, K., Sugihara, K., Asano, M., and Kitagawa, H.** (2013b). EXTL2, a member of the EXT family of tumor suppressors, controls glycosaminoglycan biosynthesis in a xylose kinase-dependent manner. *J. Biol. Chem.* **288**, 9321-9333.
- Nagai, N., Habuchi, H., Sugaya, N., Nakamura, M., Imamura, T., Watanabe, H., and Kimata, K.** (2013). Involvement of heparan sulfate 6-O-sulfation in the regulation of energy metabolism and the alteration of thyroid hormone levels in male mice. *Glycobiology* **23**, 980-992.
- Narvid, J., Gorno-Tempini, M.L., Slavotinek, A., Dearmond, S.J., Cha, Y.H., Miller, B.L., and Rankin, K.** (2009). Of brain and bone: the unusual case of Dr. A. *Neurocase* **15**, 190-205.
- Neugebauer, J.M., Cadwallader, A.B., Amack, J.D., Bisgrove, B.W., and Yost, H.J.** (2013). Differential roles for 3-OSTs in the regulation of cilia length and motility. *Development* **140**, 3892-3902.
- Ng, A., Wong, M., Viviano, B., Erlich, J.M., Alba, G., Pfleiderer, C., Jay, P.Y., and Saunders, S.** (2009). Loss of glypcan-3 function causes growth factor-dependent defects in cardiac and coronary vascular development. *Dev. Biol.* **335**, 208-215.
- Nicole, S., Davoine, C.S., Topaloglu, H., Cattolico, L., Barral, D., Beighton, P., Hamida, C.B., Hammouda, H., Cruaud, C., White, P.S. et al.** (2000). Perlecan, the major proteoglycan of basement membranes, is altered in patients with Schwartz-Jampel syndrome (chondrodystrophic myotonia). *Nat. Genet.* **26**, 480-483.
- Norton, W.H., Ledin, J., Grandel, H., and Neumann, C.J.** (2005). HSPG synthesis by zebrafish Ext2 and Extl3 is required for Fgf10 signalling during limb development. *Development* **132**, 4963-4973.
- Ogata-Iwao, M., Inatani, M., Iwao, K., Takihara, Y., Nakaishi-Fukuchi, Y., Irie, F., Sato, S., Furukawa, T., Yamaguchi, Y., and Tanihara, H.** (2011). Heparan sulfate regulates intraretinal axon pathfinding by retinal ganglion cells. *Invest. Ophthalmol. Vis. Sci.* **52**, 6671-6679.

- Ohkawara, B., Yamamoto, T.S., Tada, M., and Ueno, N.** (2003). Role of glycan 4 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* **130**, 2129-2138.
- Okamoto, K., Tokunaga, K., Doi, K., Fujita, T., Suzuki, H., Katoh, T., Watanabe, T., Nishida, N., Mabuchi, A., Takahashi, A. et al.** (2011). Common variation in GPC5 is associated with acquired nephrotic syndrome. *Nat. Genet.* **43**, 459-463.
- Olivares, G.H., Carrasco, H., Aroca, F., Carvallo, L., Segovia, F., and Larraín, J.** (2009). Syndecan-1 regulates BMP signaling and dorso-ventral patterning of the ectoderm during early *Xenopus* development. *Dev. Biol.* **329**, 338-349.
- Ostrovsyky, O., Shimoni, A., Rand, A., Vlodavsky, I., and Nagler, A.** (2010). Genetic variations in the heparanase gene (HPSE) associate with increased risk of GVHD following allogeneic stem cell transplantation: effect of discrepancy between recipients and donors. *Blood* **115**, 2319-2328.
- Otsuki, S., Hanson, S.R., Miyaki, S., Grogan, S.P., Kinoshita, M., Asahara, H., Wong, C.H., and Lotz, M.K.** (2010). Extracellular sulfatases support cartilage homeostasis by regulating BMP and FGF signaling pathways. *Proc. Natl. Acad. Sci. USA* **107**, 10202-10207.
- Paine-Saunders, S., Viviano, B.L., Zupicich, J., Skarnes, W.C., and Saunders, S.** (2000). glycan-3 controls cellular responses to Bmp4 in limb patterning and skeletal development. *Dev. Biol.* **225**, 179-187.
- Pallerla, S.R., Pan, Y., Zhang, X., Esko, J.D., and Grobe, K.** (2007). Heparan sulfate NdSt1 gene function variably regulates multiple signaling pathways during mouse development. *Dev. Dyn.* **236**, 556-563.
- Pallerla, S.R., Lawrence, R., Lewejohann, L., Pan, Y., Fischer, T., Schlomann, U., Zhang, X., Esko, J.D., and Grobe, K.** (2008). Altered heparan sulfate structure in mice with deleted NDST3 gene function. *J. Biol. Chem.* **283**, 16885-16894.
- Pan, Y., Woodbury, A., Esko, J.D., Grobe, K., and Zhang, X.** (2006). Heparan sulfate biosynthetic gene NdSt1 is required for FGF signaling in early lens development. *Development* **133**, 4933-4944.

- Pan, Y., Carbe, C., Powers, A., Zhang, E.E., Esko, J.D., Grobe, K., Feng, G.S., and Zhang, X.** (2008). Bud specific N-sulfation of heparan sulfate regulates Shp2-dependent FGF signaling during lacrimal gland induction. *Development* **135**, 301-310.
- Pan, Y., Carbe, C., Kupich, S., Pickhinke, U., Ohlig, S., Frye, M., Seelige, R., Pallerla, S.R., Moon, A.M., Lawrence, R. et al.** (2014). Heparan sulfate expression in the neural crest is essential for mouse cardiogenesis. *Matrix Biol.* **35**, 253-265.
- Pang, J., Zhang, S., Yang, P., Hawkins-Lee, B., Zhong, J., Zhang, Y., Ochoa, B., Agundez, J.A., Voelckel, M.A., Fisher, R.B. et al.** (2010). Loss-of-function mutations in HPSE2 cause the autosomal recessive urofacial syndrome. *Am. J. Hum. Genet.* **86**, 957-962.
- Pilia, G., Hughes-Benzie, R.M., MacKenzie, A., Baybayan, P., Chen, E.Y., Huber, R., Neri, G., Cao, A., Forabosco, A., and Schlessinger, D.** (1996). Mutations in GPC3, a glycan gene, cause the Simpson-Golabi-Behmel overgrowth syndrome. *Nat. Genet.* **12**, 241-247.
- Planer, D., Metzger, S., Zcharia, E., Wexler, I.D., Vlodavsky, I., and Chajek-Shaul, T.** (2011). Role of heparanase on hepatic uptake of intestinal derived lipoprotein and fatty streak formation in mice. *PLoS One* **6**, e18370.
- Poon, I.K., Goodall, K.J., Phipps, S., Chow, J.D., Pagler, E.B., Andrews, D.M., Conlan, C.L., Ryan, G.F., White, J.A., Wong, M.K. et al.** (2014). Mice deficient in heparanase exhibit impaired dendritic cell migration and reduced airway inflammation. *Eur. J. Immunol.* **44**, 1016-1030.
- Poulain, F.E. and Chien, C.B.** (2013). Proteoglycan-mediated axon degeneration corrects pretarget topographic sorting errors. *Neuron* **78**, 49-56.
- Pratt, T., Conway, C.D., Tian, N.M., Price, D.J., and Mason, J.O.** (2006). Heparan sulphation patterns generated by specific heparan sulfotransferase enzymes direct distinct aspects of retinal axon guidance at the optic chiasm. *J. Neurosci.* **26**, 6911-6923.
- Qi, Q., Menzaghi, C., Smith, S., Liang, L., de Rekeneire, N., Garcia, M.E., Lohman, K.K., Miljkovic, I., Strotmeyer, E.S., and Cummings, S.R.** (2012). Genome-wide association analysis identifies TYW3/CRYZ and NDST4 loci associated with circulating resistin levels. *Hum. Mol. Genet.* **21**, 4774-4780.
- Qiang, B., Lim, S.Y., Lekas, M., Kuliszewski, M.A., Wolff, R., Osherov, A.B., Rudenko, D., Leong-Poi, H., Noyan, H., Husain, M. et al.** (2014). Perlecan heparan sulfate proteoglycan is a

critical determinant of angiogenesis in response to mouse hind-limb ischemia. *Can. J. Cardiol.* **30**, 1444-1451.

Qu, X., Carbe, C., Tao, C., Powers, A., Lawrence, R., van Kuppevelt, T.H., Cardoso, W.V., Grobe, K., Esko, J.D., and Zhang, X. (2011). Lacrimal gland development and Fgf10-Fgfr2b

signaling are controlled by 2-O- and 6-O-sulfated heparan sulfate. *J. Biol. Chem.* **286**, 14435-14444.

Qu, X., Pan, Y., Carbe, C., Powers, A., Grobe, K., and Zhang, X. (2012). Glycosaminoglycan-dependent restriction of FGF diffusion is necessary for lacrimal gland development. *Development* **139**, 2730-2739.

Ramsbottom, S.A., Maguire, R.J., Fellgett, S.W., and Pownall, M.E. (2014). Sulf1 influences the Shh morphogen gradient during the dorsal ventral patterning of the neural tube in *Xenopus tropicalis*. *Dev. Biol.* **391**, 207-218.

Rauch, S.M., Huen, K., Miller, M.C., Chaudry, H., Lau, M., Sanes, J.R., Johanson, C.E., Stopa, E.G., and Burgess, R.W. (2011). Changes in brain β-amyloid deposition and aquaporin 4 levels in response to altered agrin expression in mice. *J. Neuropathol. Exp. Neurol.* **70**, 1124-1137.

Reijmers, R.M., Vondenhoff, M.F., Roozendaal, R., Kuil, A., Li, J.P., Spaargaren, M., Pals, S.T., and Mebius, R.E. (2010). Impaired lymphoid organ development in mice lacking the heparan sulfate modifying enzyme glucuronyl C5-epimerase. *J. Immunol.* **184**, 3656-3664.

Reijmers, R.M., Groen, R.W., Kuil, A., Weijer, K., Kimberley, F.C., Medema, J.P., van Kuppevelt, T.H., Li, J.P., Spaargaren, M., and Pals, S.T. (2011). Disruption of heparan sulfate proteoglycan conformation perturbs B-cell maturation and APRIL-mediated plasma cell survival. *Blood* **117**, 6162-6171.

Reizes, O., Lincecum, J., Wang, Z., Goldberger, O., Huang, L., Kaksonen, M., Ahima, R., Hinkes, M.T., Barsh, G.S., Rauvala, H., and Bernfield, M. (2001). Transgenic expression of syndecan-1 uncovers a physiological control of feeding behavior by syndecan-3. *Cell* **106**, 105-116.

Reizes, O., Benoit, S.C., Strader, A.D., Clegg, D.J., Akunuru, S., and Seeley, R.J. (2003). Syndecan-3 modulates food intake by interacting with the melanocortin/AgRP pathway. *Ann. NY Acad. Sci.* **994**, 66-73.

- Reuter, M.S., Musante, L., Hu, H., Diederich, S., Sticht, H., Ekici, A.B., Uebe, S., Wienker, T.F., Bartsch, O., Zechner, U. et al.** (2014). NDST1 missense mutations in autosomal recessive intellectual disability. *Am. J. Med. Genet. A.* **164A**, 2753-2763.
- Rieubland, C., Jacquemont, S., Mittaz, L., Osterheld, M.C., Vial, Y., Superti-Furga, A., Unger, S., and Bonafé, L.** (2010). Phenotypic and molecular characterization of a novel case of dyssegmental dysplasia, Silverman-Handmaker type. *Eur. J. Med. Genet.* **53**, 294-298.
- Ringvall, M., Ledin, J., Holmborn, K., van Kuppevelt, T., Ellin, F., Eriksson, I., Olofsson, A.M., Kjellen, L., and Forsberg, E.** (2000). Defective heparan sulfate biosynthesis and neonatal lethality in mice lacking N-deacetylase/N-sulfotransferase-1. *J. Biol. Chem.* **275**, 25926-25930.
- Rodgers, K.D., Sasaki, T., Aszodi, A., and Jacenko, O.** (2007). Reduced perlecan in mice results in chondrodysplasia resembling Schwartz-Jampel syndrome. *Hum. Mol. Genet.* **16**, 515-528.
- Röll, S., Seul, J., Paulsson, M., and Hartmann, U.** (2006). Testican-1 is dispensable for mouse development. *Matrix Biol.* **25**, 373-381.
- Rops, A.L., Götte, M., Baselmans, M.H., van den Hoven, M.J., Steenbergen, E.J., Lensen, J.F., Wijnhoven, T.J., Cevikbas, F., van den Heuvel, L.P., van Kuppevelt, T.H. et al.** (2007). Syndecan-1 deficiency aggravates anti-glomerular basement membrane nephritis. *Kidney Int.* **72**, 1204-1215.
- Rossi, M., Morita, H., Sormunen, R., Airenne, S., Kreivi, M., Wang, L., Fukai, N., Olsen, B.R., Tryggvason, K., and Soininen, R.** (2003). Heparan sulfate chains of perlecan are indispensable in the lens capsule but not in the kidney. *EMBO J.* **22**, 236-245.
- Rygh, C.B., Løkka, G., Heljasvaara, R., Taxt, T., Pavlin, T., Sormunen, R., Pihlajaniemi, T., Curry, F.R., Tenstad, O., and Reed, R.K.** (2014). Image-based assessment of microvascular function and structure in collagen XV- and XVIII-deficient mice. *J Physiol.* **592**, 325-336.
- Sakazume, S., Okamoto, N., Yamamoto, T., Kurosawa, K., Numabe, H., Ohashi, Y., Kako, Y., Nagai, T., and Ohashi, H.** (2007). GPC3 mutations in seven patients with Simpson-Golabi-Behmel syndrome. *Am. J. Med. Genet. A.* **143A**, 1703-1707.
- Sakimoto, T., Kim, T.I., Ellenberg, D., Fukai, N., Jain, S., Azar, D.T., and Chang, J.H.** (2008). Collagen XVIII and corneal reinnervation following keratectomy. *FEBS Lett.* **582**, 3674-3680.

- Samson, S.C., Ferrer, T., Jou, C.J., Sachse, F.B., Shankaran, S.S., Shaw, R.M., Chi, N.C., Tristani-Firouzi, M., and Yost, H.J.** (2013). 3-OST-7 regulates BMP-dependent cardiac contraction. *PLoS Biol.* **11**, e1001727.
- Samuel, M.A., Valdez, G., Tapia, J.C., Lichtman, J.W., and Sanes, J.R.** (2012). Agrin and synaptic laminin are required to maintain adult neuromuscular junctions. *PLoS One* **7**, e46663.
- Sarraj, M.A., Escalona, R.M., Umbers, A., Chua, H.K., Small, C., Griswold, M., Loveland, K., Findlay, J.K., and Stenvers, K.L.** (2010). Fetal testis dysgenesis and compromised Leydig cell function in Tgfbr3 (beta glycan) knockout mice. *Biol. Reprod.* **82**, 153-162.
- Scarpellini, A., Huang, L., Burhan, I., Schroeder, N., Funck, M., Johnson, T.S., and Verderio, E.A.** (2014). Syndecan-4 knockout leads to reduced extracellular transglutaminase-2 and protects against tubulointerstitial fibrosis. *J. Am. Soc. Nephrol.* **25**, 1013-1027.
- Schneider, V.A. and Granato, M.** (2006). The myotomal diwanka (lh3) glycosyltransferase and type XVIII collagen are critical for motor growth cone migration. *Neuron* **50**, 683-695.
- Schüring, A.N., Lutz, F., Tüttelmann, F., Gromoll, J., Kiesel, L., and Götte, M.** (2009). Role of syndecan-3 polymorphisms in obesity and female hyperandrogenism. *J. Mol. Med. (Berl.)* **87**, 1241-1250.
- Seppinen, L., Sormunen, R., Soini, Y., Elamaa, H., Heljasvaara, R., and Pihlajaniemi, T.** (2008). Lack of collagen XVIII accelerates cutaneous wound healing, while overexpression of its endostatin domain leads to delayed healing. *Matrix Biol.* **27**, 535-546.
- Serpinskaya, A.S., Feng, G., Sanes, J.R., and Craig, A.M.** (1999). Synapse formation by hippocampal neurons from agrin-deficient mice. *Dev. Biol.* **205**, 65-78.
- Serralbo, O. and Marcelle, C.** (2014). Migrating cells mediate long-range WNT signaling. *Development* **141**, 2057-2063.
- Sertié, A.L., Sossi, V., Camargo, A.A., Zatz, M., Brahe, C., and Passos-Bueno, M.R.** (2000). Collagen XVIII, containing an endogenous inhibitor of angiogenesis and tumor growth, plays a critical role in the maintenance of retinal structure and in neural tube closure (Knobloch syndrome). *Hum. Mol. Genet.* **9**, 2051-2058.

- Sgariglia, F., Candela, M.E., Huegel, J., Jacenko, O., Koyama, E., Yamaguchi, Y., Pacifici, M., and Enomoto-Iwamoto, M.** (2013). Epiphyseal abnormalities, trabecular bone loss and articular chondrocyte hypertrophy develop in the long bones of postnatal Ext1-deficient mice. *Bone* **57**, 220-231.
- Shah, M.M., Sakurai, H., Sweeney, D.E., Gallegos, T.F., Bush, K.T., Esko, J.D., and Nigam, S.K.** (2010). Hs2st mediated kidney mesenchyme induction regulates early ureteric bud branching. *Dev. Biol.* **339**, 354-365.
- Shi, W. and Filmus, J.** (2009). A patient with the Simpson-Golabi-Behmel syndrome displays a loss-of-function point mutation in GPC3 that inhibits the attachment of this proteoglycan to the cell surface. *Am. J. Med. Genet. A* **149A**, 552-554.
- Shiau, C.E., Hu, N., and Bronner-Fraser, M.** (2010). Altering Glypican-1 levels modulates canonical Wnt signaling during trigeminal placode development. *Dev. Biol.* **348**, 107-118.
- Shieh, Y.E., Wells, D.E., and Sater, A.K.** (2014). Zygotic expression of Exostosin1 (Ext1) is required for BMP signaling and establishment of dorsal-ventral pattern in Xenopus. *Int. J. Dev. Biol.* **58**, 27-34.
- Shimo, T., Gentili, C., Iwamoto, M., Wu, C., Koyama, E., and Pacifici, M.** (2004). Indian hedgehog and syndecans-3 coregulate chondrocyte proliferation and function during chick limb skeletogenesis. *Dev. Dyn.* **229**, 607-617.
- Shimokawa, K., Kimura-Yoshida, C., Nagai, N., Mukai, K., Matsubara, K., Watanabe, H., Matsuda, Y., Mochida, K., and Matsuo, I.** (2011). Cell surface heparan sulfate chains regulate local reception of FGF signaling in the mouse embryo. *Dev. Cell* **21**, 257-272.
- Shworak, N.W., HajMohammadi, S., de Agostini, A.I., and Rosenberg, R.D.** (2002). Mice deficient in heparan sulfate 3-O-sulfotransferase-1: normal hemostasis with unexpected perinatal phenotypes. *Glycoconj. J.* **19**, 355-361.
- Sipola, A., Seppinen, L., Pihlajaniemi, T., and Tuukkanen, J.** (2009). Endostatin affects osteoblast behavior in vitro, but collagen XVIII/endostatin is not essential for skeletal development in vivo. *Calcif. Tissue Int.* **85**, 412-420.
- Song, H.H., Shi, W., Xiang, Y.Y., and Filmus, J.** (2005). The loss of glycan-3 induces alterations in Wnt signaling. *J. Biol. Chem.* **280**, 2116-2125.

- Stanford, K.I., Bishop, J.R., Foley, E.M., Gonzales, J.C., Niesman, I.R., Witztum, J.L., and Esko, J.D.** (2009). Syndecan-1 is the primary heparan sulfate proteoglycan mediating hepatic clearance of triglyceride-rich lipoproteins in mice. *J. Clin. Invest.* **119**, 3236-3245.
- Stanford, K.I., Wang, L., Castagnola, J., Song, D., Bishop, J.R., Brown, J.R., Lawrence, R., Bai, X., Habuchi, H., Tanaka, M. et al.** (2010). Heparan sulfate 2-O-sulfotransferase is required for triglyceride-rich lipoprotein clearance. *J. Biol. Chem.* **285**, 286-294.
- Stenvers, K.L., Tursky, M.L., Harder, K.W., Kountouri, N., Amatayakul-Chantler, S., Grail, D., Small, C., Weinberg, R.A., Sizeland, A.M., and Zhu, H.J.** (2003). Heart and liver defects and reduced transforming growth factor beta2 sensitivity in transforming growth factor beta type III receptor-deficient embryos. *Mol. Cell. Biol.* **23**, 4371-4385.
- Stepp, M.A., Gibson, H.E., Gala, P.H., Iglesia, D.D., Pajooohesh-Ganji, A., Pal-Ghosh, S., Brown, M., Aquino, C., Schwartz, A.M., Goldberger, O. et al.** (2002). Defects in keratinocyte activation during wound healing in the syndecan-1-deficient mouse. *J. Cell. Sci.* **115**, 4517-4531.
- Stickens, D., Clines, G., Burbee, D., Ramos, P., Thomas, S., Hogue, D., Hecht, J.T., Lovett, M., and Evans, G.A.** (1996). The EXT2 multiple exostoses gene defines a family of putative tumour suppressor genes. *Nat. Genet.* **14**, 25-32.
- Stickens, D., Zak, B.M., Rougier, N., Esko, J.D., and Werb, Z.** (2005). Mice deficient in Ext2 lack heparan sulfate and develop exostoses. *Development* **132**, 5055-5068.
- Strader, A.D., Reizes, O., Woods, S.C., Benoit, S.C., and Seeley, R.J.** (2004). Mice lacking the syndecan-3 gene are resistant to diet-induced obesity. *J. Clin. Invest.* **114**, 1354-1360.
- Stum, M., Davoine, C.S., Vicart, S., Guillot-Noël, L., Topaloglu, H., Carod-Artal, F.J., Kayserili, H., Hentati, F., Merlini, L., Urtizberea, J.A. et al.** (2006). Spectrum of HSPG2 (Perlecan) mutations in patients with Schwartz-Jampel syndrome. *Hum. Mutat.* **27**, 1082-1091.
- Stum, M., Girard, E., Bangratz, M., Bernard, V., Herbin, M., Vignaud, A., Ferry, A., Davoine, C.S., Echaniz-Laguna, A., René, F. et al.** (2008). Evidence of a dosage effect and a physiological endplate acetylcholinesterase deficiency in the first mouse models mimicking Schwartz-Jampel syndrome neuromyotonia. *Hum. Mol. Genet.* **17**, 3166-3179.
- Sugar, T., Wassenhove-McCarthy, D.J., Esko, J.D., van Kuppevelt, T.H., Holzman, L., and McCarthy, K.J.** (2014). Podocyte-specific deletion of NDST1, a key enzyme in the sulfation of

heparan sulfate glycosaminoglycans, leads to abnormalities in podocyte organization in vivo.

Kidney Int. **85**, 307-318.

Suzuki, O., Kague, E., Bagatini, K., Tu, H., Heljasvaara, R., Carvalhaes, L., Gava, E., de Oliveira, G., Godoi, P., Oliva, G. et al. (2009). Novel pathogenic mutations and skin biopsy analysis in Knobloch syndrome. *Mol. Vis.* **15**, 801-809.

Suzuki, O.T., Sertié, A.L., Der Kaloustian, V.M., Kok, F., Carpenter, M., Murray, J., Czeizel, A.E., Kliemann, S.E., Rosemberg, S., Monteiro, M. et al. (2002). Molecular analysis of collagen XVIII reveals novel mutations, presence of a third isoform, and possible genetic heterogeneity in Knobloch syndrome. *Am. J. Hum. Genet.* **71**, 1320-1329.

Syu, A., Ishiguro, H., Inada, T., Horiuchi, Y., Tanaka, S., Ishikawa, M., Arai, M., Itokawa, M., Niizato, K., Iritani, S. et al. (2010). Association of the HSPG2 gene with neuroleptic-induced tardive dyskinesia. *Neuropsychopharmacology* **35**, 1155-1164.

Takahashi, I., Noguchi, N., Nata, K., Yamada, S., Kaneiwa, T., Mizumoto, S., Ikeda, T., Sugihara, K., Asano, M., Yoshikawa, T. et al. (2009). Important role of heparan sulfate in postnatal islet growth and insulin secretion. *Biochem. Biophys. Res. Commun.* **383**, 113-118.

Tang, G.L. and Weitz, K. (2015). Impaired arteriogenesis in syndecan-1(-/-) mice. *J. Surg. Res.* **193**, 22-27.

Tao, Q., Yokota, C., Puck, H., Kofron, M., Birsoy, B., Yan, D., Asashima, M., Wylie, C.C., Lin, X., and Heasman, J. (2005). Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in Xenopus embryos. *Cell* **120**, 857-871.

Thompson, W.R., Modla, S., Grindel, B.J., Czymbek, K.J., Kirn-Safran, C.B., Wang, L., Duncan, R.L., and Farach-Carson, M.C. (2011). Perlecan/Hspg2 deficiency alters the pericellular space of the lacunocanalicular system surrounding osteocytic processes in cortical bone. *J. Bone Miner. Res.* **26**, 618-629.

Topczewski, J., Sepich, D.S., Myers, D.C., Walker, C., Amores, A., Lele, Z., Hammerschmidt, M., Postlethwait, J., and Solnica-Krezel, L. (2004). The zebrafish glycan knypek controls cell polarity during gastrulation movements of convergent extension. *Dev. Cell* **1**, 251-264.

Tornberg J., Sykiotis, G.P., Keefe, K., Plummer, L., Hoang, X., Hall, J.E., Quinton, R., Seminara, S.B., Hughes, V., Van Vliet, G. et al. (2011). Heparan sulfate 6-O-sulfotransferase 1, a

- gene involved in extracellular sugar modifications, is mutated in patients with idiopathic hypogonadotropic hypogonadism. *Proc. Natl. Acad. Sci. USA* **108**, 11524-11529.
- Tran, P.K., Tran-Lundmark, K., Soininen, R., Tryggvason, K., Thyberg, J., and Hedin, U.** (2004). Increased intimal hyperplasia and smooth muscle cell proliferation in transgenic mice with heparan sulfate-deficient perlecan. *Circ. Res.* **94**, 550-558.
- Tran, T.H., Shi, X., Zaia, J., and Ai, X.** (2012). Heparan sulfate 6-O-endosulfatases (Sulfs) coordinate the Wnt signaling pathways to regulate myoblast fusion during skeletal muscle regeneration. *J. Biol. Chem.* **287**, 32651-32664.
- Trowe, T., Klostermann, S., Baier, H., Granato, M., Crawford, A.D., Grunewald, B., Hoffmann, H., Karlstrom, R.O., Meyer, S.U., Müller, B. et al.** (1996). Mutations disrupting the ordering and topographic mapping of axons in the retinotectal projection of the zebrafish, *Danio rerio*. *Development* **123**, 439-450.
- Tzeng, S.T., Tsai, M.H., Chen, C.L., Lee, J.X., Jao, T.M., Yu, S.L., Yen, S.J., and Yang, Y.C.** (2013). NDST4 is a novel candidate tumor suppressor gene at chromosome 4q26 and its genetic loss predicts adverse prognosis in colorectal cancer. *PLoS One* **8**, e67040.
- Utriainen, A., Sormunen, R., Kettunen, M., Carvalhaes, L.S., Sajanti, E., Eklund, L., Kauppinen, R., Kitten, G.T., and Pihlajaniemi, T.** (2004). Structurally altered basement membranes and hydrocephalus in a type XVIII collagen deficient mouse line. *Hum. Mol. Genet.* **13**, 2089-2099.
- Venero Galanternik, M., Kramer, K.L., and Piotrowski, T.** (2015). Heparan Sulfate Proteoglycans Regulate Fgf Signaling and Cell Polarity during Collective Cell Migration. *Cell Rep.* pii: S2211-1247(14)01096-1.
- Veugelers, M., Cat, B.D., Muyldermans, S.Y., Reekmans, G., Delande, N., Frints, S., Legius, E., Fryns, J.P., Schrander-Stumpel, C., Weidle, B. et al.** (2000). Mutational analysis of the GPC3/GPC4 glycan gene cluster on Xq26 in patients with Simpson-Golabi-Behmel syndrome: identification of loss-of-function mutations in the GPC3 gene. *Hum. Mol. Genet.* **9**, 1321-1328.
- Viviano, B.L., Silverstein, L., Pfleiderer, C., Paine-Saunders, S., Mills, K., and Saunders, S.** (2005). Altered hematopoiesis in glycan-3-deficient mice results in decreased osteoclast differentiation and a delay in endochondral ossification. *Dev. Biol.* **282**, 152-162.

- Walker, K.A., Sims-Lucas, S., Caruana, G., Cullen-McEwen, L., Li, J., Sarraj, M.A., Bertram, J.F., and Stenvers, K.L.** (2011). Betaglycan is required for the establishment of nephron endowment in the mouse. *PLoS One* **6**, e18723.
- Wang, F., Wolfson, S.N., Gharib, A., and Sagasti, A.** (2012). LAR receptor tyrosine phosphatases and HSPGs guide peripheral sensory axons to the skin. *Curr. Biol.* **22**, 373-382.
- Wang, K.S., Wang, L., Liu, X., and Zeng, M.** (2013). Association of HS6ST3 gene polymorphisms with obesity and triglycerides: gene x gender interaction. *J. Genet.* **92**, 395-402.
- Wang, L., Fuster, M., Sriramrao, P., and Esko, J.D.** (2005). Endothelial heparan sulfate deficiency impairs L-selectin- and chemokine-mediated neutrophil trafficking during inflammatory responses. *Nat. Immunol.* **6**, 902-910.
- Wang, Q., Yang, L., Alexander, C., and Temple, S.** (2012). The niche factor syndecan-1 regulates the maintenance and proliferation of neural progenitor cells during mammalian cortical development. *PLoS One* **7**, e42883.
- Wang, S., Ai, X., Freeman, S.D., Pownall, M.E., Lu, Q., Kessler, D.S., and Emerson, C.P. Jr.** (2004). QSulf1, a heparan sulfate 6-O-endosulfatase, inhibits fibroblast growth factor signaling in mesoderm induction and angiogenesis. *Proc. Natl. Acad. Sci. USA* **101**, 4833-4838.
- Wang, W., Zhong, B., Sun, J., Cao, J., Tian, J., Zhong, N., Zhao, W., Tian, L., Xu, P., Guo, D. et al.** (2011). Down-regulated HS6ST2 in osteoarthritis and Kashin-Beck disease inhibits cell viability and influences expression of the genes relevant to aggrecan metabolism of human chondrocytes. *Rheumatology (Oxford)* **50**, 2176-2186.
- Wei, K.H. and Liu, I.H.** (2014). Heparan sulfate glycosaminoglycans modulate migration and survival in zebrafish primordial germ cells. *Theriogenology* **81**, 1275-1285.
- Wilson, N.H. and Stoeckli, E.T.** (2013). Sonic hedgehog regulates its own receptor on postcrossing commissural axons in a glypican1-dependent manner. *Neuron* **79**, 478-491.
- Wiweger, M.I., Avramut, C.M., de Andrea, C.E., Prins, F.A., Koster, A.J., Ravelli, R.B., and Hogendoorn, P.C.** (2011). Cartilage ultrastructure in proteoglycan-deficient zebrafish mutants brings to light new candidate genes for human skeletal disorders. *J. Pathol.* **223**, 531-542.

- Wiweger, M.I., Zhao, Z., van Merkesteyn, R.J., Roehl, H.H., and Hogendoorn, P.C.** (2012). HSPG-deficient zebrafish uncovers dental aspect of multiple osteochondromas. *PLoS One* **7**, e29734.
- Wuyts, W., Van Hul, W., De Boulle, K., Hendrickx, J., Bakker, E., Vanhoenacker, F., Mollica, F., Lüdecke, H.J., Sayli, B.S., Pazzaglia, U.E. et al.** (1998). Mutations in the EXT1 and EXT2 genes in hereditary multiple exostoses. *Am. J. Hum. Genet.* **62**, 346-354.
- Xiao, T. and Baier, H.** (2007). Lamina-specific axonal projections in the zebrafish tectum require the type IV collagen Dragnet. *Nat. Neurosci.* **10**, 1529-1537.
- Xiong, D.H., Liu, X.G., Guo, Y.F., Tan, L.J., Wang, L., Sha, B.Y., Tang, Z.H., Pan, F., Yang, T.L., Chen, X.D. et al.** (2009). Genome-wide association and follow-up replication studies identified ADAMTS18 and TGFBR3 as bone mass candidate genes in different ethnic groups. *Am. J. Hum. Genet.* **84**, 388-398.
- Xu, J., Park, P.W., Kheradmand, F., and Corry, D.B.** (2005). Endogenous attenuation of allergic lung inflammation by syndecan-1. *J. Immunol.* **174**, 5758-5765.
- Xu, Z., Ichikawa, N., Kosaki, K., Yamada, Y., Sasaki, T., Sakai, L.Y., Kurosawa, H., Hattori, N., and Arikawa-Hirasawa, E.** (2010). Perlecan deficiency causes muscle hypertrophy, a decrease in myostatin expression, and changes in muscle fiber composition. *Matrix Biol.* **29**, 461-470.
- Yamamoto, A., Uchiyama, K., Nara, T., Nishimura, N., Hayasaka, M., Hanaoka, K., and Yamamoto, T.** (2014). Structural abnormalities of corpus callosum and cortical axonal tracts accompanied by decreased anxiety-like behavior and lowered sociability in spock3- mutant mice. *Dev. Neurosci.* **36**, 381-395.
- Yasuda, T., Mundy, C., Kinumatsu, T., Shibukawa, Y., Shibutani, T., Grobe, K., Minugh-Purvis, N., Pacifici, M., and Koyama, E.** (2010). Sulfotransferase Ndst1 is needed for mandibular and TMJ development. *J. Dent. Res.* **89**, 1111-1116.
- Ylikärppä, R., Eklund, L., Sormunen, R., Kontiola, A.I., Utriainen, A., Määttä, M., Fukai, N., Olsen, B.R., and Pihlajaniemi, T.** (2003). Lack of type XVIII collagen results in anterior ocular defects. *FASEB J.* **17**, 2257-2259.

- Zak, B.M., Schuksz, M., Koyama, E., Mundy, C., Wells, D.E., Yamaguchi, Y., Pacifici, M., and Esko, J.D.** (2011). Compound heterozygous loss of Ext1 and Ext2 is sufficient for formation of multiple exostoses in mouse ribs and long bones. *Bone* **48**, 979-987.
- Zcharia, E., Metzger, S., Chajek-Shaul, T., Aingorn, H., Elkin, M., Friedmann, Y., Weinstein, T., Li, J.P., Lindahl, U., Vlodavsky, I.** (2004). Transgenic expression of mammalian heparanase uncovers physiological functions of heparan sulfate in tissue morphogenesis, vascularization, and feeding behavior. *FASEB J.* **18**, 252-263.
- Zcharia, E., Philp, D., Edovitsky, E., Aingorn, H., Metzger, S., Kleinman, H.K., Vlodavsky, I., Elkin, M.** (2005a) Heparanase regulates murine hair growth. *Am. J. Pathol.* **166**, 999-1008.
- Zcharia, E., Zilka, R., Yaar, A., Yacoby-Zeevi, O., Zetser, A., Metzger, S., Sarid, R., Naggi, A., Casu, B., Ilan, N. et al.** (2005b). Heparanase accelerates wound angiogenesis and wound healing in mouse and rat models. *FASEB J.* **19**, 211-221.
- Zcharia, E., Jia, J., Zhang, X., Baraz, L., Lindahl, U., Peretz, T., Vlodavsky, I., Li, J.P.** (2009). Newly generated heparanase knock-out mice unravel co-regulation of heparanase and matrix metalloproteinases. *PLoS One* **4**, e5181.
- Zhang, B., Xiao, W., Qiu, H., Zhang, F., Moniz, H.A., Jaworski, A., Condac, E., Gutierrez-Sanchez, G., Heiss, C., Clugston, R.D. et al.** (2014). Heparan sulfate deficiency disrupts developmental angiogenesis and causes congenital diaphragmatic hernia. *J. Clin. Invest.* **124**, 209-221.
- Zhang, X., Wu, C., Song, J., Götte, M., Sorokin, L.** (2013). Syndecan-1, a cell surface proteoglycan, negatively regulates initial leukocyte recruitment to the brain across the choroid plexus in murine experimental autoimmune encephalomyelitis. *J. Immunol.* **191**, 4551-4556.
- Zhang, Y., Li, J., Partovian, C., Sellke, F.W., Simons, M.** (2003). Syndecan-4 modulates basic fibroblast growth factor 2 signaling in vivo. *Am. J. Physiol. Heart. Circ. Physiol.* **284**, H2078-2082.
- Zheng, Q., Zhu, J., Shanabrough, M., Borok, E., Benoit, S.C., Horvath, T.L., Clegg, D.J., Reizes, O.** (2010). Enhanced anorexigenic signaling in lean obesity resistant syndecan-3 null mice. *Neuroscience* **171**, 1032-1040.
- Zoeller, J.J., McQuillan, A., Whitelock, J., Ho, S.Y., Iozzo, R.V.** (2008). A central function for perlecan in skeletal muscle and cardiovascular development. *J. Cell Biol.* **181**, 381-394.

Zoeller, J.J., Whitelock, J.M., Iozzo, R.V. (2009). Perlecan regulates developmental angiogenesis by modulating the VEGF-VEGFR2 axis. *Matrix Biol.* **28**, 284-291.

Zuberi, R.I., Ge, X.N., Jiang, S., Bahaei, N.S., Kang, B.N., Hosseinkhani, R.M., Frenzel, E.M., Fuster, M.M., Esko, J.D., Rao, S.P., Sriramaraao, P. (2009). Deficiency of endothelial heparan sulfates attenuates allergic airway inflammation. *J. Immunol.* **183**, 3971-3979.