

RESEARCH ARTICLE

Translation repression by maternal RNA binding protein Zar1 is essential for early oogenesis in zebrafish

Liyun Miao^{1,2,3,4}, Yue Yuan^{1,3,4}, Feng Cheng^{1,*}, Junshun Fang¹, Fang Zhou¹, Weirui Ma^{1,‡}, Yan Jiang⁵, Xiahe Huang¹, Yingchun Wang¹, Lingjuan Shan¹, Dahua Chen⁶ and Jian Zhang^{1,3,4,§}

ABSTRACT

A large amount of maternal RNA is deposited in oocytes and is reserved for later development. Control of maternal RNA translation during oocyte maturation has been extensively investigated and its regulatory mechanisms are well documented. However, translational regulation of maternal RNA in early oogenesis is largely unexplored. In this study, we generated zebrafish *zar1* mutants that result in early oocyte apoptosis and fully penetrant male development. Loss of p53 suppresses the apoptosis in *zar1* mutants and restores oocyte development. *zar1* immature ovaries show upregulation of proteins implicated in endoplasmic reticulum (ER) stress and the unfolded protein response (UPR). More importantly, loss of Zar1 causes marked upregulation of zona pellucida (ZP) family proteins, while overexpression of ZP proteins in oocytes causes upregulation of stress-related *activating transcription factor 3* (*atf3*), arguing that tightly controlled translation of ZP proteins is essential for ER homeostasis during early oogenesis. Furthermore, Zar1 binds to ZP gene mRNAs and represses their translation. Together, our results indicate that regulation of translational repression and de-repression are essential for precisely controlling protein expression during early oogenesis.

KEY WORDS: Zar1, Zona pellucida, Oogenesis, p53, Zebrafish, Translational control, ER stress, Unfolded protein response, Apoptosis, Sex differentiation

INTRODUCTION

In contrast to spermatogenesis, oogenesis is associated with increasing cell volume and accumulating maternal molecules such as RNAs and proteins, which are essential for oogenesis itself and early embryogenesis. Maternal gene expression is temporally and spatially controlled (Becalska and Gavis, 2009; Curtis et al., 1995; Evans and Hunter, 2005; Li et al., 2010; Richter and Lasko, 2011). In most species, maternal RNAs play central roles during early embryogenesis owing to the absence of zygotic transcription at initial embryonic stages. To accomplish their functions during early embryogenesis, some of the maternal RNAs are extensively regulated post-transcriptionally during

oocyte maturation and early embryogenesis. For example, maternal mRNA translation can be regulated by cytoplasmic polyadenylation (Richter, 2007; Richter and Lasko, 2011). Maternal RNA stability can also be regulated by micro RNAs, 3' untranslated region (UTR) length and even codon usage (Barckmann and Simonelig, 2013; Mishima and Tomari, 2016; Norbury, 2013), which can further temporally control protein expression.

In addition to cytoplasmic components, oocytes also maintain complex cell surface structures including the glycoprotein-rich zona pellucida (ZP), which surrounds oocyte plasma membrane and is vital for oocyte integrity. At fertilization in mouse, ZP proteins (such as ZP2 and ZP3) are required to bind to spermatozoa and to initiate acrosome reaction (Conner et al., 2005). FIG α activates transcription of the mouse genes *Zp1*, *Zp2* or *Zp3* through an E-box in their promoters (Liang et al., 1997; Soyal et al., 2000). In zebrafish, the vitelline envelope (equivalent to mammalian ZP) starts to form in stage IB follicles. ZP glycoproteins are encoded by multicopy ZP genes whose expression is tightly regulated at transcriptional and translational levels, yet their regulators remain unclear (Mold et al., 2009).

Zebrafish oocyte development is closely associated with sex determination (Liew and Orban, 2013). Adult zebrafish possess only testes or ovaries, but juveniles are initially bipotential. Undifferentiated gonads are ovary-like with early stage oocytes (Maack and Segner, 2003; Takahashi, 1977). These oocytes grow to full-size in females, but degenerate and are replaced by spermatogenesis in males (Uchida et al., 2002). Zebrafish without germ cells develop into male adults, indicating germ cells are essential for female development (Campbell et al., 2015; Draper et al., 2007; Houwing et al., 2007; Siegfried and Nüsslein-Volhard, 2008; Slanchev et al., 2005). When oocytes are depleted in juveniles, zebrafish also develop into males (Dranow et al., 2013; Hartung et al., 2014; Houwing et al., 2008; Rodríguez-Marí et al., 2010, 2011; Shive et al., 2010; White et al., 2011). Attenuation of apoptosis pathways by p53 deficiency restores ovarian development in otherwise all male *fancl* and *brca2* mutants (Rodríguez-Marí et al., 2010, 2011; Shive et al., 2010). In addition, germ cells and oocytes are also required to maintain female phenotype in adult zebrafish (Dranow et al., 2016, 2013). Separately, activation of the estrogen pathway suppresses apoptosis in mouse and fish ovaries (Janz and Van Der Kraak, 1997; Kim et al., 2009). Estrogen is also essential for zebrafish female development (Dranow et al., 2016; Guiguen et al., 2010). Furthermore, juvenile fish exposed to estrogen are likely to become females (Larsen et al., 2009; Örn et al., 2003).

Maternal genes have been studied in a number of model organisms. *Zar1* (*zygotic arrest 1*) was initially identified as a maternal gene in mouse. *Zar1* null female mice generate fully grown oocytes and the eggs can be fertilized, but the resulting embryos fail to develop beyond the 2-cell stage, suggesting that *Zar1* is required for very early embryogenesis (Wu et al., 2003a). The molecular

¹State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China. ²University of Chinese Academy of Sciences, Beijing 100101, China.

³Center for Life Sciences; School of Life Sciences, Yunnan University, Kunming 650500, China. ⁴State Key Laboratory for Conservation and Utilization of Bio-Resources, Kunming 650500, China. ⁵Jilin University, Changchun 130012, China.

⁶Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China. *Present address: University of California, San Francisco, CA 94143, USA.

[‡]Present address: Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA.

§Author for correspondence (jianzhang84@genetics.ac.cn)

id J.Z., 0000-0002-5867-9808

regulatory mechanism of ZAR1 in mouse oocytes is largely unknown, but *in vitro* results suggest that its *Xenopus* homolog may function as a RNA binding protein to regulate RNA translation (Yamamoto et al., 2013). ZAR1 proteins are conserved in vertebrates and contain an atypical plant homeodomain (PHD) zinc finger (Znf) domain in the C-terminus (Wu et al., 2003b). Here, we show that Zar1 functions by repressing ZP gene translation, thereby preventing overload of ZP proteins in early oocytes. Our results indicate that Zar1 is essential for early oogenesis.

RESULTS

Zebrafish *zar1* null mutants exclusively develop into adult males

To study the function of Zar1, we first examined *zar1* mRNA expression. Zebrafish *zar1* is expressed in ovaries but absent in testes (Fig. 1A). *zar1* mRNA is highly expressed in primary growth (stage I) and cortical alveolus (stage II) oocytes (oocyte staging according to Selman et al., 1993) (Fig. 1A). Zar1 protein was also detected in Stage I and II oocytes but not in Stage III oocytes (Fig. 1B). We generated several *zar1* mutant lines using TALEN (Fig. 1C–E) and CRISPR/Cas9 (Fig. S1) technology (Chang et al., 2013; Huang et al., 2011; Li et al., 2011). We focused on the *zar1*^{gd5} (Fig. 1E) and *zar1*^{gd6} mutants (Fig. S1C). Immunoblotting results indicate that Zar1 protein is absent in both *zar1*^{gd5/gd5} (Fig. 1F) and *zar1*^{gd6/gd6} homozygotes (Fig. S1D).




Fig. 1. Generation of a *zar1* mutant with TALENs in zebrafish. (A) *In situ* hybridization on cryosections of wild-type gonads with *zar1* antisense probe and sense probe. Scale bar: 200 μ m. (B) Immunoblotting to detect Zar1 protein in stage I, stage II and stage III oocytes. Ten oocytes were lysed for each stage. (C) The TALEN sequences for *zar1* mutant generation. (D) Digestion of PCR products from wild-type (WT) and *zar1* TALEN mRNA injected embryos (TLN) with *NdeI* restriction enzyme. (E) DNA sequences of the *zar1*^{gd5} mutant fish line. A premature stop codon was generated. (F) Western blot of Zar1 in gonads from WT, heterozygotes (hetero) and homozygotes (homo) at 25 dpf. DNA sequences highlighted in yellow are the start codon; red, *NdeI* recognition sites; green, premature stop codon.

We intercrossed *zar1*^{gd5/+} heterozygotes to obtain *zar1*^{gd5/gd5} homozygotes. No morphological difference was observed between *zar1*^{gd5/gd5} homozygotes and siblings during embryonic and juvenile stages (data not shown). To our surprise, we could not detect adult female *zar1* homozygotes based on external morphology. In contrast, sex ratios in sibling wild-type and heterozygotes were normal (Fig. 2A). Furthermore, histological analysis of adult gonads indicated no ovary in *zar1* homozygotes (Fig. 2B–D). It should be noted that all *zar1*^{gd5/gd5} homozygous males were fertile. There is no histological difference in testes between the homozygotes and siblings (Fig. 2C,D). Analysis of *zar1*^{gd6/gd6} homozygotes further suggests that loss of Zar1 causes the all-male phenotype (Fig. S1E). Results in this study were generated using *zar1*^{gd5}, unless otherwise indicated.

To confirm that the all-male phenotype was due to Zar1 deficiency, we used an oocyte-specific *zp3b* (*zpc*) promoter (Onichtchouk et al., 2003) to drive *zar1* transgene expression in oocytes. *EGFP* driven by the heart-specific *cmlc2* promoter was inserted into the transgenic construct *Tg(zp3b:zar1,cmlc2:EGFP)* to facilitate screening of transgenic zebrafish (Fig. 2E). *EGFP*-positive embryos, indicating embryos carrying *Tg(zp3b:zar1,cmlc2:EGFP)*, were selected at 48–72 h post fertilization (Fig. S2A–A'). *EGFP*-positive *zar1* homozygous males were crossed with *EGFP*-positive *zar1* heterozygous females. As expected, all *EGFP*-negative *zar1*^{-/-} homozygotes were male. In contrast, females were recovered from *EGFP*-positive *zar1*^{-/-} homozygotes (Fig. 2F). Accordingly Zar1 expression, albeit at relatively low level, was detected in *EGFP* positive *zar1*^{-/-} homozygotes (Fig. S2B). Ovaries of the rescued *zar1* homozygous females were morphologically similar to those of the heterozygous females (Fig. 2G,H). Together, these results confirm that the all-male phenotype results from Zar1 deficiency.

Loss of Zar1 results in female-to-male sex reversal

Two possibilities could account for the all-male phenotype of *zar1* mutants: *zar1* homozygous females might have died during development or *zar1* homozygous females could have reversed to males. To resolve this issue, we crossed *zar1* heterozygous females with *zar1* homozygous males. If the all-male phenotype is caused by female lethality, the number of homozygotes should be about 50% of the number of heterozygotes, assuming the allele segregates in a Mendelian manner. If it is caused by sex-reversal, the numbers of homozygotes and heterozygotes should be similar. The observed survival rate was 91.6% (174 out of 190), indicating that the all-male phenotype in *zar1* mutants is not caused by female lethality. The number of *zar1* homozygotes was similar to that of *zar1* heterozygotes (Fig. 3A). Among heterozygotes, about 50% were female. As expected, no females were observed among *zar1* homozygotes (Fig. 3A). These results indicate that the all-male phenotype in *zar1* mutants is caused by female-to-male sex reversal.

To better understand the sex reversal process in *zar1* homozygotes, we examined their gonads at different developmental stages. Undifferentiated gonads in *zar1* homozygotes resembled those in control siblings (heterozygotes and wild type) at 22 days post fertilization (dpf) (Fig. 3B,C). Only stage I oocytes were detected in these undifferentiated gonads. At 33 dpf, sex is determined, but gonads are immature (Rodríguez-Marí et al., 2010). At 33 dpf, similar to the control gonads, *zar1* mutant gonads were either immature ovaries (Fig. 3D,E) or immature testes (Fig. 3F,G). From 22 dpf to 33 dpf, oocyte volume increased similarly both in *zar1* mutants and in control siblings. Most oocytes in *zar1* homozygotes resembled the control oocytes. However aberrant vesicles started to appear in *zar1* mutant oocytes (Fig. 3E, arrows), suggesting that




Fig. 2. Loss of *Zar1* causes all-male phenotype in zebrafish. (A) Analysis of genders of *zar1* homozygotes (homo), heterozygotes (hetero) and wild-type siblings. (B-D) Histological analysis of gonads of *zar1* homozygotes and sibling controls by H&E staining. About half of the gonads (20/37) from wild-type and *zar1* heterozygotes were ovaries (B) while the other half (17/37) were testes (C). All the 13 gonads from *zar1* homozygotes were testes (D). (E) Schematic diagram of the *Tg(zp3b:zar1)* transgenic construct. *egfp* coding sequences were placed under control of the *cmlc2* promoter to help visually identify transgenic fish. (F) Gender analysis of *zar1*^{-/-} homozygotes with or without EGFP signal, indicating *Tg(zp3b:zar1)* transgene. Females were recovered only from *zar1*^{-/-} homozygotes with the *Tg(zp3b:zar1)* transgene. (G,H) H&E staining of ovaries of *zar1*^{+/-} heterozygotes and *zar1*^{-/-} homozygotes on the *Tg(zp3b:zar1)* transgenic backgrounds. Ovaries from *zar1*^{-/-} homozygotes rescued with the *Tg(zp3b:zar1)* transgene are normal histologically. *Tg(zp3b:zar1)*, *Tg(zp3b:zar1)*, *cmlc2:EGFP*; sg, spermatogonia; sc, spermatocytes; sp, sperm; I,II,III, oocyte stage I, II or III. Scale bars: 40 μ m.

ovarian development defects occurred as early as 33 dpf. At 39 dpf, stage II oocytes appeared in *zar1* heterozygous ovaries and wild-type ovaries (Fig. 3H). In contrast, oocyte growth appeared arrested in the mutants (Fig. 3I) with obvious aberrant vesicles (Fig. 3I, arrows). The vesicles resemble cortical granules (CGs). To identify the nature of the vesicles, we used *Maclura pomifera* agglutinin (MPA), which recognizes CG contents (Becker and Hart, 1999). The aberrant vesicles in the mutant oocytes were indeed MPA positive (Fig. 3L), indicating they are CGs. The CGs in the mutants are larger than those in sibling controls. In addition, the CGs are found in smaller oocytes in the mutants, suggesting that they accumulate prematurely. We also noticed that CG number in mutant oocytes is far less than that in sibling oocytes (Fig. 3M).

At 50 dpf, *zar1* mutant ovaries became ovotestis, containing both oocytes and spermatocytes (Fig. 3N,O). At 60 dpf, when oocytes reached stage III in control ovaries, spermatogenesis dominated in the *zar1* mutant ovotestis with a few residual stage I oocytes (Fig. 3P,Q). Testis development in *zar1* mutants resembled that in sibling controls (Fig. 3F,G,J,K, and Fig. S3). The percentage of immature ovaries at 33-41 dpf in *zar1* homozygotes (54.5%) is similar to that in *zar1* heterozygous controls (56.9%) (Fig. S4). These results confirm that the all-male phenotype in *zar1* mutants is due to female-to-male sex reversal.

Apoptosis in *zar1* mutants is mediated through the p53 pathway

We hypothesized that degeneration of *zar1* homozygous oocytes is mediated by apoptosis. To test this, we used the TUNEL assay to examine apoptosis in immature ovaries at 37-40 dpf. Few apoptotic

cells could be seen in immature ovaries of *zar1* heterozygotes but apoptosis was readily detected in *zar1* homozygous ovaries (Fig. 4A-C). p53-mediated germ cell apoptosis causes gonad transformation in several zebrafish mutants (Rodríguez-Mari et al., 2010, 2011; Shive et al., 2010). To test whether apoptosis in *zar1* homozygotes is mediated by p53 (also known as Tp53), we crossed *zar1*^{-/-} males with *p53*^{M214K/M214K} (*p53*^{-/-}) females and then crossed *p53*^{+/-}; *zar1*^{+/-} F1 progenies. Of the *p53*^{-/-}; *zar1*^{-/-} double mutant offspring, 8 out of 14 were female while all the other 28 mutant siblings (*p53*^{+/-}; *zar1*^{-/-} or *p53*^{+/-}; *zar1*^{-/-}) were male (Fig. 4D), suggesting oogenesis arrest in *zar1*^{-/-} mutants is suppressed by p53 deficiency. The ovaries of the *p53*^{-/-}; *zar1*^{-/-} double mutant were indistinguishable from control ovaries histologically (Fig. 4E,F), but the chorions of *p53*^{-/-}; *zar1*^{-/-} double mutant eggs failed to lift upon activation (Fig. S5). As expected, apoptosis was also blocked in *p53*^{-/-}; *zar1*^{-/-} double mutant ovaries (Fig. 4G-J). Thus, p53-mediated apoptosis contributes to oocyte degeneration in the *zar1*^{-/-} mutants.

Estrogen treatment restores oogenesis in *zar1* mutants

Estrogen is required for zebrafish female development (Dranow et al., 2016; Guiguen et al., 2010). Furthermore, juvenile fish exposed to 17 α -ethinylestradiol (EE2), a synthetic estrogen agonist, usually develop into females (Örn et al., 2003). To explore whether estrogen can affect *zar1* mutant ovarian development, we used EE2 (10 ng/l, 20-60 dpf) to treat *zar1* homozygous juveniles. Females were recovered from EE2-treated *zar1* homozygous mutants, although the percentage of female in *zar1* homozygotes is not as high as that in *zar1* heterozygotes (Fig. 5A). No obvious




Fig. 3. Zar1 deficiency causing all-male phenotype is due to female-to-male sex reversal. (A) Analysis of genders of *zar1* homozygotes (homo) and heterozygous siblings (hetero). *zar1*^{+/-} heterozygous females were crossed with *zar1*^{-/-} homozygous males and genders of their progenies were analyzed. (B-K,N-Q) Gonad development of *zar1* homozygotes and control siblings analyzed by H&E staining. At 22 dpf, zebrafish gonads are undifferentiated. Gonads of *zar1* homozygotes and control siblings are indistinguishable histologically (B,C). At 33 dpf, WT gonads differentiate into immature ovaries and immature testes. Minor developmental abnormalities are observed in *zar1* homozygotes. Oocytes in *zar1* homozygotes are similar to those in WT in size and morphology, but aberrant vesicles (arrow indicated) are observed in ooplasm of a few *zar1* homozygous oocytes (D,E). Testis development in *zar1* homozygotes is normal compared with that in control siblings (F,G). At 39 dpf, ovarian developmental abnormality in *zar1* homozygotes becomes more pronounced (H,I), while testis development in *zar1* homozygotes is similar to the controls (J,K). Immature ovaries in control siblings contain stage I and stage II oocytes (H) while oocytes in *zar1* homozygotes resemble stage I oocytes, indicating oogenesis arrest in *zar1* homozygotes. In addition, aberrant vesicles (arrow) in *zar1* homozygotes increase significantly in size and number. (L) MPA (*Maclura pomifera* agglutinin) staining of ovary sections. Juveniles at 37-40 dpf fixed with 4% PFA were embedded in paraffin and sections stained with MPA; 0.5 M D-galactose inhibited MPA staining. MPA specifically stains CGs in control siblings. The aberrant vesicles in *zar1* mutant ovaries are MPA positive (arrows). (M) Comparison of CG numbers (mean±s.e.m.) between *zar1* mutants (*n*=11) and siblings (*n*=11). At 50 dpf, stage II oocytes are observed in heterozygous and wild-type ovaries (N), but large numbers of oocytes underwent atresia in *zar1* homozygotes, and spermatogonia and spermatocytes appear among oocytes, indicating transitional ovaries (ovotestis) (O). At 60 dpf, stage III oocytes were seen in the control ovaries (P); in contrast, very few oocytes remain in ovotestis of *zar1* mutants and spermatogenesis has progressed further (Q). At 22 dpf, six juveniles were analyzed for each genotype. At 33 dpf, 39 dpf, 50 dpf and 60 dpf, 20 juveniles per stage were analyzed for each genotype. Stars indicate degenerating oocytes. sg, spermatogonia; sc, spermatocytes; sp, sperm. Scale bar: 40 μm.

histological difference could be detected between EE2-treated *zar1* mutant ovaries and control ovaries (Fig. 5B,C), yet the chorions of eggs from EE2-treated *zar1* mutants failed to lift properly upon activation (Fig. S6).

Estrogen treatment has been shown to suppress apoptosis in mouse and fish ovaries (Janz and Van Der Kraak, 1997; Kim et al., 2009). Similarly, in human breast cancer cells, estrogen downregulates the P53 target genes, including *ATF3*, *BGT2* and *TRAF4*, which are

involved in P53-mediated apoptosis (Bailey et al., 2012). We found that at 33-34 dpf, *btg2* mRNA was downregulated while *traf4a* mRNA had similar levels of expression in *zar1* mutant ovaries and control sibling ovaries (Fig. 5D). Only *atf3* was upregulated in *zar1* mutant ovaries (Fig. 5E). Moreover, EE2 treatment was able to repress the *atf3* upregulation in *zar1* mutant ovaries (Fig. 5E). These results suggest that estrogen functions through suppression of stress-related genes to restore oogenesis.




Fig. 4. Zar1 deficiency causing all-male phenotype is due to p53-mediated apoptosis. (A-C) TUNEL staining of ovary sections of *zar1* homozygotes and control siblings at 37-40 dpf. Obvious apoptotic cells are observed in immature ovaries in *zar1* homozygotes (A), but not in sibling controls (B). Quantification of apoptotic cells is shown in C; 18 sections from 6 juveniles were counted for each genotype. (D) Gender analysis of *zar1*^{-/-} homozygotes on different *p53* genotype backgrounds (*p53*^{+/+}, *p53*^{+/-} and *p53*^{-/-}). Females are only observed in *p53*^{-/-}; *zar1*^{-/-} double homozygous mutants. (E,F) H&E staining of sections of *p53*^{-/-}; *zar1*^{+/-} and *p53*^{-/-}; *zar1*^{-/-} adult ovaries. *p53*^{-/-}; *zar1*^{-/-} ovaries are morphologically normal. (G-J) TUNEL staining of ovary sections of *p53*^{-/-}; *zar1*^{-/-} ovaries at 37-40 dpf with *p53*^{-/-}; *zar1*^{+/-} and *p53*^{+/-}; *zar1*^{-/-} ovaries as controls. (J) Quantification of apoptotic cells. Nine sections from three juveniles were counted for each genotype. Scale bar: 20 μ m.

Loss of Zar1 triggers ER stress and the unfolded protein response (UPR)

The ATF4-ATF3-CHOP cascade has been implicated in ER stress and the UPR (Jiang et al., 2004; Schröder and Kaufman, 2005). ER is involved in protein folding, post-translational modification and secretory activities. ER homeostasis is essential for normal cell functions (Wang and Kaufman, 2012). ER homeostasis can be disrupted by misfolded proteins and abnormally elevated secretory protein synthesis. Under ER stress, cells activate the UPR to alleviate ER burden by reducing protein translation, increasing protein degradation and generating additional chaperones to assist protein folding. When the UPR fails to restore ER homeostasis, cells may undergo apoptosis (Breckenridge et al., 2003; Shore et al., 2011; Szegezdi et al., 2006). The UPR functions through three major pathways, initiated by three ER-localized transmembrane proteins, to restore ER homeostasis. One of them is initiated by protein kinase RNA-like ER kinase (PERK). Activation of PERK contributes to the accumulation of activating transcription factor 4 (ATF4) (Bettigole and Glimcher, 2015; Sano and Reed, 2013), which upregulates ATF3 and CHOP expression. We checked the expression level of the CHOP-encoding gene *ddit3* and found that, like *atf3*, *ddit3* was upregulated in *zar1* mutants (Fig. 6A). These results suggest that loss of Zar1 causes ER stress in ovaries.

The lack of specific antibodies against zebrafish antigens prevents us from using immunological methods to analyze ER stress and the UPR targets at the protein level. To further study cellular stress in *zar1* mutant ovaries, we quantitatively compared proteomes of *zar1* homozygous ovaries with that of *zar1* heterozygous ones using isobaric tags for relative and absolute quantitation (iTRAQ) technology. We chose ovaries at 33 dpf based on the following considerations: (1) prior to 33 dpf, ovaries and testes are nearly indistinguishable morphologically; (2) at 33 dpf, oocyte sizes are similar in both groups yet aberrant cortical granules start to appear in the mutants, indicating initial oocyte defects. More than 5300 proteins were identified in ovaries from the two genotypes (Table S3), 325 proteins show differential expression ($P < 0.05$) (Table S4). A total of 42 proteins were increased or decreased by more than twofold, with five proteins, including Zar1, downregulated and 37 proteins upregulated in homozygous *zar1*^{-/-} ovaries. We examined seven upregulated proteins and analyzed their mRNA expression (Fig. S7). Five of them were also transcriptionally upregulated. RNA levels of the other two were similar between the two groups, suggesting that protein upregulation occurs at the post-transcriptional level.

Among the 37 upregulated proteins are ER stress- and UPR-related proteins, such as Dnajc3a, Vapb and Pdla4 (Fig. 6B and




Fig. 5. EE2 treatment restores ovarian development in *zar1* homozygous mutant. (A) Statistics for sex ratio of *zar1* homozygous and heterozygous mutants after EE2 treatment. (B,C) H&E staining of ovarian tissue in EE2-treated *zar1* homozygotes and heterozygotes. Similar to *zar1* heterozygous ovaries, the treated homozygous mutant ovaries possess oocytes at all stages. (D) *btg2* and *traf4a* expression analyzed by qPCR in *zar1* homozygotes ($n=6$) and heterozygotes ($n=6$). (E) *atf3* expression analyzed by qPCR in *zar1* homozygotes ($n=7$) and heterozygotes ($n=8$) with or without EE2 treatment. Gene expression was normalized to expression of *elongation factor 1 alpha* (*ef1a*). hetero, *zar1* heterozygotes; homo, *zar1* homozygotes; EE2-hetero, EE2-treated *zar1* heterozygotes ($n=6$); EE2-homo, EE2 treated *zar1* homozygotes ($n=6$). Data are mean \pm s.e.m. ** $P<0.01$, *** $P<0.001$; ns, not significant. Scale bar: 20 μ m.

Table S1), indicating that ER stress and the UPR occurred. If the PERK-ATF4-ATF3-CHOP UPR cascade is a major contributor for oogenesis arrest, we would expect to restore female development by inhibiting PERK activity. To this end, we used PERK inhibitors (GSK2606414 and ISRIB) to treat *zar1*^{-/-} homozygous juveniles. In both treatments, females were recovered in *zar1* mutants (Fig. 6C,D), suggesting that ER stress and the UPR play an important role in *zar1* oogenesis arrest.

Zar1 regulates translation of ZP proteins

In addition to the three upregulated ER stress-related proteins (Fig. 6A and Table S1), there are another 34 upregulated proteins, among which 23 proteins have been annotated in various databases. Of these, 17 can be classified into two categories (Table S1). Seven proteins are ZP glycoproteins (Fig. 7A), and ten proteins have been implicated in immune regulation (Fig. S8). We speculated that upregulation of ER stress, the UPR and immunity-related proteins are secondary events. We hypothesize that overexpression of ZP glycoproteins in the *zar1* mutants may overwhelm translational and post-translational machineries in early oocytes and cause ER stress and the UPR. To test this hypothesis, we overexpressed Zp3b protein in zebrafish oocytes by injection of *zp3b-flag* mRNA (Fig. 7B). Indeed, *zp3b-flag* injected oocytes showed higher *atf3* expression compared with oocytes injected with control *rfp-flag* mRNA (Fig. 7C), suggesting that ZP overexpression causes the UPR, possibly by overwhelming ER capacity.

We noticed that mRNA levels of the seven ZP genes were not upregulated in *zar1* homozygous mutants, with the *zp211* transcript level actually downregulated (Fig. 7D), indicating that upregulation of ZP protein occurs post-transcriptionally. One possible role for Zar1 to fit into this model is to function as a translational repressor. As a first step to test the hypothesis, we examined whether Zar1 protein is associated with ZP mRNAs. We first confirmed that affinity-purified Zar1 antibodies can specifically immunoprecipitate endogenous Zar1 in juvenile ovaries (Fig. 8A). We compared mRNAs precipitated by the antibody between *zar1* mutants and heterozygous siblings. The results indicate that all seven ZP mRNAs are significantly enriched in the Zar1 protein precipitate, whereas control mRNAs (*ef1a* and *gapdh*) recovered similarly between the two lysates (Fig. 8B). Furthermore, in a yeast three-hybrid assay (Bernstein et al., 2002; Chen et al., 2014), Zar1 was shown to activate *lacZ* reporter expression in the presence of either *zp211* or *zp3b* RNAs (Fig. 8C,D), suggesting direct protein-RNA interaction. To check whether the Zar1 Znf motif is required for the binding, we mutated all of the eight cysteines in the Znf domain to alanines (Fig. S9). The Zar1 Znf domain mutant (Zar1-mu) failed to activate the reporter, suggesting the Znf domain is required for the binding (Fig. 8C,D). Next, we tested whether Zar1 could repress ZP mRNA translation in zebrafish oocytes. Using a




Fig. 6. ER stress and the UPR may be responsible for deprivation of female development of *zar1* homozygous mutants. (A) Transcriptional upregulation of *ddit3*, examined with qPCR, in mutant ovaries at 33-34 dpf (hetero, $n=6$; homo, $n=6$). Internal control: *ef1a*. (B) ER stress- and UPR-related proteins are upregulated in *zar1* homozygous (homo) ovaries compared with heterozygous (hetero) ovaries at 33-34 dpf as determined by iTRAQ. (C,D) PERK inhibitor treatment restores female development in *zar1* homozygous mutants. Juveniles were treated with two PERK inhibitors from 22 dpf to 60 dpf. Final concentration: GSK2606414, 6 nM; ISRIB, 50 nM. ** $P<0.01$, *** $P<0.001$.




Fig. 7. Translational upregulation of zona pellucida proteins may cause ER stress. (A) ZP proteins are upregulated in *zar1* homozygous (homo) ovaries compared with heterozygous (hetero) ovaries at 33-34 dpf determined by iTRAQ. (B,C) Overexpression of *zp3b* in oocytes causes upregulation of *atf3*. Stage IV oocytes were injected with 600 pg *rfp* or *zp3b* mRNAs. Injected oocytes were cultured in OCM for 4 h and collected for RNA and protein extraction with TRIzol reagent. (B) Western blot to confirm the translation of injected mRNAs. (C) qPCR to detect *atf3* expression in injected oocytes. (D) qPCR to detect ZP gene transcription (hetero, $n=6$; homo, $n=6$). Internal control: *ef1a*. ** $P<0.01$, *** $P<0.001$; ns, not significant. Data are mean \pm s.e.m.

ZP-luciferase reporter, we showed that wild-type Zar1, but not Zar1- μ , downregulated ZP gene translation (Fig. 8E-G).

Next, using immunoprecipitation and mass spectrometry (MS), we identified four Zar1-interacting candidates: vitellogenin 4 (Vtg4), poly(A) binding protein cytoplasmic 1-like (Pabpc11), eukaryotic translation initiation factor 4E transporter (Eif4enif1/4E-T) and cytoplasmic polyadenylation element-binding protein 1 (Cpeb1) (Fig. S10 and Table S5). Results from a STRING database search (Franceschini et al., 2013) indicate that Zar1, Pabpc11 (ePAB), 4E-T and Cpeb1 (CPEB) share an association network (Fig. S10A). ePAB, 4E-T and CPEB are all involved in translational regulation and ovarian development (Gray et al., 2000; Guzeloglu-Kayisli et al., 2012; Kamenska et al., 2014; Kasippillai et al., 2013; Voeltz et al., 2001). The interaction of Zar1 with CPEB and ePAB has been reported in *Xenopus* (Cook and Charlesworth, 2015). We confirmed the interaction between Zar1 and 4E-T both *in vivo* and *in vitro* (Fig. S10B-D). These results suggest that Zar1 is likely to interact with translational regulatory factors to function in a common complex that regulates oogenesis.

Taken together, these results suggest that in early oocytes (stage I and II), Zar1 is required to negatively regulate ZP protein expression through direct interactions between ZP mRNA and Zar1. The lack of

Zar1 in early oocytes causes excess ZP mRNA translation, and the resulting ZP proteins overwhelm ER capacity and cause the UPR and ultimately, apoptosis (Fig. 8H).

DISCUSSION

Maternal effect genes are relatively less studied compared with somatic genes in vertebrates. Previously, maternal ZAR1 was shown to be essential for early embryogenesis and was proposed to regulate protein translation (Wu et al., 2003a; Yamamoto et al., 2013). In this study, we reveal that zebrafish Zar1 is essential for very early oogenesis. Loss of Zar1 causes early oogenesis arrest and female-to-male sex reversal. Both genetic ablation of *p53* and estrogen treatment restore oogenesis and female fertility. Mechanistically, Zar1 binds to ZP mRNAs and represses their translation, whereas ZP protein overexpression in oocytes may cause ER stress and the UPR. Our results reveal previously unappreciated functions of Zar1 during early oogenesis.

Zar1 proteins are conserved in vertebrates, but their *in vivo* targets and molecular functions were largely unknown prior to this study. There are clear functional differences between zebrafish Zar1 and its mouse homolog. In mouse, *Zar1* mutants develop normally, yet loss of maternal ZAR1 results in two-cell arrest




Fig. 8. Zar1 represses translation of ZP proteins in zebrafish oocytes. (A-D) Zar1 binds to ZP gene mRNAs. (A,B) RNA immunoprecipitation with anti-Zar1 antibody from *zar1* homozygous ovaries (homo) and heterozygous ovaries (hetero) at 33-34 dpf. (A) Zar1 protein was precipitated by anti-Zar1 antibody in heterozygous ovaries. (B) qPCR analysis of ZP gene mRNAs immunoprecipitated from heterozygous ovaries with anti-Zar1 antibodies. Relative level of immunoprecipitated *ef1a* mRNA compared with input *ef1a* mRNA was assigned as 1. (C) Analyzing interaction of zebrafish Zar1 and ZP mRNAs with a yeast three-hybrid system. (D) Western blot analysis indicates expression of Zar1 and Zar1-mu in yeast. (E-G) Zar1 represses translation of Zp proteins in oocytes. (E) Statistics of relative luciferase activity. *zar1* mRNA injection represses Zp3b translation. Mutation of Zar1 Znf domain abolishes its translational repression activity. (F) RT-PCR of injected RNA reporters. (G) Western blot analysis shows expression of RFP-Flag, Zar1 and Zar1-mu. (H) A model of translational regulation of ZP mRNAs. In early oocytes (stage I and II), Zar1 binds to ZP gene mRNAs and negatively regulates their translation. Later, Zar1 proteins are gradually degraded and the translational repression of ZP mRNAs is removed, allowing enough ZP proteins to be used in the vitelline membrane development. The lack of Zar1 in early oocytes causes excess ZP gene translation and the resulting ZP proteins overwhelm the ER, which causes the UPR and apoptosis. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; ns, not significant;

(Wu et al., 2003a). In contrast, zebrafish *zar1* mutants show oogenesis arrest and female-to-male sex reversal. We also noticed that even a very low level of *zar1* transgene expression in the homozygous mutants was sufficient to restore ovarian development. Like the *fancl* mutants, the *zar1* mutants showed p53-mediated apoptosis and female-to-male sex reversal. In both *zar1* and *fancl* mutants, p53 deficiency can restore ovarian development. The *zar1*^{-/-};*p53*^{-/-} double mutants also showed a chorion elevation defect. In contrast, it is unknown whether a similar chorion phenotype occurs in *fancl*^{-/-};*p53*^{-/-} double homozygous females. Oocyte clearing-out time is also different between *fancl* mutants (32 dpf) and *zar1* mutants (50 dpf). The difference could be due to a requirement for Fancl in overall

genome stability (Rodríguez-Marí et al., 2011), while Zar1 is involved in translational control of a limited numbers of RNAs.

Undifferentiated gonads in *zar1* homozygotes are indistinguishable from wild-type gonads, indicating that the lack of Zar1 does not affect early gonad development prior to sex differentiation. In *zar1* homozygotes, oocyte size at 33 dpf is much larger than that at 22 dpf, indicating that loss of Zar1 does not impair initial gonad differentiation into ovaries. Furthermore, immature ovary to immature testis ratio (at 33-41 dpf) in *zar1* homozygotes is approximately 1:1, which is similar to that in *zar1* heterozygotes, suggesting that Zar1 is not required for initial sex differentiation. The appearance of aberrant cortical granules in *zar1* oocytes at 33 dpf is the initial sign of defective oogenesis, but the mutant

oocytes could last more than 2 weeks before they were cleared out, suggesting that detrimental factors accumulate gradually in the oocytes. Several studies support the hypothesis that oocytes are essential for ovarian development and may suppress testis development (Dranow et al., 2016, 2013; Hartung et al., 2014; Houwing et al., 2008; Rodríguez-Marí et al., 2010, 2011; Shive et al., 2010; White et al., 2011). We hypothesize that gradual loss of oocytes leads to testis development in *zar1* homozygous females and eventually causes female-to-male sex reversal.

p53-mediated apoptosis is attributed to oocyte clearance in zebrafish *fancl* and *brca2* mutants (Rodríguez-Marí et al., 2010, 2011; Rodríguez-Marí and Postlethwait, 2011; Shive et al., 2010). Apoptosis in *zar1* mutants is also mediated by p53 (Fig. 4). In *zar1* mutant ovaries, upregulation of ER stress- and the UPR-related genes suggests that loss of Zar1 causes ER stress and the UPR. This hypothesis was partially supported by *atf3* upregulation upon overexpression of ZP proteins in oocytes (Fig. 7C). Both EE2 treatment and p53 deficiency restore ovarian development and female development in *zar1* homozygous females. Previous studies suggest that in some organisms, estrogen can suppress oocyte apoptosis (Janz and Van Der Kraak, 1997; Kim et al., 2009). Hydroxysteroid (17- β) dehydrogenase 1, an enzyme known to catalyze estrogen synthesis (Mindnich et al., 2004), is less abundant in *zar1* mutant ovaries, suggesting that loss of Zar1 may affect estrogen synthesis. EE2 represses the upregulation of *atf3* in *zar1* mutants, suggesting some crosstalk between the UPR and estrogen pathways, although additional functions of EE2 cannot be ruled out.

In *Xenopus laevis*, Zar1 binds to the translational control sequence (TCS) of *Wee1* and *Mos* mRNAs and represses their translation in immature oocytes (Yamamoto et al., 2013). However, we did not recover *Wee1* and *Mos* in the iTRAQ analysis. This may be due to the detection limitation of iTRAQ or *Wee1* and *Mos* may not be expressed during early oogenesis in zebrafish. ZP mRNAs are highly enriched in Zar1 immunoprecipitates in early ovaries (Fig. 8B). Furthermore, Zar1 probably binds to ZP mRNAs directly, as shown in the yeast three-hybrid assay (Fig. 8C). The finding that Zar1 with a mutated Znf domain failed to bind to ZP mRNAs to repress its translation in oocytes suggests that the Znf domain mediates the binding of Zar1 to ZP mRNAs. The exact molecular mechanism by which Zar1 represses ZP mRNA translation remains to be investigated. There are a number of ways that RNA binding proteins can exert their translational inhibitory functions towards their RNA targets. In general, depolyadenylation is considered to be a mechanism to repress mRNA translation, while polyadenylation is used to initiate and enhance translation (Richter, 2007). We noticed that all seven ZP mRNAs have relatively short 5'UTRs and 3'UTRs (most of them are under 100 bp), suggesting that Zar1 may not use ZP mRNA UTRs to regulate translation. However, there might be common binding motifs/secondary structures in ZP mRNAs for Zar1 binding. Bioinformatics and experimental analyses of ZP mRNA structures are warranted. More recently, Amon and colleagues showed that the RNA binding protein Rim4 can form amyloid-like aggregates that translationally repress its target RNAs in gametogenesis (Berchowitz et al., 2015). This mode of mRNA-specific repression of translation may be used in zebrafish oogenesis. Zar1 contains a putative low complexity/disordered region, which is usually found in amyloid-like proteins or proteins capable of forming hydrogel (Courchaine et al., 2016; Kato et al., 2012). Whether Zar1 can form amyloid-like structures or hydrogel will be examined experimentally.

Zar1 was reported to be associated with known translation factors, such as CPEB and ePAB (Cook and Charlesworth, 2015).

Similarly, we found that zebrafish Zar1 coimmunoprecipitated with Cpeb1, ePAB, and 4E-T. ePAB and CPEB have been shown to control mRNA translation and oogenesis (Gray et al., 2000; Guzeloglu-Kayisli et al., 2012; Voeltz et al., 2001). Human 4E-T controls mRNA decay and represses translation of bound mRNAs (Kamenska et al., 2014). Moreover, 4E-T mutations are implicated in human primary ovarian insufficiency (Kasipillai et al., 2013). Previous studies in *Xenopus laevis* show that CPEB forms an RNP complex and interacts with Pabpc11 and 4E-T (Kim and Richter, 2007; Minshall et al., 2007; Standart and Minshall, 2008). We propose that Zar1, as a component of a maternal translational complex, may recruit other translational regulators and repress ZP mRNA translation in early oocytes. Future studies will examine how the translational complex is regulated in zebrafish oocytes.

MATERIALS AND METHODS

Zebrafish and maintenance

All animal studies in this report were approved by the Institutional Review Board of the Institute of Genetics and Developmental Biology, Chinese Academy of Sciences. Zebrafish husbandry and manipulations were performed as described (Westerfield, 2000). Zebrafish *zar1* mutants were generated using the Tübingen strain with TALEN system or CRISPR/Cas9 system using primers as detailed in the supplementary Materials and methods. Genotyping for the *p53* mutant was performed as reported (Berghmans et al., 2005).

Generation of *Tg(zp3:zar1,cmlc2:EGFP)* transgenic zebrafish

The 412 bp promoter of zebrafish *zp3b* (*zpc*) (Onichtchouk et al., 2003) was amplified by PCR with reported primers and cloned in the upstream of *zar1* coding sequence. The *egfp* sequence under the control of *cmlc2* promoter was cloned into *zar1* construct in different directions. The plasmid containing *cmlc2* promoter was reported previously (Chen et al., 2010). The transgene was introduced into zebrafish genome with the *Tol2* transposon system (Kawakami et al., 2004).

qRT-PCR and statistical analysis

Total RNA was extracted from the isolated gonads using TRIzol reagent (Thermo Fisher Scientific). cDNA was synthesized from 2 μ g total RNA with M-MLV reverse transcriptase. qPCR was performed using SsoFast EvaGreen Supermix (Bio-Rad) in 10 μ l reactions. *ef1a* and *gapdh* were amplified with reported primers (McCurley and Callard, 2008). Additional primers are listed in Table S2. Ovaries from at least six fish were analyzed for each genotype and treatment. The two-tailed unpaired *t*-test was used to determine whether the difference between the two groups was significant.

RNA *in situ* hybridization on cryosections

Animals were sacrificed with standard operations (Westerfield, 2000). Ovaries and testes were isolated and fixed in 4% paraformaldehyde (PFA) overnight at 4°C. Cryosection *in situ* hybridization was performed as described (Simmons et al., 2007). The full-length *zar1* coding sequences were used to synthesize DIG-labeled probe (Roche).

Histology, immunocytochemistry and TUNEL assay

For juveniles at 22–60 dpf, heads and tails were removed and the middle body parts containing gonads were fixed in Bouin's solution (Sigma) overnight at 4°C. For adults, ovaries and testes were isolated and fixed in Bouin's solution overnight at 4°C. Fixed tissues were embedded in paraffin and sectioned at 10 μ m. Hematoxylin and Eosin (H&E) staining was then performed on the sections. Samples for TUNEL assay (Roche) were fixed in 4% PFA. A list of antibodies and sources is provided in the supplementary Materials and methods.

RNA immunoprecipitation (RIP)

Ovaries were homogenized and lysed with RIP buffer (20 mM Tris-HCl, pH 8.0, 137 mM NaCl, 10% glycerol, 0.5% NP-40, 2 mM EDTA, RNase inhibitor and protease inhibitors were added fresh) on ice for 30 min.

Antibodies were added to the supernatant and incubated for 4 h with gentle rotation at 4°C. Protein A beads were then added, incubated for 1.5 h and washed four times (10 min each). The suspension was divided into two parts for protein and RNA preparation.

Juvenile treatment with EE2 and inhibitors

EE2 (Sigma, E4876), GSK2606414 (Selleck, S7307) and ISRIB (Selleck, S7400) were diluted to working concentration with system water. Thirty juveniles were put in a 10 liter tank containing the working solution. Fish water was renewed by dripping 20 liters of working solution per day. The juveniles were treated from 20 dpf or 22 dpf to 60 dpf and then transferred to zebrafish housing system and their gender was determined 1 month later.

Oocyte isolation and microinjection

Ovaries were isolated from adult females and transferred into fish oocyte culture medium (OCM; 20 mM HEPES, 0.2 mg/ml of BSA, 75% L-15 medium, pH 7.5) (Mold et al., 2009). Oocytes were dissociated by gentle pipetting. Oocytes at desired stage were collected, microinjected and cultured in OCM. Stage II oocytes were used to test the function of Zar1. As oocytes earlier than stage IV are easily stressed during *in vitro* manipulation, only healthy stage IV oocytes were used to test the relationship between ZP gene overexpression and ER stress.

Luciferase assay

zp3b was cloned downstream of Firefly luciferase encoding sequences (*firefly*) in pCS2 vector. *firefly-zp3b*, *rfp-flag*, *zar1* and *zar1-mu* (Znf domain mutated) mRNAs were synthesized *in vitro* using the SP6 Transcription Kit (Thermo, AM1340). *Renilla* mRNAs were synthesized *in vitro* using T7 Transcription Kit (Thermo, AM1344). *firefly-zp3b* mRNAs (200 ng/μl) and *Renilla* mRNAs (70 ng/μl) were mixed with *rfp-flag* mRNAs (210 ng/μl), *zar1* mRNAs (260 ng/μl) or *zar1-mu* mRNAs (260 ng/μl). The mixed mRNAs were injected into stage II oocytes (0.2 nl per oocyte). The injected oocytes were cultured in OCM for 4 h. Luciferase assay was performed using the Dual-Luciferase Reporter Assay System (Promega, E1910).

Quantitative proteomic analysis by iTRAQ

At 33 dpf, 10 ovaries for each genotype (*zar1^{-/-}* and *zar1^{+/-}*) were pooled and homogenized in denaturing buffer (1%SDS, 1 mM sodium orthovanadate, 1 mM sodium fluoride, protease inhibitors) on ice. The iTRAQ analysis was performed as described (Bi et al., 2014; Udeshi et al., 2013) (see supplementary Materials and methods for details). The UniProt proteome sequences for *Danio rerio* were used for the database searching.

Acknowledgements

We thank Drs Bo Zhang and Jingwei Xiong for their assistance with TALEN and CRISPR/Cas9 systems. We thank Drs Anming Meng, Jinrong Peng, Jun Chen, Zhaohui Wang, and Xiushan Wu for reagents and help.

Competing interests

The authors declare no competing or financial interests.

Author contributions

L.M. and J.Z. designed the experiments and wrote the paper; L.M., Y.Y., F.C., J.F., F.Z. W.M., Y.J., X.H. and L.S. performed the experiments; L.M., X.H., Y.W., D.C. and J.Z. analyzed the data.

Funding

This work has been supported by grants from National Natural Science Foundation of China (31590830, 31471359), Ministry of Science and Technology of the People's Republic of China (2013CB945000) and the Chinese Academy of Sciences (XDA01010108).

Data availability

Raw data of the iTRAQ proteome analyses are available in Figshare under accession number 4253888 (available at: https://figshare.com/articles/Translation_Repression_by_Maternal_RNA_Binding_Protein_Zar1_is_Essential_for_Early_Oogenesis_in_Zebrafish_rar/4253888).

Supplementary information

Supplementary information available online at <http://dev.biologists.org/lookup/doi/10.1242/dev.144642.supplemental>

References

- Bailey, S. T., Shin, H., Westerling, T., Liu, X. S. and Brown, M. (2012). Estrogen receptor prevents p53-dependent apoptosis in breast cancer. *Proc. Natl. Acad. Sci. USA* **109**, 18060-18065.
- Barckmann, B. and Simonelig, M. (2013). Control of maternal mRNA stability in germ cells and early embryos. *Biochim. Biophys. Acta* **1829**, 714-724.
- Becalska, A. N. and Gavis, E. R. (2009). Lighting up mRNA localization in *Drosophila* oogenesis. *Development* **136**, 2493-2503.
- Becker, K. A. and Hart, N. H. (1999). Reorganization of filamentous actin and myosin-II in zebrafish eggs correlates temporally and spatially with cortical granule exocytosis. *J. Cell Sci.* **112**, 97-110.
- Berchowitz, L. E., Kabachinski, G., Walker, M. R., Carlile, T. M., Gilbert, W. V., Schwartz, T. U. and Amon, A. (2015). Regulated formation of an amyloid-like translational repressor governs gametogenesis. *Cell* **163**, 406-418.
- Berghmans, S., Murphey, R. D., Wienholds, E., Neuberg, D., Kutok, J. L., Fletcher, C. D. M., Morris, J. P., Liu, T. X., Schulte-Merker, S., Kanki, J. P. et al. (2005). tp53 mutant zebrafish develop malignant peripheral nerve sheath tumors. *Proc. Natl. Acad. Sci. USA* **102**, 407-412.
- Bernstein, D. S., Buter, N., Stumpf, C. and Wickens, M. (2002). Analyzing mRNA-protein complexes using a yeast three-hybrid system. *Methods* **26**, 123-141.
- Bettigole, S. E. and Glimcher, L. H. (2015). Endoplasmic reticulum stress in immunity. *Annu. Rev. Immunol.* **33**, 107-138.
- Bi, J., Wang, W., Liu, Z., Huang, X., Jiang, Q., Liu, G., Wang, Y. and Huang, X. (2014). Seipin promotes adipose tissue fat storage through the ER Ca(2+)-ATPase SERCA. *Cell Metab.* **19**, 861-871.
- Breckenridge, D. G., Germain, M., Mathai, J. P., Nguyen, M. and Shore, G. C. (2003). Regulation of apoptosis by endoplasmic reticulum pathways. *Oncogene* **22**, 8608-8618.
- Campbell, P. D., Heim, A. E., Smith, M. Z. and Marlow, F. L. (2015). Kinesin-1 interacts with Bucky ball to form germ cells and is required to pattern the zebrafish body axis. *Development* **142**, 2996-3008.
- Chang, N., Sun, C., Gao, L., Zhu, D., Xu, X., Zhu, X., Xiong, J.-W. and Xi, J. J. (2013). Genome editing with RNA-guided Cas9 nuclease in Zebrafish embryos. *Cell Res.* **23**, 465-472.
- Chen, T. F., Luo, N., Xie, H. P., Wu, X. S. and Deng, Y. (2010). Construction and expression analysis of the zebrafish heart-specific transgenic vector based on Tol2 transposable element. *Chin. J. Biotech.* **26**, 230-236.
- Chen, D., Wu, C., Zhao, S., Geng, Q., Gao, Y., Li, X., Zhang, Y. and Wang, Z. (2014). Three RNA binding proteins form a complex to promote differentiation of germline stem cell lineage in *Drosophila*. *PLoS Genet.* **10**, e1004797.
- Conner, S. J., Lefievre, L., Hughes, D. C. and Barratt, C. L. (2005). Cracking the egg: increased complexity in the zona pellucida. *Hum. Reprod.* **20**, 1148-1152.
- Cook, J. and Charlesworth, A. (2015). The developmentally important RNA-binding protein, Zygote arrest (Zar), Regulates mRNA translation. *FASEB J.* **29** (1 Supplement), 711.18.
- Courchaine, E. M., Lu, A. and Neugebauer, K. M. (2016). Droplet organelles? *EMBO J.* **35**, 1603-1612.
- Curtis, D., Lehmann, R. and Zamore, P. D. (1995). Translational regulation in development. *Cell* **81**, 171-178.
- Dranow, D. B., Tucker, R. P. and Draper, B. W. (2013). Germ cells are required to maintain a stable sexual phenotype in adult zebrafish. *Dev. Biol.* **376**, 43-50.
- Dranow, D. B., Hu, K., Bird, A. M., Lawry, S. T., Adams, M. T., Sanchez, A., Amatruza, J. F. and Draper, B. W. (2016). Bmp15 is an oocyte-produced signal required for maintenance of the adult female sexual phenotype in zebrafish. *PLoS Genet.* **12**, e1006323.
- Draper, B. W., McCallum, C. M. and Moens, C. B. (2007). nanos1 is required to maintain oocyte production in adult zebrafish. *Dev. Biol.* **305**, 589-598.
- Evans, T. C. and Hunter, C. P. (2005). Translational control of maternal RNAs. *WormBook Nov* **10**, 1-11.
- Franceschini, A., Szklarczyk, D., Frankild, S., Kuhn, M., Simonovic, M., Roth, A., Lin, J., Minguez, P., Bork, P., von Mering, C. et al. (2013). STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res.* **41**, D808-D815.
- Gray, N. K., Collier, J. M., Dickson, K. S. and Wickens, M. (2000). Multiple portions of poly(A)-binding protein stimulate translation *in vivo*. *EMBO J.* **19**, 4723-4733.
- Guiguen, Y., Fostier, A., Piferrer, F. and Chang, C.-F. (2010). Ovarian aromatase and estrogens: a pivotal role for gonadal sex differentiation and sex change in fish. *Gen. Comp. Endocrinol.* **165**, 352-366.
- Guzeloglu-Kayisli, O., Lalioti, M. D., Aydinler, F., Sasson, I., Ilbay, O., Sakkas, D., Lowther, K. M., Mehlmann, L. M. and Seli, E. (2012). Embryonic poly(A)-binding protein (EPAB) is required for oocyte maturation and female fertility in mice. *Biochem. J.* **446**, 47-58.
- Hartung, O., Forbes, M. M. and Marlow, F. L. (2014). Zebrafish vasa is required for germ-cell differentiation and maintenance. *Mol. Reprod. Dev.* **81**, 946-961.

- Houwing, S., Kamminga, L. M., Berezikov, E., Cronembold, D., Girard, A., van den Elst, H., Filipov, D. V., Blaser, H., Raz, E., Moens, C. B. et al. (2007). A role for Piwi and piRNAs in germ cell maintenance and transposon silencing in zebrafish. *Cell* **129**, 69-82.
- Houwing, S., Berezikov, E. and Ketting, R. F. (2008). Zili is required for germ cell differentiation and meiosis in zebrafish. *EMBO J.* **27**, 2702-2711.
- Huang, P., Xiao, A., Zhou, M., Zhu, Z., Lin, S. and Zhang, B. (2011). Heritable gene targeting in zebrafish using customized TALENs. *Nat. Biotechnol.* **29**, 699-700.
- Janz, D. M. and Van Der Kraak, G. (1997). Suppression of apoptosis by gonadotropin, 17beta-estradiol, and epidermal growth factor in rainbow trout preovulatory ovarian follicles. *Gen. Comp. Endocrinol.* **105**, 186-193.
- Jiang, H.-Y., Wek, S. A., McGrath, B. C., Lu, D., Hai, T., Harding, H. P., Wang, X., Ron, D., Cavener, D. R. and Wek, R. C. (2004). Activating transcription factor 3 is integral to the eukaryotic initiation factor 2 kinase stress response. *Mol. Cell. Biol.* **24**, 1365-1377.
- Kamenska, A., Lu, W.-T., Kubacka, D., Broomhead, H., Minshall, N., Bushell, M. and Standart, N. (2014). Human 4E-T represses translation of bound mRNAs and enhances microRNA-mediated silencing. *Nucleic Acids Res.* **42**, 3298-3313.
- Kasippillai, T., MacArthur, D. G., Kirby, A., Thomas, B., Lambalk, C. B., Daly, M. J. and Welt, C. K. (2013). Mutations in eIF4ENIF1 are associated with primary ovarian insufficiency. *J. Clin. Endocrinol. Metab.* **98**, E1534-E1539.
- Kato, M., Han, T. W., Xie, S., Shi, K., Du, X., Wu, L. C., Mirzaei, H., Goldsmith, E. J., Longgood, J., Pei, J. et al. (2012). Cell-free formation of RNA granules: low complexity sequence domains form dynamic fibers within hydrogels. *Cell* **149**, 753-767.
- Kawakami, K., Takeda, H., Kawakami, N., Kobayashi, M., Matsuda, N. and Mishina, M. (2004). A transposon-mediated gene trap approach identifies developmentally regulated genes in zebrafish. *Dev. Cell* **7**, 133-144.
- Kim, J. H. and Richter, J. D. (2007). RINGO/cdk1 and CPEB mediate poly(A) tail stabilization and translational regulation by ePAB. *Genes Dev.* **21**, 2571-2579.
- Kim, H., Nakajima, T., Hayashi, S., Chambon, P., Watanabe, H., Iguchi, T. and Sato, T. (2009). Effects of diethylstilbestrol on programmed oocyte death and induction of polyovular follicles in neonatal mouse ovaries. *Biol. Reprod.* **81**, 1002-1009.
- Larsen, M. G., Bilberg, K. and Baatrup, E. (2009). Reversibility of estrogenic sex changes in zebrafish (*Danio rerio*). *Environ. Toxicol. Chem.* **28**, 1783-1785.
- Li, L., Zheng, P. and Dean, J. (2010). Maternal control of early mouse development. *Development* **137**, 859-870.
- Li, T., Huang, S., Jiang, W. Z., Wright, D., Spalding, M. H., Weeks, D. P. and Yang, B. (2011). TAL nucleases (TALNs): hybrid proteins composed of TAL effectors and FokI DNA-cleavage domain. *Nucleic Acids Res.* **39**, 359-372.
- Liang, L., Soyak, S. M. and Dean, J. (1997). FIGalpha, a germ cell specific transcription factor involved in the coordinate expression of the zona pellucida genes. *Development* **124**, 4939-4947.
- Liew, W. C. and Orban, L. (2013). Zebrafish sex: a complicated affair. *Brief. Funct. Genomics* **13**, 172-187.
- Maack, G. and Segner, H. (2003). Morphological development of the gonads in zebrafish. *J. Fish Biol.* **62**, 895-906.
- McCurley, A. T. and Callard, G. V. (2008). Characterization of housekeeping genes in zebrafish: male-female differences and effects of tissue type, developmental stage and chemical treatment. *BMC Mol. Biol.* **9**, 102.
- Mindnich, R., Deluca, D. and Adamski, J. (2004). Identification and characterization of 17 beta-hydroxysteroid dehydrogenases in the zebrafish, *Danio rerio*. *Mol. Cell. Endocrinol.* **215**, 19-30.
- Minshall, N., Reiter, M. H., Weil, D. and Standart, N. (2007). CPEB interacts with an ovary-specific eIF4E and 4E-T in early *Xenopus* oocytes. *J. Biol. Chem.* **282**, 37389-37401.
- Mishima, Y. and Tomari, Y. (2016). Codon usage and 3' UTR length determine maternal mRNA stability in zebrafish. *Mol. Cell* **61**, 874-885.
- Mold, D. E., Diniz, A. E. and Sambandan, D. R. (2009). Regulation of zebrafish zona pellucida gene activity in developing oocytes. *Biol. Reprod.* **81**, 101-110.
- Norbury, C. J. (2013). Cytoplasmic RNA: a case of the tail wagging the dog. *Nat. Rev. Mol. Cell Biol.* **14**, 643-653.
- Onichtchouk, D., Aduroja, K., Belting, H.-G., Gnügge, L. and Driever, W. (2003). Transgene driving GFP expression from the promoter of the zona pellucida gene *zpc* is expressed in oocytes and provides an early marker for gonad differentiation in zebrafish. *Dev. Dyn.* **228**, 393-404.
- Örn, S., Holbech, H., Madsen, T. H., Norrgren, L. and Petersen, G. I. (2003). Gonad development and vitellogenin production in zebrafish (*Danio rerio*) exposed to ethinylestradiol and methyltestosterone. *Aquatic Toxicol.* **65**, 397-411.
- Richter, J. D. (2007). CPEB: a life in translation. *Trends Biochem. Sci.* **32**, 279-285.
- Richter, J. D. and Lasko, P. (2011). Translational control in oocyte development. *Cold Spring Harb. Perspect. Biol.* **3**, a002758.
- Rodríguez-Marí, A. and Postlethwait, J. H. (2011). The role of Fanconi anemia/BRCA genes in zebrafish sex determination. *Methods Cell Biol.* **105**, 461-490.
- Rodríguez-Marí, A., Cañestro, C., BreMiller, R. A., Nguyen-Johnson, A., Asakawa, K., Kawakami, K. and Postlethwait, J. H. (2010). Sex reversal in zebrafish fanl mutants is caused by Tp53-mediated germ cell apoptosis. *PLoS Genet.* **6**, e1001034.
- Rodríguez-Marí, A., Wilson, C., Titus, T. A., Cañestro, C., BreMiller, R. A., Yan, Y.-L., Nanda, I., Johnston, A., Kanki, J. P., Gray, E. M. et al. (2011). Roles of *brca2* (*fancl1*) in oocyte nuclear architecture, gametogenesis, gonad tumors, and genome stability in zebrafish. *PLoS Genet.* **7**, e1001357.
- Sano, R. and Reed, J. C. (2013). ER stress-induced cell death mechanisms. *Biochim. Biophys. Acta* **1833**, 3460-3470.
- Schröder, M. and Kaufman, R. J. (2005). The mammalian unfolded protein response. *Annu. Rev. Biochem.* **74**, 739-789.
- Selman, K., Wallace, R. A., Sarka, A. and Qi, X. (1993). Stages of oocyte development in the zebrafish, *Brachydanio rerio*. *J. Morphol.* **218**, 203-224.
- Shive, H. R., West, R. R., Embree, L. J., Azuma, M., Sood, R., Liu, P. and Hickstein, D. D. (2010). *brca2* in zebrafish ovarian development, spermatogenesis, and tumorigenesis. *Proc. Natl. Acad. Sci. USA* **107**, 19350-19355.
- Shore, G. C., Papa, F. R. and Oakes, S. A. (2011). Signaling cell death from the endoplasmic reticulum stress response. *Curr. Opin. Cell Biol.* **23**, 143-149.
- Siegfried, K. R. and Nüsslein-Volhard, C. (2008). Germ line control of female sex determination in zebrafish. *Dev. Biol.* **324**, 277-287.
- Simmons, D. G., Fortier, A. L. and Cross, J. C. (2007). Diverse subtypes and developmental origins of trophoblast giant cells in the mouse placenta. *Dev. Biol.* **304**, 567-578.
- Slanchev, K., Stebler, J., de la Cueva-Mendez, G. and Raz, E. (2005). Development without germ cells: the role of the germ line in zebrafish sex differentiation. *Proc. Natl. Acad. Sci. USA* **102**, 4074-4079.
- Soyak, S. M., Amleh, A. and Dean, J. (2000). FIGalpha, a germ cell-specific transcription factor required for ovarian follicle formation. *Development* **127**, 4645-4654.
- Standart, N. and Minshall, N. (2008). Translational control in early development: CPEB, P-bodies and germinal granules. *Biochem. Soc. Trans.* **36**, 671-676.
- Szegezdi, E., Logue, S. E., Gorman, A. M. and Samali, A. (2006). Mediators of endoplasmic reticulum stress-induced apoptosis. *EMBO Rep.* **7**, 880-885.
- Takahashi, H. (1977). Juvenile hermaphroditism in the zebrafish, *Brachydanio rerio*. *Bull. Fac. Fish Hokkaido Univ.* **28**, 57-65.
- Uchida, D., Yamashita, M., Kitano, T. and Iguchi, T. (2002). Oocyte apoptosis during the transition from ovary-like tissue to testes during sex differentiation of juvenile zebrafish. *J. Exp. Biol.* **205**, 711-718.
- Udeshi, N. D., Svinkina, T., Mertins, P., Kuhn, E., Mani, D. R., Qiao, J. W. and Carr, S. A. (2013). Refined preparation and use of anti-diglycine remnant (K-epsilon-GG) antibody enables routine quantification of 10,000s of ubiquitination sites in single proteomics experiments. *Mol. Cell. Proteomics* **12**, 825-831.
- Voeltz, G. K., Ongkasuwan, J., Standart, N. and Steitz, J. A. (2001). A novel embryonic poly(A) binding protein, ePAB, regulates mRNA deadenylation in *Xenopus* egg extracts. *Genes Dev.* **15**, 774-788.
- Wang, S. and Kaufman, R. J. (2012). The impact of the unfolded protein response on human disease. *J. Cell Biol.* **197**, 857-867.
- Westerfield, M. (2000). *The Zebrafish Book: A Guide for the Laboratory Use of Zebrafish (Danio rerio)*. Eugene, OR: University of Oregon Press.
- White, Y. A. R., Woods, D. C. and Wood, A. W. (2011). A transgenic zebrafish model of targeted oocyte ablation and de novo oogenesis. *Dev. Dyn.* **240**, 1929-1937.
- Wu, X., Viveiros, M. M., Eppig, J. J., Bai, Y., Fitzpatrick, S. L. and Matzuk, M. M. (2003a). Zygote arrest 1 (*Zar1*) is a novel maternal-effect gene critical for the oocyte-to-embryo transition. *Nat. Genet.* **33**, 187-191.
- Wu, X., Wang, P., Brown, C. A., Zilinski, C. A. and Matzuk, M. M. (2003b). Zygote arrest 1 (*Zar1*) is an evolutionarily conserved gene expressed in vertebrate ovaries. *Biol. Reprod.* **69**, 861-867.
- Yamamoto, T. M., Cook, J. M., Kotter, C. V., Khat, T., Silva, K. D., Ferreyros, M., Holt, J. W., Knight, J. D. and Charlesworth, A. (2013). *Zar1* represses translation in *Xenopus* oocytes and binds to the TCS in maternal mRNAs with different characteristics than *Zar2*. *Biochim. Biophys. Acta* **1829**, 1034-1046.

Supplementary information

Translation Repression by Maternal RNA Binding Protein Zar1 is Essential for Early Oogenesis in Zebrafish

Liyun Miao^{1,2,4,5}, Yue Yuan^{1,4,5}, Feng Cheng^{1,7}, Junshun Fang¹, Fang Zhou¹, Weirui Ma^{1,8}, Yan Jiang⁶, Xiahe Huang¹, Yingchun Wang¹, Lingjuan Shan¹, Dahua Chen³, Jian Zhang^{1,4,5*}

Materials and Methods

Mutant generation and genotyping

The zebrafish *zar1* mutants were generated using the Tübingen strain with TALEN system or CRISPR/Cas9 system. The following primers were used to identify *zar1* mutants generated with TALEN system: for mutant screening a pair of primers (forward primer: 5'-CTTTCCCAAACCTCGAAAATCGT-3'; reverse primer: 5'-GGGGTGAGATTTGGGTTGATCTG-3') were used; for *zar1*^{gd5} mutant line specific genotyping two forward primers (5'-GCGAAAATGGCTACATAT-3' to detect the WT allele and 5'-GCGAAAATGGCTACATGG-3' to detect the mutant allele) and a reverse primer (5'-TCCGGGATTCTACTGGGGAGTAA-3') were used

Antibodies

Zar1 antibody was generated with full-length recombinant Zar1 protein purified from *E.coli* by immunizing rabbits and affinity purified with SulfoLink Immobilization Trial Kit (Thermo, 20325). Zar1 antibody was diluted 1:1000. Other antibodies and beads were purchased: 4E-T antibody (Abcam, ab6034, 1:500); tubulin antibody (Sigma, T6074, 1:4000); FLAG antibody (Sigma, F3165, 1:2000); anti-FLAG M2 affinity gel (Sigma, A2220); and protein A beads (Millipore, 16-156).

Quantitative proteomic analysis by iTRAQ

The ovary lysates were boiled for 10 min and centrifuged (12000 rpm, 10 min). The supernatant (100 µg protein / sample) were reduced (10 mM DTT, 37°C, 1 h), alkylated (55 mM iodoacetamide, room temperature, 1 h, in the dark), and transferred to the Microcon YM-30 centrifugal filter units (EMD Millipore Corporation) where the lysis buffer was replaced with iTRAQ dissolution buffer. Samples were digested with trypsin (sequencing grade, 1:50, 37 °C, overnight). The resulting peptides were labeled, according to the manufacturer's manual with slight modifications, with iTRAQ Reagents (AB Sciex Inc.). Briefly, the peptides were incubated with the ethanol-dissolved iTRAQ reagents (2 h, room temperature), and then terminated by adding H₂O to 30% of the total reaction volume. The labeled samples were mixed together with equal ratios in amount and fractionated by reverse phase HPLC. The mixed peptides were resuspended in 0.5 ml of RP-HPLC solvent A (2% ACN, 5 mM ammonium formate, pH 10), and separated into 10 fractions using the Gemini-NX 5u c18 110A (P/No: 00G-4454-Y0 3*250 mm length, 5 µm particle size, Phenomenex) on a Waters e2695 separations module system with a flow rate of 0.4 ml/min. A 97 min basic RP LC method was utilized for offline fractionation. The gradient consisted of an initial increase to 8% solvent B (1.1%B/min) (90% ACN/5 mM ammonium formate, pH 10) followed by a 38 min linear gradient (0.5% B/min) from 8% solvent B to 27% B and successive ramps to 31% B (1%B/min), 39% B (0.5%B/min), and 60% B (3%B/min). The separated peptides were further desalted with C18 StageTips, concentrated with a SpeedVac, resuspended with 0.1% formic acid.

Figures




Fig. S1. *zar1* mutant generation with CRISPR/Cas9 system

(A) Selected gRNA sequence. DNA sequences highlighted in red are the protospacer-adjacent motif (PAM). (B) PCR fragments (amplified with the following primers: primer-F: 5'-AGGTCACAGAGACGGTTGACAGC-3' and primer-R: 5'-ACTTTCCCACCGTAGGTTGCAGT-3') were digested with T7 endonuclease I to test the gRNA. (C) DNA sequences of the *zar1^{gd6}* mutant line. Green highlighted DNA sequences are the premature stop codon. Two *zar1^{gd6}* mutant line specific forward primers (5'-CCAGTGAAAACCGAACA-3' to detect the WT allele and 5'-CCAGTGAAAACCGAATGA-3' to detect the mutant allele) were used together with the reverse primer (primer-R) mentioned above to do genotyping. (D) Western blot of Zar1 in gonads at 25 dpf from wild-type (WT), *zar1^{gd6/+}* heterozygotes (hetero) and *zar1^{gd6/gd6}* homozygotes (homo). Gonads from 8 fish were mixed together and lysed for each genotype. (E) Analysis of the gender of *zar1^{gd6/gd6}* homozygotes (homo), *zar1^{gd6/+}* heterozygotes (hetero) and WT siblings.




Fig. S2. Zar1 is expressed in *zar1* homozygous females with the *Tg(zp3b:zar1,cmlc2:EGFP)* transgene.

(A, A') Heart specific EGFP signal (arrow) seen in transgenic embryos. (B) Western blot to detect Zar1 protein in adult ovaries. Zar1 was expressed in the transgene rescued adult *zar1* homozygous ovaries (arrowhead). Compared with Zar1 expression level in *zar1* heterozygous mutant ovaries (hetero), Zar1 expression level in transgene rescued *zar1* homozygous (homo) mutant ovaries was low. The smaller bands are non-specific.




Fig. S3. Spermatogenesis is not affected in *zar1* mutant males.

Histology analysis of testes of *zar1* homozygous males and control siblings at 50 dpf and 60 dpf. Spermatogenesis in *zar1* homozygous males resembles that in control siblings. sg: spermatogonia; sc: spermatocytes; sp: sperm; Scale bar: 0.04 mm.




Fig. S4. Sex ratio in juveniles at 33-41 dpf is not affected in *zar1* mutants.

Sex ratios in *zar1* homozygotes (homo) and *zar1* heterozygotes (hetero) at 33 dpf-41 dpf. Sex of *zar1* homozygotes and *zar1* heterozygotes was determined at 33-41 dpf according to H&E staining results.




Fig. S5. The chorions of eggs from $p53^{-/-};zar1^{-/-}$ failed to elevate normally.

Five $p53^{-/-};zar1^{+/-}$ females and five $p53^{-/-};zar1^{-/-}$ females were crossed with wild-type males. The chorions of eggs from $p53^{-/-};zar1^{+/-}$ lifted normally (A). The chorions of eggs from $p53^{-/-};zar1^{-/-}$ failed to lift. Cleaved embryos (B) and uncleaved embryos (C); P-female: genotypes of the mothers




Fig. S6. The chorions of $zar1$ mutant eggs failed to elevate normally.

EE2 treated females were crossed with wild-type males. Embryos from 6 heterozygous females (EE2-hetero) and 6 homozygous females (EE2-homo) were analyzed. The chorions of $zar1$ homozygotes failed to lift properly compared with those of $zar1$ heterozygotes. P-female: genotypes of the mothers.




Fig. S7. qPCR to confirm the iTRAQ result.

Statistical analysis of mRNA levels of upregulated genes in *zar1* homozygotes (homo), according to iTRAQ result. Ovaries were isolated from *zar1* homozygotes (homo) and *zar1* heterozygotes (hetero) at 33 dpf. Seven out of the first 25 upregulated genes were tested. Internal control: *efla*; ns: not significant; *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$.




Fig. S8. Immunity related proteins are upregulated in *zar1* mutants.

Statistical analysis of immune related proteins upregulated in *zar1* homozygotes (homo) compared to heterozygotes (hetero) determined by iTRAQ. *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$.

Zar1 SAYVWCVQG¹TNKVYFKQFC²RTCCQKSFNPYRVEDIAC³QTCKKARCTCSVKSRHVDPKRPHRQDL⁴CGRCKGKRLSCDSTFSFKYII
 Zar1-mu SAYVWCVQG¹TNKVYFKQF²ARTAQKSFNPYRVEDIAA³QTAKKARATASVKSRHVDPKRPHRQDL⁴AGRAK⁵GKRLSCDSTFSFKYII

Fig. S9. Mutation of the Zar1 Znf domain.

The 8 cysteines in Zar1 Znf domain are mutated to alanines.




Fig. S10. Interaction between Zar1 and translation initiation factor 4E transporter.

(A) Association analysis of Zar1 with the immunoprecipitated proteins by searching STING database (evidence view). (B) 4E-T was immunoprecipitated by the Zar1 antibody from *zar1* heterozygous ovaries (hetero, *p53*^{-/-};*zar1*^{+/-}) but not from *zar1* homozygous ovaries (homo, *p53*^{-/-};*zar1*^{-/-}). (C) Zar1 was immunoprecipitated from ovaries by a 4E-T antibody but not by a control antibody. (D) Zar1 was immunoprecipitated by the Flag antibody from HEK293 cells co-transfected with *zar1* and *eif4enif1-flag* plasmids but not from cells transfected with *zar1* plasmid alone.

Tables

Table S1. The three categories of upregulated proteins (> 2 fold over controls) in *zar1* mutants

| Category | Accession | Protein Name | Gene symbol |
|-----------------------------------|-----------|---|-------------------------|
| ER stress & the UPR related | Q6P0U6 | DnaJ (Hsp40) homolog, subfamily C, member 3a | <i>dnajc3a</i> |
| | Q6P2B0 | Vesicle-associated membrane protein-associated protein B/C | <i>vapb</i> |
| Zona pellucida proteins | Q6P3I1 | Protein disulfide-isomerase A4 | <i>pdia4</i> |
| | Q5TYP2 | Zona pellucida glycoprotein 3a, tandem duplicate 2 | <i>zp3a.2</i> |
| | A7MBW8 | Zona pellucida glycoprotein 2, like 1 | <i>zp2l1</i> |
| | A8WG31 | pellucida sperm-binding protein 3 | <i>LOC100148225</i> |
| | F1R4N4 | Egg envelope glycoprotein-like | <i>si:dkey-19b23.11</i> |
| | A7MBS3 | Zona pellucida glycoprotein 3b | <i>zp3b</i> |
| | B0R0H4 | ZPC domain containing protein 1 | <i>si:ch211-14a17.7</i> |
| | Q5TYX2 | ZPA domain containing protein | <i>si:dkeyp-50f7.2</i> |
| Immunity related | Q24JW2 | Lysozyme | <i>lyz</i> |
| | Q1LV07 | Metallothionein | <i>mt2</i> |
| | F1QYN0 | Complement C3-like | <i>LOC100331428</i> |
| | F1QLN6 | Complement component c3a, duplicate 5 | <i>c3a.5</i> |
| | Q803G1 | Solute carrier family 3, member 2b | <i>slc3a2b</i> |
| | E7FCS3 | Alpha-2-macroglobulin-liked, duplicate 1 | <i>a2m1</i> |
| | F1QF63 | Alpha-2-macroglobulin-liked, duplicate 2 | <i>a2m2</i> |
| | F1Q6K5 | Lectin, galactoside-binding, soluble, 3 binding protein b | <i>lgals3bpb</i> |
| | A2VD28 | Complement factor B | <i>cfb</i> |
| | Q6PH75 | Cathepsin B | <i>ctspa</i> |

Table S2. Sequences of primers used in qPCR and RT-PCR.

| <i>gene</i> | primers |
|-----------------------|--|
| <i>atf3</i> | F: 5'-TCAATGGCTACTGAATTCCAAC TG-3' R: 5'-TTCTTGTGGCATGTTATGTGGAC-3' |
| <i>ddit3</i> | F: 5'-AGTTGGAGGCGTGGTATGAAGAC-3' R: 5'-GTCAACCAGGTGAGCGAACAG-3' |
| <i>btg2</i> | F: 5'-CATTAGAAACCAGACAAATCCTCGT-3' R: 5'-GGAGCGGTGCTGTGGTTAAG-3' |
| <i>traf4a</i> | F: 5'-GAACTCTGGACTCGACATGCTCA-3' R: 5'-ATTGCACAAGGCTCATCTTCCTC-3' |
| <i>LOC566745</i> | F: 5'-GAGTTGCAACTTGAGAAGCTCTTC-3' R: 5'-TCCTCCAACCCAAACCCAGATC-3' |
| <i>lyz</i> | F: 5'-AGAATTTGTGCAAAGTGGCCTGT-3' R: 5'-AAGAATCCCAGGTTTCCCATGAT-3' |
| <i>si:dkey-7i4.24</i> | F: 5'-CCGCTCTGAAACAAAAGCAGTGG-3' R: 5'-CGCTTAGATGGATGGGGTCAGGT-3' |
| <i>mt2</i> | F: 5'-GACTGGA ACTTGCAACTGTGGTG-3' R: 5'-GGGCAGCAAGAACAACA ACTCTT-3' |
| <i>serpinb5</i> | F: 5'-TGATACAGGCAATGGTGGAGTTG-3' R: 5'-CTGTGAGGAATCTGCTCCTAGC-3' |
| <i>zp21l</i> | F: 5'-TTTCTGACTTTGGGTCGTTGTTG-3' R: 5'-GGGAACAAAGTGGTCAGGTAACG-3' |
| <i>LOC100148225</i> | F: 5'-GCCTTCATGTTCCAGGATACACC-3' R: 5'-CCCAGTAAGTTGGCCTTCAACAC-3' |
| <i>dnajc3a</i> | F: 5'-AGACA ACTTCCAGGATGCAGAGG-3' R: 5'-GTCCAGAACCAAACGGATTGAAC-3' |
| <i>zp3b</i> | F: 5'-TTGTGTTTCATCGACTGGTGTGTG-3' |

| | |
|-------------------------|----------------------------------|
| | R: 5'-ACTCCTGCTATCTGCAAGACATC-3' |
| | F: 5'-GGAGTGATGTCGTAGCCCTCATT-3' |
| <i>Slc3a2b</i> | R: 5'-GAGGAGCAAGTCCACACCAGTTT-3' |
| | F: 5'-ACCCAGGACTGCATTCCTACTGC-3' |
| <i>zp3a.2</i> | R: 5'-ACAAGTTCAGGAGAACTCTAATC-3' |
| | F: 5'-TATTCCTCTGACGCAAAGCCAGT-3' |
| <i>si:dkey-19b23.11</i> | R: 5'-GTCCACTGAAAGGGTTTCAAGGT-3' |
| | F: 5'-GCAGTGTTTCATGCTGAATTTGG-3' |
| <i>si:ch211-14a17.7</i> | R: 5'-GAGACCACTCTTTTGGCCTTTCA-3' |
| | F: 5'-CAGGCCTAAATTCAGCAATGACC-3' |
| <i>si:dkeyp-50f7.2</i> | R: 5'-GCGGTTGTGAGGTAAACGAAATC-3' |
| | F: 5'-GTTGACCGCCTGAAGTCTCTGAT-3' |
| <i>firefly</i> | R: 5'-CCACGATCTCTTTTTCCGTCATC-3' |
| | F: 5'-ATTGAATCGGACCCAGGATTCTT-3' |
| <i>renilla</i> | R: 5'-TTTCATCAGGTGCATCTTCTTGC-3' |

Table S3. Proteomic comparison between *zar1* homozygotes and *zar1* heterozygotes.

Proteomic analysis with iTRAQ to compare *zar1* homozygous ovaries (homo) with *zar1* heterozygous ovaries (hetero). 10 ovaries for each genotype were isolated at 33 dpf and analyzed with iTRAQ. Two replicates were performed for each genotype. The UniProt proteome sequences for *Danio rerio* were used for the database searching.

[Click here to Download Table S3](#)

Table S4. Summary of differentially expressed proteins based on the iTRAQ result.

Analysis of differentially expressed proteins between *zar1* homozygous ovaries (homo) and *zar1* heterozygous ovaries (hetero) determined by iTRAQ. 325 proteins show differential expression ($P < 0.05$). 42 proteins changed by more than 2 folds, among which 37 proteins were upregulated and 5 proteins were down-regulated in homozygous ovaries compared with the control ovaries.

[Click here to Download Table S4](#)

Table S5. Mass Spectrometry (MS) analysis of proteins precipitated by the Zar1 antibody.

MS analysis of immunoprecipitated proteins from *zar1* heterozygotes and *zar1* homozygotes. Immunoprecipitation with the Zar1 antibody was performed using *zar1* heterozygous mutant ovaries ($p53^{-/-};zar1^{+/-}$) and *zar1* homozygous mutant ovaries ($p53^{-/-};zar1^{-/-}$) followed by MS analysis. 59 proteins from *zar1* heterozygous ovaries and 44 proteins from *zar1* homozygous ovaries were precipitated.

[Click here to Download Table S5](#)