

Fibrillin-1 directly regulates osteoclast formation and function by a dual mechanism

Kerstin Tiedemann^{1,2}, Iris Boraschi-Diaz^{1,2}, Irina Rajakumar^{1,*}, Jasvir Kaur³, Peter Roughley², Dieter P. Reinhardt^{1,3,‡} and Svetlana V. Komarova^{1,2,3,‡,§}

¹Faculty of Dentistry, McGill University, 3640 rue University, Montreal, Quebec, Canada, H3A 0C7

²Shriners Hospital for Children – Canada, 1529 Cedar Ave, Montreal, Quebec, Canada, H3G 1A6

³Department of Anatomy and Cell Biology, Faculty of Medicine, McGill University, 3640 rue University, Montreal, Quebec, Canada, H3A 0C7

*Present address: London Health Science Center, London, ON, Canada

‡These authors contributed equally to this work

§Author for correspondence (svetlana.komarova@mcgill.ca)

Accepted 25 June 2013

Journal of Cell Science 126, 4187–4194

© 2013. Published by The Company of Biologists Ltd

doi: 10.1242/jcs.127571

Summary

Mutations in the fibrillin-1 gene give rise to a number of heritable disorders, which are all characterized by various malformations of bone as well as manifestations in other tissues. However, the role of fibrillin-1 in the development and homeostasis of bone is not well understood. Here, we examined the role of fibrillin-1 in regulating osteoclast differentiation from primary bone-marrow-derived precursors and monocytic RAW 264.7 cells. The soluble N-terminal half of fibrillin-1 (rFBN1-N) strongly inhibited osteoclastogenesis, whereas the C-terminal half (rFBN1-C) did not. By contrast, when rFBN1-N was immobilized on calcium phosphate, it did not affect osteoclastogenesis but modulated osteoclast resorptive activity, which was evident by a larger number of smaller resorption pits. Using a panel of recombinant sub-fragments spanning rFBN1-N, we localized an osteoclast inhibitory activity to the 63 kDa subfragment rF23 comprising the N-terminal region of fibrillin-1. Osteoclastic resorption led to the generation of small fibrillin-1 fragments that were similar to those identified in human vertebral bone extracts. rF23, but not rFBN1-N, was found to inhibit the expression of cathepsin K, matrix metalloproteinase 9 and Dcstamp in differentiating osteoclasts. rFBN1-N, but not rF23, exhibited interaction with RANKL. Excess RANKL rescued the inhibition of osteoclastogenesis by rFBN1-N. By contrast, rF23 disrupted RANKL-induced Ca²⁺ signaling and activation of transcription factor NFATc1. These studies highlight a direct dual inhibitory role of N-terminal fibrillin-1 fragments in osteoclastogenesis, the sequestration of RANKL and the inhibition of NFATc1 signaling, demonstrating that osteoclastic degradation of fibrillin-1 provides a potent negative feedback that limits osteoclast formation and function.

Key words: Calcium signaling, Fibrillin, NFATc1, Osteoclastogenesis, RANKL

Introduction

Mutations in the fibrillin-1 gene cause a range of connective tissue disorders characterized by variable clinical symptoms in the skeletal, cardiovascular and ocular systems. These disorders are termed type I fibrillinopathies and include, among others, Marfan syndrome (MFS), dominant Weill-Marchesani syndrome, geleophysic dysplasia and acromicric dysplasia (Hubmacher and Reinhardt, 2011). The skeletal symptoms in these disorders show a remarkable range, from long bone overgrowth, arachnodactyly and loose joints in MFS to short stature and brachydactyly in the other type I fibrillinopathies, suggesting that fibrillin-1 plays a major role in regulating skeletal homeostasis. Osteopenia is a hallmark of MFS and may be present in Weill–Marchesani syndrome (Giordano et al., 1997; Pyeritz, 2000).

This study is motivated by a largely unexplained low bone mass present particularly in individuals with MFS that likely increases the risk of fractures as the patient age increases due to successful treatment of cardiovascular aneurysms (Kohlmeier et al., 1995). Children with MFS have significantly lower bone mineral density compared with healthy height-matched controls (Kohlmeier et al., 1995; Moura et al., 2006), which was shown to persist in both women (Kohlmeier et al., 1995; Le Parc et al., 1999) and men (Carter et al., 2000; Le Parc et al., 1999) with

MFS. While no appropriately powered study was performed yet on the fracture risk in older MFS patients, studies with small numbers of young patients indicate increased peripheral traumatic fractures (Grahame and Pyeritz, 1995; Kohlmeier et al., 1995).

The fibrillin super-family of large cysteine-rich glycoproteins consists of fibrillin-1, -2 and -3 and the latent transforming growth factor (TGF) β -binding proteins (LTBPs). Fibrillins constitute the backbone of extracellular microfibrils, which confer structural and functional integrity to all tissues including the skeleton (Hubmacher et al., 2006; Ramirez et al., 2007). In the skeletal system, the fibrillins are expressed in long bones, ribs, vertebral bodies and cartilage (Arteaga-Solis et al., 2001; Keene et al., 1991; Kitahama et al., 2000; Quondamatteo et al., 2002; Sabatier et al., 2011; Zhang et al., 1995). Fibrillins and fibrillin-containing microfibrils regulate the bioavailability of growth factors of the TGF β superfamily including TGF β and bone morphogenetic proteins (BMPs) (Neptune et al., 2003; Nistala et al., 2010b; Sengle et al., 2008). In turn, TGF β and BMPs are important regulators of bone growth and remodeling (Giannoudis et al., 2007; Guo et al., 2008; Janssens et al., 2005; Quinn et al., 2001). Elevated TGF β levels are a major disease-causing factor in MFS and other fibrillinopathies (Doyle et al., 2012; Le Goff and Cormier-Daire, 2012).

Changes in bone are regulated by the activities of osteoblasts, the cells responsible for new bone formation, and osteoclasts that remove mineralized tissue. In mouse models of progressively severe MFS, *Fbn1*^{mgR/mgR} (Nistala et al., 2010a), as well as *Fbn1*^{-/-} (Nistala et al., 2010c), osteoclastogenesis was reported to increase in a TGFβ- and osteoblast-dependent manner. Importantly, in the *Fbn1*^{mgR/mgR} mice, treatment with alendronate, a bisphosphonate that directly targets osteoclasts, but not with an angiotensin receptor I antagonist that lowers TGFβ signaling, resulted in significant improvement of bone quality (Nistala et al., 2010a). These data suggest that osteoclasts are critical, but unexplored, players in the skeletal pathophysiology of MFS and possibly other type I fibrillinopathies. In this study, we addressed the direct role of fibrillin-1 in regulating osteoclast differentiation and function.

Results

Fibrillin-1 and fibrillin-2 are produced and secreted by differentiating osteoblasts

First we demonstrated that fibrillin-1 and fibrillin-2 are expressed and secreted by differentiating osteoblasts (supplementary material Fig. S1). Both, *Fbn1* and *Fbn2* mRNA were strongly induced during the first week of osteoblast differentiation, and the development of fibrillin-1-containing microfibrils was apparent by immunofluorescence during the first week of differentiation. Fibrillin-1 was undetectable in cells of osteoclast lineage (supplementary material Fig. S1D).

Fibrillin-1 affects function and differentiation of osteoclasts

Since full-length fibrillin-1 is not amenable for sufficient recombinant production due to its propensity to aggregate (Lin

et al., 2002), we utilized recombinant N-terminal (rFBN1-N) and C-terminal (rFBN1-C) halves of fibrillin-1 (Jensen et al., 2001). rFBN1-N and rFBN1-C were immobilized on calcium phosphate, and osteoclast differentiation from bone marrow cells was induced with MCSF (50 ng/ml) and RANKL (50 ng/ml). The presence of fibrillin-1 on calcium phosphate did not affect osteoclast formation (Fig. 1A top, B). Although the total area resorbed per osteoclast remained the same under all conditions, on the calcium phosphate plates coated with rFBN1-N the osteoclasts formed 6-fold more resorption pits, which were significantly smaller in size than control pits (Fig. 1A bottom, C). We next examined how the presence of soluble rFBN1-N or rFBN1-C in the incubation medium affects osteoclast formation. Bone marrow-derived precursors were cultured with MCSF and RANKL or RAW 264.7 cells were cultured with RANKL (50 ng/ml) for 5–6 days with or without 50 μg/ml rFBN1-N or rFBN1-C, and osteoclast differentiation was quantified. There was no difference in the appearance or numbers of osteoclasts formed in the presence of rFBN1-C compared with positive control (Fig. 1D,E). However, when rFBN1-N was added to the culture medium, osteoclast formation was significantly decreased (Fig. 1D,E). Albumin (50 μg/ml) used as a control to coat calcium phosphate or as a soluble ligand did not affect osteoclast differentiation (supplementary material Fig. S2).

A small N-terminal fragment of fibrillin-1 acts as an inhibitor of osteoclastogenesis

We used a panel of recombinant sub-fragments spanning rFBN1-N (Fig. 2A) as soluble ligands during osteoclast differentiation from bone marrow or RAW 264.7 cells. The N-terminal half rFBN1-N and rF23, that represents the most N-terminal region of




Fig. 1. The N-terminal half of fibrillin-1 affects osteoclastic resorption and inhibits osteoclast differentiation. (A–C) Bone marrow cells were plated on calcium phosphate that was uncoated (positive control, PC) or coated with 50 μg/ml rFBN1-N (1N) or rFBN1-C (1C) and differentiation was induced with MCSF (50 ng/ml) and RANKL (50 ng/ml). After 5–6 days the samples were fixed and stained for TRAP. Osteoclasts were quantified, removed and resorption was assessed. (A) Representative images of osteoclasts (top) and resorption pits (bottom) formed under the indicated conditions. Scale bars apply to all images in the corresponding rows. (B,C) Average numbers of differentiated osteoclasts (B, top), total area resorbed per osteoclast (B, bottom), average pit size (C, top), and numbers of resorption pits per osteoclast (C, bottom). Data are means ± s.e.m., $n=3$ experiments; * $P<0.05$ assessed by Student's *t*-test. (D,E) Bone marrow cells [white bars, all conditions treated with MCSF (50 ng/ml)] or RAW 264.7 (black bars) were cultured for 5 days without RANKL (negative control, NC); with RANKL (50 ng/ml, positive control, PC) or with RANKL and soluble rFBN1-N or rFBN1-C, fixed and stained for TRAP. (D) Representative images of osteoclastic cells formed from RAW 264.7 under the indicated conditions. Scale bars apply to all images. (E) Average numbers of osteoclasts. Data are means ± s.e.m., $n=4$ experiments for bone marrow cells, $n=5–6$ experiments for RAW 264.7; ** $P<0.01$, *** $P<0.001$ assessed by Student's *t*-test.




Fig. 2. A small N-terminal subfragment of fibrillin-1 inhibits osteoclastogenesis. (A) Schematic of recombinant N-terminal (rFBN1-N) and C-terminal (rFBN1-C) halves of fibrillin-1, and recombinant sub-fragments of rFBN1-N – rF23, rF51, rF1F and rF18. (B–D) Bone marrow cells (white bars) and RAW 264.7 (black bars) were cultured for 5 days untreated (negative control, NC), treated with RANKL only (50 ng/ml, positive control, PC) or treated with RANKL and 50 μ g/ml of soluble fibrillin-1 fragments rFBN1-N, rFBN1-C, rF23, rF51, rF1F or rF18. (B) Average numbers of osteoclasts formed under the indicated conditions. Data are means \pm s.e.m., $n=3-8$ experiments; * $P<0.05$, ** $P<0.01$, *** $P<0.001$ compared with PC assessed by Student's t -test. (C,D) mRNA was isolated and the expression of cathepsin K (*Ctsk*, C) and *Dcstamp* (D) was quantified. Data are means \pm s.e.m., $n=3-9$ experiments; * $P<0.05$, *** $P<0.001$, compared with PC assessed by Student's t -test.

fibrillin-1, significantly inhibited osteoclastogenesis (Fig. 2B; supplementary material Fig. S3A). Expression of RANKL-induced osteoclast marker genes *Ctsk*, *Mmp9* and *Dcstamp* was significantly inhibited by rF23, but not by other fibrillin-1 fragments (Fig. 2C,D; supplementary material Fig. S3B). The expression of *Trap*, *Rank* and *Itgb3* (integrin $\beta 3$) was not affected by fibrillin-1 fragments (supplementary material Fig. S3B). rFBN1-N and rF23 inhibited osteoclastogenesis in a dose-dependent manner (supplementary material Fig. S3C) and were most effective when applied for the duration of the experiment (supplementary material Fig. S3D). The inhibitory potential of fibrillin-1 fragments was specific to osteoclasts, since none of the fibrillin-1 fragments affected differentiation of osteoblasts (supplementary material Fig. S4).

Fibrillin-1 fragments are released during resorption and inhibit osteoclasts *in vivo*

Biotinylated rFBN1-N was coated on calcium phosphate and osteoclastogenesis from bone marrow cells was induced with RANKL and MCSF. Mature osteoclasts were cultured for additional 24 hours in serum free medium, which was then collected and the degradation pattern of fibrillin-1 was visualized by immunoblotting (Fig. 3A). rFBN1-N as well as multiple fibrillin-1 degradation products in the range of 35–85 kDa were released by osteoclasts (Fig. 3A, lane OC). No spontaneous release of biotinylated material was observed when coated wells were incubated without cells (Fig. 3A, lane M). To explore if fibrillin-1 and its degradation products are present in the human bone matrix, we extracted proteins from vertebral bones of healthy individuals. We identified fibrillin-1 multimers, full length fibrillin-1 as well as a number of smaller fibrillin-1 degradation products in the bone tissue (Fig. 3B). To assess if fibrillin-1 degradation fragments can inhibit osteoclast formation in a complex *in vivo* environment, we used a small sub-fragment of rF23, the 32 kDa rF31 (Fig. 3C). We confirmed that similar to rF23, rF31 inhibits osteoclast formation from RAW 264.7 or bone marrow precursors (Fig. 3D). Next, we injected healthy 6-week-old male balb/c mice with rF31 (50 μ g/kg)

intra-peritoneally every other day for 1 week, and examined 1 week after the last injection osteoclasts in histological sections. In mice injected with rF31 the osteoclast number per bone perimeter was decreased by 50% compared with vehicle-injected animals (Fig. 3E).

TGF β does not affect osteoclast formation in the cell culture models used

TGF β plays a key role in fibrillin-mediated cellular effects in vasculature and bone (Neptune et al., 2003; Nistala et al., 2010b). However, differentiation of RAW 264.7 cells into osteoclasts was not affected by addition of TGF $\beta 1$ (1–50 ng/ml) or by inhibition of TGF β function using a pan-specific TGF β -neutralizing antibody or an inhibitor of T β RI (supplementary material Fig. S5A,B), indicating that the osteoclastogenesis system used in our study is not sensitive to TGF β . To exclude that TGF β co-purified with fibrillin fragments (Kaur and Reinhardt, 2012) may inhibit osteoclastogenesis, we added TGF β neutralizing antibodies or T β RI inhibitor in the differentiation assays with fibrillin-1 fragments (supplementary material Fig. S5C). Reducing TGF β levels or signaling did not prevent osteoclast inhibition by rFBN1-N or rF23. Thus, the osteoclast-inhibitory activity exerted by fibrillin-1 fragments in our experimental setup is not mediated by TGF β .

The N-terminal half of fibrillin-1 binds and sequesters RANKL

We hypothesized that direct binding of RANKL to fibrillin-1 fragments may mediate their osteoclast inhibitory effects. We performed solid phase binding assays, using RANKL as immobilized and rFBN1-N, rFBN1-C or rF23 as soluble ligands. We confirmed that rFBN1-N and rF23 are recognized by the detector antibody with similar efficiencies (supplementary material Fig. S6). rFBN1-N, but not rFBN1-C or rF23, interacted with RANKL in a Ca^{2+} -dependent manner (Fig. 4A). Furthermore, inhibition of osteoclastogenesis by rFBN1-N was rescued by increasing RANKL levels (Fig. 4B). This strategy was not effective in rescuing osteoclast inhibition induced by rF23 (Fig. 4C).




Fig. 3. Fragments of fibrillin-1 are released during resorption and inhibit osteoclasts *in vivo*. (A) Bone marrow cells were plated on calcium phosphate coated with biotinylated rFBN1-N (80–155 $\mu\text{g}/\text{cm}^2$) and after mature osteoclasts developed (5–7 days), the cultures were maintained for an additional 24 hours in serum free medium. Lane B, Biotinylated rFBN1-N before coating. Lane OC, conditioned medium from wells cultured with osteoclasts. Lane M, conditioned medium from wells cultured identically, but without cells. Arrowheads indicate degradation products in the range of 35–85 kDa. (B) Proteins were extracted from human vertebral bone and immunoblotted for fibrillin-1. Arrows indicate multimeric (FBN1 mult.) and monomeric full length fibrillin-1 (FBN1); arrowheads indicate fibrillin-1 degradation products. (C) Schematic of recombinant sub-fragments rF23 and rF31 (compare with Fig. 2 for position within full length fibrillin-1). (D) Bone marrow cells (white bars) and RAW 264.7 (black bars) were cultured in parallel for 5 days with RANKL only (50 ng/ml, positive control, PC), or with RANKL and 50 $\mu\text{g}/\text{ml}$ of rF23 or rF31. Data are means \pm s.e.m., $n=2-3$ experiments; * $P<0.05$, compared with PC assessed by Student's *t*-test. (E) Healthy mice were injected with vehicle (PBS) or recombinant rF31 (50 $\mu\text{g}/\text{kg}$) intra-peritoneally every other day for 1 week and osteoclast numbers were assessed 1 week after the last injection. Left, representative images of upper tibial metaphyseal area of vehicle- or rF31-treated mice. Right, osteoclast number (OCN) per bone perimeter (BP) in vehicle- and rF31-injected mice. Data are means \pm s.d., $n=11$ mice for vehicle-injected and $n=4$ mice for rF31-injected; * $P<0.05$, assessed by Student's *t*-test.

rF23 impairs NFATc1 translocation and Ca^{2+} signaling in osteoclast precursors

Since *Ctsk* mRNA expression is regulated by the transcription factor nuclear factor of activated T-cells (NFATc1) (Balkan et al., 2009; Ishida et al., 2002; Takayanagi et al., 2002), we hypothesized that rF23 affects NFATc1 activation. NFATc1 protein expression is induced by RANKL (Ishida et al., 2002; Takayanagi et al., 2002). Inactive hyper-phosphorylated NFATc1




Fig. 4. The N-terminal half of fibrillin-1 binds and sequesters RANKL. (A) rFBN1-N (circles), but not rFBN1-C (squares) or rF23 (triangles), binds RANKL. RANKL (20 $\mu\text{g}/\text{ml}$) was immobilized and rFBN1-N, rFBN1-C or rF23 (0–150 $\mu\text{g}/\text{ml}$) were added as soluble ligands, in Ca^{2+} -containing (open symbols) or Ca^{2+} -free (filled symbols) buffer, and bound proteins were detected. Non-specific binding to milk proteins without coated RANKL was subtracted from all values. Data are representative of three independent experiments. (B,C) RAW 264.7 cells were incubated with 50 $\mu\text{g}/\text{ml}$ of rFBN1-N (B) or rF23 (C) and increasing concentrations of RANKL (0.1, 1, 3 and 10 $\mu\text{g}/\text{ml}$), and osteoclast numbers were assessed. Data in B are means \pm s.e.m., $n=3-5$ experiments, * $P<0.05$ and ** $P<0.01$ indicate significance compared with PC; # $P<0.05$ indicates significance compared with the rFBN1-N treated sample including 0.1 $\mu\text{g}/\text{ml}$ RANKL, as analyzed by ANOVA. Data in C are means \pm s.e.m., $n=4$ experiments; ** $P<0.01$ compared with PC, as assessed by ANOVA.

resides in the cytosol. Calcium-dependent phosphatase calcineurin dephosphorylates NFATc1 allowing its nuclear translocation. Osteoclastogenesis from RAW 264.7 cells was induced in the absence or presence of fibrillin-1 fragments and after 3 days NFATc1 levels were monitored. None of the fibrillin-1 fragments affected NFATc1 protein levels (supplementary material Fig. S7). Using immunofluorescence we examined the subcellular localization of NFATc1 (Fig. 5A,B). Nuclear translocation of NFATc1 was prevented by rF23, but was not affected by rFBN1-C or rF18. In the presence of rFBN1-N, a trend of decreased nuclear localization of NFATc1 was observed. To investigate the consequences of fibrillin-1 fragments on cytosolic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$), RAW 264.7 cells were cultured with RANKL and rFBN1-N, rFBN1-C or rF23 for 3 days, and basal $[\text{Ca}^{2+}]_i$ of Fura-2 loaded osteoclast precursors was examined. The $[\text{Ca}^{2+}]_i$ in RANKL-treated precursors demonstrated characteristic oscillatory fluctuations (Fig. 5C, top), which were prevented by rF23 (Fig. 5C, bottom). For each cell we quantified the average $[\text{Ca}^{2+}]_i$ over 120 seconds of measurement, and the variation in $[\text{Ca}^{2+}]_i$ assessed as standard deviation from the average. Plotting average $[\text{Ca}^{2+}]_i$ and variation in $[\text{Ca}^{2+}]_i$ in the order of ascending basal $[\text{Ca}^{2+}]_i$ demonstrated that the cells with higher average $[\text{Ca}^{2+}]_i$ also exhibited higher levels of baseline variation (Fig. 5D). Treatment with rF23 resulted in significant decrease in average $[\text{Ca}^{2+}]_i$ and in variation of $[\text{Ca}^{2+}]_i$ (Fig. 5D,E). rFBN1-C did not affect calcium signaling in osteoclast precursors, but rFBN1-N significantly decreased both average and variation in $[\text{Ca}^{2+}]_i$ in osteoclast precursors (Fig. 5E).

Discussion

In this study, we discovered a novel role of fibrillin-1 in the regulation of osteoclast differentiation. Osteoclastogenesis was found to be negatively regulated by the N-terminal half of fibrillin-1 (rFBN1-N) and by a small 63 kDa fibrillin-1 N-terminal sub-fragment (rF23). This function was specific for osteoclasts and restricted to the application of soluble fibrillin-1




Fig. 5. Fibrillin-1 fragments act through the Ca^{2+} -NFATc1 pathway. RAW 264.7 cells were cultured untreated (negative control, NC), treated with RANKL (positive control, PC) or RANKL and 50 $\mu\text{g}/\text{ml}$ of rFBN1-N (1N), rFBN1-C (1C), rF23 (23) or rF18 (18) for 48 hours. (A) NFATc1 localization was determined by immunofluorescence (green). Nuclei were counterstained using DAPI (blue). Scale bar: 100 μm . (B) The percentage of cells exhibiting nuclear NFATc1 of the total number of cells. Data are means \pm s.d.; $n=220$ cells for positive control, $n=137$ cells for rFBN1-N-treated, $n=282$ cells for rFBN1-C-treated, $n=135$ cells for rF23-treated and $n=117$ cells for rF18-treated samples; $*P<0.05$ assessed by Student's *t*-test. (C–E) Osteoclast precursors were loaded with Fura-2AM, and $[\text{Ca}^{2+}]_i$ was monitored for 120 seconds. (C) Representative traces demonstrate changes in $[\text{Ca}^{2+}]_i$ in eight individual cells per experimental condition. (D) For each cell, the average $[\text{Ca}^{2+}]_i$ (top) and variation in baseline $[\text{Ca}^{2+}]_i$, measured as a standard deviation of basal levels (bottom) were determined and presented in an ascending order of the average $[\text{Ca}^{2+}]_i$; $n=59$ cells for positive control, $n=65$ cells for rF23-treated samples. (E) Average $[\text{Ca}^{2+}]_i$ (top) and variation in $[\text{Ca}^{2+}]_i$ (bottom). Data are means \pm s.d.; $n=59$ cells for PC, $n=65$ for rFBN1-C-treated cells, $n=85$ for rFBN1-N-treated cells, $n=65$ for rF23-treated cells; $*P<0.5$, $**P<0.01$ indicates significance compared with PC, as assessed by Student's *t*-test.

fragments to the apical cell surface. Similar fibrillin-1 fragments were found to be solubilized by the resorptive activity of differentiated osteoclasts. We have identified two differential mechanisms by which fibrillin-1 fragments can inhibit osteoclastogenesis. First, rFBN1-N sequestered RANKL. Second, rF23 prevented calcium fluctuations and NFATc1 translocation in differentiating osteoclasts. The data indicate that osteoclastic degradation of fibrillin-1 might provide a potent negative feedback that limits osteoclast formation and function.

In line with previous studies (Arteaga-Solis et al., 2001; Keene et al., 1991; Kitahama et al., 2000; Quondamatteo et al., 2002; Zhang et al., 1995), we have found that fibrillin-1 and fibrillin-2 are expressed during osteoblast differentiation and secreted into the extracellular matrix. Importantly, we demonstrate that fibrillin-1 is also proteolytically processed during osteoclastic resorption, resulting in several specific degradation products, which were identified in cell culture, as well as in the extracts of human vertebral bone. The significance of our study is that distinct fibrillin-1 fragments are capable of powerful direct inhibition of osteoclastogenesis. The osteoclast inhibitory properties were associated with two different fibrillin-1 sub-fragments, rFBN1-N and rF23. When added as soluble ligands, these proteins strongly inhibited osteoclast differentiation. In contrast, when rFBN1-N was coated on calcium phosphate, it did not affect osteoclast formation, but modulated the resorbing activity. These data suggest that coating fibrillin-1 on the calcium phosphate surface masks the osteoclast-inhibitory epitopes, which become exposed following substrate degradation and solubilization during osteoclastic resorption.

One of the prominent functions of fibrillins is their role in regulating TGF β and BMPs (Neptune et al., 2003; Nistala et al., 2010b; Sengle et al., 2008). However, we have found that the osteoclastogenesis system used in our study is not sensitive to

TGF β , and that interference with TGF β levels or signaling minimally affected the osteoclast-inhibitory efficacy of fibrillin-1 fragments. We hypothesized that fibrillin-1 may modulate the bioavailability of RANKL. We have found that rFBN1-N but not rF23 or rFBN1-C, interacts with RANKL in a Ca^{2+} -dependent manner, and that inhibition of osteoclastogenesis by rFBN1-N was rescued by increasing RANKL. Based on the differential binding of RANKL to rFBN1-N but not to rF23, the binding site must be located in the region outside of rF23 between EGF4 and cbEGF22. While most fibrillin-1-interacting proteins bind at the N-terminal region comprising rF23, proteoglycans, including perlecan and versican, and heparin/heparan sulfate interact with the central region between EGF4 and cbEGF22 (Isogai et al., 2002; Tiedemann et al., 2001; Tiedemann et al., 2005). In addition to a direct interaction with fibrillin-1, RANKL may also interact with fibrillin-1-bound heparan sulfate, which in turn contributes to RANKL sequestration (Ariyoshi et al., 2008).

We have identified other differences in osteoclastogenesis inhibiting mechanisms between rFBN1-N and rF23, despite the fact that rF23 is fully contained within rFBN1-N. Whereas RANKL-induced gene expression was not affected by rFBN1-N, rF23 selectively inhibited the mRNA expression of proteases cathepsin K and MMP-9, as well as fusion factor Dcstamp. Moreover, calcium/NFATc1 signaling was disturbed to a larger degree by rF23 compared with rFBN1-N. Calcium signaling, which leads to NFATc1 nuclear translocation, is generated by activation of co-stimulatory immune receptors linked to immunoreceptor tyrosine-based activation motif (ITAM)-harboring adaptors (Shinohara et al., 2008). Two ITAM adapter proteins, Fc receptor common c subunit γ (FcR γ) and DNAX-activating protein 12 (DAP12), were identified to be essential for osteoclastogenesis (Koga et al., 2004; Mócsai et al., 2004), and to be coupled to activation of osteoclast-associated immunoglobulin-like

receptor (OSCAR) and the triggering receptor expressed in myeloid cells-2 (TREM-2) (Koga et al., 2004; Negishi-Koga and Takayanagi, 2009). Interestingly, OSCAR has been recently identified as a receptor for extracellular matrix proteins, including triple helical collagen I, II and III (Barrow et al., 2011). While we have not identified the osteoclast receptor for rF23 in this study, it may be possible that OSCAR-induced positive osteoclastogenic signaling is inhibited by rF23.

Modulation of the negative feedback described in our study may contribute to the development of low bone mass in mouse models of MFS, *Fbn1*^{mgR/mgR} (Nistala et al., 2010a), and *Fbn1*^{-/-} (Nistala et al., 2010c), as well as to osteopenia in MFS patients and potentially other type I fibrillinopathies (Giordano et al., 1997). Irrespective of the underlying molecular pathogenetic mechanisms in MFS, including haploinsufficiency, enhanced proteolytic degradation, delayed secretion, and altered proprotein processing, the net effect is a reduced level of fully functional microfibrils in the extracellular matrix and enhanced TGF β levels (Hubmacher and Reinhardt, 2011). Decreased levels of microfibrils would result in a reduction in the absolute amount of fibrillin-1 degradation fragments generated in bone turnover. This in turn would release the negative feedback controlled by N-terminal fibrillin-1 fragments resulting in enhanced osteoclast formation and activity. MFS mutations typically render fibrillin-1 more susceptible to proteolysis (Kirschner et al., 2011; McGettrick et al., 2000; Reinhardt et al., 2000; Vollbrandt et al., 2004). Cathepsin K cleaves rFBN1-N exclusively in the linker regions between domains producing N-terminal fragments almost identical to the inhibitory rF23 through cleavage sites at Val⁴⁴⁴ and Leu⁴⁴⁵ (Kirschner et al., 2011). In contrast, most of the tested mutations resulted in enhanced degradation within disulfide-bonded loops of individual cbEGF or TB domains. Mutation-mediated cleavage events may also modulate interdomain processing by long-range structural effects that have been reported previously (Kirschner et al., 2011; Vollbrandt et al., 2004). In this regard it is important to note that the rF1F fragment tested in our study fully contains rF23, but lacks osteoclast inhibitory properties, which could be potentially explained by different folding of various length fibrillin-1 fragments. Thus it would be of interest to further validate if MFS mutation-mediated proteolysis of fibrillin-1 may result in degradation products with different conformations, which would in turn cause inappropriate stimulation of osteoclastogenesis.

Taken together with previously published data (Nistala et al., 2010c; Nistala et al., 2010d), our findings suggest a paradigm in which normal bone homeostasis is maintained by fibrillin-1 both indirectly, through regulating TGF β and RANKL levels, as well as directly by affecting osteoclast formation and function. In MFS and potentially other type I fibrillinopathies, the following consequences may occur: (1) TGF β levels produced by osteoblasts increase and induce RANKL expression (Nistala et al., 2010d), (2) compromised sequestration of RANKL by mutated fibrillin-1 additionally increases RANKL levels, and (3) production of small N-terminal inhibitory fragments is decreased either due to overall decrease in fibrillin-1 or due to its abnormal degradation. All these additive mechanisms result in elevated osteoclastogenesis and osteopenia.

Materials and Methods

Recombinant proteins, cytokines and growth factors

Recombinant fragments of human fibrillin-1, rFBN1-N (rF16), rFBN1-C (rF6H), rF18, rF23, rF1F, rF31 and rF51, were produced as described previously (El-Hallous et al., 2007; Jensen et al., 2001; Keene et al., 1997; Reinhardt et al., 1996a;

Reinhardt et al., 1996b), diluted at 50 μ g/ml in 50 mM Tris-HCl, pH 7.4, 150 mM NaCl (TBS) including 2 mM CaCl₂. TGF β 1 (1–50 ng/ml) was from Invitrogen (PHG9204), TGF β neutralizing antibody (15 μ g/ml) was from R&D (AB-100-NA) and T β RI inhibitor (5 μ M) was from Sigma (SB 431542). Recombinant GST-RANKL was purified from clones kindly provided by Dr M. F. Manolopoulos (University of Toronto) (Manolopoulos et al., 2003). MCSF was from Peprotech (300-25). For biotinylation, rFBN1-N was dialyzed in 20 mM HEPES, pH 7.4, 500 mM NaCl, 2.5 mM CaCl₂ and biotinylated using EZ-Link Sulfo-NHS-LC-Biotin (Thermo Scientific; 21335).

Osteoclast cultures

The mouse monocytic cell line RAW 264.7 obtained from the American Tissue Culture Collection (ATCC; TIB-71) was cultured in DMEM (Wisent; 319-020-CL) with 10% fetal bovine serum (FBS) (Wisent; 080152), 1 mM pyruvate (Wisent; 600-110-EL), and 1% penicillin, streptomycin, L-glutamine solution (Wisent; 450-202-EL). To generate osteoclasts, RAW 264.7 cells were plated at 5×10^3 cells/cm² and grown with RANKL (50 ng/ml) for 5–7 days with one medium change on day 3.

Studies were compliant with McGill University guidelines established by the Canadian Council on Animal Care. Mouse bone marrow was isolated as described previously (Hussein et al., 2011). Bone marrow cells were incubated overnight in α MEM (Wisent; 310-022-CL) supplemented with 10% FBS, 1 mM pyruvate, 1% penicillin, streptomycin, L-glutamine solution and 50 ng/ml MCSF. Non-adherent cells were harvested, plated at 5×10^4 cells/cm² and grown with MCSF (50 ng/ml) and RANKL (50 ng/ml) for 5–7 days with one medium change on day 3.

Cultures were fixed with 10% formalin and stained for TRAP (Sigma; 387A-KT). Osteoclasts were identified as TRAP-positive cells with three or more nuclei.

Extraction of human bone

Vertebrae from a 16-year-old female were obtained with consent through the Transplant Quebec Organ Donation Program, and stored at -70°C . Proteins were extracted from the mineralized bone matrix as described previously (Dickson and Roughley, 1993). Briefly, the bone was powdered, treated with 4 M guanidinium chloride to remove proteins present in soft connective tissues, then bone powder was treated with 4 M guanidinium chloride containing 0.2 M EDTA to extract proteins from the mineralized matrix. Proteins in the extract were precipitated with 9 volumes of ethanol and used directly for analysis by SDS-PAGE. Fibrillin-1 was immunoblotted using primary polyclonal antibody against rFBN1-N (1:100) (Lin et al., 2002).

Osteoclastic resorption

Osteoassay plates (Corning; 3989) were preincubated with complete medium and 50 μ g/ml of rFBN1-N, rFBN1-C or BSA overnight at 37°C. To quantify resorption, the cells were removed with 10% NaClO, the wells were washed with water and air dried. Resorption pits were measured using PixelINK Capture SE software.

Animal model

Studies were compliant with McGill University guidelines established by the Canadian Council on Animal Care. Male Balb/c mice (6 weeks) were injected intra-peritoneally with vehicle (PBS, $n=11$) or 50 μ g/kg of rF31 ($n=4$) every other day for 1 week. The mice were euthanized 1 week after the last injection, and hind limbs were dissected immediately. Tibias were decalcified in EDTA, embedded in paraffin, sectioned (5 μ m thickness) and stained for TRAP in the histology platform of the Centre for Bone and Periodontal Research, McGill University. TRAP-positive cells were counted 100 μ m below the growth plate, in two fields per mouse using Osteomeasure 3.0 (Osteometrics).

Quantitative real-time PCR (qPCR)

Total RNA was isolated as described previously (Guo et al., 2008; Hussein et al., 2011), quantified with a Quant-iT instrument (Invitrogen; #Q32860) and reverse transcribed using the High Capacity cDNA Reverse Transcription Kit (Life Technologies; #4368814). PCR was performed with a 7500 Applied Biosystem instrument using TaqMan probes with the universal PCR Master Mix (Life Technologies), or SYBR Green primers and PCR Master Mix (Hoffmann-La Roche Ltd) in a total volume of 25 μ l including 0.9 μ M of reverse and forward primers from Life Technologies. The following probes were used: Taqman: *Ctsk* (Cathepsin K): Mm00484039_m1; *Mmp9* (Matrix metalloproteinase 9): Mm00600163_m1; *Fbn1* (Fibrillin-1): Mm00514908_m1; *Fbn2* (Fibrillin-2): Mm00515742_m1; *Tnfrsf11a* (Rank): Mm00437132_m1; *Acp5* (Trap): Mm00475698_m1; *Sp7* (Osterix): Mm00504574_m1; *Colla1* (Collagen I alpha 1): Mm00801666_g1; *Gapdh* (Glyceraldehyde 3-phosphate dehydrogenase): Mm99999915_g1. SYBR Green (forward, reverse): *Itgb3* (Integrin β 3): (TCCAGACCCTGGGTACCAAG, GCCAATCCGAAGTTGCTAG); *Dcstamp* (Dendritic cell-specific transmembrane protein): (CTTCCGTGGGCCAGAAGTT, AGGCCAGTCTGACTAGGATGA); *Opn* (Osteopontin): (GTGGACTC-GGATGAATCTG, TCGACTGTAGGGACGATTG); *Runx2* (Runt-related

transcription factor 2): (TGGCTTGGGTTTCAGGTTAG, TCGGTTTCTTAGG-GTCTTGGGA); *Gapdh*: (CAAGTATGATGACATCAAGAAGGTGG, GGAAGA-GTGGGAGTTGCTGTTG).

Immunoblotting

Whole cell lysates were prepared as described previously (Tiedemann et al., 2009). Briefly, the cells were lysed using a RIPA buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Nonidet P-40, 1 mM EDTA, 10 mg/ml aprotinin, 1 mg/ml leupeptin, 0.1 M phenylmethylsulfonyl fluoride, 0.5 M sodium fluoride and 1 mM sodium orthovanadate). The protein concentrations were determined using a Quant-iT protein assay kit (Invitrogen). Conditioned media were collected and filtered using a 0.22 µm filter, aliquots were kept at -80°C. Proteins (50 µg) were precipitated in 10% trichloroacetic acid, redissolved in 1× SDS sample buffer, separated by 7.5% gel electrophoresis under reducing conditions, transferred to a nitrocellulose membrane (Bio-Rad, #162-0115; 0.45 µm) using 10 mM sodium borate, pH 8.9, blocked in 5% non-fat milk in TBST buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.05% Tween 20) for 1 h at 4°C followed by overnight incubation at 4°C with the primary antibodies: NFATc1 (1:100, sc-7294, Santa Cruz Biotechnology), tubulin (1:5000, T6793, Sigma), hexahistidine tag (1:5,000, R930-25, Invitrogen); rFBN1-N (1:100), rFBN1-C (1:250), rFBN2-C (1:250) (Lin et al., 2002; Tiedemann et al., 2001; Tiedemann et al., 2005); then with horseradish peroxidase-conjugated secondary antibodies (1:200–1:10,000, Bio-Rad; goat anti-mouse, #170-5047; goat anti-rabbit #170-5046) or NeutrAvidin horseradish peroxidase conjugate (Thermo Scientific, #31001, 1:800) and visualized using a chemiluminescence substrate (Thermo Scientific, #32106).

Immunofluorescence

Immunostaining for NFATc1 was performed as described previously (Guo et al., 2008). Briefly, osteoclasts on glass coverslips were fixed with 10% formalin, incubated with mouse anti-NFATc1 antibody (1:100, Santa Cruz Biotechnology; #SC-7294), followed by a biotinylated goat anti-mouse IgG (1:200, Invitrogen; #A10519) and Alexa-Fluor-488-conjugated streptavidin (1:500, Invitrogen; #S-11223). Nuclei were counterstained using 4',6-diamidino-2-phenylindole dihydrochloride (DAPI; 1:10,000, Invitrogen; #D1306). For each experiment, five random images per condition were recorded. NFATc1 localization was considered nuclear when the fluorescence intensity of nuclei exceeded that of the cytoplasm.

Osteoblasts were fixed with 10% formalin, and immunofluorescence of fibrillin-1 and fibrillin-2 was performed as described previously with primary antibodies against rFBN1-C (1:100) and rFBN2-C (1:100) (Lin et al., 2002; Tiedemann et al., 2001; Tiedemann et al., 2005).

Solid-phase binding assay

Solid-phase binding assays were performed as described previously (Lin et al., 2002). Briefly, 96-well plates (MaxiSorp, Nalge Nunc International) were coated overnight in 4°C with RANKL (20 µg/ml) in TBS or TBS alone, and blocked for 1 hour with 5% (w/v) non-fat milk in TBS. rFBN1-N, rFBN1-C or rF23 were incubated as soluble ligands (0–150 µg/ml) in TBS with 5% non-fat milk, with either 5 mM CaCl₂ or 10 mM EDTA at room temperature, and washed with TBS containing 0.05% (v/v) Tween 20. The wells were incubated for 1 hour with the primary antibody against rFBN1-N [1:250, (Lin et al., 2002)], followed by horseradish-peroxidase-conjugated secondary goat anti-rabbit antibody (1:800 diluted; Bio-Rad) for 1 hour. Color development was performed with 1 mg/ml 5-aminosalicylic acid (Sigma) in 20 mM phosphate buffer, pH 6.8, including 0.1% H₂O₂ for 3–5 minutes, stopped with 2 M NaOH, and measured at 492 nm using a Microplate reader. OD measured in control wells (blocked with non-fat milk, but without immobilized ligand) were subtracted from all measured values.

Microspectrofluorometry

Cells plated onto glass-bottom 35 mm dishes (MatTek.) were loaded with Fura-2-AM (Invitrogen; F1221) at room temperature for 40 minutes and changes in [Ca²⁺]_i were measured for 120 seconds, as described previously (Tiedemann et al., 2009). [Ca²⁺]_i values were calculated using a Fura-2-AM calcium imaging calibration kit (Invitrogen; F6774).

Statistics

Data are presented as means ± standard error of the mean (s.e.m.) with sample size (*n*) indicating the number of independent experiments, or as means ± standard deviation (s.d.) with sample size (*n*) indicating the number of samples. Differences were assessed by ANOVA or Student's *t*-test and accepted as statistically significant at *P*<0.05.

Acknowledgements

We are grateful to Dr Morris F. Manolson for providing GST-RANKL clones and to Dr Lisbet Haglund for preparing the human bone sample, and to Ms Amelie Pagliuzza for performing a control

ELISA and Ms Meng Lan Li for helping with histomorphometric analysis.

Author contributions

K.T., D.P.R. and S.V.K. designed the experiments, interpreted the results and prepared the manuscript. K.T. performed the experiments. I.B.D. contributed to experimental design and performed the experiments. I.R. performed key preliminary experiments. J.K. prepared recombinant fibrillin fragments. P.R. provided human bone samples. All authors read and approved the manuscript.

Funding

This work was supported by the Canadian Institutes of Health Research [grant number MOP-106494 to D.P.R. and MOP-77643 to S.V.K.]; the Réseau de recherche en santé buccodentaire et osseuse; the Natural Sciences and Engineering Research Council of Canada [grant number RGPIN 375738-09]. D.P.R. holds a Canada Research Chair in Cell-Matrix Biology and S.V.K. holds a Canada Research Chair in Osteoclast Biology.

Supplementary material available online at

<http://jcs.biologists.org/lookup/suppl/doi:10.1242/jcs.127571/-/DC1>

References

- Ariyoshi, W., Takahashi, T., Kanno, T., Ichimiya, H., Shinmyouzu, K., Takano, H., Koseki, T. and Nishihara, T. (2008). Heparin inhibits osteoclastic differentiation and function. *J. Cell. Biochem.* **103**, 1707-1717.
- Arteaga-Solis, E., Gayraud, B., Lee, S. Y., Shum, L., Sakai, L. and Ramirez, F. (2001). Regulation of limb patterning by extracellular microfibrils. *J. Cell Biol.* **154**, 275-281.
- Balkan, W., Martinez, A. F., Fernandez, I., Rodriguez, M. A., Pang, M. and Troen, B. R. (2009). Identification of NFAT binding sites that mediate stimulation of cathepsin K promoter activity by RANK ligand. *Gene* **446**, 90-98.
- Barrow, A. D., Raynal, N., Andersen, T. L., Slatter, D. A., Bihan, D., Pugh, N., Cella, M., Kim, T., Rho, J., Negishi-Koga, T. et al. (2011). OSCAR is a collagen receptor that costimulates osteoclastogenesis in DAP12-deficient humans and mice. *J. Clin. Invest.* **121**, 3505-3516.
- Carter, N., Duncan, E. and Wordsworth, P. (2000). Bone mineral density in adults with Marfan syndrome. *Rheumatology (Oxford)* **39**, 307-309.
- Dickson, I. R. and Roughley, P. J. (1993). The effects of vitamin D deficiency on proteoglycan and hyaluronate constituents of chick bone. *Biochim. Biophys. Acta* **1181**, 15-22.
- Doyle, J. J., Gerber, E. E. and Dietz, H. C. (2012). Matrix-dependent perturbation of TGFβ signaling and disease. *FEBS Lett.* **586**, 2003-2015.
- El-Hallous, E., Sasaki, T., Hubmacher, D., Getie, M., Tiedemann, K., Brinckmann, J., Bätge, B., Davis, E. C. and Reinhardt, D. P. (2007). Fibrillin-1 interactions with fibulins depend on the first hybrid domain and provide an adaptor function to tropoelastin. *J. Biol. Chem.* **282**, 8935-8946.
- Giannoudis, P. V., Kanakaris, N. K. and Einhorn, T. A. (2007). Interaction of bone morphogenetic proteins with cells of the osteoclast lineage: review of the existing evidence. *Osteoporos. Int.* **18**, 1565-1581.
- Giordano, N., Senesi, M., Battisti, E., Mattii, G. and Gennari, C. (1997). Weill-Marchesani syndrome: report of an unusual case. *Calcif. Tissue Int.* **60**, 358-360.
- Grahame, R. and Pyeritz, R. E. (1995). The Marfan syndrome: joint and skin manifestations are prevalent and correlated. *Br. J. Rheumatol.* **34**, 126-131.
- Guo, Y., Tiedemann, K., Khalil, J. A., Russo, C., Siegel, P. M. and Komarova, S. V. (2008). Osteoclast precursors acquire sensitivity to breast cancer derived factors early in differentiation. *Bone* **43**, 386-393.
- Hubmacher, D. and Reinhardt, D. P. (2011). Microfibrils and fibrillin. In *Biology of Extracellular Matrix*, (ed. R. P. Mecham), pp. 233-265. New York, NY: Springer.
- Hubmacher, D., Tiedemann, K. and Reinhardt, D. P. (2006). Fibrillins: from biogenesis of microfibrils to signaling functions. *Curr. Top. Dev. Biol.* **75**, 93-123.
- Hussein, O., Tiedemann, K. and Komarova, S. V. (2011). Breast cancer cells inhibit spontaneous and bisphosphonate-induced osteoclast apoptosis. *Bone* **48**, 202-211.
- Ishida, N., Hayashi, K., Hoshijima, M., Ogawa, T., Koga, S., Miyatake, Y., Kumegawa, M., Kimura, T. and Takeya, T. (2002). Large scale gene expression analysis of osteoclastogenesis in vitro and elucidation of NFAT2 as a key regulator. *J. Biol. Chem.* **277**, 41147-41156.
- Isogai, Z., Asperger, A., Keene, D. R., Ono, R. N., Reinhardt, D. P. and Sakai, L. Y. (2002). Versican interacts with fibrillin-1 and links extracellular microfibrils to other connective tissue networks. *J. Biol. Chem.* **277**, 4565-4572.
- Janssens, K., ten Dijke, P., Janssens, S. and Van Hul, W. (2005). Transforming growth factor-beta1 to the bone. *Endocr. Rev.* **26**, 743-774.
- Jensen, S. A., Reinhardt, D. P., Gibson, M. A. and Weiss, A. S. (2001). Protein interaction studies of MAGP-1 with tropoelastin and fibrillin-1. *J. Biol. Chem.* **276**, 39661-39666.

- Kaur, J. and Reinhardt, D. P. (2012). Immobilized metal affinity chromatography copurifies TGF- β 1 with histidine-tagged recombinant extracellular proteins. *PLoS ONE* **7**, e48629.
- Keene, D. R., Sakai, L. Y. and Burgeson, R. E. (1991). Human bone contains type III collagen, type VI collagen, and fibrillin: type III collagen is present on specific fibers that may mediate attachment of tendons, ligaments, and periosteum to calcified bone cortex. *J. Histochem. Cytochem.* **39**, 59-69.
- Keene, D. R., Jordan, C. D., Reinhardt, D. P., Ridgway, C. C., Ono, R. N., Corson, G. M., Fairhurst, M., Sussman, M. D., Memoli, V. A. and Sakai, L. Y. (1997). Fibrillin-1 in human cartilage: developmental expression and formation of special banded fibers. *J. Histochem. Cytochem.* **45**, 1069-1082.
- Kirschner, R., Hubmacher, D., Iyengar, G., Kaur, J., Fagotto-Kaufmann, C., Brömme, D., Bartels, R. and Reinhardt, D. P. (2011). Classical and neonatal Marfan syndrome mutations in fibrillin-1 cause differential protease susceptibilities and protein function. *J. Biol. Chem.* **286**, 32810-32823.
- Kitahama, S., Gibson, M. A., Hatzinikolas, G., Hay, S., Kuliwaba, J. L., Evdokiou, A., Atkins, G. J. and Findlay, D. M. (2000). Expression of fibrillins and other microfibril-associated proteins in human bone and osteoblast-like cells. *Bone* **27**, 61-67.
- Koga, T., Inui, M., Inoue, K., Kim, S., Suematsu, A., Kobayashi, E., Iwata, T., Ohnishi, H., Matozaki, T., Kodama, T. et al. (2004). Costimulatory signals mediated by the ITAM motif cooperate with RANKL for bone homeostasis. *Nature* **428**, 758-763.
- Kohlmeier, L., Gasner, C., Bachrach, L. K. and Marcus, R. (1995). The bone mineral status of patients with Marfan syndrome. *J. Bone Miner. Res.* **10**, 1550-1555.
- Le Goff, C. and Cormier-Daire, V. (2012). From tall to short: the role of TGF β signaling in growth and its disorders. *Am. J. Med. Genet. C. Semin. Med. Genet.* **160C**, 145-153.
- Le Parc, J. M., Plantin, P., Jondeau, G., Goldschild, M., Albert, M. and Boileau, C. (1999). Bone mineral density in sixty adult patients with Marfan syndrome. *Osteoporos. Int.* **10**, 475-479.
- Lin, G., Tiedemann, K., Vollbrandt, T., Peters, H., Batge, B., Brinckmann, J. and Reinhardt, D. P. (2002). Homo- and heterotypic fibrillin-1 and -2 interactions constitute the basis for the assembly of microfibrils. *J. Biol. Chem.* **277**, 50795-50804.
- Manolson, M. F., Yu, H., Chen, W., Yao, Y., Li, K., Lees, R. L. and Heersche, J. N. (2003). The $\alpha 3$ isoform of the 100-kDa V-ATPase subunit is highly but differentially expressed in large (≥ 10 nuclei) and small (≤ 10 nuclei) osteoclasts. *J. Biol. Chem.* **278**, 49271-49278.
- McGettrick, A. J., Knott, V., Willis, A. and Handford, P. A. (2000). Molecular effects of calcium binding mutations in Marfan syndrome depend on domain context. *Hum. Mol. Genet.* **9**, 1987-1994.
- Mócsai, A., Humphrey, M. B., Van Ziffle, J. A., Hu, Y., Burghardt, A., Spusta, S. C., Majumdar, S., Lanier, L. L., Lowell, C. A. and Nakamura, M. C. (2004). The immunomodulatory adapter proteins DAP12 and Fc receptor gamma-chain (FcRgamma) regulate development of functional osteoclasts through the Syk tyrosine kinase. *Proc. Natl. Acad. Sci. USA* **101**, 6158-6163.
- Moura, B., Tubach, F., Sulpice, M., Boileau, C., Jondeau, G., Muti, C., Chevallier, B., Ounnoughene, Y., Le Parc, J. M.; Multidisciplinary Marfan Syndrome Clinic Group (2006). Bone mineral density in Marfan syndrome. A large case-control study. *Joint Bone Spine* **73**, 733-735.
- Negishi-Koga, T. and Takayanagi, H. (2009). Ca²⁺-NFATc1 signaling is an essential axis of osteoclast differentiation. *Immunol. Rev.* **231**, 241-256.
- Neptune, E. R., Frischmeyer, P. A., Arking, D. E., Myers, L., Bunton, T. E., Gayraud, B., Ramirez, F., Sakai, L. Y. and Dietz, H. C. (2003). Dysregulation of TGF-beta activation contributes to pathogenesis in Marfan syndrome. *Nat. Genet.* **33**, 407-411.
- Nistala, H., Lee-Arteaga, S., Carta, L., Cook, J. R., Smaldone, S., Siciliano, G., Rifkin, A. N., Dietz, H. C., Rifkin, D. B. and Ramirez, F. (2010a). Differential effects of alendronate and losartan therapy on osteopenia and aortic aneurysm in mice with severe Marfan syndrome. *Hum. Mol. Genet.* **19**, 4790-4798.
- Nistala, H., Lee-Arteaga, S., Siciliano, G., Smaldone, S. and Ramirez, F. (2010b). Extracellular regulation of transforming growth factor beta and bone morphogenetic protein signaling in bone. *Ann. N. Y. Acad. Sci.* **1192**, 253-256.
- Nistala, H., Lee-Arteaga, S., Smaldone, S., Siciliano, G., Carta, L., Ono, R. N., Sengle, G., Arteaga-Solis, E., Levasseur, R., Ducy, P. et al. (2010c). Fibrillin-1 and -2 differentially modulate endogenous TGF- β and BMP bioavailability during bone formation. *J. Cell Biol.* **190**, 1107-1121.
- Nistala, H., Lee-Arteaga, S., Smaldone, S., Siciliano, G. and Ramirez, F. (2010d). Extracellular microfibrils control osteoblast-supported osteoclastogenesis by restricting TGFbeta stimulation of RANKL production. *J. Biol. Chem.* **285**, 34126-34133.
- Pyeritz, R. E. (2000). The Marfan syndrome. *Annu. Rev. Med.* **51**, 481-510.
- Quinn, J. M., Itoh, K., Udagawa, N., Hausler, K., Yasuda, H., Shima, N., Mizuno, A., Higashio, K., Takahashi, N., Suda, T. et al. (2001). Transforming growth factor beta affects osteoclast differentiation via direct and indirect actions. *J. Bone Miner. Res.* **16**, 1787-1794.
- Quondamatteo, F., Reinhardt, D. P., Charbonneau, N. L., Pophal, G., Sakai, L. Y. and Herken, R. (2002). Fibrillin-1 and fibrillin-2 in human embryonic and early fetal development. *Matrix Biol.* **21**, 637-646.
- Ramirez, F., Sakai, L. Y., Rifkin, D. B. and Dietz, H. C. (2007). Extracellular microfibrils in development and disease. *Cell. Mol. Life Sci.* **64**, 2437-2446.
- Reinhardt, D. P., Keene, D. R., Corson, G. M., Pöschl, E., Bächinger, H. P., Gambree, J. E. and Sakai, L. Y. (1996a). Fibrillin-1: organization in microfibrils and structural properties. *J. Mol. Biol.* **258**, 104-116.
- Reinhardt, D. P., Sasaki, T., Dzamba, B. J., Keene, D. R., Chu, M. L., Göhring, W., Timpl, R. and Sakai, L. Y. (1996b). Fibrillin-1 and fibulin-2 interact and are colocalized in some tissues. *J. Biol. Chem.* **271**, 19489-19496.
- Reinhardt, D. P., Ono, R. N., Notbohm, H., Müller, P. K., Bächinger, H. P. and Sakai, L. Y. (2001). Mutations in calcium-binding epidermal growth factor modules render fibrillin-1 susceptible to proteolysis. A potential disease-causing mechanism in Marfan syndrome. *J. Biol. Chem.* **275**, 12339-12345.
- Sabatier, L., Miosge, N., Hubmacher, D., Lin, G., Davis, E. C. and Reinhardt, D. P. (2011). Fibrillin-3 expression in human development. *Matrix Biol.* **30**, 43-52.
- Sengle, G., Charbonneau, N. L., Ono, R. N., Sasaki, T., Alvarez, J., Keene, D. R., Bächinger, H. P. and Sakai, L. Y. (2008). Targeting of bone morphogenetic protein growth factor complexes to fibrillin. *J. Biol. Chem.* **283**, 13874-13888.
- Shinohara, M., Koga, T., Okamoto, K., Sakaguchi, S., Arai, K., Yasuda, H., Takai, T., Kodama, T., Morio, T., Geha, R. S. et al. (2008). Tyrosine kinases Btk and Tec regulate osteoclast differentiation by linking RANK and ITAM signals. *Cell* **132**, 794-806.
- Takayanagi, H., Kim, S., Koga, T., Nishina, H., Isshiki, M., Yoshida, H., Saiura, A., Isobe, M., Yokochi, T., Inoue, J. et al. (2002). Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev. Cell* **3**, 889-901.
- Tiedemann, K., Bätge, B., Müller, P. K. and Reinhardt, D. P. (2001). Interactions of fibrillin-1 with heparin/heparan sulfate, implications for microfibrillar assembly. *J. Biol. Chem.* **276**, 36035-36042.
- Tiedemann, K., Sasaki, T., Gustafsson, E., Göhring, W., Bätge, B., Notbohm, H., Timpl, R., Wedel, T., Schlötzer-Schrehardt, U. and Reinhardt, D. P. (2005). Microfibrils at basement membrane zones interact with perlecan via fibrillin-1. *J. Biol. Chem.* **280**, 11404-11412.
- Tiedemann, K., Hussein, O., Sadvakassova, G., Guo, Y., Siegel, P. M. and Komarova, S. V. (2009). Breast cancer-derived factors stimulate osteoclastogenesis through the Ca²⁺/protein kinase C and transforming growth factor-beta/MAPK signaling pathways. *J. Biol. Chem.* **284**, 33662-33670.
- Vollbrandt, T., Tiedemann, K., El-Hallous, E., Lin, G., Brinckmann, J., John, H., Bätge, B., Notbohm, H. and Reinhardt, D. P. (2004). Consequences of cysteine mutations in calcium-binding epidermal growth factor modules of fibrillin-1. *J. Biol. Chem.* **279**, 32924-32931.
- Zhang, H., Hu, W. and Ramirez, F. (1995). Developmental expression of fibrillin genes suggests heterogeneity of extracellular microfibrils. *J. Cell Biol.* **129**, 1165-1176.

Kerstin Tiedemann, Iris Boraschi-Diaz, Irina Rajakumar, Jasvir Kaur, Peter Roughley, Dieter P. Reinhardt and Svetlana V. Komarova

Supplemental Figure S1.




Fig. S1. Osteoblasts produce fibrillin-1 and -2. Mouse bone marrow cells were treated with ascorbic acid (AA; 50 $\mu\text{g/ml}$). (A) Osteoblast differentiation was monitored by the expression of alkaline phosphatase (ALP). (B) The expression of osteoblast marker genes including Runt-related transcription factor 2 (*Runx2*), osteopontin (*Opn*), and osterix (*Sp7*) was assessed during the differentiation period. *Runx2*, *Opn* and *Sp7* reached maximum expression after 2 weeks of culture and remained up-regulated during the third week of differentiation. (C) The expression of the extracellular matrix proteins by osteoblastic cells. Collagen type IA (*Col1a1*) gradually increased over the first two weeks of culture and remained up-regulated thereafter. Both fibrillin-1 (*Fbn1*) and fibrillin-2 (*Fbn2*) mRNA were strongly induced during the first week of osteoblast differentiation. In the subsequent weeks, *Fbn1* mRNA remained up-regulated, but the expression of *Fbn2* mRNA decreased to levels similar to undifferentiated cells. Data are means \pm SEM, $n = 3-5$ experiments, * $p < 0.05$ indicates significance compared to week 0 assessed by ANOVA. (D) RAW 264.7 cells (RAW) or differentiated osteoclasts (OC) do not express *Fbn1*. Mature osteoblasts (OB) were used as a positive control. (E) Differentiated osteoblasts (OB) secreted fibrillin-1 and -2 into the conditioned medium. In contrast, untreated bone marrow cells (negative control, NC) did not secrete detectable amounts of fibrillin-1 or -2. (F) Development of fibrillin-1-containing microfibrils was apparent by immunofluorescence during the first week (W1) of osteoblast differentiation. (G) Only diffuse staining was observed for fibrillin-2 in osteoblast cultures.

Kerstin Tiedemann, Iris Boraschi-Diaz, Irina Rajakumar, Jasvir Kaur, Peter Roughley, Dieter P. Reinhardt and Svetlana V. Komarova

Supplemental Figure S2.




Fig. S2. Bovine serum albumin (BSA) does not affect osteoclast formation. RAW 264.7 cells (black bars) and mouse bone marrow cells (white bars) were cultured for 5 days untreated (negative control, NC), treated with RANKL only (50 ng/ml, positive control, PC), or treated with RANKL and BSA (50 μ g/ml), or soluble fibrillin-1 fragments rFBN1-N or rFBN1-C (50 μ g/ml). Samples were fixed and stained for TRAP. Shown are representative images of osteoclasts formed in bone marrow cultures under the indicated conditions. Scale bar of 250 μ m applies to all images. The addition of BSA did not affect osteoclast formation, only addition of rFBN1-N resulted in significant decrease in osteoclast numbers. Data are means \pm SEM, n = 3-5 independent experiments, *p<0.05, ***p<0.001 compared to PC.

Kerstin Tiedemann^{1,2}, Iris Boraschi-Diaz^{1,2}, Irina Rajakumar^{1,#}, Jasvir Kaur³, Peter Roughley², Dieter P. Reinhardt^{1,3,†} and Svetlana V. Komarova^{1,2,3,†*}

Supplemental Information S3.




Fig. S3. Effect of fibrillin-1 fragments on osteoclast formation and gene expression. (A) Representative images of osteoclasts formed from RAW 264.7 monocytes after addition of soluble rFBN1-N, rFBN1-C, rF23, rF51, rF1F, or rF18 (50 µg/ml) for the duration of the experiment. Scale bar applies to all images. (B) RAW 264.7 cells were cultured for 5 days untreated, treated with RANKL only (50 ng/ml), or treated with RANKL and soluble rFBN1-N, rFBN1-C, rF23, rF51, rF1F, or rF18 (50 µg/ml), and mRNA expression of tartrate resistant acid phosphatase (*Acp5*), receptor activator of nuclear factor B (*Tnfrsf11a*), metalloproteinase 9 (*Mmp9*) and integrin β3 (*Itgb3*) was assessed. RANKL treatment alone significantly increased gene expression of *Acp5* and *Mmp9*. rF23, rF51, and rF1F significantly decreased gene expression of *Mmp9*. Data are means ± SEM, n = 3 experiments, ***p < 0.001, compared to PC as assessed by t test; # p < 0.05, compared to NC as assessed by t-test. (C) Bone marrow cells were cultured with 50 ng/ml MCSF and RANKL for 4 days with or without rF23 and rFBN1-N in concentrations of 6.25, 12.5, 25 and 50 µg/ml. (D) Bone marrow cells were cultured with 50 ng/ml MCSF and RANKL for 4 days with or without rF23 and rFBN1-N added for the duration of the experiment (day 1-4, D1-4), for the first 2 days only (D1-2), or for the last 2 days only (D3-4). Data are means ± SEM, n = 3 experiments normalized to osteoclast numbers observed in positive control (PC), *p < 0.05, **p < 0.01, ***p < 0.001 compared to positive control as assessed by t-test; # p < 0.05, compared to negative control (NC) as determined by t-test.

Fibrillin-1 directly regulates osteoclast formation and function by a dual mechanism

Kerstin Tiedemann^{1,2}, Iris Boraschi-Diaz^{1,2}, Irina Rajakumar^{1,#}, Jasvir Kaur³, Peter Roughley², Dieter P. Reinhardt^{1,3,¶} and Svetlana V. Komarova^{1,2,3,¶*}

Supplemental Figure S4.




Fig. S4. Fibrillin-1 has no effect on osteoblast differentiation. Osteoblast differentiation of mouse bone marrow cells was induced with ascorbic acid (50 $\mu\text{g}/\text{ml}$) in the absence (positive control, PC) or presence of vehicle (Tris-based buffer) or soluble fibrillin-1 fragments (50 $\mu\text{g}/\text{ml}$), rFBN1-N, rFBN1-C, rF23 (23), rF51 (51), rF1F (1F) and rF18 (18). After 1 week, cells were fixed and stained for ALP (red). Shown are representative images of osteoblastic cells formed under different conditions. Scale bar applies to all images.

Kerstin Tiedemann, Iris Boraschi-Diaz, Irina Rajakumar, Jasvir Kaur, Peter Roughley, Dieter P. Reinhardt and Svetlana V. Komarova

Supplemental Figure S5.




Fig. S5. TGFβ does not affect osteoclastogenesis in the cell culture model used. (A) RAW 264.7 cells were cultured for 5 days untreated or treated with RANKL (50 ng/ml) and recombinant TGFβ1 (1-50 ng/ml) and osteoclast numbers were quantified. (B) Mouse bone marrow cells were treated with MCSF (50 ng/ml) and RANKL (50 ng/ml) in the presence of TGFβ receptor I inhibitor (TβRI inh, 5 μM) or neutralizing antibody against TGFβ1 (TGFβ ab, 15 μg/ml) and osteoclast numbers were determined. For A and B, data are means ± SEM, n=3-5 experiments, no significant differences between RANKL-treated conditions. (C) Inhibition of TGFβ minimally affects anti-osteoclastogenic effect of fibrillin-1 fragments. Bone marrow cells were cultured for 5 days with MCSF and RANKL without (positive control) or with 50 μg/ml of rFBN1-N or rF23 in the absence or presence of TβRI inhibitor (5 μM) or TGFβ neutralizing antibody (15 μg/ml), and osteoclast numbers were counted. Data are means ± SEM, n = 3-5 experiments, *p < 0.001 indicates significance compared to positive control and #p < 0.05 indicates significance compared to the samples without inhibitors, as determined by t-test.

Fibrillin-1 directly regulates osteoclast formation and function by a dual mechanism

Kerstin Tiedemann^{1,2}, Iris Boraschi-Diaz^{1,2}, Irina Rajakumar^{1,#}, Jasvir Kaur³, Peter Roughley², Dieter P. Reinhardt^{1,3,¶} and Svetlana V. Komarova^{1,2,3,¶*}

Supplemental Figure S6.




Fig. S6. Efficiency of the detector antibody for rFBN1-N and its fragments rF18 and rF23. ELISA demonstrates similar titers for rFBN1-N and the tested fragments. Background values without coated proteins were ≤ 0.075 . Data are means \pm SD, $n = 3$ for rFBN1-N and rF23, and $n = 2$ for rF18.

Supplemental Figure S7.




Fig. S7. NFAT protein levels are not affected by fibrillin-1 fragments. RAW 264.7 cells were either untreated (NC) or treated with RANKL (50 ng/ml, PC) alone or RANKL and rFBN1-C (1C), rFBN1-N (1N), rF23 (23), rF51 (51), rF1F (1F), rF18 (18) or BSA (50 μ g/ml) for 3 days. Cell lysates were collected and NFATc1 protein levels were assessed using immunoblotting, α -tubulin was used as a loading control. Fibrillin-1 fragments did not affect protein level of NFATc1. Data are representative of three independent experiments.