

Pbx homeodomain proteins direct Myod activity to promote fast-muscle differentiation

Lisa Maves^{1,*}, Andrew Jan Waskiewicz², Biswajit Paul¹, Yi Cao¹, Ashlee Tyler¹, Cecilia B. Moens³ and Stephen J. Tapscott¹

The basic helix-loop-helix (bHLH) transcription factor Myod directly regulates gene expression throughout the program of skeletal muscle differentiation. It is not known how a Myod-driven myogenic program is modulated to achieve muscle fiber-type-specific gene expression. Pbx homeodomain proteins mark promoters of a subset of Myod target genes, including myogenin (*Myog*); thus, Pbx proteins might modulate the program of myogenesis driven by Myod. By inhibiting Pbx function in zebrafish embryos, we show that Pbx proteins are required in order for Myod to induce the expression of a subset of muscle genes in the somites. In the absence of Pbx function, expression of *myog* and of fast-muscle genes is inhibited, whereas slow-muscle gene expression appears normal. By knocking down Pbx or Myod function in combination with another bHLH myogenic factor, *Myf5*, we show that Pbx is required for Myod to regulate fast-muscle, but not slow-muscle, development. Furthermore, we show that Sonic hedgehog requires Myod in order to induce both fast- and slow-muscle markers but requires Pbx only to induce fast-muscle markers. Our results reveal that Pbx proteins modulate Myod activity to drive fast-muscle gene expression, thus showing that homeodomain proteins can direct bHLH proteins to establish a specific cell-type identity.

KEY WORDS: Myod, Pbx, Muscle, Zebrafish

INTRODUCTION

Basic helix-loop-helix (bHLH) transcription factors are crucial regulators, in some instances referred to as ‘master regulators’, of cell-type identities. Among the bHLH proteins, the Myod family drives skeletal myogenesis, similar to the regulation of neurogenesis by NeuroD proteins and lymphocyte development by E proteins (Tapscott, 2005; Lee, 1997; Quong et al., 2002). Homeodomain transcription factors have also been termed ‘master regulators’, but typically of organ or positional identities. For example, the Hox proteins control segmental identities, and Pax6 controls eye development (Pearson et al., 2005; Gehring, 1996). How are positional-identity programs linked with cell-type-identity programs, so that neurons in the eye acquire a different phenotype to neurons in the brain, or that neurons in one hindbrain segment differ from those in another segment? Although the mechanism is not known, one possibility is that positional-identity factors modulate the activity of cell-type-identity factors. As suggested by Westerman et al. (Westerman et al., 2003), homeodomain proteins might act in their region of expression to directly modulate the activity of bHLH proteins and thus confer specific characteristics on a cell-type identity. Although synergistic interactions of bHLH and homeodomain proteins have been described at some promoters (Westerman et al., 2003), the model in which homeodomain proteins can instruct the bHLH proteins to modulate a cell-type program to generate, for example, a particular type of neuron or muscle cell has not yet been demonstrated.

Previous demonstrations of molecular interactions between the bHLH Myod protein and the homeodomain proteins Pbx and Meis suggest that skeletal myogenesis might be a good system to directly test the role of homeodomain proteins in modulating bHLH-driven cell-type programs (Knoepfler et al., 1999; Berkes et al., 2004). Furthermore, myogenesis is a model system in which to study the acquisition of cellular phenotype diversity; for example, the generation of different fiber types. Skeletal myogenesis in vertebrate embryos is coordinated by the bHLH transcription factors Myod, Myf5, Myog and Mrf4 (Buckingham, 2001). Myod is sufficient to convert fibroblasts and other non-muscle cells into skeletal muscle (Weintraub et al., 1989). Myod directly activates the expression of multiple additional transcription factors, including Myog, and acts in a feed-forward mechanism in cooperation with those factors to directly activate muscle genes expressed later in the differentiation program (Penn et al., 2004; Cao et al., 2006). At a portion of these later target genes, Myod appears necessary to initiate chromatin remodeling before other transcription factors, such as Myog, can bind to the promoters (Cao et al., 2006). Therefore, Myod has a crucial function of identifying and remodeling the target genes that will subsequently be available for activation by other factors. Because Myod directly regulates a broad suite of muscle differentiation genes, one question that arises is how does Myod correctly regulate promoters used in different skeletal muscle types, such as fast- or slow-twitch muscle?

Mutations of the histidine- and cysteine-rich domain or the C-terminal helix III domain of Myod prevent full activation of a subset of Myod target genes and also prevent cooperative DNA binding with Pbx and Meis proteins on specific Myod target promoters, including that of *Myog* (Berkes et al., 2004). Because Pbx and Meis proteins are present on some of these promoters prior to Myod expression, we suggested that a Pbx-Meis complex might mark a subset of genes for Myod activation (Berkes et al., 2004; de la Serna et al., 2005). However, mutations in the Myod protein might alter interactions with additional factors, and a requirement for Pbx-Meis in skeletal myogenesis has not yet been demonstrated. Thus, it

¹Division of Human Biology, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA. ²Department of Biological Sciences, University of Alberta, Edmonton, Alberta, T6G2E9, Canada. ³Howard Hughes Medical Institute and Division of Basic Science, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA.

*Author for correspondence (e-mail: lmaves@fhcrc.org)

remains to be determined whether a Pbx-Meis complex is necessary for Myod activity at *Myog* or other specific genes in vivo and whether this subset of genes regulates a specific aspect of muscle development.

Zebrafish embryos provide an attractive biological system in which to test the role of Pbx proteins in skeletal myogenesis. Pbx (and Meis) proteins are TALE (three amino acid loop extension)-class homeodomain proteins that are best characterized as cofactors for Hox proteins. In mice, overlapping expression and functional redundancy among the Pbx genes complicates the analysis of their requirements, and skeletal muscle defects have not been reported for knock-outs of individual Pbx genes (Moens and Selleri, 2006). Zebrafish have five Pbx genes, but only *pbx2* and *pbx4* are expressed during the period of early myogenesis (Pöpperl et al., 2000; Waskiewicz et al., 2002). Knock-down of both *pbx2* and *pbx4* in zebrafish results in severe hindbrain-patterning defects that reflect the role of Pbx proteins as Hox cofactors (Waskiewicz et al., 2002), but skeletal muscle development was not assessed. The Pbx-Meis binding site in the mouse *Myog* promoter is conserved in the zebrafish *myog* gene (Berkes et al., 2004), providing an opportunity to test the developmental role of Pbx in myogenesis. Here, we demonstrate that Pbx proteins are necessary for the timely activation of *myog* by Myod, demonstrating that Pbx has a necessary role in marking the *myog* gene for efficient activation during development. Furthermore, we demonstrate that Pbx proteins facilitate Myod activity to drive the expression of a subset of genes necessary for fast-muscle differentiation, thus showing that homeodomain proteins can direct bHLH proteins to establish a specific cell-type identity.

MATERIALS AND METHODS

Zebrafish stocks

Zebrafish (*Danio rerio*) were raised and staged as previously described (Westerfield, 2000). Time (hpf) refers to hours post-fertilization at 28.5°C. In some cases, embryos were raised for periods at room temperature, at approximately 25°C. The wild-type stock used was an AB/WIK hybrid. For microarray analyses, the AB stock was used. The *pbx4* mutant line has been described (Pöpperl et al., 2000).

Morpholino and mRNA injections

Morpholino injections were performed, and working concentrations were determined, as previously described (Maves et al., 2002). Morpholinos (shown 5' to 3') were used at the following working concentrations: *pbx2*-MO2 (Waskiewicz et al., 2002), CCGTTGCCTGTGATGGGCTGCTGCG, 0.25 mg/ml; *pbx2*-MO3, GCTGCAACATCCTGAGCACTACATT, 0.5 mg/ml; *pbx2*-MO3 five-base mismatch control (mismatched bases shown in lowercase), GCTcCAAgATCCTcAcCAGTACATT, 0.5 mg/ml; *pbx4*-MO1, AATACTTTTGAGCCGAATCTCTCCG, 0.5 mg/ml; *pbx4*-MO2, CGCCGCAAACCAATGAAAGCGTGTT, 0.5 mg/ml; *myod*-splMO E111, AATAAGTTTCTCACAATGCCATCAG, 5 mg/ml; *myod*-splMO E212, TTTCGAGCAAACCTACCATTGGTG, 2.5 mg/ml; *myod*-MO1, GCAAGAAATGTACTTGAATGTTTCC, 0.5 mg/ml; *myod*-MO2, GGAATAGTAAGACAAAGTCCTCAG, 5 mg/ml; *myf5*-MO1, GATTGGTTTGGTGTGAAGGTTTCT, 0.25 mg/ml; *myf5*-MO2, GATCTGGGATGTGGAGAATACGTCC, 0.25 mg/ml. *pbx2*-MO2, *pbx2*-MO3, *pbx4*-MO1 and *pbx4*-MO2 were combined, *myod*-MO1 and *myod*-MO2 were combined, and *myf5*-MO1 and *myf5*-MO2 were combined in order to knock down their respective gene products. Embryos that are knocked-down for *pbx2* and *pbx4* function show a developmental delay of approximately two somites during somitogenesis stages, comparable to maternal-zygotic *pbx4*^{-/-}; *pbx2*-MO embryos (see Fig. S2 in the supplementary material; C.B.M., unpublished observations). We somite-stage-matched sibling control and MO-treated embryos when collecting embryos for staining or for RNA or protein analyses.

Injections of *exd* or *shh* (*shha* – ZFIN) mRNA were performed as previously described (Pöpperl et al., 2000; Barresi et al., 2000).

RNA in situ hybridization and immunocytochemistry

RNA in situ hybridizations were performed as previously described (Maves et al., 2002). The following cDNA probes were used: *myod* (Weinberg et al., 1996); *krox-20* [*egr2b* – Zebrafish Information Network (Oxtoby and Jowett, 1993)]; *myog* (Weinberg et al., 1996); *myhc4* (EST fb27a08); *aldh1a2* (Begemann et al., 2001); *mylz2* (Xu et al., 2000); *chrna1* (Sepich et al., 1998); *tmem161a* (IMAGE:7149790); *ttnl* (EST eu247); *atp2a1* (EST fc22f07); *srl* (EST fb94b10); *vmhc* (Yelon et al., 1999); *smyhc1* (Bryson-Richardson et al., 2005); and *cxcr4a* (EST cb824, Zebrafish International Resource Center). We subcloned the *desmin* EST cb290 (Zebrafish International Resource Center) into pCRII-TOPO, which was linearized with *Bam*HI and transcribed with T7 to make the antisense *desmin* probe.

Whole-embryo immunostaining was performed with the following primary antibodies: anti-pan zebrafish Pbx, 1:500 (Pöpperl et al., 2000); anti-Myf5, 1:100 [Santa Cruz, C-20, sc-302 (Hammond et al., 2007)]; anti-myosin heavy chain F59, 1:10 [supernatant (Devoto et al., 1996)]; anti-myosin heavy chain MF20, 1:10 [supernatant (Bader et al., 1982)]; and anti-fast myosin light chain F310, 1:10 [supernatant (Hamade et al., 2006)]. F59, F310 and MF20 antibodies were developed by F. E. Stockdale and D. A. Fischman and were obtained from the Developmental Studies Hybridoma Bank, maintained by the University of Iowa. Stainings were performed as previously described (Feng et al., 2006) with the following modifications: for anti-Pbx staining, embryos were fixed in 4% PFA at 4°C for 4 hours and methanol dehydration was omitted; for anti-Myf5 (Myod) staining, embryos were fixed in 4% PFA for 1 hour at room temperature, methanol dehydration was omitted, and washes and incubations were done in PBS+1% Triton-X; for F59, F310 and MF20 staining, secondaries (Southern Biotech) used were goat anti-mouse IgG1-FITC (1:100, F59 and F310) and goat anti-mouse IgG2b-TRITC (1:100, MF20); for SYTOX Green staining, embryos were rinsed in PBS following antibody staining, and then were incubated in a 1:10,000 dilution of SYTOX Green (Invitrogen) overnight at 4°C.

Embryos were photographed using a Zeiss Axioplan2 microscope, a SPOT RT digital camera (Diagnostic Instruments) and MetaMorph software or imaged using a Zeiss Pascal confocal microscope. Images were assembled using Adobe Photoshop.

Western analysis

Western analysis was performed as previously described (Waskiewicz et al., 2001). Samples were run on NuPage 4-12% BIS-TRIS gels (Invitrogen). The equivalent of approximately one embryo was loaded per lane. Antibodies were used at the following dilutions: anti-pan zebrafish Pbx (Pöpperl et al., 2000), 1:2000; and anti-Histone H4 (Upstate Biotechnology), 1:1000.

Quantitative real-time RT-PCR

Dechorionated embryos were homogenized in TRIzol using a 1 cc insulin syringe, and RNA was isolated following the TRIzol protocol (Invitrogen). 1 µg of RNA plus random hexamers were used in a reverse-transcriptase (RT) reaction with SuperScriptII Reverse Transcriptase (Invitrogen). Real-time PCRs were performed using an Applied Biosystems 7900HT System according to the manufacturer's instructions. The relative expression levels were normalized to those of *ornithine decarboxylase 1* (*odc1*) (Draper et al., 2001). We used the following probe and primer sets (sequences given are 5' to 3'): *myog*L1 forward, CTGGGGTGTCTCCTCTAGT; *myog*R1 reverse, TCGTCTGTTACAGAGATCCT; *myog*P1 probe, TGGAGCAGC-GCGTCTGATCA; *myod*L2 forward, TCAGACGAGAAGACGGAACA; *myod*R2 reverse, CACGATGCTGGACAGACAAT; *myod*P2 probe, CACCAAATGCTGACGCACGG; *odc*L2 forward, GACTTTGACTTCGCC-TTCT; *odc*R2 reverse, GAGGTGCTTCTCAGGACATC; *odc*P1 probe; CGCGCATATCGTGGAGCAG; *desmin*L1 forward, GAGGCCGAG-GACTGGTATAA; *desmin*R1 reverse, GGTGTAGGACTGGAGCTGGT; *desmin*P1 probe, AGCCAAGCAGGAGACCATGCAA.

cDNA microarray analysis

pbx2-MO; *pbx4*-MO embryos as well as their control siblings (*pbx2*-MO3 mismatch control) were collected at the 10 somite (10s) or 18s stage from three independent sets of injections. A subset of embryos from each set of injections were used for in situ hybridization to confirm somite staging

(using *myod*) as well as the loss of *krox-20* expression and downregulation of *myog* expression in *pbx2*-MO; *pbx4*-MO embryos (see Fig. 1M,N for example). Total RNA was harvested using TRIzol followed by Qiagen RNeasy clean-up according to the Affymetrix recommendations. cRNA labeling and hybridization to Affymetrix Zebrafish Genome arrays were performed using the manufacturer's protocols. The perfect match (PM) probe intensities were corrected by robust multi-array average, normalized by quantile normalization and summarized by medianpolish using the Affymetrix package of Bioconductor. The gene expression profiles of control versus *pbx2*-MO; *pbx4*-MO were compared using the LIMMA package of Bioconductor. Array data are available at NCBI GEO (www.ncbi.nlm.nih.gov/geo/), accession GSE8428.

EMSA

Electrophoretic mobility shift assays (EMSAs) were performed as previously described (Berkes et al., 2004). Pbx4 and Meis3 pCS2 expression constructs were as previously described (Pöpperl et al., 2000; Vlachakis et al., 2000).

RESULTS

Pbx and Myod are required for activation of *myog* expression

To address the role of Pbx in myogenesis, we used antisense morpholino oligos (MOs) to knock down the maternal and zygotic expression of Pbx2 and Pbx4 (Waskiewicz et al., 2002). In control embryos, a pan-zebrafish-Pbx antibody (Pöpperl et al., 2000) identified ubiquitous nuclear Pbx expression in early paraxial mesoderm, and ubiquitous Pbx expression was maintained during somite development and early muscle differentiation (see Fig. S1A-C in the supplementary material), consistent with prior reports of broad *pbx* mRNA expression during early zebrafish development (Pöpperl et al., 2000; Vlachakis et al., 2000; Waskiewicz et al., 2002), and confirming that Pbx proteins are present at the right time and place to potentially regulate early muscle development. By contrast, Pbx protein expression was essentially absent in *pbx2*-MO; *pbx4*-MO embryos (see Fig. S1D in the supplementary material). Western analysis confirmed the loss of Pbx2 and Pbx4 expression (see Fig. S1E in the supplementary material), and the loss of hindbrain expression of *krox-20*, a Pbx-dependent gene (Waskiewicz et al., 2002), confirmed the abrogation of Pbx function in *pbx2*-MO; *pbx4*-MO embryos (see Fig. 1).

We first wanted to test whether Pbx is required for the activation of *myog* expression. During somitogenesis stages in control embryos, *myod* was expressed adjacent to the notochord in adaxial cells (presumptive slow-muscle cells) as well as in more-lateral cells in the posterior of each somite (presumptive fast-muscle cells). *myog* showed a similar pattern of expression, which was delayed relative to *myod* (Weinberg et al., 1996; Groves et al., 2005) (Fig. 1A,D,G,J). In comparison to control embryos, *myod* expression was slightly increased in *pbx2*-MO; *pbx4*-MO embryos, based on in situ hybridization (Fig. 1A,B,G-H,M,N), and this slight increase was confirmed by quantitative real-time reverse transcriptase (RT)-PCR (qRT-PCR; Fig. 1S). Despite the presence of increased levels of *myod* in *pbx2*-MO; *pbx4*-MO embryos, expression of *myog* was severely reduced at the 6-10 somite (s) stages (Fig. 1D,E). As somitogenesis proceeded, *pbx2*-MO; *pbx4*-MO embryos exhibited a delayed pattern of *myog* expression (Fig. 1J,K,M,N). The reduction and delay of *myog* expression were confirmed by qRT-PCR (Fig. 1T). By approximately 24 hours post fertilization (hpf), *myog* expression approached its normal expression pattern but remained at somewhat reduced levels (Fig. 1T and data not shown). These results show that Pbx function is required for the efficient initiation of *myog* expression.

To demonstrate the specificity of the *pbx* MOs, we injected mRNA for the *Drosophila* *pbx* gene ortholog, *extradenticle* (*exd*), into *pbx2*-MO; *pbx4*-MO embryos. *exd* mRNA previously was shown to rescue *krox-20* expression in *pbx4*^{-/-} embryos (Pöpperl et al., 2000). We found that *exd* mRNA rescued *krox-20* expression and rescued the reduced and delayed *myog* expression in *pbx2*-MO; *pbx4*-MO embryos (see Fig. S2 in the supplementary material). Additionally, knock-down of either *pbx2* or *pbx4* had little or no effect on *myog* expression, and a five-base mismatch control MO for *pbx2* caused no defects in *myog* expression (data not shown). Therefore, the delay and reduction in *myog* expression in *pbx2*-MO; *pbx4*-MO embryos is specific and dependent upon the loss of function of both *pbx* genes.

We then addressed whether the requirements for Myod are similar to those for Pbx in the activation of *myog* expression. We used two types of MOs to knock down Myod: translation-blocking MOs (*myod*-MO) to knock down all Myod activity, and splice-blocking MOs (*myod*-splMO) to knock down the helix III-containing exon 3 of zebrafish *myod*, because the helix III domain of Myod is necessary for cooperative binding with Pbx in vitro (Berkes et al., 2004). *myod*-splMOs induced aberrantly spliced cDNAs that are predicted to generate truncated proteins lacking the helix III domain but retaining the activation and bHLH domains of Myod, whereas the *myod*-MOs caused the loss of Myod protein (see Fig. S3 in the supplementary material). If the delay in *myog* expression in *pbx2*-MO; *pbx4*-MO embryos was due to requirements for an interaction between Pbx and Myod, then we expect that knock-down of Myod would also result in a delay in *myog* expression. Similar to *pbx2*-MO; *pbx4*-MO embryos, *myog* expression was reduced and delayed with knock-down of Myod helix III (Fig. 1F,L,T). Knock-down of Myod protein caused a similar delay in *myog* activation and, at later stages, *myod*-MO embryos maintained a slightly more severe reduction of *myog* expression compared with *pbx2*-MO; *pbx4*-MO embryos (Fig. 1O; see also Fig. 4). These results show that Myod is indeed required for the proper expression of *myog* in zebrafish and, in particular, that it has similar requirements as Pbx for the timely initiation of *myog* expression. These results thus support our hypothesis that interaction with Pbx is necessary for Myod to properly initiate *myog* expression.

To test whether Pbx or Myod knock-down cause a delay in all muscle gene expression, we analyzed the expression of *desmin* (*desm*), one of the earliest-expressed muscle-specific genes in zebrafish (Xu et al., 2000) and a known direct transcriptional target of Myod that does not depend on Myod helix III in cultured mouse cells (Bergstrom et al., 2002; Berkes et al., 2004). In control embryos, *desm* is expressed subsequent to *myod*, mainly in adaxial cells (Xu et al., 2000) (Fig. 1P). *desm* expression appeared largely unaffected in *pbx2*-MO; *pbx4*-MO, *myod*-splMO or *myod*-MO embryos, either by RNA in situ hybridizations (Fig. 1P-R; see also Figs 2 and 4) or by qRT-PCR (Fig. 1U). We demonstrate below (see Fig. 4) that zebrafish *desm* is indeed Myod-regulated. These results show that Pbx is not required for the proper initiation of all muscle gene expression.

Microarray analysis identifies Pbx-dependent somite and muscle gene expression

To identify the subset of muscle genes that require Pbx for normal developmental expression, we used Affymetrix expression microarrays to compare gene expression in embryos lacking Pbx function (*pbx2*-MO; *pbx4*-MO embryos) to control MO-treated embryos at the 10s and 18s stages. We chose the 10s stage because that was when *myog* expression showed a strong reduction in *pbx2*-

MO; *pbx4*-MO embryos (Fig. 1) and when early muscle-specific genes are beginning to be expressed (Xu et al., 2000). We chose the 18s stage because many muscle differentiation genes are not expressed until mid- or late-stage somitogenesis in zebrafish (Xu et al., 2000). Using the cut-offs of >1.5-fold change and a false discovery rate of <0.05, our array analysis identified 188 Pbx-

dependent genes at 10s and 258 genes at 18s, including all of the previously known Pbx-dependent genes expressed in the hindbrain, such as *krox-20* (Waskiewicz et al., 2002) (see Tables S1 and S2 in the supplementary material). From these groups, we selected the genes that are known to be expressed in wild-type somites and early skeletal muscle (Thisse et al., 2001; ZFIN, 2006) (Table 1). *myog* is




Fig. 1. Pbx and Myod are required for the proper initiation of *myog* expression. (A-R) RNA in situ expression of (A-C,G-I) *myod* and *krox-20*, (D-F,J-L) *myog* and *krox-20*, (M-O) *myog* (blue) and *myod* (red), or (P-R) *desm* and *krox-20* in (A,D,G,J,M,P) wild-type control, (B,E,H,K,N,Q) *pbx2*-MO; *pbx4*-MO, (C,F,I,L,R) *myod*-splMO, or (O) *myod*-MO embryos. Similar phenotypes were observed for both *myod* splice-blocking MOs (*myod*-splMOs). Somite (s) staging was confirmed by counting somites using Nomarski optics. All embryos are shown in dorsal view, anterior towards the left. (A) Arrowheads point to adaxial cells; arrows point to lateral somite cells; and r3 and r5 indicate *krox-20* expression in hindbrain rhombomeres. (S-U) Graphs of real-time reverse-transcriptase (RT)-PCR quantities for (S) *myod*, (T) *myog* and (U) *desm*. (T, bottom) Enlarged view of 6s stage *myog* expression and the key for S-U. x-axes are developmental stage and y-axes are relative mRNA expression level. Quantities were normalized to *odc1* expression. Error bars represent standard deviation. *myodE212*-splMO targets the exon 2-intron 2 boundary (see Fig. S3 in the supplementary material). The *myod* real-time primers measure only mRNA with properly spliced intron 2. Quantitative (q)RT-PCR shows that less than about 10-15% of normally spliced *myod* transcripts remain in *myod*-splMO embryos (Fig. 1S and see Fig. S3 in the supplementary material; and data not shown). Similar quantities for *myog* and *desmin* were observed for both *myod*-splMOs.

Table 1. Microarray-identified Pbx-dependent genes expressed in somites and early muscle

Gene symbol	Affymetrix ID number	Fold change at 10s (controlMO: <i>pbx2</i> -MO; <i>pbx4</i> -MO)	Expression pattern at 10s	RNA in situ expression in <i>pbx2</i> -MO; <i>pbx4</i> -MO at 10s	RNA in situ expression in <i>myod</i> -MO at 10s
<i>si:dkey-119m7.4</i>	Dr26487.1.S1_at	3.85	Segmental plate	ND	ND
<i>angptl7</i>	Dr10141.1.A1_at	2.62	Broad through lateral somites	Very reduced	Upregulated
<i>hoxa3a</i>	Dr24774.1.S2_at	2.25	Anterior cells of anterior somites; central nervous system (CNS)	All domains reduced	Not affected
<i>myog</i>	Dr2499.1.S1_at	2.04	Adaxial cells and lateral somites	Very reduced	Very reduced
<i>egfl6</i>	Dr1807.1.A1_at	1.95	Segmental plate; CNS	All domains reduced	Not affected
<i>bhik</i>	DrAffx.1.2.S1_at	1.92	Somites; lateral plate mesoderm; tail bud; polster	ND	ND
<i>fabp3</i>	Dr6814.1.S1_at	1.74	Posterior cells in anterior somites; CNS; eye	All domains reduced	Somite expression reduced
<i>pwp2h</i>	Dr3590.1.S1_at	1.61	Segmental plate; CNS	All domains reduced	Not affected
<i>ppp1r3b</i>	Dr4453.1.S1_at	1.59	Ubiquitous	ND	ND
<i>acta1</i>	AFFX-Dr-acta1-5_at	1.59	Basal ubiquitous expression; stronger in adaxial cells	Reduced	Reduced
<i>meis3</i>	Dr1680.1.S1_at	1.56	Anterior somites; head mesenchyme; CNS	All domains reduced	Not affected
<i>ttn</i>	Dr4681.1.A1_at	1.53	Adaxial cells; lateral somites; heart primordium	Heart domain reduced	Not affected
<i>chrna1</i>	Dr123.1.S1_at	1.52	Adaxial cells	Reduced	Reduced
<i>gstm</i>	Dr1754.1.S1_at	1.52	Diffuse in somites, CNS and eye	All domains reduced	Somite expression reduced
<i>ttnl</i>	Dr1662.1.S1_at	1.52	Adaxial cells; lateral somites; segmental plate	Reduced	Very reduced
Gene symbol	Affymetrix ID number	Fold change at 18s (controlMO: <i>pbx2</i> -MO; <i>pbx4</i> -MO)	Expression pattern at 18s	RNA in situ expression in <i>pbx2</i> -MO; <i>pbx4</i> -MO at 18s	RNA in situ expression in <i>myod</i> -MO at 18s
<i>vmhc</i>	Dr10607.1.A1_at	3.20	Adaxial cells; heart	Heart domain reduced	Not affected
<i>myhz1</i>	Dr23067.1.S1_at	2.99	Adaxial cells and lateral somites	ND	ND
<i>myhc4</i>	Dr4812.1.S1_s_at	2.51	Adaxial and lateral somites	Reduced	Very reduced
<i>eng2a</i>	Dr496.1.S1_at	2.30	Muscle pioneers; CNS	CNS expression reduced	Not affected
<i>aldh1a2</i>	Dr5206.1.S1_at	2.26	Somites; lateral plate mesoderm; eye	All domains reduced, but increased expression in tail bud	Not affected
<i>mylz2</i>	Dr2914.1.S1_at	2.08	Adaxial cells and lateral somites	Very reduced	Very reduced
<i>chrna1</i>	Dr123.1.S1_at	2.06	Adaxial cells and lateral somites	Very reduced	Very reduced
<i>ppp1r3b</i>	Dr4453.1.S1_at	1.97	Ubiquitous	ND	ND
<i>angptl7</i>	Dr10141.1.A1_at	1.96	Lateral somites	Reduced	Upregulated
<i>ttn</i>	Dr4681.1.A1_at	1.88	Adaxial cells; lateral somites; heart	Reduced	Reduced
<i>bhik</i>	DrAffx.1.2.S1_at	1.84	Anterior somites; tail bud	ND	ND
<i>tmem161a</i>	Dr5047.1.A1_at	1.80	Adaxial cells and lateral somites	Reduced	Very reduced
<i>ttnl</i>	Dr1662.1.S1_at	1.78	Adaxial cells and lateral somites	Reduced	Very reduced
<i>dhrs3</i>	Dr2644.1.A1_at	1.76	Anterior cells of somites; ventral mesoderm; spinal cord	All domains reduced	Somite expression expanded
<i>atp2a1</i>	Dr20010.7.S1_at	1.76	Adaxial cells and lateral somites	Reduced	Very reduced
<i>acta1</i>	AFFX-Dr-acta1-5_at	1.75	Basal ubiquitous expression, stronger in adaxial cells and somites	Reduced	Reduced
<i>srl</i>	Dr21216.1.A1_at	1.72	Adaxial cells and lateral somites	Reduced	Very reduced
<i>si:dkey-119m7.4</i>	Dr26487.1.S1_at	1.66	Posterior segmental plate	ND	ND
<i>myog</i>	Dr2499.1.S1_at	1.63	Adaxial cells and lateral somites	Reduced	Reduced
<i>fabp3</i>	Dr6814.1.S1_at	1.54	Posterior-ventral somite expression; CNS; eye	Upregulated in somites; reduced in CNS	Upregulated in somites
<i>atp1a2a</i>	Dr10438.1.S1_at	1.52	Lateral somites	Reduced	Reduced
<i>cmyc</i>	Dr1.1.S1_at	1.51	Lateral somites; CNS; eye	CNS and eye reduced	Not affected

10s/18s, 10/18 somites; ND, not determined.

one of these somite-expressed genes and provides an initial validation of our array analysis (Table 1). Because some of these genes were expressed in domains in addition to the somites, we used RNA in situ hybridization to confirm that most of these genes indeed show reduced somite expression in *pbx2*-MO; *pbx4*-MO embryos (Table 1 and Fig. 2A-P). However, there were some exceptions. For example, *vmhc* was expressed both in adaxial cells and the heart primordium, but only the heart expression was reduced in *pbx2*-MO; *pbx4*-MO embryos (Fig. 2Q,R). Furthermore, *desm* was not significantly changed on the arrays and showed normal expression in *pbx2*-MO; *pbx4*-MO embryos at 18s (Fig. 2S,T). These results indicate that Pbx is indeed required for a subset of skeletal muscle gene expression.

We tested whether these microarray-identified Pbx-dependent genes were also Myod-dependent and found that, of the 20 genes whose somite/skeletal muscle expression was Pbx-dependent, 13 also showed reduced somite/muscle expression in *myod*-MO embryos (Table 1). At the 18s stage, these genes generally exhibited a delayed pattern of expression in *pbx2*-MO; *pbx4*-MO embryos, which was slightly more reduced in *myod*-MO embryos (Table 1), similar to that observed for *myog*. We noted that many of the Pbx- and Myod-dependent genes identified in our 18s arrays are predicted to play roles in fast-muscle differentiation, in particular *myhz1*, *myhc4*, *mylz2*, *atp2a1*, *srl* and *atp1a2a* (Feng et al., 2006; Groves et al., 2005; Bárány, 1967; Reiser et al., 1985; Ohlendieck et al., 1999).

Previous work showed that knock-down of Myod in zebrafish causes reduced *mylz2* expression and, therefore, a defect in fast-muscle development (Hammond et al., 2007). We examined *mylz2* expression at early stages and found that, like other ‘fast-muscle’ genes in zebrafish (Bryson-Richardson et al., 2005; Xu et al., 2000) (data not shown), *mylz2* was initially expressed mainly in adaxial cells, but also in lateral somite cells, and that both of these domains were reduced and delayed in *pbx2*-MO; *pbx4*-MO and *myod*-MO embryos (Fig. 2U-W). Similar results were observed for *myhc4*, *atp2a1*, *ttnl* and *srl* (data not shown). Thus, we have identified a set of genes that are both Pbx- and Myod-dependent. Based on the predicted functions of these genes and their expression defects, our data suggest that Pbx might facilitate Myod-activation of a set of genes that function in fast-muscle differentiation.

Pbx function is required for efficient fast-muscle differentiation in zebrafish embryos

To further address the requirements for Pbx and Myod in the development of fast and slow muscle in zebrafish, we examined additional markers that label fast- or slow-muscle precursors. *smyhc1*, which encodes a slow myosin heavy chain and is expressed specifically in adaxial cells/slow-muscle precursors (Bryson-Richardson et al., 2005), was expressed normally in *pbx2*-MO; *pbx4*-MO and in *myod*-MO embryos (Fig. 2X-Z). Engrailed staining labels muscle pioneers – an early-differentiating subset of the slow-




Fig. 2. Validation of microarray-identified Pbx-dependent genes. (A-Z) RNA in situ

expression of genes from Table 1 in (A,C,E,G,I,K,M,O,Q,S,U,X) wild-type control, (B,D,F,H,J,L,N,P,R,T,V,Y) *pbx2*-MO; *pbx4*-MO, or (W,Z) *myod*-MO embryos. (S,T) Control gene, *desm*; (X-Z) slow-muscle gene, *smyhc1*. Although *pbx2*-MO; *pbx4*-MO embryos appear delayed relative to controls, all embryos are stage-matched at (A-T) 18-somite stage (18s), (U-W) 16s or (X-Z) 14s. *krox-20* was included in all in situs to control for the *pbx* MOs. (A) r3 and r5 indicate *krox-20* expression in hindbrain rhombomeres. (Q) Arrow marks *vmhc* expression in the heart primordium. Embryos are shown in (A-P,S-T) left-side view or (Q-R,U-Z) dorsal view, anterior towards the left.

muscle cell population (Devoto et al., 1996; Hatta et al., 1991) – which appeared to develop normally in *pbx2*-MO; *pbx4*-MO embryos (data not shown). The F59 anti-myosin heavy chain antibody, which specifically labels slow-muscle precursors during somitogenesis stages (Devoto et al., 1996), was expressed normally in *pbx2*-MO; *pbx4*-MO, in *myod*-splMO and in *myod*-MO embryos (Fig. 3A,B,E). The F310 anti-myosin light chain antibody labels fast-muscle myosins in zebrafish (Hamada et al., 2006). In contrast to F59 staining, F310 staining was delayed and reduced in *pbx2*-MO; *pbx4*-MO, in *myod*-splMO and in *myod*-MO embryos (Fig. 3C,D,F and see Fig. S4 in the supplementary material). *myod*-MO embryos showed slightly more reduced fast-muscle development than *pbx2*-MO; *pbx4*-MO or *myod*-splMO embryos (Fig. 3F and see Fig. S4 in the supplementary material). Expression of a pan-muscle myosin marker MF20 (Bader et al., 1982) initiated on time, and its slow-muscle expression appeared normal in *pbx2*-MO; *pbx4*-MO, in *myod*-splMO and in *myod*-MO embryos (Fig. 3A-F and see Fig. S4 in the supplementary material). Thus, upon loss of Pbx or Myod function, we observed delayed and reduced fast-muscle differentiation, using multiple markers, and we observed no change, including no delay, reduction, or increase, in slow-muscle-differentiation marker expression. These results, combined with our microarray findings, show that Pbx function is needed to promote efficient fast-muscle differentiation, as is Myod.

Multiple populations of fast-muscle fibers have been described in zebrafish. Primary fast muscle comprises medial fast fibers, which express low levels of Engrailed in response to Hedgehog signaling (Wolff et al., 2003), and lateral fast fibers, which are dependent on Fgf signaling (Groves et al., 2005). A secondary population of fast-muscle precursors, initially marked by *pax3* and *pax7* expression, contributes fast-muscle fibers to the growing myotomes, beginning at around 24 hpf (Hollway et al., 2007; Stellabotte et al., 2007).

Because Engrailed expression appeared normal in *pbx2*-MO; *pbx4*-MO embryos (data not shown) and because fast-muscle differentiation in *pbx2*-MO; *pbx4*-MO embryos appeared to recover by about 24 hpf (Fig. 3), we conclude that Pbx is necessary for the efficient differentiation of primary lateral fast fibers.

We next looked for evidence that lateral somite cells are maintaining a prolonged undifferentiated state in the absence of Pbx. Decreased fast-muscle differentiation and loss of Myod function are associated with increased and prolonged *pax3* and *pax7* expression, markers of undifferentiated myogenic precursors (Groves et al., 2005; Hammond et al., 2007). We could not identify a significant upregulation of *pax3* or *pax7* in *pbx2*-MO; *pbx4*-MO embryos (data not shown). However, our microarray analysis identified the upregulation of *cxcr4a* in *pbx2*-MO; *pbx4*-MO embryos (see Table S2 in the supplementary material). *cxcr4a*, the expression of which in the anterior-lateral cells of each somite is initially very similar to *pax7*, is normally downregulated in all but the most posterior somites at the 18s stage (Hollway et al., 2007) (see Fig. S5 in the supplementary material). We observed prolonged maintenance of *cxcr4a* in trunk somites in *pbx2*-MO; *pbx4*-MO embryos as well as in *myod*-MO embryos (see Fig. S5 in the supplementary material). These results suggest that Pbx, as well as Myod, are important for the efficient differentiation of lateral somite cells.

Pbx functions with Myod to promote fast-muscle, but not slow-muscle, differentiation

Previous work has shown that *myod* functions redundantly with *myf5* in slow-muscle development, because loss of *myod* together with *myf5* causes loss of slow-muscle myosin heavy chain expression (Hammond et al., 2007). Redundancy with Myf5 might thus mask requirements for Pbx acting with Myod. In particular, if Pbx is necessary for Myod to activate slow-muscle gene expression, then




Fig. 3. Pbx is required for the proper initiation of fast-muscle differentiation. (A-D) Antibody staining of slow muscle with F59 antibody (green; A,B), or fast muscle with F310 antibody (green; C,D), in (A,C) wild-type control or (B,D) *pbx2*-MO; *pbx4*-MO embryos. Antibody staining of muscle with MF20 antibody (red; A-D) is included to control for somite staging. Embryos are shown in left-side view, anterior towards the left. (E,F) Quantification of (E) F59/slow- and (F) F310/fast-muscle marker expression. MF20 expression is included to control for somite (s) staging. Graphs show number of somites expressing a muscle marker at different developmental stages. Similar phenotypes were observed for both *myod* splice-blocking MOs (*myod*-splMOs). Error bars represent standard deviation. *P* values (Mann-Whitney test; normalized to somite number): **P*<0.0001 compared to same-stage control; except for *pbx2*-MO; *pbx4*-MO F310 compared to control at 24 hpf, *P*<0.03.




Fig. 4. Pbx functions with Myod in fast-muscle, but not slow-muscle, differentiation. RNA in situ expression at the 10-somite (10s) stage of (A-G) *myog* and *krox-20*, (H-N) *desm* and *krox-20* or (O-U) *smyhc1* and *krox-20*, or at 16s of (V-BB) *mylz2* and *krox-20* in (A,H,O,V) wild-type control, (B,I,P,W) *pbx2*-MO; *pbx4*-MO, (C,J,Q,X) *myod*-MO, (D,K,R,Y) *myf5*-MO, (E,L,S,Z) *myf5*-MO; *myod*-MO, (F,M,T,AA) *myf5*-MO; *pbx2*-MO; *pbx4*-MO, or (G,N,U,BB) *myod*-MO; *pbx2*-MO; *pbx4*-MO embryos. Embryos are shown in dorsal view, anterior towards the left. Somite staging was confirmed by counting somites using Nomarski optics. (A) r3 and r5 indicate *krox-20* expression in hindbrain rhombomeres.

knock-down of *Myf5* together with *Pbx* should cause slow-muscle defects. Alternatively, because the helix III *Pbx*-interacting domain of *Myod* is conserved in *Myf5* (Bergstrom and Tapscott, 2001), knock-down of *Myod* together with *Pbx* might cause slow-muscle defects. We found that *myf5*-MO embryos showed no defects in *myog* or *desm* expression (Fig. 4D,K), whereas *myf5*-MO; *myod*-MO embryos showed complete loss of *myog* and *desm* expression (Fig. 4E,L), confirming that our *myf5* and *myod* MOs were functioning. *myf5*-MO; *pbx2*-MO; *pbx4*-MO embryos resembled *pbx2*-MO; *pbx4*-MO embryos, in that *myog* expression was delayed in lateral somite cells and *desm* expression was unaffected (Fig. 4F,M). *myod*-MO; *pbx2*-MO; *pbx4*-MO embryos, however, showed almost complete loss of *myog* expression at 10s (Fig. 4G), whereas *desm* expression was unaffected (Fig. 4N). By 18s, in *myod*-MO; *pbx2*-MO; *pbx4*-MO embryos, *myog* expression was still much reduced compared with *myod*-MO embryos (data not shown). These results suggest that *Pbx* proteins are needed for both *Myod* and *Myf5* to activate *myog* expression.

We then assessed slow- and fast-muscle differentiation in these MO combinations using *smyhc1*, F59, *mylz2* and F310. As previously reported, all slow-muscle differentiation was lost in *myf5*-MO; *myod*-MO embryos (Hammond et al., 2007) (see Fig. 4S). All

fast-muscle differentiation was also lost in *myf5*-MO; *myod*-MO embryos (Fig. 4Z and data not shown), revealing that *myf5* acts with *myod* in fast muscle. *myf5*-MO; *pbx2*-MO; *pbx4*-MO embryos resembled *pbx2*-MO; *pbx4*-MO embryos (Fig. 4T,AA and data not shown). Importantly, slow-muscle differentiation was not delayed in *myf5*-MO; *pbx2*-MO; *pbx4*-MO embryos (Fig. 4T and data not shown). These results show that *Pbx* is not necessary for *Myod* to activate slow-muscle gene expression. We also observed that slow-muscle differentiation was not delayed in *myod*-MO; *pbx2*-MO; *pbx4*-MO embryos (Fig. 4U and data not shown). However, similar to *myog* expression, *myod*-MO; *pbx2*-MO; *pbx4*-MO embryos showed severely reduced activation of *mylz2* (Fig. 4BB) and very reduced F310 staining even at 24 hpf (data not shown). Taken together, these results show that *Pbx* homeodomain proteins are necessary to modulate the activity of *Myod* and, probably, *Myf5*, in driving fast-muscle-specific differentiation.

Pbx is required downstream of Shh signaling for *Myod* to induce fast-muscle, but not slow-muscle, gene expression


To further demonstrate that *Pbx* directs *Myod* to drive fast-muscle, but not slow-muscle, development, we tested whether *Pbx* and *Myod* are required for the ability of Sonic hedgehog (*Shh*) signaling to

Fig. 5. Pbx is required downstream of Shh signaling to induce fast-muscle, but not slow-muscle, gene expression.

(A-L) RNA in situ expression of (A-F) *smyh1* (blue) and *myod* (red) at the 14-somite (14s) stage or (G-L) *myl2* (blue) and *myod* (red) at 16s in embryos injected with (A,G) *pbx2*-MO mismatch control and *gfp* mRNA, (B,H) *pbx2*-MO mismatch control and *shh* mRNA, (C,I) *myod*-MO and *gfp* mRNA, (D,J) *myod*-MO and *shh* mRNA, (E,K) *pbx2*-MO; *pbx4*-MO and *gfp* mRNA, or (F,L) *pbx2*-MO; *pbx4*-MO and *shh* mRNA. The most-anterior 8-10 somites of each embryo are shown, in dorsal view, anterior towards the left.

Control embryos injected with *shh* mRNA (control+*shh*) show increased expression of *smyh1* (B, 106/135 embryos; 29/135 show normal *smyh1* expression) or increased expression of *myl2* (H, 88/121; 33/121 show normal *myl2* expression). *myod*-MO+*shh* embryos show less-frequent and less-extensive induction of *smyh1* compared with control+*shh* (46/90 embryos resemble D; 44/90 show normal *smyh1* expression) and fail to upregulate *myl2* expression beyond that seen in *myod*-MO+*gfp* embryos (84/88 embryos resemble J; 4/88 show slightly increased *myl2* expression).

pbx2-MO; *pbx4*-MO+*shh* embryos show strongly increased expression of *smyh1* (77/79 embryos resemble F; 2/79 have normal *smyh1* expression). Approximately half (43/102; L) of *pbx2*-MO; *pbx4*-MO+*shh* embryos show expansion of weak levels of *myl2* across the somites; the other half (59/102) resemble *pbx2*-MO; *pbx4*-MO+*gfp* embryos. Similar results were observed in multiple experiments.



drive slow- or fast-muscle gene expression. Previous studies have shown that injecting *shh* mRNA into wild-type embryos induces the upregulation of *myod* and slow-muscle gene expression, such as *smyh1*, causing expanded slow-muscle differentiation across the lateral somites (Currie and Ingham, 1996; Blagden et al., 1997; Du et al., 1997) (Fig. 5B). We tested whether this slow-muscle expansion requires Myod and found that *shh*-induced upregulation of *smyh1* was partially inhibited in *myod*-MO embryos (Fig. 5D). However, *shh* mRNA readily induced increased *myod* and *smyh1* expression in *pbx2*-MO; *pbx4*-MO embryos (Fig. 5F), showing that Pbx is not needed for Shh/Myod-induced slow-muscle development. Whereas *shh* induces repression of fast-muscle markers at late stages (30 hpf) (Blagden et al., 1997; Ju et al., 2003) (data not shown), we found that, at earlier stages, *shh* can induce the expansion of fast-muscle markers, such as *myl2*, in control embryos (Fig. 5H and data not shown). Shh-induced upregulation of fast-muscle markers was inhibited in *myod*-MO embryos, even though *myod* expression was upregulated (Fig. 5J and data not shown). In contrast to *smyh1*, Shh/Myod induction of fast-muscle markers was inhibited in *pbx2*-MO; *pbx4*-MO embryos (Fig. 5L and data not shown). Thus, our experiments here show that the slow- or fast-muscle-inducing effects of Shh are at least in part mediated by Myod, but that Pbx is required only for the induction of fast-muscle markers. Taken together, these experiments further demonstrate that Pbx functions specifically in regulating Myod activation of fast-muscle gene expression.

Pbx cooperates with Myod to regulate fast-muscle gene expression independent of Myog or retinoic acid and might directly regulate *myl2*

We next wanted to address how directly Pbx and Myod function in the activation of fast-muscle-specific genes. There are several reasons why Myog could be the main direct target of Pbx-Myod. First, the *myog* promoter contains the best-characterized binding site for a Pbx-Myod complex (Berkes et al., 2004). Second, *myog* expression is affected strongly and early in *pbx2*-MO; *pbx4*-MO

embryos and in *myod*-MO embryos (Fig. 1). Also, work in mammalian cell culture has demonstrated that Myog is a key downstream effector of Myod (Cao et al., 2006). If Myog is the main effector of Pbx-Myod in zebrafish, then knocking down Myog function should cause similar defects as those seen in *pbx2*-MO; *pbx4*-MO embryos and *myod*-MO embryos. We confirmed that Myog protein is almost completely abolished in *myog*-MO embryos (data not shown). However, we observed no delays or defects in fast-muscle gene expression in *myog*-MO embryos (data not shown). We also analyzed additional Pbx-dependent genes and found that *acta1*, *ttn*, *chrna1* and *ttnl* expression all appeared to be normal in *myog*-MO embryos (data not shown). These results reveal that Pbx-Myod must have direct targets, in addition to *myog*, to mediate the defects in fast-muscle gene expression.

Our microarray analysis identified the expression of *aldh1a2*, which encodes an enzyme that is crucial for the synthesis of retinoic acid (RA) (Begemann et al., 2001; Grandel et al., 2002), as Pbx-dependent (Table 1; Fig. 2D). Because RA promotes fast-muscle development in zebrafish (Hamade et al., 2006), *aldh1a2*/RA signaling could be an effector of Pbx. Incubating wild-type embryos in an RA bath induces the upregulation of *myod* and *myog* expression, and precocious fast-muscle differentiation (Hamade et al., 2006). We found that, although RA is still able to strongly increase *myod* expression in *pbx2*-MO; *pbx4*-MO embryos, RA did not rescue *myog* expression nor the defect in fast-muscle development in *pbx2*-MO; *pbx4*-MO embryos (data not shown). Therefore, Pbx is required for the efficient activation of fast-muscle genes by Myod in a manner that is independent of its role in RA synthesis. Taken together, our findings are consistent with the hypothesis that Pbx not only recruits Myod to the *myog* promoter (Berkes et al., 2004) but also to additional fast-muscle gene promoters.

To determine whether Pbx directly regulates fast-muscle gene expression, we have initiated an analysis of fast-muscle gene promoters. We identified a near-consensus Pbx-Meis binding site in

the *myl2* promoter and determined that in vitro-translated Pbx-Meis heterodimers bind to this site in electrophoretic mobility shift assays (see Fig. S6 in the supplementary material). Although additional studies are necessary to test the biological relevance of this site, the presence of a Pbx-binding site in the *myl2* promoter suggests a direct role for Pbx in the regulation of fast-muscle gene expression.

DISCUSSION

Previous studies suggested that Pbx marks a subset of genes for activation by Myod but they did not demonstrate a necessary or biological role for this function (Berkes et al., 2004). We set out to address two specific hypotheses of Pbx function: that it is required for *myog* expression and that it is required for a biologically relevant subset of Myod-regulated gene expression. Our analysis of zebrafish embryos lacking Pbx function demonstrates that Pbx is indeed necessary in order for Myod to accurately regulate the expression of *myog* and that the subset of Myod-regulated genes that also require Pbx confer the fast-skeletal-muscle phenotype in zebrafish embryos. More generally, our results provide the initial support for an instructional role of homeodomain proteins in modulating the activity of bHLH proteins in driving a specific cellular phenotype. Previous work has shown that Myod, Neurod and other bHLH proteins drive a broad program of gene expression to establish a general cell type, such as muscle or neuron, whereas our study demonstrates that homeodomain proteins modulate this broad program to establish the regional identity of that cell type by marking certain subsets of genes for activation.

Our work demonstrates that Pbx is needed for the activity of a non-homeodomain protein. Pbx proteins are most-characterized for their role as Hox cofactors, but Pbx proteins also physically and functionally interact with other homeodomain proteins, including with Pdx1 (Peers et al., 1995; Kim et al., 2002) and Engrailed (Peltenburg and Murre, 1996; Kobayashi et al., 2003; Erickson et al., 2007). Our demonstration that Pbx proteins modulate the set of genes regulated by the bHLH protein Myod suggests that Pbx proteins could have more-widespread interactions than previously appreciated. Indeed, the increasing data on the requirements for Pbx in mice and zebrafish underscore the expanding roles for Pbx beyond that of Hox cofactors (Moens and Selleri, 2006; Erickson et al., 2007) (this work). Although previous studies in worms have shown that Pbx homologs are required for muscle development via their roles in mesoderm development (Liu and Fire, 2000), our study is the first to show that Pbx is required for vertebrate myogenesis.

Although we had hypothesized that Pbx would be required for a subset of Myod activity and therefore for the differentiation of a subset of muscle cell types, it is unexpected that Pbx is needed specifically for fast-muscle differentiation. Pbx proteins are expressed in presumptive slow- (adaxial) and fast- (lateral somite) muscle cells along with Myod. The Pbx- and Myod-dependent genes identified in our study are expressed in both of these cell populations. Adaxial cells thus express both Pbx-dependent and non-Pbx-dependent genes. Therefore, Pbx and Myod appear to be acting at the level of promoter activity and not in specific cell types. If Pbx proteins are ubiquitously expressed, how does Pbx help target Myod to specific promoters? Pbx and Myod directly bind the mouse *Myog* promoter (Berkes et al., 2004), and we found that Pbx can bind the zebrafish *myog* (A.T. and L.M., unpublished observations) and *myl2* promoters, suggesting a direct role for Pbx in recruiting Myod to fast-muscle gene promoters. An alternative, although not necessarily mutually exclusive, hypothesis is that additional Pbx-dependent intermediates might exist that modulate the activity of Myod on fast-muscle gene regulatory regions. In support of this

hypothesis, our microarray analysis identified at least seven genes whose expression was Pbx-dependent but not Myod-dependent (Table 1), although the functions of these genes in muscle development is unknown. Whether acting directly or indirectly, Pbx is necessary for the efficient expression of *myog* and fast-muscle genes, but not for the entire myogenic program. Therefore, our results provide the first evidence that homeodomain proteins directly modulate the activity of a bHLH protein in directing the phenotype of a general differentiation program.

What happens to cells that are delayed in differentiating into fast muscle in the absence of Pbx? We did not find any evidence for an increase in slow-muscle differentiation in the absence of Pbx, and we did not find a severe loss of cells, because *myod* was still expressed strongly in the somites. Thus, these cells might remain in a prolonged undifferentiated state. In the absence of Fgf8 signaling, lateral somite cells fail to properly activate *myod*, *myog* and fast-muscle gene expression but show increased expression of *pax3* and *pax7*, suggesting prolonged maintenance of an undifferentiated state (Groves et al., 2005; Hammond et al., 2007). Loss of Myod and Myf5 function also leads to increased *pax3* and *pax7* expression (Hammond et al., 2007). *pax3* and *pax7* expression does not normally decline in the somites until after *pbx2*-MO; *pbx4*-MO embryos have started to recover fast-muscle differentiation; thus, we have not been able to identify a significant upregulation of *pax3* or *pax7* in *pbx2*-MO; *pbx4*-MO embryos. We did, however, see prolonged maintenance of *cxcr4a* expression in *pbx2*-MO; *pbx4*-MO embryos and in *myod*-MO embryos. Thus, Pbx function might be important for the efficient progression of fast-muscle differentiation. Perhaps the upregulation of *aldh1a2* expression in the tail of *pbx2*-MO; *pbx4*-MO embryos (Fig. 2D) further reflects a defect in differentiation.

The delayed differentiation of a specific muscle type in *pbx2*-MO; *pbx4*-MO embryos is reminiscent of that observed for *Myod* or *Myf5* knock-outs in mice. Whereas loss of *Myod* and *Myf5* together in mice causes a loss of all skeletal muscle development (Rudnicki et al., 1993), loss of *Myod* alone results in delayed hypaxial muscle differentiation, and loss of *Myf5* alone results in delayed epaxial muscle differentiation (Braun et al., 1994; Kablar et al., 1997). Such studies analyzing muscle-differentiation delays have revealed that *Myod* and *Myf5* can play unique roles, and, thus, have had significant impacts on our understanding of muscle development.

Our demonstration that Pbx is needed for the activity of a bHLH protein, in addition to previous examples of homeodomain-bHLH interactions, suggests that homeodomain modulation of bHLH activity could be a widespread mechanism to modulate cell-type diversity. Previous studies have shown interactions between homeodomain proteins and bHLH proteins at specific promoters (Westerman et al., 2003). In particular, the bHLH protein NeuroD1, along with its bHLH heterodimer partner E47, binds with the homeodomain protein Pdx1 to synergistically activate the insulin promoter in B cells within the pancreas (Glick et al., 2000). Mouse *Pbx1*^{-/-} embryos have defects in pancreas differentiation, and Pbx1 interacts with Pdx1 (Kim et al., 2002), but it is not known whether Pbx modulates bHLH activity in the pancreas. In addition to their biochemical interactions, genetic interactions have suggested that bHLH and homeodomain proteins can modulate the activity of each other. Interactions between the homeodomain Nkx2-2 and Olig bHLH proteins in the mouse spinal cord regulate a cell-fate decision between an interneuron fate and a glial fate (Sun et al., 2001). This study suggested that the activity of Olig is dependent on its context with a homeodomain protein. Also, a Hairy-related bHLH protein can inhibit the activity of a Hox protein in regulating epidermal cell

fates in *Caenorhabditis elegans* (Alper and Kenyon, 2001). However, in these cases, the mechanism by which these modulations occur is not known.

Our results provide a novel demonstration of how interactions between different types of 'master regulatory' factors control cell-type diversity. Homeodomain factors control positional identities, whereas bHLH proteins control cell-type identities. We propose that these identity programs merge together such that homeodomain proteins modulate the set of bHLH-activated genes to achieve region-specific cellular phenotypes. Recent studies are continuing to underscore the role of homeodomain proteins in modulating cellular diversity within a cell type, because Hox proteins are now understood to modulate skin-cell phenotype in different regions of the body (Rinn et al., 2006). We propose that homeodomain proteins, by instructing bHLH proteins to regulate a subset of their target genes, provide competence for a cell to execute a region-specific differentiation program.

We thank the following colleagues for generously providing reagents: Peter Currie, Stephen Devoto, Zhiyuan Gong, Randall Moon, Heike Pöpperl, Charles Sagerstrom, Monte Westerfield and Debbie Yelon. The Zebrafish International Resource Center (supported by grants RR12546 and RR15402-01 from the NIH) provided cDNA clones. We thank Daniel Osborn and Simon Hughes for advice on use of the anti-Myf5 antibody. We also thank Erwin Analau and Sean Rhodes for technical assistance; and Stephen Devoto, Clarissa Henry, Estelle Hirsinger and members of the Moens and Tapscott labs, especially Laurie Snider, for advice. B.P. was supported by an HHMI Undergraduate Research Internship and by the Washington NASA Space Grant Consortium. C.B.M. is an investigator with the Howard Hughes Medical Institute. This work was supported by a Research Development Grant from the Muscular Dystrophy Association (L.M.) and by NIH AR45113 (S.J.T.).

Supplementary material

Supplementary material for this article is available at <http://dev.biologists.org/cgi/content/full/134/18/3371/DC1>

References

- Alper, S. and Kenyon, C. (2001). REF-1, a protein with two bHLH domains, alters the pattern of cell fusion in *C. elegans* by regulating Hox protein activity. *Development* **128**, 1793-1804.
- Bader, D., Masaki, T. and Fischman, D. A. (1982). Immunocytochemical analysis of myosin heavy chain during avian myogenesis in vivo and in vitro. *J. Cell Biol.* **95**, 763-770.
- Bárány, M. (1967). ATPase activity of myosin correlated with speed of muscle shortening. *J. Gen. Physiol. Suppl.* **50**, 197-216.
- Barresi, M. J. F., Stickney, H. L. and Devoto, S. H. (2000). The zebrafish *slow-muscle-omitted* gene product is required for Hedgehog signal transduction and the development of slow muscle identity. *Development* **127**, 2189-2199.
- Begemann, G., Schilling, T. F., Rauch, G. J., Geisler, R. and Ingham, P. W. (2001). The zebrafish neckless mutation reveals a requirement for raldh2 in mesodermal signals that pattern the hindbrain. *Development* **128**, 3081-3094.
- Bergstrom, D. A. and Tapscott, S. J. (2001). Molecular distinction between specification and differentiation in the myogenic basic helix-loop-helix transcription factor family. *Mol. Cell Biol.* **21**, 2404-2412.
- Bergstrom, D. A., Penn, B. H., Strand, A., Perry, R. L., Rudnicki, M. A. and Tapscott, S. J. (2002). Promoter-specific regulation of MyoD binding and signal transduction cooperate to pattern gene expression. *Mol. Cell* **9**, 587-600.
- Berkes, C. A., Bergstrom, D. A., Penn, B. H., Seaver, K. J., Knoepfler, P. S. and Tapscott, S. J. (2004). Pbx marks genes for activation by MyoD indicating a role for a homeodomain protein in establishing myogenic potential. *Mol. Cell* **14**, 465-477.
- Blagden, C. S., Currie, P. D., Ingham, P. W. and Hughes, S. M. (1997). Notochord induction of zebrafish slow muscle mediated by Sonic hedgehog. *Genes Dev.* **11**, 2163-2175.
- Braun, T., Bober, E., Rudnicki, M. A., Jaenisch, R. and Arnold, H.-H. (1994). MyoD expression marks the onset of skeletal myogenesis in homozygous Myf-5 mutant mice. *Development* **120**, 3083-3092.
- Bryson-Richardson, M. J., Dagggett, D. F., Cortes, F., Neyt, C., Keenan, D. G. and Currie, P. D. (2005). Myosin heavy chain expression in zebrafish and slow muscle composition. *Dev. Dyn.* **233**, 1018-1022.
- Buckingham, M. (2001). Skeletal muscle formation in vertebrates. *Curr. Opin. Genet. Dev.* **11**, 440-448.
- Cao, Y., Kumar, R. M., Penn, B. H., Berkes, C. A., Kooperberg, C., Boyer, L. A., Young, R. A. and Tapscott, S. J. (2006). Global and gene-specific analyses show distinct roles for MyoD and MyoG at a common set of promoters. *EMBO J.* **25**, 502-511.
- Currie, P. D. and Ingham, P. W. (1996). Induction of a specific muscle cell type by a hedgehog-like protein in zebrafish. *Nature* **382**, 452-455.
- de la Serna, I. L., Ohkawa, Y., Berkes, C. A., Bergstrom, D. A., Dacwag, C. S., Tapscott, S. J. and Imbalzano, A. N. (2005). MyoD targets chromatin remodeling complexes to the myogenin locus prior to forming a stable DNA-bound complex. *Mol. Cell Biol.* **25**, 3997-4009.
- Devoto, S. H., Melancon, E., Eisen, J. S. and Westerfield, M. (1996). Identification of separate slow and fast muscle precursor cells in vivo, prior to somite formation. *Development* **121**, 439-451.
- Draper, B. W., Morcos, P. A. and Kimmel, C. B. (2001). Inhibition of zebrafish fgf8 pre-mRNA splicing with morpholino oligos: a quantifiable method for gene knockdown. *Genesis* **30**, 154-156.
- Du, S. J., Devoto, S. H., Westerfield, M. and Moon, R. T. (1997). Positive and negative regulation of muscle cell identity by members of the hedgehog and TGF-beta gene families. *J. Cell Biol.* **139**, 145-156.
- Erickson, T., Scholpp, S., Brand, M., Moens, C. B. and Waskiewicz, A. J. (2007). Pbx proteins cooperate with Engrailed to pattern the midbrain-hindbrain and diencephalic-mesencephalic boundaries. *Dev. Biol.* **301**, 504-517.
- Feng, X., Adiarte, E. G. and Devoto, S. H. (2006). Hedgehog acts directly on the zebrafish dermomyotome to promote myogenic differentiation. *Dev. Biol.* **300**, 736-746.
- Gehring, W. J. (1996). The master control gene for morphogenesis and evolution of the eye. *Genes Cells* **1**, 11-15.
- Glick, E., Leshkowitz, D. and Walker, M. D. (2000). Transcription factor BETA2 acts cooperatively with E2A and PDX1 to activate the insulin gene promoter. *J. Biol. Chem.* **275**, 2199-2204.
- Grandel, H., Lun, K., Rauch, G. J., Rhinn, M., Piotrowski, T., Houart, C., Sordino, P., Kuchler, A. M., Schulte-Merker, S., Geisler, R. et al. (2002). Retinoic acid signalling in the zebrafish embryo is necessary during pre-segmentation stages to pattern the anterior-posterior axis of the CNS and to induce a pectoral fin bud. *Development* **129**, 2851-2865.
- Groves, J. A., Hammond, C. L. and Hughes, S. M. (2005). Fgf8 drives myogenic progression of a novel lateral fast muscle fibre population in zebrafish. *Development* **132**, 4211-4222.
- Hamade, A., Deries, M., Begemann, G., Bally-Cuif, L., Genêt, C., Sabatier, F., Bonniou, A. and Cousin, X. (2006). Retinoic acid activates myogenesis in vivo through Fgf8 signalling. *Dev. Biol.* **289**, 127-140.
- Hammond, C. L., Hinitz, Y., Osborn, D. P. S., Minchin, J. E. N., Tettamanti, G. and Hughes, S. M. (2007). Signals and myogenic regulatory factors restrict *pax3* and *pax7* expression to dermomyotome-like tissue in zebrafish. *Dev. Biol.* **302**, 504-521.
- Hatta, K., Bremiller, R., Westerfield, M. and Kimmel, C. B. (1991). Diversity of expression of engrailed-like antigens in zebrafish. *Development* **112**, 821-832.
- Hollway, G. E., Bryson-Richardson, R. J., Berger, S., Cole, N. J., Hall, T. E. and Currie, P. D. (2007). Whole-somite rotation generates muscle progenitor cell compartments in the developing zebrafish embryo. *Dev. Cell* **12**, 207-219.
- Ju, B., Chong, S. W., He, J., Wang, X., Xu, Y., Wan, H., Tong, Y., Yan, T., Korzh, V. and Gong, Z. (2003). Recapitulation of fast skeletal muscle development in zebrafish by transgenic expression of GFP under the *mylz2* promoter. *Dev. Dyn.* **227**, 14-26.
- Kablar, B., Krastel, K., Ying, C., Asakura, A., Tapscott, S. J. and Rudnicki, M. A. (1997). MyoD and Myf-5 differentially regulate the development of limb versus trunk skeletal muscle. *Development* **124**, 4729-4738.
- Kim, S. K., Selleri, L., Lee, J. S., Zhang, A. Y., Gu, X., Jacobs, Y. and Cleary, M. L. (2002). Pbx1 inactivation disrupts pancreas development and in *lpf1*-deficient mice promotes diabetes mellitus. *Nat. Genet.* **30**, 430-435.
- Knoepfler, P. S., Bergstrom, D. A., Uetsuki, T., Dac-Korytko, I., Sun, Y. H., Wright, W. E., Tapscott, S. J. and Kamps, M. P. (1999). A conserved motif N-terminal to the DNA-binding domains of myogenic bHLH transcription factors mediates cooperative DNA binding with Pbx-Meis1/Prep1. *Nucleic Acids Res.* **27**, 3752-3761.
- Kobayashi, M., Fujioka, M., Tolkunova, E. N., Deka, D., Abu-Shaar, M., Mann, R. S. and Jaynes, J. B. (2003). Engrailed cooperates with extradenticle and homothorax to repress target genes in *Drosophila*. *Development* **130**, 741-751.
- Lee, J. E. (1997). Basic helix-loop-helix genes in neural development. *Curr. Opin. Neurobiol.* **7**, 13-20.
- Liu, J. and Fire, A. (2000). Overlapping roles of two Hox genes and the *exd* ortholog *ceh-20* in diversification of the *C. elegans* postembryonic mesoderm. *Development* **127**, 5179-5190.
- Maves, L., Jackman, W. and Kimmel, C. B. (2002). FGF3 and FGF8 mediate a rhombomere 4 signaling activity in the zebrafish hindbrain. *Development* **129**, 3825-3837.
- Moens, C. B. and Selleri, L. (2006). Hox cofactors in vertebrate development. *Dev. Biol.* **291**, 193-206.
- Ohlendieck, K., Frömming, G. R., Murray, B. E., Maguire, P. B., Leisner, E.,

- Traub, I. and Pette, D.** (1999). Effects of chronic low-frequency stimulation on Ca²⁺-regulatory membrane proteins in rabbit fast muscle. *Pflugers Arch.* **438**, 700-708.
- Oxtoby, E. and Jowett, T.** (1993). Cloning of the zebrafish krox-20 gene (*krx-20*) and its expression during hindbrain development. *Nucleic Acids Res.* **21**, 1087-1095.
- Pearson, J. C., Lemons, D. and McGinnis, W.** (2005). Modulating Hox gene functions during animal body patterning. *Nat. Rev. Genet.* **6**, 893-904.
- Peers, B., Sharma, S., Johnson, T., Kamps, M. and Montminy, M.** (1995). The pancreatic islet factor STF-1 binds cooperatively with Pbx to a regulatory element in the somatostatin promoter: importance of the FPWMK motif and of the homeodomain. *Mol. Cell. Biol.* **15**, 7091-7097.
- Peltenburg, L. T. and Murre, C.** (1996). Engrailed and Hox homeodomain proteins contain a related Pbx interaction motif that recognizes a common structure present in Pbx. *EMBO J.* **15**, 3385-3393.
- Penn, B. H., Bergstrom, D. A., Dilworth, F. J., Bengal, E. and Tapscott, S. J.** (2004). A MyoD-generated feed-forward circuit temporally patterns gene expression during skeletal muscle differentiation. *Genes Dev.* **18**, 2348-2353.
- Pöpperl, H., Rikhof, H., Chang, H., Haffter, P., Kimmel, C. B. and Moens, C. B.** (2000). Lazarus is a novel pbx gene that globally mediates hox gene function in zebrafish. *Mol. Cell* **6**, 255-267.
- Quong, M. W., Romanow, W. J. and Murre, C.** (2002). E protein function in lymphocyte development. *Annu. Rev. Immunol.* **20**, 301-322.
- Reiser, P. J., Moss, R. L., Giulian, G. G. and Greaser, M. L.** (1985). Shortening velocity in single fibers from adult rabbit soleus muscles is correlated with myosin heavy chain composition. *J. Biol. Chem.* **260**, 9077-9080.
- Rinn, J. L., Bondre, C., Gladstone, H. B., Brown, P. O. and Chang, H. Y.** (2006). Anatomic demarcation by positional variation in fibroblast gene expression programs. *PLoS Genet.* **2**, e119.
- Rudnicki, M. A., Schnegelsberg, P. N. J., Stead, R. H., Braun, T., Arnold, H.-H. and Jaenisch, R.** (1993). MyoD or Myf-5 is required for the formation of skeletal muscle. *Cell* **75**, 1351-1359.
- Sepich, D. S., Wegner, J., O'Shea, S. and Westerfield, M.** (1998). An altered intron inhibits synthesis of the acetylcholine receptor α -subunit in the paralyzed zebrafish mutant *nic1*. *Genetics* **148**, 361-372.
- Stellabotte, F., Dobbs-McAuliffe, B., Fernandez, D. A., Feng, X. and Devoto, S. H.** (2007). Dynamic somite cell rearrangements lead to distinct waves of myotome growth. *Development* **134**, 1253-1257.
- Sun, T., Echelard, Y., Lu, R., Yuk, D., Kaing, S., Stiles, C. D. and Rowitch, D. H.** (2001). Olig bHLH proteins interact with homeodomain proteins to regulate cell fate acquisition in progenitors of the ventral neural tube. *Curr. Biol.* **11**, 1413-1420.
- Tapscott, S. J.** (2005). The circuitry of a master switch: MyoD and the regulation of skeletal muscle gene transcription. *Development* **132**, 2685-2695.
- Thisse, B., Pflumio, S., Fürthauer, M., Loppin, B., Heyer, V., Degrave, A., Woehl, R., Lux, A., Steffan, T., Charbonnier, X. Q. et al.** (2001). Expression of the zebrafish genome during embryogenesis (NIH R01 RR15402). ZFIN Direct Data Submission. ZFIN ID: ZDB-PUB-010810-1.
- Vlachakis, N., Ellstrom, D. R. and Sagerstöm, C. G.** (2000). A novel pbx family member expressed during early zebrafish embryogenesis forms trimeric complexes with Meis3 and Hoxb1b. *Dev. Dyn.* **217**, 3227-3239.
- Waskiewicz, A. J., Rikhof, H. A., Hernandez, R. E. and Moens, C. B.** (2001). Zebrafish Meis functions to stabilize Pbx proteins and regulate hindbrain patterning. *Development* **128**, 4139-4151.
- Waskiewicz, A. J., Rikhof, H. A. and Moens, C. B.** (2002). Eliminating zebrafish pbx proteins reveals a hindbrain ground state. *Dev. Cell* **3**, 723-733.
- Weinberg, E. S., Allende, M. L., Kelly, C. S., Abdelhamid, A., Murakami, T., Andermann, P., Doerre, O. G., Grunwald, D. J. and Riggelman, B.** (1996). Developmental regulation of zebrafish MyoD in wild-type, no tail and spadetail embryos. *Development* **122**, 271-280.
- Weintraub, H., Tapscott, S. J., Davis, R. L., Thayer, M. J., Adam, M. A., Lassar, A. B. and Miller, A. D.** (1989). Activation of muscle-specific genes in pigment, nerve, fat, liver, and fibroblast cell lines by forced expression of MyoD. *Proc. Natl. Acad. Sci. USA* **86**, 5434-5438.
- Westerfield, M.** (2000). *The Zebrafish Book. A Guide for the Laboratory use of Zebrafish (Danio rerio)* (4th edn). Eugene: University of Oregon Press.
- Westerman, B. A., Murre, C. and Oudejans, C. B.** (2003). The cellular Pax-Hox-helix connection. *Biochim. Biophys. Acta* **1629**, 1-7.
- Wolff, C., Roy, S. and Ingham, P. W.** (2003). Multiple muscle cell identities induced by distinct levels and timing of hedgehog activity in the zebrafish embryo. *Curr. Biol.* **13**, 1169-1181.
- Xu, Y., He, J., Wang, X., Lim, T. M. and Gong, Z.** (2000). Asynchronous activation of 10 muscle-specific genes during zebrafish somitogenesis. *Dev. Dyn.* **219**, 201-215.
- Yelon, D., Horne, S. A. and Stainier, D. Y. R.** (1999). Restricted expression of cardiac myosin genes reveals regulated aspects of heart tube assembly in zebrafish. *Dev. Biol.* **214**, 23-37.
- ZFIN** (2006). Zebrafish Information Network. <http://www.zfin.org/>.

Table S1. Microarray analysis of control MO versus pbx2-MO; pbx4-MO at 10-somite stage (normalized expression levels are in log2 scale)													
Affymetrix ID	Gene	False discovery rate	Control lmean	pbx-MO Mean	Control l1	Control l2	Control l3	pbx-MO 1	pbx-MO 2	pbx-MO 3	Fold Change	RefID	
AFFX-Dr-acta1-5_at	acta1	0.0023	8.76	8.091	8.805	8.615	8.859	8.156	8.058	8.06	0.669	NM_212971 /// NM_213221	
Dr10085.2.S1_a_at	ccni	0.00024	9.862	9.059	9.842	9.851	9.893	9.066	9.009	9.103	0.803	NM_684760	
Dr10102.1.A1_at	sb:cb528	0.00249	8.746	7.87	8.548	8.752	8.939	7.985	7.874	7.75	0.876	NM_684760	
Dr10102.2.S1_at	sb:cb528	0.00256	9.39	8.423	9.238	9.352	9.579	8.608	8.444	8.216	0.967	NM_001006073	
Dr10141.1.A1_at	zgc:101687	2.00E-04	7.526	6.139	7.439	7.386	7.754	6.218	6.045	6.155	1.387	NM_131844	
Dr10168.2.S1_at	maf	0.04611	6.289	6.956	6.064	6.354	6.448	6.678	6.836	7.354	-0.667	NM_131602	
Dr10244.1.S1_at	dcd1	0.00207	8.356	8.96	8.294	8.426	8.349	8.966	8.881	9.034	-0.604	NM_131510	
Dr10334.1.S1_at	casp8	0.00693	7.431	8.197	7.654	7.481	7.158	8.161	8.146	8.286	-0.766	---	
Dr10958.1.A1_at	---	0.00633	6.498	7.172	6.37	6.522	6.601	6.967	7.258	7.291	-0.674	---	
Dr11077.1.A1_at	---	0.00291	7.716	7.076	7.835	7.569	7.744	7.114	7.076	7.039	0.64	---	
Dr1121.1.A1_at	---	0.02911	4.845	5.483	4.699	5.075	4.762	5.216	5.498	5.734	-0.638	---	
Dr11316.1.S1_at	nap11	0.00792	10.738	10.087	10.829	10.889	10.498	10.127	10.112	10.021	0.651	NM_201318	
Dr11374.1.A1_at	alcy12	0.01297	8.012	8.643	7.738	8.168	8.129	8.578	8.693	8.656	-0.631	NM_201340	
Dr11374.2.A1_at	LOC555165	0.00101	8.718	9.319	8.695	8.742	8.717	9.323	9.356	9.279	-0.601	NM_697408	
Dr1139.1.A1_at	---	0.00072	6.766	5.826	6.767	6.697	6.833	5.838	5.978	5.663	0.94	---	
Dr11479.1.A1_at	LOC561833	0.01474	6.986	7.734	7.21	6.954	6.795	7.582	7.612	8.008	-0.748	NM_685232	
Dr11520.2.S1_a_at	LOC563637	0.00337	10.479	9.87	10.478	10.537	10.422	10.008	9.828	9.774	0.609	NM_200917 /// NM_684021 /// NM_690444 /// NM_704135 /// NM_704136 /// NM_704137	
Dr11532.1.S1_at	LOC567152	0.01696	6.506	7.166	6.384	6.635	6.499	6.873	7.269	7.356	-0.66	NM_131445	
Dr123.1.S1_at	chrna1	0.02794	6.297	5.693	6.008	6.393	6.491	5.761	5.507	5.811	0.604	NM_200605	
Dr12459.1.A1_at	zgc:63561	0.02068	7.18	7.907	6.938	7.112	7.489	7.678	8.005	8.037	-0.727	NM_001020560	
Dr12544.1.A1_at	zgc:110044	0.0087	7.771	8.39	7.78	7.847	7.686	8.451	8.544	8.175	-0.619	NM_131851	
Dr12573.1.S1_at	sox32	0.00021	5.772	6.679	5.77	5.84	5.704	6.712	6.607	6.719	-0.907	NM_688507	
Dr12669.1.A1_at	wu:fi72h12	0.00312	7.679	8.278	7.784	7.584	7.671	8.295	8.199	8.338	-0.599	---	
Dr12753.1.A1_at	---	0.00055	7.671	8.398	7.655	7.717	7.64	8.365	8.36	8.468	-0.727	---	
Dr12868.1.A1_at	---	0.00792	6.725	7.526	6.793	6.633	6.75	7.612	7.217	7.749	-0.801	---	
Dr12920.1.S1_at	zgc:103764	0.00063	10.461	9.804	10.459	10.478	10.447	9.816	9.833	9.764	0.657	---	
Dr12947.1.A1_at	---	0.0037	6.429	7.036	6.289	6.532	6.465	6.98	7.024	7.105	-0.607	---	
Dr13338.1.S1_at	gbx1	0.01237	7.77	7.122	7.679	7.712	7.918	6.998	7	7.369	0.648	NM_174861	

Dr19562.1.S1_at	--	0.0064	9.424	8.242	9.748	9.568	8.957	8.239	8.37	8.117	1.182	--
Dr19567.1.S1_at	zgc:73377	0.00063	6.989	6.176	7.057	6.965	6.944	6.186	6.069	6.273	0.813	NM_200810
Dr19794.1.A1_at	--	0.02068	7.338	8.121	7.515	7.513	6.987	7.867	8.258	8.239	-0.783	--
Dr1999.1.S1_at	zgc:109940	0.01188	7.231	6.224	6.823	7.285	7.585	6.296	6.349	6.026	1.007	NM_001020532
Dr20125.1.A1_s_at	zgc:111983	0.01216	7.692	7.007	7.493	7.736	7.847	6.987	7.212	6.822	0.685	NM_001017803
Dr20131.2.A1_at	clu:///	0.00644	6.01	6.703	5.864	5.964	6.202	6.548	6.826	6.736	-0.693	NM_200802 /// XM_695015
Dr20214.1.A1_at	tuba8l3	0.01641	8.252	7.488	8.133	8.042	8.583	7.587	7.593	7.283	0.764	NM_001003558
Dr2054.1.A1_at	wu:fb78e03	9.00E-04	7.243	7.89	7.224	7.24	7.266	7.826	7.888	7.956	-0.647	--
Dr20954.1.S1_at	kns2	0.00256	9.875	9.26	9.904	9.765	9.957	9.211	9.228	9.342	0.615	NM_199954
Dr20962.1.S1_at	hoxb2a	0.00017	8.354	7.5	8.404	8.329	8.329	7.479	7.513	7.507	0.854	NM_131116
Dr20990.2.S1_at	ttn	0.01473	8.491	7.896	8.602	8.421	8.45	8.143	7.781	7.765	0.595	XM_679005
Dr2177.1.A1_at	--	0.00337	8.808	8.178	8.853	8.749	8.822	8.25	8.261	8.021	0.63	--
Dr21859.1.A1_at	wu:fc74e04	0.00574	7.735	6.857	7.539	8.014	7.651	6.996	6.894	6.682	0.878	--
Dr21919.1.A1_at	zgc:86773	0.00121	5.136	6.031	5	5.122	5.287	5.98	5.95	6.164	-0.895	NM_001002129
Dr21935.1.A1_at	LOC558348	0.00172	6.038	7.04	6.052	5.943	6.119	6.798	7.261	7.06	-1.002	XM_697991
Dr22027.1.A1_at	LOC559444	0.00017	6.333	5.159	6.195	6.362	6.444	5.249	5.095	5.133	1.174	XM_682782
Dr22306.1.S1_at	--	0.02642	6.873	8.484	6.786	6.545	7.288	7.665	8.555	9.234	-1.611	--
Dr23191.1.A1_at	--	0.00043	11.143	10.378	11.215	11.11	11.103	10.413	10.35	10.372	0.765	--
Dr23348.1.A1_at	bmp3	0.00671	9.276	9.996	9.365	9.055	9.407	10.092	9.831	10.066	-0.72	--
Dr24018.1.A1_at	--	0.00901	6.134	5.525	6.303	5.946	6.153	5.64	5.451	5.485	0.609	--
Dr2408.2.S1_at	mimp2	0.00278	9.041	9.632	9.036	9.076	9.012	9.657	9.725	9.512	-0.591	NM_198067
Dr24244.2.S1_a_at	pax6a /// LOC566915	0.00508	8.735	8.104	8.724	8.678	8.805	8.261	8.113	7.939	0.631	NM_131304 /// XM_690200
Dr24246.1.S1_at	h1fx	0.00172	6.809	7.565	6.74	6.982	6.704	7.616	7.513	7.566	-0.756	NM_199276
Dr2426.1.S1_at	zgc:66321	2.00E-04	8.049	6.867	8.11	7.916	8.121	6.749	6.865	6.987	1.182	NM_201118
Dr24774.1.S1_at	hoxa3a	7.00E-05	8.2	7.147	8.143	8.224	8.233	7.104	7.194	7.144	1.053	NM_131534
Dr24774.1.S2_at	hoxa3a	8.00E-05	6.25	5.082	6.192	6.268	6.291	4.993	5.181	5.074	1.168	NM_131534
Dr2499.1.S1_at	myog	0.01023	8.331	7.301	8.17	8.075	8.749	7.479	7.34	7.085	1.03	NM_131006
Dr25422.1.S1_s_at	hbae1 /// LOC572729	0.01893	8.29	7.59	8.17	8.104	8.597	7.775	7.42	7.574	0.7	NM_182940 /// XM_678451 /// XM_679920 /// XM_680614 /// XM_680782 /// XM_701993 /// XM_702647 /// XM_702998 /// XM_703072
Dr25425.1.A1_at	sdf2	0.00579	8.495	9.12	8.306	8.549	8.632	9.052	9.19	9.119	-0.625	NM_200039
Dr25450.1.A1_at	--	0.00882	8.518	9.234	8.258	8.598	8.698	9.125	9.35	9.226	-0.716	--
Dr25497.1.S1_at	szl	0.0057	8.426	7.641	8.611	8.448	8.218	7.745	7.456	7.723	0.785	NM_181663
Dr25527.1.A1_at	LOC572080	0.00508	6.625	5.996	6.582	6.801	6.493	6.07	5.932	5.986	0.629	XM_695747
Dr25527.1.A1_x_at	LOC572080	0.00193	6.623	6.001	6.645	6.694	6.53	6.074	5.971	5.957	0.622	XM_695747
Dr25555.1.A1_at	--	0.00172	6.682	7.693	6.534	6.597	6.914	7.748	7.527	7.805	-1.011	--
Dr25683.10.S1_at	ctslb	0.00193	8.521	7.854	8.538	8.403	8.622	7.865	7.911	7.787	0.667	NM_131198
Dr25770.3.A1_at	zgc:113283	0.01996	6.415	7.033	6.497	6.137	6.612	6.956	6.964	7.179	-0.618	NM_001017556

Dr9272.1.A1_at	zgc:112278 wu:fc21h08 ///	0.00147	9	8.35	9.099	8.949	8.953	8.418	8.311	8.321	0.65	NM_001029958
Dr9288.1.S1_at	LOC572929	0.00334	7.048	7.68	6.966	6.992	7.187	7.727	7.724	7.59	-0.632	NM_001001815 ///XM_682622
Dr9353.1.S1_at	zgc:55733	0.00046	9.676	8.79	9.648	9.632	9.749	8.913	8.748	8.707	0.886	NM_200993
Dr9654.1.A1_at	ppia	0.00085	8.465	9.838	8.653	8.229	8.513	10.059	9.615	9.84	-1.373	NM_212758
DrAffx.1.2.S1_at	bhik	0.00725	10.022	9.083	10.036	10.28	9.751	9.065	9.31	8.874	0.939	NM_131803
DrAffx.1.42.S1_s_at	cugbp2	0.0023	7.667	7.046	7.574	7.781	7.647	7.019	7.105	7.014	0.621	NM_194401
DrAffx.1.74.S1_at	sepw2b	0.00613	9.107	8.451	9.219	9.154	8.949	8.611	8.413	8.33	0.656	NM_194418
DrAffx.2.59.S1_at	LOC555198	0.00133	8.148	7.163	8.217	8.064	8.163	7.334	6.931	7.225	0.985	XM_677650
DrAffx.2.62.S1_at	cyp26b1	0.00046	7.264	6.262	7.38	7.23	7.181	6.406	6.179	6.2	1.002	NM_212666

Table S2. Microarray analysis of control MO versus pbx2-MO; pbx4-MO at 18-somite stage (normalized expression levels are in log2 scale)

Affymetrix ID	Gene	False discovery rate	Control mean	pbx-MO mean	Control 1	Control 2	Control 3	pbx-MO 1	pbx-MO 2	pbx-MO 3	Fold Change	RefID
AFFX-Dr-acta1-5_at	acta1	0.00229	9.651	8.848	9.763	9.59	9.601	9.049	8.63	8.864	0.803	NM_131591
Dr1.1.S1_at	cmyc	0.00639	7.628	7.031	7.536	7.607	7.741	7.039	6.86	7.195	0.597	NM_131412
Dr10141.1.A1_at	zgc:101687	0.00066	7.058	6.086	7.156	7.098	6.919	6.177	5.905	6.177	0.972	NM_001006073
Dr10295.1.A1_at	--	0.00286	6.599	7.212	6.645	6.707	6.445	7.174	7.274	7.188	-0.613	--
Dr10326.1.S1_at	junb	0.01457	6.418	7.28	6.216	6.41	6.627	6.872	7.332	7.637	-0.862	NM_213556
Dr10438.1.S1_at	atp1a2a /// LOC557474	0.00434	6.842	6.239	6.898	6.861	6.767	6.39	6.069	6.258	0.603	NM_131683 /// XM_680554
Dr10457.1.S2_at	vent	0.01438	4.61	5.287	4.414	4.716	4.701	4.99	5.349	5.522	-0.677	NM_131700
Dr10602.1.A1_at	LOC556764	0.00158	8.849	9.519	8.713	8.9	8.935	9.517	9.568	9.471	-0.67	XM_677901 /// XM_701698 /// XM_701699 /// XM_701700 /// XM_701701 /// XM_701702 /// XM_701703 /// XM_701704 /// XM_701705
Dr10607.1.A1_at	vmhc	0.00021	6.358	4.68	6.679	6.395	6.001	4.711	4.693	4.636	1.678	XM_679302 /// XM_702387 /// XM_702388 /// XM_702389 /// XM_702390 /// XM_702391 /// XM_702392
Dr1081.1.S1_at	eef1b2	0.00115	12.231	11.616	12.274	12.22	12.199	11.655	11.591	11.603	0.615	NM_199949
Dr11035.1.A1_at	wu:fj35f12	0.00317	5.049	5.71	5.012	5.086	5.048	5.525	5.719	5.885	-0.661	--
Dr11059.1.S1_at	zgc:109987	0.00376	5.35	5.939	5.304	5.337	5.409	5.773	6.018	6.027	-0.589	--
Dr11072.1.A1_at	zgc:73075	0.00239	3.492	4.121	3.462	3.497	3.518	4.005	4.276	4.081	-0.629	NM_200825
Dr11316.1.S1_at	nap1l1	0.00325	10.004	9.3	10.2	10.03	9.782	9.295	9.32	9.285	0.704	NM_201318
Dr11316.2.A1_at	wu:fb33e04	0.00325	10.569	9.936	10.668	10.613	10.427	9.872	9.865	10.072	0.633	--
Dr11520.3.S1_a_at	sich211-51e12.7 /// LOC556109 /// LOC563637	0.00292	10.45	9.799	10.385	10.521	10.443	9.904	9.876	9.616	0.651	XM_678095 /// XM_679914 /// XM_681132 /// XM_701815 /// XM_703223 /// XM_703224 /// XM_703225 /// XM_704710 /// XM_704711 /// XM_704712
Dr11661.1.S1_at	ypel3	0.03215	6.79	7.452	6.536	6.795	7.039	7.192	7.37	7.794	-0.662	NM_212790
Dr12107.1.A1_at	ndrg1	0.01214	6.255	6.929	6.067	6.497	6.201	6.691	7.065	7.032	-0.674	NM_213348
Dr12134.2.A1_at	LOC566559	0.00126	9.134	10.017	9.156	9.314	8.934	9.997	10.119	9.935	-0.883	XM_687419 /// XM_704560 /// XM_704561
Dr123.1.S1_at	chrna1	0.00079	6.865	5.826	6.888	6.929	6.778	5.824	5.602	6.051	1.039	NM_131445
Dr12387.1.S1_at	efna3	0.00265	6.526	5.926	6.548	6.4	6.63	5.867	5.993	5.919	0.6	NM_131854

Dr12459.1.A1_at	zgc:63561	0.00151	7.005	7.638	6.976	6.958	7.08	7.629	7.565	7.72	-0.633	NM_200605
Dr12467.1.A1_at	--	0.00654	6.804	7.399	6.672	6.972	6.769	7.337	7.543	7.316	-0.595	--
Dr12544.1.A1_at	zgc:110044	0.02092	6.679	7.339	6.719	6.893	6.427	7.389	7.562	7.066	-0.66	NM_001020560
Dr12573.1.S1_at	sox32	0.00159	4.87	5.909	4.813	4.623	5.174	5.938	6.003	5.787	-1.039	NM_131851
Dr12868.1.A1_at	--	0.00115	5.533	6.172	5.479	5.504	5.616	6.176	6.153	6.188	-0.639	--
Dr13284.1.A1_at	LOC557600 /// LOC558499	0.01636	4.408	5.384	4.184	4.608	4.432	5.097	5.929	5.126	-0.976	XM_681807 /// XM_682586 /// XM_703468 /// XM_703726 /// XM_703727 /// XM_703728
Dr13338.1.S1_at	gbx1	0.00011	6.843	5.661	6.713	7.001	6.814	5.636	5.725	5.623	1.182	NM_174861
Dr13544.1.S1_at	pdcd6 /// LOC573822	0.0018	6.874	7.608	6.693	6.958	6.97	7.621	7.521	7.683	-0.734	NM_200950 /// XM_694060
Dr13750.1.S1_at	--	0.0371	6.727	5.895	7.094	6.83	6.256	6.257	5.832	5.955	0.832	--
Dr13796.1.S1_at	upf3b	0.01465	6.352	7.049	6.588	6.301	6.167	7.179	7.217	6.753	-0.697	NM_200954
Dr13796.2.S1_at	upf3b	0.00723	5.39	6.007	5.53	5.228	5.412	6.187	5.965	5.869	-0.617	NM_200954
Dr13796.2.S1_x.at	upf3b	0.00928	6.031	6.62	6.154	5.843	6.096	6.775	6.612	6.474	-0.589	NM_200954
Dr138.1.A1_at	zgc:65774	0.00115	5.238	4.282	5.014	5.422	5.278	4.175	4.264	4.407	0.956	NM_199924
Dr13843.1.S1_at	bhlhb5	0.00012	5.659	4.008	5.483	5.765	5.728	3.95	3.794	4.28	1.651	NM_200955
Dr1388.1.S1_at	gspt1	0.0011	11.223	10.59	11.178	11.28	11.211	10.582	10.577	10.611	0.633	NM_198806
Dr13933.1.A1_at	dlnmt6	3.00E-05	6.319	4.843	6.266	6.483	6.209	4.828	4.783	4.917	1.476	NM_001018140
Dr13971.1.A1_at	--	0.00419	4.175	4.922	4.186	4.132	4.208	4.818	5.208	4.741	-0.747	--
Dr14127.1.S1_at	gfap	0.00234	8.751	8.129	8.776	8.858	8.62	8.065	8.134	8.189	0.622	NM_131373
Dr14140.1.S1_at	LOC569795	0.0015	8.462	9.454	8.468	8.571	8.347	9.425	9.72	9.218	-0.992	XM_693202
Dr14317.1.A1_at	--	0.0011	5.743	4.836	5.888	5.521	5.819	4.8	4.791	4.916	0.907	--
Dr14668.1.S1_at	gch	0.00722	5.157	4.432	5.281	5.264	4.925	4.212	4.468	4.614	0.725	NM_131667
Dr14787.1.A1_at	s100b	0.00527	4.509	3.855	4.604	4.661	4.261	3.831	3.888	3.846	0.654	--
Dr14950.1.A1_at	zgc:123068	0.03099	5.971	5.293	6.23	6.046	5.637	5.489	5.403	4.987	0.678	NM_001007132
Dr15055.1.S1_at	cxcr4a	0.00145	8.24	8.976	8.347	8.261	8.111	8.89	9.09	8.949	-0.736	NM_131882
Dr15056.1.S1_at	mab21l2	0.00204	8.385	9.124	8.401	8.475	8.278	9.295	9.102	8.975	-0.739	NM_152975
Dr15174.1.A1_at	--	0.00409	7.056	6.434	6.94	7.013	7.215	6.499	6.497	6.305	0.622	--
Dr15236.1.A1_at	--	0.00099	5.825	6.714	5.781	5.895	5.799	6.604	6.921	6.616	-0.889	--
Dr15287.1.S1_at	LOC553758	0.00375	9.906	8.94	9.774	9.765	10.178	8.977	8.666	9.179	0.966	XM_686123
Dr15413.1.A1_at	--	0.0206	5.249	5.877	5.439	5.034	5.273	5.629	6.128	5.874	-0.628	--
Dr15714.1.S1_at	hoxd9a	8.00E-05	6.312	7.408	6.361	6.328	6.246	7.34	7.373	7.509	-1.096	NM_131126
Dr15717.1.S1_at	hoxb1a	0.00047	7.146	6.19	7.024	7.1	7.314	6.268	6.207	6.093	0.956	NM_131115
Dr15849.1.S1_at	zgc:55604	0.00654	4.734	5.404	4.634	4.948	4.62	5.294	5.334	5.586	-0.67	NM_201002
Dr15964.1.A1_at	--	0.00646	7.031	7.658	7.102	7.141	6.85	7.81	7.649	7.514	-0.627	--
Dr16035.1.S1_at	zgc:101776	0.00184	6.334	7.035	6.455	6.361	6.186	6.948	7.038	7.119	-0.701	NM_001007404
Dr16111.1.S1_at	tshz1	0.0018	8.461	7.735	8.304	8.529	8.55	7.739	7.628	7.838	0.726	--
Dr16193.1.S1_at	--	0.00132	4.396	4.988	4.448	4.374	4.367	4.981	4.976	5.009	-0.592	--
Dr16230.1.A1_at	zgc:154090	0.008	7.548	6.941	7.485	7.525	7.635	6.797	6.842	7.182	0.607	--
Dr16375.1.A1_at	zgc:136476	0.01296	6.28	6.903	6.437	6.388	6.015	7.042	6.926	6.742	-0.623	NM_001040394
Dr16377.1.A1_at	--	0.00375	6.36	6.998	6.316	6.346	6.418	6.878	6.917	7.199	-0.638	--
Dr16392.1.A1_at	c6	0.00689	5.074	4.401	5.3	5.02	4.903	4.557	4.264	4.381	0.673	NM_200638

Dr16427.1.A1_at	--	0.00491	5.06	5.701	4.844	5.236	5.101	5.644	5.711	5.747	-0.641	--
Dr16456.1.S1_at	mtmr6	0.00394	6.314	6.952	6.486	6.183	6.274	7.073	6.887	6.897	-0.638	NM_199791
Dr16499.1.A1_at	zgc:113038	0.00544	7.446	8.09	7.512	7.28	7.547	8.109	8.238	7.923	-0.644	NM_001020711
Dr1662.1.S1_at	ttnl	0.00434	8.97	8.137	9.028	8.96	8.922	8.479	7.951	7.98	0.833	XM_677962 /// XM_678057 /// XM_678144
Dr16960.1.A1_at	--	0.00334	5.234	5.832	5.182	5.386	5.135	5.909	5.82	5.766	-0.598	--
Dr17100.1.A1_at	LOC558276	0.00066	8.521	7.695	8.513	8.555	8.494	7.781	7.764	7.541	0.826	XM_681478
Dr17108.1.S1_at	--	0.00407	9.02	8.364	8.919	9.248	8.894	8.37	8.331	8.392	0.656	--
Dr1725.1.A1_at	--	0.01901	8.555	7.857	8.438	8.489	8.739	7.659	8.231	7.681	0.698	--
Dr1726.1.A1_at	no16	0.00149	9.274	8.653	9.298	9.188	9.336	8.67	8.69	8.599	0.621	XM_685738
Dr17431.1.S1_at	ica	0.00294	4.905	5.647	4.899	5.053	4.763	5.582	5.839	5.52	-0.742	NM_201587
Dr17549.1.A1_at	egr2a	0.00617	5.539	4.928	5.456	5.45	5.71	5.08	4.892	4.81	0.611	NM_183341
Dr17767.2.A1_at	--	0.00684	6.951	6.352	7.057	6.729	7.066	6.269	6.401	6.386	0.599	--
Dr17767.2.A1_x_at	--	0.00716	7.396	6.722	7.452	7.27	7.466	6.46	6.811	6.896	0.674	--
Dr17971.1.S1_at	--	0.00229	6.96	6.193	6.983	7.013	6.884	6.254	6.354	5.97	0.767	--
Dr180.1.A1_at	cyp26b1	2.00E-05	7.719	6.262	7.639	7.787	7.732	6.392	6.228	6.165	1.457	NM_212666
Dr18158.1.A1_at	--	0.00286	5.129	4.471	5.181	5.094	5.112	4.338	4.661	4.415	0.658	--
Dr18211.1.A1_at	--	0.00234	6.374	7.018	6.268	6.521	6.334	6.96	7.097	6.997	-0.644	--
Dr18293.1.S1_at	LOC572973	1.00E-05	7.769	6.181	7.774	7.86	7.674	6.164	6.15	6.229	1.588	XM_678098 /// XM_701820
Dr18315.1.S1_at	eef1g	0.00132	12.787	12.164	12.848	12.767	12.745	12.135	12.13	12.226	0.623	NM_173263
Dr18514.1.S1_at	zgc:101025	0.00291	7.852	7.169	7.909	7.736	7.912	7.122	7.036	7.349	0.683	NM_001012315
Dr1899.3.A1_at	--	0.01157	9.068	8.384	8.835	9.117	9.252	8.362	8.61	8.18	0.684	--
Dr19144.1.A1_at	p4ha2	0.03256	8.186	8.862	7.915	8.162	8.48	8.559	8.872	9.153	-0.676	NM_001007285
Dr19395.1.A1_at	LOC564042	0.00108	6.232	6.942	6.325	6.23	6.141	6.883	6.996	6.948	-0.71	XM_687402
Dr19416.1.S1_at	otomp	0.00384	5.943	5.253	5.907	5.792	6.131	5.331	5.105	5.322	0.69	XM_689087
Dr19518.1.S1_at	LOC567197	0.00718	5.449	6.125	5.635	5.35	5.361	5.894	6.235	6.245	-0.676	XM_690489
Dr19562.1.S1_at	--	0.00016	8.452	7.233	8.533	8.543	8.279	7.346	7.246	7.108	1.219	--
Dr19888.1.S1_at	ved	0.0011	7.478	8.171	7.455	7.464	7.514	8.061	8.192	8.258	-0.693	NM_183074
Dr19965.1.S1_at	fn1	0.00853	8.703	9.561	8.679	8.813	8.617	9.715	9.817	9.15	-0.858	NM_131520
Dr1999.1.S1_at	zgc:109940	0.0018	7.717	6.81	7.791	7.616	7.744	6.905	6.54	6.985	0.907	NM_001020532
Dr20010.7.S1_at	atp2a1	0.04258	9.398	8.583	9.581	9.382	9.231	8.993	7.989	8.767	0.815	NM_001007029
Dr20131.2.A1_at	clu /// LOC573883	0.0011	5.661	6.825	5.409	5.68	5.892	6.641	6.822	7.012	-1.164	NM_200802 /// XM_695015
Dr20181.1.S1_at	anp32e	0.00428	10.045	9.357	10.031	10.155	9.948	9.506	9.432	9.133	0.688	NM_213277
Dr2054.1.A1_at	wu:fb78e03	0.00419	6.091	6.73	6.138	6.106	6.028	6.584	6.939	6.668	-0.639	--
Dr20850.1.S1_at	fabp7a	0.04631	6.855	6.116	6.781	6.549	7.234	6.056	5.777	6.516	0.739	NM_131605
Dr20962.1.S1_at	hoxb2a	0.00311	7.477	6.724	7.483	7.646	7.303	6.834	6.794	6.545	0.753	NM_131116
Dr20990.2.S1_at	ttn	0.00849	8.763	7.877	9.086	8.405	8.797	8.049	7.686	7.896	0.886	XM_679005
Dr21031.1.S1_at	hoxa4a	0.00412	7.314	6.721	7.274	7.348	7.32	6.595	6.67	6.899	0.593	NM_131535
Dr21216.1.A1_at	wu:fb94b10	0.00132	7.776	6.995	7.822	7.838	7.669	7.164	6.882	6.94	0.781	--
Dr2132.1.A1_at	hao1	0.02787	7.063	6.124	7.17	6.989	7.03	5.652	5.97	6.749	0.939	XM_685763
Dr21613.1.A1_at	--	0.00171	7.235	7.91	7.212	7.308	7.186	7.782	8.026	7.923	-0.675	--
Dr21618.1.A1_at	--	0.00325	7.127	7.716	7.18	7.216	6.984	7.79	7.648	7.71	-0.589	--
Dr21755.1.A1_at	wu:fc51h05	0.00115	6.641	5.901	6.761	6.573	6.588	5.794	5.94	5.968	0.74	--

Dr21808.1.A1_at	wu:fc44g03	0.02111	8.145	8.786	8.223	8.022	8.19	8.755	8.488	9.115	-0.641	--
Dr21859.1.A1_at	wu:fc74e04	0.01747	6.608	5.984	6.546	6.72	6.559	6.026	6.25	5.676	0.624	--
Dr2193.1.A1_at	si:ch211-199g17.7	0.00352	4.532	5.202	4.574	4.572	4.451	5.274	5.341	4.993	-0.67	--
Dr22027.1.A1_at	LOC559444	0.00267	6.274	5.499	6.058	6.399	6.365	5.349	5.612	5.536	0.775	XM_682782
Dr22510.1.S1_at	zgc:112473	0.0018	5.399	4.682	5.396	5.385	5.416	4.624	4.554	4.869	0.717	NM_001017676
Dr2259.1.S1_at	wu:fc58c10	0.00132	6.35	7.303	6.135	6.39	6.524	7.202	7.234	7.474	-0.953	--
Dr22620.1.S1_at	zgc:65964	0.00229	5.704	6.353	5.626	5.803	5.682	6.219	6.415	6.426	-0.649	NM_200584
Dr22831.1.A1_at	map3k5	0.00854	6.378	7.062	6.487	6.394	6.254	7.097	7.292	6.798	-0.684	XM_678295 ///
Dr23067.1.S1_at	myhz1	0.01721	8.126	6.548	8.126	8.137	8.116	6.638	5.624	7.382	1.578	XM_678389
Dr23191.1.A1_at	--	5.00E-05	9.966	8.897	9.994	9.966	9.938	8.903	8.946	8.841	1.069	XM_703824
Dr23348.1.A1_at	bmp3	0.0018	8.142	8.779	8.259	8.046	8.12	8.833	8.747	8.756	-0.637	--
Dr23993.1.A1_at	sb:cb412	0.01066	5.673	6.527	5.615	5.69	5.714	6.126	6.907	6.548	-0.854	--
Dr240.1.A1_at	timp2	0.00132	5.852	6.556	5.817	5.95	5.789	6.471	6.526	6.672	-0.704	NM_182874
Dr24246.1.S1_at	h1fx	0.00021	6.412	8.034	6.267	6.533	6.437	7.99	8.354	7.758	-1.622	NM_199276
Dr2426.1.S1_at	zgc:66321	0.0011	7.328	6.412	7.41	7.331	7.241	6.415	6.212	6.607	0.916	NM_201118
Dr2452.1.A1_at	c9	0.02193	6.669	6.075	6.628	6.649	6.729	5.97	5.842	6.413	0.594	NM_001024435
Dr2452.2.A1_x_at	c9	0.00487	6.429	5.818	6.523	6.362	6.401	5.794	5.665	5.996	0.611	NM_001024435
Dr24527.1.S1_at	--	0.01289	8.058	7.365	8.021	8.033	8.118	7.51	7.581	7.003	0.693	--
Dr24685.1.S1_at	zgc:55970	0.00291	11.07	10.413	11.127	11.102	10.982	10.524	10.483	10.232	0.657	NM_201176
Dr24774.1.S1_at	hoxa3a	0.00026	7.902	6.931	7.792	7.897	8.017	7.005	6.835	6.952	0.971	NM_131534
Dr24774.1.S2_at	hoxa3a	0.00039	5.556	4.511	5.403	5.564	5.7	4.636	4.387	4.509	1.045	NM_131534
Dr24930.1.S1_at	c20orf45	0.00689	8.883	8.109	8.852	8.864	8.932	7.962	7.906	8.46	0.774	NM_199734
Dr2499.1.S1_at	myog	0.00257	9.406	8.702	9.511	9.394	9.312	8.884	8.567	8.654	0.704	NM_131006
Dr25060.3.A1_at	wu:fb53f04	0.00115	8.278	7.429	8.101	8.385	8.35	7.452	7.529	7.307	0.849	--
Dr25187.3.S1_at	zgc:101883	0.00426	9.814	10.443	9.764	9.973	9.707	10.336	10.584	10.411	-0.629	NM_001007375
Dr25214.1.A1_at	--	0.01731	4.187	4.788	3.899	4.228	4.434	4.671	4.824	4.867	-0.601	--
Dr2527.1.A1_at	wu:fb92a12	0.00308	7.753	8.453	7.864	7.872	7.523	8.5	8.44	8.418	-0.7	--
Dr25356.1.A1_at	wu:fc60e10	0.00272	7.765	7	7.524	7.837	7.935	6.918	7.008	7.073	0.765	--
Dr25442.1.A1_at	elovl4	0.00737	6.384	7.072	6.159	6.592	6.402	6.953	7.014	7.247	-0.688	NM_199972
Dr25548.1.S1_at	hsp90b1	0.02754	8.434	9.135	8.361	8.306	8.636	8.983	8.868	9.552	-0.701	NM_198210
Dr25683.10.S1_at	ctslb	0.00241	7.208	6.445	7.078	7.236	7.309	6.243	6.493	6.598	0.763	NM_131198
Dr25683.8.S1_at	LOC567263	0.0011	11.351	10.73	11.364	11.333	11.356	10.734	10.767	10.689	0.621	XM_701148 ///
Dr26.1.S1_at	egr2b	0	7.809	4.485	7.754	7.726	7.947	4.704	4.516	4.235	3.324	XM_704541 ///
Dr26174.1.A1_at	--	0.00079	5.334	6.006	5.331	5.299	5.371	6.053	5.973	5.992	-0.672	XM_704542 ///
Dr26342.1.A1_at	cyp26b1	0	6.725	4.433	6.609	6.826	6.738	4.389	4.35	4.56	2.292	XM_704543 ///
Dr26383.1.A1_at	zgc:113159	0.00245	7.596	8.202	7.593	7.552	7.644	8.063	8.295	8.248	-0.606	XM_704544 ///
Dr2644.1.A1_at	dhrr3	0.00145	6.272	5.455	6.255	6.425	6.134	5.326	5.445	5.594	0.817	NM_001003477
Dr26487.1.S1_at	si:dkey-119m7.4	0.0018	6.387	5.66	6.534	6.202	6.426	5.699	5.642	5.639	0.727	NM_205587
Dr26498.1.A1_at	upf3b	0.00132	8.169	9.032	8.342	8.142	8.024	9.133	9.079	8.884	-0.863	NM_200954
Dr2654.1.A1_at	zgc:91808	0.00229	8.624	9.327	8.513	8.56	8.798	9.221	9.395	9.365	-0.703	NM_001013278
Dr2710.2.A1_at	TOM1	0.00229	6.846	6.214	6.919	6.78	6.84	6.256	6.08	6.306	0.632	--

Dr2813.1.S1_at	zg:c:55461	3.00E-04	9.678	8.798	9.693	9.655	9.685	8.923	8.715	8.755	0.88	NM_213490
Dr2850.1.S1_at	mat2a	0.00603	9.999	9.398	10.102	10.079	9.816	9.417	9.276	9.499	0.601	NM_212637
Dr2914.1.S1_at	mylzz /// LOC572658	0.04547	9.27	8.212	9.406	9.083	9.322	8.533	7.375	8.73	1.058	NM_131188 /// XM_678881
Dr2988.1.A1_at	--	0.00055	9.357	8.442	9.333	9.515	9.222	8.407	8.526	8.393	0.915	--
Dr3010.1.S1_at	pbx2	7.00E-05	8.144	9.771	7.997	8.204	8.23	9.881	9.915	9.518	-1.627	NM_131612
Dr3061.1.A1_at	wu:ff44b09	0.00187	5.888	5.216	5.862	5.864	5.937	5.289	5.055	5.304	0.672	--
Dr318.1.A1_at	asc11b	0.00419	7.271	6.669	7.316	7.38	7.116	6.572	6.664	6.773	0.602	NM_131231
Dr3205.1.S1_at	LOC566096	0.00317	10.868	10.04	10.854	10.784	10.966	10.117	10.248	9.754	0.828	XM_689364
Dr3216.1.A1_at	np	0.00315	7.967	8.571	7.894	8.049	7.957	8.435	8.591	8.687	-0.604	NM_213311
Dr3224.1.S1_at	wu:fc33f12	0.00021	4.7	3.262	4.883	4.835	4.382	3.284	3.215	3.287	1.438	--
Dr3332.1.A1_at	--	0.0011	6.954	6.085	7.131	6.84	6.892	6.205	5.98	6.069	0.869	--
Dr3483.1.A1_at	wu:fc49d09	3.00E-04	9.316	10.101	9.296	9.298	9.354	10.073	10.151	10.078	-0.785	--
Dr3514.1.S1_at	cox4nb	0.0018	7.955	8.568	7.968	7.897	7.998	8.488	8.663	8.552	-0.613	NM_213370
Dr356.1.S1_at	gata2	0.00196	8.101	8.797	8.095	7.956	8.253	8.782	8.87	8.739	-0.696	NM_131233
Dr3681.1.S1_at	zg:c:110380	0.00132	11.025	11.746	10.877	11.055	11.141	11.695	11.774	11.768	-0.721	NM_001017660
Dr3817.1.S1_at	pdia4	0.01027	7.956	8.721	7.857	7.885	8.124	8.506	8.587	9.07	-0.765	NM_199779
Dr3932.1.S1_at	p4ha1	0.00462	8.715	9.586	8.509	9.058	8.579	9.461	9.713	9.584	-0.871	NM_214691
Dr4060.1.S1_at	hadhb	0.01563	8.445	7.844	8.419	8.424	8.491	7.914	7.541	8.077	0.601	NM_200019
Dr4096.1.S1_at	zg:c:65861	0.00055	10.915	11.818	10.896	11.078	10.772	11.835	11.857	11.763	-0.903	NM_199552
Dr4164.1.A1_at	fbx15	0.00243	7.942	8.762	7.85	8.07	7.905	8.603	9.01	8.672	-0.82	XM_689952
Dr421.1.A1_at	dusp22	0.00132	5.479	6.245	5.31	5.628	5.5	6.276	6.213	6.245	-0.766	NM_001002514
Dr4212.1.S1_at	ldha	0.0369	6.691	7.514	6.542	7.08	6.45	7.41	7.97	7.16	-0.823	NM_131246
Dr4227.1.S1_at	otop1	0.00132	7.205	7.899	7.203	7.192	7.22	7.75	7.981	7.966	-0.694	NM_198803
Dr4233.1.A1_at	wu:fk35g07	0.00916	7.012	6.387	7.063	6.978	6.996	6.561	6.487	6.113	0.625	--
Dr4286.1.A1_at	im:689552	0.00023	8.948	9.871	9.008	8.983	8.852	9.933	9.892	9.789	-0.923	XM_690536
Dr4305.1.A1_at	wsb1	0.00495	8.975	9.686	8.792	9.066	9.069	9.5	9.875	9.682	-0.711	NM_199633
Dr4453.1.S1_at	ppp1r3b	0.00018	8.957	7.978	8.966	9.01	8.896	7.986	7.878	8.072	0.979	NM_212709
Dr463.1.S1_at	her5	0.00168	5.741	4.811	5.69	5.571	5.961	4.634	4.842	4.958	0.93	NM_131077
Dr4681.1.A1_at	LOC556025	0.00229	8.656	7.743	8.737	8.666	8.564	8.043	7.654	7.532	0.913	XM_678716
Dr4794.2.S1_at	zg:c:66117	0.0088	9.425	8.733	9.412	9.33	9.534	8.773	8.453	8.973	0.692	NM_199560
Dr4797.1.S1_at	zg:c:123103	0.00108	9.239	8.519	9.193	9.363	9.162	8.546	8.532	8.48	0.72	NM_001037120
Dr4812.1.S1_s_at	myhz1 /// myhz2 /// myhc4	0.04833	9.074	7.746	9.152	9.034	9.036	8.043	6.662	8.532	1.328	NM_001020485 /// NM_152982 /// XM_703824
Dr4887.1.A1_at	wu:fb22f10	0.00197	10.343	10.937	10.399	10.388	10.242	10.974	10.883	10.954	-0.594	--
Dr4907.1.S1_at	fgg /// LOC566316	0.04856	8.381	7.794	8.466	8.218	8.458	7.811	7.416	8.156	0.587	NM_213054 /// XM_689588
Dr4925.1.S1_x_at	cygb	0.01438	7.068	7.668	7.105	6.998	7.101	7.934	7.406	7.665	-0.6	NM_152952
Dr496.1.S1_at	eng2a	0.00184	6.552	5.351	6.556	6.66	6.439	5.013	5.305	5.734	1.201	NM_131044
Dr4965.1.A1_at	--	0.04905	6.378	7.139	5.806	6.727	6.599	6.923	7.326	7.167	-0.761	--
Dr498.1.S1_at	eng2b	0.00257	7.775	6.491	7.669	7.785	7.872	6.324	6.158	6.99	1.284	NM_131040
Dr5022.1.A1_at	cpsf1	0.00066	9.676	8.867	9.662	9.77	9.595	8.891	8.766	8.944	0.809	XM_688537
Dr5038.1.S1_at	cldna	0.00654	4.444	3.792	4.312	4.366	4.655	3.706	3.712	3.957	0.652	NM_131762
Dr5047.1.A1_at	wu:fc01e05	0.00159	8.047	7.197	8.002	7.979	8.16	7.046	7.421	7.123	0.85	--
Dr5167.1.A1_at	--	0.00325	10.15	9.386	10.332	9.943	10.176	9.539	9.365	9.255	0.764	--

Dr5206.1.S1_at	aldh1a2	4.00E-05	8.387	7.212	8.466	8.392	8.302	7.265	7.192	7.18	1.175	NM_131850
Dr5214.1.S1_at	LOC553298	0.0019	7.665	7.062	7.767	7.591	7.635	7.008	7.099	7.078	0.603	XM_680794
Dr5231.1.S1_at	hist2h2l	0.00985	8.912	9.625	9.008	8.865	8.862	9.603	9.343	9.928	-0.713	NM_200117
Dr5372.9.S1_s_at	her4.1	0.00883	6.227	5.516	5.944	6.362	6.375	5.651	5.56	5.339	0.711	NM_131090
Dr538.1.S1_at	rx1 /// LOC555024 /// LOC567319 /// LOC567374	0.00132	7.566	6.803	7.718	7.519	7.461	6.908	6.779	6.723	0.763	NM_131225 /// XM_690614 /// XM_690667 /// XM_696870
Dr5423.1.A1_at	bat1	0.00412	8.267	8.893	8.241	8.425	8.136	8.96	8.953	8.766	-0.626	NM_200943
Dr5434.1.S3_at	plp1a	0.01029	8.019	7.378	7.904	8.318	7.836	7.397	7.411	7.325	0.641	NM_175576
Dr5438.1.S1_at	arf1l	0.00151	9.909	9.318	9.957	9.934	9.837	9.338	9.319	9.298	0.591	NM_201480
Dr546.1.S1_at	vsx2	0.00026	5.765	4.162	5.759	5.699	5.838	3.955	3.985	4.545	1.603	NM_131462
Dr5464.1.S1_at	riok3	0.00115	8.427	9.327	8.542	8.413	8.326	9.486	9.366	9.129	-0.9	NM_001003614 NM_130920 /// XM_677632 /// XM_701595
Dr5479.1.S1_at	rbp4 /// LOC572824	0.00126	10.775	9.943	10.819	10.842	10.663	9.985	9.763	10.082	0.832	NM_212641 /// XM_680694 /// XM_703016 /// XM_703017
Dr5506.1.A1_at	pa2g4b /// LOC558579	0.00265	11.114	10.521	11.13	11.134	11.077	10.595	10.591	10.378	0.593	---
Dr5510.1.A1_at	wu:fc09b09	0.00019	8.635	7.124	8.591	8.608	8.706	7.086	6.866	7.422	1.511	---
Dr5513.1.S1_at	slc3a2	0.00195	8.754	9.401	8.692	8.719	8.852	9.294	9.5	9.409	-0.647	NM_131601
Dr5531.1.S1_a_at	LOC553371	0.00077	5.945	5.219	6.036	5.899	5.899	5.256	5.162	5.239	0.726	---
Dr5758.1.S1_at	hoxd10a	0.00184	7.833	8.423	7.869	7.855	7.774	8.359	8.504	8.404	-0.59	NM_131166
Dr5772.1.S1_at	hoxa2b	0.00128	7.721	6.915	7.617	7.663	7.883	6.968	6.788	6.989	0.806	NM_131106
Dr578.1.A1_at	tef	0.00229	8.207	7.6	8.161	8.183	8.279	7.627	7.691	7.482	0.607	NM_131400
Dr578.2.S1_a_at	tef	0.00111	7.877	7.225	7.831	7.968	7.832	7.225	7.224	7.227	0.652	NM_131400
Dr582.1.S1_a_at	cx43	0.00042	7.537	6.319	7.449	7.752	7.411	6.132	6.341	6.483	1.218	NM_131038
Dr5846.1.A1_at	atf4	0.00032	10.017	11.101	10.014	9.878	10.158	11.182	10.939	11.181	-1.084	NM_213233
Dr5878.1.A1_at	---	0.00323	7.522	8.348	7.324	7.661	7.581	8.274	8.582	8.189	-0.826	---
Dr5932.1.A1_at	LOC566384	0.00095	7.487	8.233	7.591	7.426	7.443	8.232	8.31	8.156	-0.746	XM_682814 /// XM_703885 /// XM_703886
Dr5962.1.A1_at	nfyA	0.00229	4.875	5.471	4.791	5	4.835	5.438	5.513	5.462	-0.596	XM_679202 /// XM_702327 /// XM_702328 /// XM_702329 /// XM_702330 /// XM_702331 /// XM_702332
Dr6065.1.S1_at	cecr1	0.00145	6.775	7.398	6.737	6.822	6.767	7.312	7.423	7.461	-0.623	NM_001033717
Dr608.1.A1_at	mafB	0.00016	7.599	6.236	7.549	7.57	7.678	6.427	5.995	6.287	1.363	NM_131015
Dr616.1.S1_at	six3a /// LOC573784	0.00235	7.732	8.404	7.72	7.756	7.722	8.215	8.521	8.476	-0.672	NM_131362 /// XM_693508
Dr6278.1.A1_at	---	0.00829	6.927	6.185	6.711	6.942	7.128	5.992	6.131	6.431	0.742	---
Dr642.1.S1_at	myl7	0.005	7.078	6.151	7.161	7.162	6.91	6.085	5.862	6.507	0.927	NM_131329
Dr6453.1.A1_at	LOC573786	0.00704	7.413	6.773	7.5	7.397	7.343	6.597	6.703	7.02	0.64	XM_693540
Dr6460.1.S1_at	zgc:92606	0.0011	7.192	7.993	7.065	7.334	7.176	7.89	8.072	8.018	-0.801	NM_001002707
Dr656.1.S1_at	iclN	0.00286	8.047	7.447	8.034	7.985	8.124	7.38	7.376	7.586	0.6	NM_131424

Dr6680.1.S1_at	zg:c:55587	0.00279	5.978	6.627	6.016	5.955	5.964	6.438	6.72	6.723	-0.649	NM_212723
Dr6814.1.S1_at	fabp3	0.00229	11.354	10.729	11.319	11.414	11.327	10.793	10.585	10.81	0.625	NM_152961
Dr6818.1.S1_at	mpz /// grb10	0.0018	6.839	6.01	6.703	7.015	6.799	6.143	5.845	6.043	0.829	NM_001004287 ///NM_194361
Dr6848.1.A1_at	wu:fd51a05	0.00047	5.267	6.296	5.157	5.453	5.191	6.196	6.427	6.266	-1.029	---
Dr7022.1.A1_at	wu:fd9h01	0.00413	5.994	6.625	6.134	5.947	5.901	6.462	6.708	6.705	-0.631	---
Dr7111.1.S1_a_at	LOC556764	0.00229	7.498	8.153	7.393	7.666	7.437	8.16	8.113	8.186	-0.655	XM_677901 /// XM_701698 /// XM_701699 /// XM_701700 /// XM_701701 /// XM_701702 /// XM_701703 /// XM_701704 /// XM_701705
Dr7118.1.A1_at	zg:c:101644	0.00325	8.371	7.627	8.171	8.456	8.486	7.751	7.678	7.453	0.744	NM_001008643
Dr7118.2.A1_a_at	zg:c:101644	0.00158	8.503	7.76	8.369	8.449	8.692	7.742	7.756	7.782	0.743	NM_001008643
Dr7124.1.S1_at	mtbfd1 /// LOC554706	2.00E-05	9.928	8.666	9.906	9.967	9.91	8.657	8.608	8.733	1.262	NM_199529 /// XMI_696949
Dr7733.1.A1_at	wu:fa20f04	0.02754	6.989	6.351	6.765	6.941	7.261	6.077	6.381	6.596	0.638	---
Dr7737.1.S1_at	LOC554697	0.00737	4.976	6.247	4.806	5.069	5.054	5.961	6.886	5.894	-1.271	XM_697473
Dr77365.1.A1_at	wu:fj13h01	0.02337	6.291	6.939	6.15	6.58	6.142	7.009	7.141	6.666	-0.648	---
Dr77508.1.A1_at	wu:fd08h09	0.03634	6.845	7.509	6.695	7.105	6.733	7.654	7.762	7.111	-0.664	---
Dr7612.1.A1_at	spi1	0.00294	5.465	6.223	5.47	5.407	5.518	6.222	6.447	5.998	-0.758	NM_198062 XM_679747 /// XM_702597
Dr7912.1.A1_at	LOC557677	0.00217	5.325	6.087	5.187	5.246	5.541	6.007	6.154	6.1	-0.762	---
Dr796.1.S1_at	sb:cb492	0.00046	11.014	10.218	11.067	10.979	10.997	10.312	10.199	10.145	0.796	---
Dr796.1.S1_x_at	sb:cb492	0.00026	10.936	10.106	11.016	10.917	10.876	10.123	10.083	10.111	0.83	---
Dr8018.1.S1_at	zg:c:56140	0.01481	7.398	8.013	7.441	7.213	7.539	8.201	7.757	8.082	-0.615	NM_213005
Dr8109.1.S1_at	lcp1	0.00482	5.743	4.623	5.358	6.212	5.66	4.516	4.741	4.611	1.12	NM_131320
Dr8176.1.S1_at	pax5	0.00132	5.594	4.796	5.725	5.473	5.584	4.879	4.643	4.865	0.798	---
Dr8208.1.S1_at	lft1	0.01026	6.705	6.023	6.75	6.813	6.552	6.317	5.885	5.866	0.682	NM_130960
Dr8209.1.S2_at	foxo5	0.0064	8.49	7.764	8.418	8.779	8.272	7.764	7.844	7.684	0.726	NM_131085
Dr8328.1.S1_at	hand2	0.00099	8.308	9.073	8.283	8.243	8.398	9.048	9.19	8.982	-0.765	NM_131626
Dr8379.1.S1_at	LOC573558	7.00E-05	7.136	5.281	7.332	7.171	6.904	5.301	5.051	5.492	1.855	XM_689827
Dr8497.1.A1_at	LOC573237	0.00196	7.337	7.934	7.365	7.396	7.251	7.884	8.008	7.909	-0.597	XM_686280
Dr879.1.S1_at	prmt1	0.01301	11.742	11.129	11.785	11.767	11.676	11.293	11.269	10.824	0.613	NM_200650
Dr9236.2.A1_at	LOC559906	0.00222	6.329	5.53	6.366	6.309	6.314	5.776	5.349	5.463	0.799	XM_683302 NM_001001815 ///XM_682622
Dr9288.1.S1_at	wu:fc21h08 /// LOC572929	0.00069	6.493	7.249	6.578	6.483	6.418	7.327	7.216	7.206	-0.756	---
Dr9353.1.S1_at	zg:c:55733	0.00229	8.452	7.747	8.449	8.55	8.357	7.766	7.583	7.89	0.705	NM_200993
Dr9457.1.A1_at	egln3	0.03723	6.185	7.72	5.59	6.222	6.745	6.775	8.495	7.892	-1.535	NM_213310
Dr956.1.S1_at	zg:c:92631	0.01214	6.869	8.005	6.828	7.292	6.486	7.605	8.405	8.005	-1.136	NM_001002695
Dr9654.1.A1_at	ppia	0.00042	7.681	8.713	7.787	7.636	7.62	8.543	8.711	8.885	-1.032	NM_212758
Dr967.1.S1_at	mmp9	0.0017	6.205	6.87	6.196	6.279	6.142	6.843	6.993	6.775	-0.665	NM_213123
Dr9848.1.A1_at	wu:fk60g08	0.00916	6.453	7.161	6.308	6.358	6.693	7.07	7.015	7.398	-0.708	---
Dr9953.1.S1_at	LOC569927	0.00069	8.742	9.875	8.543	9.031	8.652	9.858	9.941	9.826	-1.133	XM_693341

DrAffx.1.2.S1_at	bhik	0.00278	7.93	7.051	7.842	7.998	7.951	7.087	7.304	6.763	0.879	NM_131803
DrAffx.1.52.S1_at	a2bp1l	0.01289	8.969	9.595	9.008	8.854	9.044	9.603	9.333	9.85	-0.626	NM_214775
DrAffx.2.105.S1_at	--	0.01408	7.945	7.335	7.638	8.066	8.132	7.369	7.371	7.266	0.61	--
DrAffx.2.59.S1_at	LOC555198	0.01264	7.082	6.317	7.366	6.702	7.176	6.399	6.185	6.368	0.765	XM_677650
DrAffx.2.62.S1_at	cyp26b1	0.0011	6.887	6.056	6.97	6.719	6.973	6.16	6.033	5.974	0.831	NM_212666