

Pax6 is essential for lens fiber cell differentiation

Ohad Shaham¹, April N. Smith², Michael L. Robinson³, Makoto M. Taketo⁴, Richard A. Lang² and Ruth Ashery-Padan^{1,*}

The developing ocular lens provides an excellent model system with which to study the intrinsic and extrinsic cues governing cell differentiation. Although the transcription factors Pax6 and Sox2 have been shown to be essential for lens induction, their later roles during lens fiber differentiation remain largely unknown. Using Cre/loxP mutagenesis, we somatically inactivated Pax6 and Sox2 in the developing mouse lens during differentiation of the secondary lens fibers and explored the regulatory interactions of these two intrinsic factors with the canonical Wnt pathway. Analysis of the Pax6-deficient lenses revealed a requirement for Pax6 in cell cycle exit and differentiation into lens fiber cells. In addition, Pax6 disruption led to apoptosis of lens epithelial cells. We show that Pax6 regulates the Wnt antagonist Sfrp2 in the lens, and that Sox2 expression is upregulated in the Pax6-deficient lenses. However, our study demonstrates that the failure of differentiation following loss of Pax6 is independent of β -catenin signaling or Sox2 activity. This study reveals that Pax6 is pivotal for initiation of the lens fiber differentiation program in the mammalian eye.

KEY WORDS: Pax6, Sox2, Lens, Crystallin, Wnt, β -catenin, Mouse

INTRODUCTION

Lens development is a complex process in which a single epithelial layer undergoes several stages of competence, induction and differentiation, ultimately forming a highly specialized organ (Grainger et al., 1997; Lovicu and Robinson, 2004; Ogino and Yasuda, 2000). The vertebrate lens comprises only two types of cells: an anterior lens epithelium (LE) and the derived lens fiber cells (LFCs). This, along with its morphological isolation from surrounding tissues, makes the lens an ideal model for the study of tissue growth and differentiation (Bhat, 2001; Lovicu and Robinson, 2004).

The transcription factor (TF) Pax6 is essential for eye development in vertebrates and invertebrates (Ashery-Padan and Gruss, 2001; Gehring, 1996; Grindley et al., 1995; Hogan et al., 1988; Walther et al., 1991). Interestingly, normal development of the mammalian eye is dependent on normal *Pax6* dosage, as heterozygotes suffer from pan-ocular disorders such as aniridia in humans and *Small eye (Sey)* in mice (Glaser et al., 1994; Glaser et al., 1990).

Pax6 is expressed throughout all stages of lens development, except in terminally differentiated LFCs. During early organogenesis, Pax6 is detected in both the lens-inducing optic vesicles, and in the lens-forming surface ectoderm (SE) (Walther et al., 1991). Expression of Pax6 in the SE is required to render it competent for induction into a lens (Fujiwara et al., 1994; Quinn et al., 1996). Inductive signals from the optic vesicle (OV) trigger a thickening of the SE known as the lens placode (LP), which is intrinsically dependent on Pax6 expression (Ashery-Padan et al., 2000). Between embryonic day (E) 10 and E11, the LP invaginates and detaches from the SE to form the lens vesicle. The anterior cells

of the lens vesicle remain as the undifferentiated LE and retain high expression of Pax6. By contrast, the posterior cells elongate and differentiate into primary LFCs and lose Pax6 expression. In the fetal and postnatal stages of development, at the transitional zone between LE and LFCs, the equatorial epithelial cells undergo proliferation, cell cycle exit, migration and elongation, and finally mature into secondary fiber cells deposited around a nucleus of primary LFCs, all this in a distinct spatial order (Bassnett and Beebe, 1992; Beebe et al., 1982; Rafferty and Rafferty, 1981). Although Pax6 expression is maintained in the LE and in the equatorial transitional zone, its role in maintaining an epithelial phenotype or in LFC differentiation remains largely unknown.

LFCs express high levels of lens-specific crystallins (Bassnett and Beebe, 1992; Beebe et al., 1982; Beebe and Piatigorsky, 1976; Rafferty and Rafferty, 1981). Crystallins have distinct expression patterns, making them the definitive markers of lens differentiation. The crystallins of the $\beta\gamma$ supergroup are highly expressed in developing LFCs and are used as markers for LFC differentiation, whereas α -crystallins are differentially expressed between LE and LFCs (Robinson and Overbeek, 1996). Pax6 activation and its binding to promoters of crystallin genes have been studied extensively (Cvekl et al., 2004; Duncan et al., 1996; Muta et al., 2002; Wawrousek et al., 1990; Yang et al., 2006). Based on its expression pattern and regulation of crystallins, it has been suggested that *Pax6* plays a dual role, acting as both an activator and a repressor of crystallin expression (Duncan et al., 1998). Furthermore, in vitro and chromatin-binding assays indicate that Pax6 co-operates with the Sox2 TF on specific crystallin enhancers during early stages of lens formation (Kamachi et al., 2001; Kondoh et al., 2004). However, the roles of Pax6 and Sox2 in the control of crystallin gene expression during LFC differentiation have not been studied in vivo.

Along with TFs, many growth factors and signaling pathways have been reported to be involved in LFC differentiation (Lovicu and McAvoy, 2005). Most notably, FGF signaling is differentially activated along the anterior-posterior axis of the lens, with increased activity at the posterior side of the lens equator (de Jongh et al., 1997; Garcia et al., 2005; Robinson, 2006). By contrast, Wnt signaling is believed to be antagonistic to LFC differentiation. Wnt receptors, co-receptors and downstream proteins are expressed in the LE (Ang

¹Department of Human Molecular Genetics and Biochemistry, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv 69978, Israel. ²Department of Developmental Biology, Children's Hospital Research Foundation, Cincinnati, OH 45229, USA.

³Department of Zoology, Miami University, Oxford, OH 45056, USA. ⁴Department of Pharmacology, Graduate School of Medicine, Kyoto University, Yoshida-Konoé-cho, Sakyo, Kyoto 606-8501, Japan.

*Author for correspondence (e-mail: ruthash@post.tau.ac.il)

et al., 2004; Chen et al., 2004; Chen et al., 2006). When the Wnt co-receptor *Lrp6* was deleted in mice, aberrant LFCs appeared in the anterior pole of the lens (Stump et al., 2003). Upon inactivation of the canonical Wnt effector β -catenin, LE markers, proliferation and differentiation were disrupted (Cain et al., 2008). These findings suggest a role for Wnt signaling in LE cell fate.

Owing to the severe ocular phenotype of *Pax6* mutants, the later developmental roles of *Pax6* could only be extrapolated from in vitro research. Herein, we introduce the first in vivo loss-of-function model of *Pax6* and its presumed transcriptional partner *Sox2*. We show that the loss of *Pax6* prevents LFC differentiation and results in cell death and in an increase in *Sox2*. However, conditional deletion of *Sox2* reveals that it is dispensable for LFC differentiation. Furthermore, overexpression of β -catenin results in a differentiation failure that is similar to, but independent of, that observed following *Pax6* loss. These findings place *Pax6* upstream in the cascade of events leading to the differentiation of LE into lens fibers in mammals.

MATERIALS AND METHODS

Mouse lines

Mouse lines employed in this study were: *Pax6^{lox}* (Ashery-Padan et al., 2000), *Mlr10* (Zhao et al., 2004), *Catnb^{lox(ex3)}* (Harada et al., 1999) and *BATlacZ* (Nakaya et al., 2005) and are described in Fig. S1 in the supplementary material. The *Sox2^{loxP}* line (see Fig. 7A) contains two loxP sites inserted around the single exon of the murine *Sox2* gene using conventional gene-targeting methods (Joyner, 1995). In the gene-targeting vector, *loxP-frt-pMC1neopA-frt*, the *neo* gene is flanked by *frt* sites. Flp recombinase activity within the *B6.SJL-Tg(CTFLPe)9205Dym/J* mouse line (Rodriguez et al., 2000) was used to delete the *neo* selection cassette.

Histology, immunofluorescence analysis, BrdU, TUNEL and X-Gal assays

Paraffin sections (10 μ m) were stained with Hematoxylin and Eosin (H&E) using standard procedures. Immunofluorescence analysis was performed on paraffin sections as previously described (Ashery-Padan et al., 2000) using the following primary antibodies: rabbit anti-Pax6 (1:1000, Chemicon), mouse anti-Ap2 α (1:50, Santa Cruz), rabbit anti-cleaved caspase 3 (1:100, Cell Signaling), goat anti- α A-crystallin (1:1000, Santa Cruz), goat anti- α B-crystallin (1:100, Santa Cruz), rabbit anti- β B1-crystallin (1:250, Santa Cruz), rabbit anti- γ F-crystallin (1:50, Santa Cruz), rabbit anti-cyclin D1 (1:250, Thermo Scientific), rat anti-Ki67 (1:100, Dako), goat anti-p57^{Kip2} (1:100, Santa Cruz), rabbit anti-Prox1 (1:50, Acris) and rabbit anti-Sox2 (1:500, Chemicon). Secondary antibodies were conjugated to RRX or Cy2 (Jackson ImmunoResearch). Nuclei were visualized with DAPI (0.1 μ g/ml, Sigma). For cell cycle quantification, BrdU (10 μ l/g of 14 mg/ml) was injected 1.5 hours before sacrifice. Slides were stained with anti-phosphohistone H3 (1:500, Santa Cruz), fixed for 10 minutes in 4% paraformaldehyde, then stained with mouse anti-BrdU (1:100, Chemicon) as described (Marquardt et al., 2001). Five eyes were used from each genotype, and the percentage of marker-positive nuclei was calculated from total DAPI-positive nuclei. A two-tailed Student's *t*-test was used for statistical analysis. X-Gal staining on cryosections was performed as described (Liu et al., 2003).

Confocal quantification of Sox2 expression

Images of E14.5 lenses were taken using a confocal microscope CLSM410 (Zeiss) and the signal was measured within the linear range using the Range Indicator application (Zeiss LSM Imager). Five nuclei each from the extreme anterior, equator and posterior of lens sections were measured for intensity (pixel values 0-255) and divided by the retinal nucleus intensity for the same section, using ImageJ software (NIH).

In situ hybridization (ISH)

ISH was performed using DIG-labeled RNA probes (Yaron et al., 2006). The *Prox1* probe was produced from a 947 bp PCR fragment (forward, 5'-CAGATGCCTAGTTCACAGACC-3'; reverse, 5'-AGAGCGTTGCA-

ATCTCTACTCG-3'). Other ISH probes used were: *Cryaa*, *Cryab* and *cMaf* (Robinson and Overbeek, 1996), *Sox1*, *Sfrp2* (Leimeister et al., 1998) and *Six3* (Oliver et al., 1995). All analyses presented in this study were conducted on at least five eyes of each genotype, from at least two different litters.

RESULTS

Somatic mutation of Pax6 in the lens results in small eyes due to lens defects

To study the role of *Pax6* in the lens after the lens vesicle stage, we employed the *Mlr10* transgene (Zhao et al., 2004) and the *Pax6^{lox}* allele (Ashery-Padan et al., 2000) and established *Pax6^{lox/lox};Mlr10* somatic mutants. *Pax6^{lox/lox}* littermates were used as controls. *Pax6^{lox/lox};Mlr10* eyes were significantly smaller than those of controls (Fig. 1B,D,F; 65% of circumference, $P < 0.001$). This reduction in size was attributed to the decrease in lens tissue, which appeared opaque and shapeless (Fig. 1D). In a previous study, *Pax6* was shown to have lens-autonomous dosage requirements at the LP stage (Davis-Silberman et al., 2005). To determine whether there is haploinsufficiency at later stages of lens development, we investigated the phenotype of the heterozygous *Pax6^{lox/+};Mlr10* littermates. We did not identify any differences in lens weight, size or opacity between the *Pax6^{lox/+};Mlr10* animals and controls (Fig. 1E,F). This result implies that a diploid dose of *Pax6* is not required after the lens structure has formed.

Somatic inactivation of Pax6 by E14.5 leads to failure in LFC differentiation

To characterize the morphological defects of *Pax6^{lox/lox};Mlr10* mice and to determine the onset of *Pax6* inactivation, we conducted a histological analysis of eyes from embryonic and postnatal stages and monitored *Pax6* loss by immunostaining (Fig. 2) and by activity of human alkaline phosphatase (hAP) from the *Z/AP* reporter (see Fig. S2 in the supplementary material) (Lobe et al., 1999). In controls, *Pax6* expression was high in the anterior LE and equator of the lens, and diminished as the cells underwent differentiation (Fig. 2K-M). At E13.5, prior to the loss of *Pax6* protein from the LE (Fig. 2K,N), the *Pax6^{lox/lox};Mlr10* and control lenses appeared similar in size (Fig. 2A,F). After *Pax6* loss at E14.5 (Fig. 2O), *Pax6^{lox/lox};Mlr10* lenses were slightly more elongated, and a few small nucleated cells were detected in the posterior part of the lens (Fig. 2G, arrow). The cornea of E14.5 mutants (Fig. 2G, double arrow) was thicker than in the controls (Fig. 2B), which suggests failure of the lens to induce mesenchymal condensation (Sevel and Isaacs, 1988). By E15.5, the mutant lenses appeared significantly smaller than the controls (Fig. 2H; 76% of circumference, $n=8$, $P < 0.002$). In E15.5 controls, the nuclei of the equatorial transitional zone, in which LE cells undergo differentiation, were organized in a characteristic bow pattern (Fig. 2C,D). By contrast, in *Pax6^{lox/lox};Mlr10* lenses, transitional zone cells were disorganized (Fig. 2H,I). At later stages, the *Pax6^{lox/lox};Mlr10* lenses remained significantly smaller than controls, as lens fiber formation seemed to be arrested from \sim E14.5 (Fig. 2; see Fig. S3 in the supplementary material). By P4, the *Pax6^{lox/lox};Mlr10* lens seemed to be mostly composed of epithelial cells surrounding a few fiber cells (Fig. 2J). In the adult [postnatal day (P) 30], only remnants of lens tissue were detected in the mutant (Fig. 1D). These morphological defects suggest that *Pax6*-deficient LE cells fail to differentiate and instead accumulate at the lens equator and in the posterior lens, and that the LFCs detected in the *Pax6^{lox/lox};Mlr10* lenses probably originate prior to *Pax6* loss.




Fig. 1. Microphthalmia in *Pax6^{lox/lox};Mir10* mice.

(A-E) Eyes (A,B) and isolated lenses (C-E) from P30 control (A,C), *Pax6^{lox/lox};Mir10* (B,D) and *Pax6^{lox/+};Mir10* (E) mice. (F) Quantification of eye circumference (light-gray bars), lens circumference (dark-gray bars) and dry mass (black bars) of controls, *Pax6^{lox/+};Mir10* and *Pax6^{lox/lox};Mir10* mice. Lens size and mass could not be measured in *Pax6^{lox/lox};Mir10* mice (D,F). ** $P=0.004$. Scale bars: 500 μm .

Pax6-deficient LE cells fail to exit the cell cycle at the lens equator

The first step in LFC differentiation is cell cycle exit (Rafferty and Rafferty, 1981). Pax6 is expressed in proliferating LE cells, as it co-localized with Ki67, a marker of actively proliferating cells (Fig. 3A) (Endl et al., 1997). However, Pax6 was also expressed after the cells had undergone cell cycle exit (Fig. 3A', arrow). This pattern of expression suggests that either Pax6 is involved in maintaining the proliferation capacity of the anterior LE, or that it is required for cell cycle exit in the LE of the equatorial zone. To distinguish between these possibilities, we characterized the distribution of Ki67 in *Pax6^{lox/lox};Mir10* embryos. The loss of Pax6 was accompanied by a change in the distribution of Ki67. In *Pax6^{lox/lox};Mir10* lenses, Ki67⁺ cells were detected posterior to the lens equator, a region which is normally devoid of proliferating cells (Fig. 3A,E). To further determine the cell cycle stage of Pax6-deficient cells at E14.5, we quantified the percentage of cells in the S and M phases using BrdU incorporation and phosphorylated histone H3 (PH3) immunostaining, respectively. Both markers were only detected anterior to the equator in control lenses (Fig. 3B), whereas in *Pax6^{lox/lox};Mir10* mutants, proliferating cells were abundant in the transitional zone and in the posterior lens (Fig. 3F' and arrowheads). In these regions, BrdU was detected in $46.2 \pm 3.5\%$ (\pm s.d.) of the nuclei, and PH3 was detected in $32.8 \pm 6.2\%$ of the nuclei. We therefore concluded that Pax6 is required for the cell cycle exit of LE cells at the lens equator.

To determine whether Pax6 loss alters the cell cycle dynamics in the anterior LE itself, we quantitatively analyzed BrdU⁺ and PH3⁺ cells in the LE of control and Pax6-deficient lenses (Fig. 3C). A significant increase in the BrdU incorporation index was observed in the LE following Pax6 loss ($70.7 \pm 5.15\%$), as compared with the controls ($52.8 \pm 1.4\%$, $P < 0.001$). The proportion of PH3⁺ cells was similar between the genotypes ($14.3 \pm 6.2\%$ in *Pax6^{lox/lox};Mir10* and $14.7 \pm 4.0\%$ in *Pax6^{lox/lox}*). This suggests a prolonged S phase in the Pax6-deficient LE, which is reminiscent of the phenotype reported in Pax6-deficient cerebral cortex (Estivill-Torrus et al., 2002).

Apoptosis in the Pax6-deficient lens

Although cells in *Pax6^{lox/lox};Mir10* lenses continued to proliferate, the overall size of the lens was reduced (Fig. 2). To test whether this tissue loss was due to apoptosis, we performed TUNEL analysis, which demonstrated an increase in apoptotic cells in the *Pax6^{lox/lox};Mir10* lenses (not shown). To perform a quantitative analysis, we detected the cleaved form of caspase 3 (cCas3, Fig. 3D,G). However, the number of cCas3⁺ cells in the *Pax6^{lox/lox};Mir10* lenses was low (2.6 ± 2.0 per section, $n=8$), suggesting that apoptosis is only partially responsible for the significant reduction in lens size, which is instead primarily due to the arrest in LFC differentiation. Interestingly, cCas3⁺ cells were never BrdU⁺ (not shown). Thus, the longer S phase observed in the Pax6⁻ LE is probably not an immediate trigger of apoptosis.

Pax6 is not required for the regulation of crystallin genes at late stages of lens development

The α A-crystallin (*Cryaa*) promoter has been shown to bind and to be activated by Pax6 in vitro (Cvekl et al., 1995; Yang and Cvekl, 2005; Yang et al., 2006). Accordingly, Pax6 was found to be required in vivo for the onset of *Cryaa* expression during early stages of lens development (Ashery-Padan et al., 2000; Cvekl et al., 1995). To examine whether Pax6 regulates crystallin expression during secondary LFC differentiation, we characterized the distribution of crystallin transcripts and proteins in the Pax6-deficient lenses. Cryaa protein was detected in both the LE and LFCs of control lenses, with elevated expression in the latter (Robinson and Overbeek, 1996) (Fig. 4A). Intriguingly, in *Pax6^{lox/lox};Mir10* lenses, Cryaa protein was maintained in the LE and in the posterior aberrant cells, similar to its expression in controls (Fig. 4F; see also Fig. S4 in the supplementary material). As crystallins are ultra-stable proteins (Jaenicke, 1996), Cryaa might be detected because of its low turnover, rather than continued expression. We therefore examined *Cryaa* transcripts by in situ hybridization (ISH). *Cryaa* transcripts were detected in the LE




Fig. 2. Pax6 ablation in the lens epithelium coincides with failure of lens fiber cell differentiation and morphological defects in the lens. (A-J) Hematoxylin and Eosin (H&E) staining of control (A-E) and *Pax6^{lox/lox};Mlr10* (F-J) eyes in E13.5 (A,F), E14.5 (B,G), E15.5 (C,D,H,I) and P4 (E,J) mice. D and I are higher magnifications of the boxed regions from C and H, respectively. Arrows mark posterior nucleated cells. Double arrows demonstrate the thickness of presumptive cornea of controls (A,B) and mutants (F,G). Asterisks mark fixation artifacts. (K-P) Pax6 protein in *Pax6^{lox/lox}* controls (K-M) and *Pax6^{lox/lox};Mlr10* mutants (N-P) at E13.5 (K,N), E14.5 (L,O) and E15.5 (M,P). White arrows indicate a few cells that retain some Pax6 activity owing to the mosaic nature of Cre activity. co, cornea; re, retina. Scale bars: 100 μ m.

of E14.5, E15.5 and E18.5 *Pax6^{lox/lox};Mlr10* lenses in a similar distribution to that of Cryaa protein (Fig. 4B,G), confirming that *Pax6* is not required for the low-level *Cryaa* expression in the LE at these stages (Fig. 4G and data not shown).

Unlike *Cryaa*, α B-crystallin (*Cryab*) is strongly expressed in the LE of the E12.5-15.5 developing lens and is reduced in LFCs, overlapping with *Pax6* expression (Robinson and Overbeek, 1996) (Fig. 4C). This, together with the results of extensive in vitro research (Gopal-Srivastava et al., 1996; Yang et al., 2004), suggest that *Pax6* is an important regulator of *Cryab* expression. However, *Cryab* expression was maintained in both control and *Pax6^{lox/lox};Mlr10* lenses, with high expression in the LE (Fig. 4C,H). This suggests that *Pax6* is not required for *Cryab* expression during the later stages of lens development.

β - and γ -crystallins are expressed throughout lens development exclusively in LFCs. Specifically, β B1-crystallin (*Crybb1*) expression is initiated precisely when lens fibers begin to elongate (Brahma, 1988; Duncan et al., 1996), making it an ideal marker for LFC differentiation. To examine LFC differentiation in *Pax6^{lox/lox};Mlr10* mutants, an antibody against *Crybb1* that identifies most β -crystallin (Cryb) proteins was employed. Cryb was detected in the LFCs of control and *Pax6^{lox/lox};Mlr10* lenses in a similar pattern (Fig. 4D,I). *Cryb* was not detected in the LE, transitional zone and aberrant posterior cells of the *Pax6*-deficient lenses (Fig. 4I), confirming the undifferentiated state of these cells. Previous in vitro studies suggested that the high level of Pax6 in the LE suppresses *Crybb1* (Duncan et al., 1996). However, removal of *Pax6* from the *Pax6^{lox/lox};Mlr10* lenses was not sufficient to induce upregulation of *Cryb* in the LE in vivo (Fig. 4I).

Similar to *Cryb*, γ -crystallins (*Cryg*) are expressed in mature LFCs and are possible targets for Pax6 regulation based on in vitro studies (Kralova et al., 2002; Yang et al., 2004). *Cryg* was not detected in the LE, transitional zone or aberrant posterior cells of *Pax6^{lox/lox};Mlr10* lenses (Fig. 4J).

Taken together, these results demonstrate that Pax6 is not required for the expression of α -crystallins or for the maintenance of an undifferentiated fate in the LE by inhibiting LFC-specific crystallins. Importantly, cells at the equator and on the posterior side of *Pax6^{lox/lox};Mlr10* lenses do not express any crystallin LFC marker (Fig. 4I,J, arrowheads). Therefore, *Pax6* is primarily required for the normal differentiation of LFCs and this activity does not depend on its regulation of crystallin expression.

Pax6 requirement for LFC differentiation is not mediated through Prox1, Sox1 or cMaf

Several TFs have been shown to be essential for LFC differentiation in vivo, namely *Prox1*, *Sox1* and *cMaf* (*Maf* – Mouse Genome Informatics). To determine whether the lack of LFC differentiation observed in the *Pax6^{lox/lox};Mlr10* mice is mediated through one of these TFs, we characterized their expression in Pax6-deficient *Pax6^{lox/lox};Mlr10* lenses.

Prox1 is essential for the elongation of primary LFCs, exit from the cell cycle and the expression of several γ -crystallins (Wigle et al., 1999). *Prox1* expression in the LP is dependent on Pax6 activity (Ashery-Padan et al., 2000). At E14.5, *Prox1* transcripts were detected in both control and *Pax6^{lox/lox};Mlr10* lenses (Fig. 5A,D). As *Prox1* protein is differentially localized during lens development (Duncan et al., 2002), we examined its spatial distribution at E14.5 by immunolabeling. In both control and *Pax6^{lox/lox};Mlr10* lenses, *Prox1* protein was detected in the nuclei and cytoplasm of LE cells, whereas in the equator and in differentiating LFCs it was mainly nuclear (Fig. 5B,E). The level of *Prox1* expression varied among *Pax6^{lox/lox};Mlr10* cells (Fig. 5E,E', asterisk). However, most nuclei maintained *Prox1* expression at E14.5 and during later stages (E15.5; data not shown). In accordance with the maintenance of *Prox1* in *Pax6^{lox/lox};Mlr10* lenses, expression of its downstream targets – cell cycle inhibitory genes *p57^{Kip2}* (*Cdkn1c*) and *p27^{Kip1}* (*Cdkn1b*) (Wigle et al., 1999) – was detected in the equatorial region of both control and Pax6-




Fig. 3. Pax6-deficient lens epithelium fails to exit the cell cycle at the lens equator and undergoes apoptosis. (A-B',E-F') Antibody labeling of control (A,B) and Pax6^{lox/lox};Mlr10 (E,F) lenses from E14.5 mouse embryos. (A,E) Expression of Pax6 (red) and Ki67 (green) with co-expression in the lens epithelium (LE) of Ki67 and Pax6 (A, yellow). Some Ki67⁻ cells express Pax6 (A', arrowhead). In Pax6^{lox/lox};Mlr10, Pax6⁻ Ki67⁺ cells are detected in the transitional zone (E') and in the posterior lens (E, arrowhead). (B,F) Phosphohistone H3 (PH3, green) and BrdU (B,F, red) were detected in the Pax6^{lox/lox} LE up to the lens equator (arrow, B) but also posterior to the lens equator in the Pax6^{lox/lox};Mlr10 lenses (F,F') and in the posterior lens (F, arrowheads). (C) Quantitative analysis reveals a significant increase in the percentage of BrdU⁺ cells in Pax6^{lox/lox};Mlr10 lenses (70.7 ± 5.1% s.d.) as compared with the control LE (52.8 ± 1.4%, **P < 0.001). The percentage of PH3⁺ cells is not altered in the Pax6⁻ LE (control 14.3 ± 6.2%; mutant 14.7 ± 4.1%). (D,G,G') Cleaved caspase 3 (cCas3)-positive cells detected in the LE of Pax6^{lox/lox};Mlr10 lenses (G,G', red, arrowhead) but not in the control E14.5 mouse embryo (D). Counterstaining is with DAPI (blue). co, cornea; re, retina. Scale bar: 100 μm.

deficient lenses (Fig. 5C,F and data not shown). These results show that during secondary LFC differentiation, Pax6 does not regulate the expression of Prox1 or of its cell cycle inhibiting targets, but is still essential for cell cycle arrest.

TFs of the Sox family are expressed during, and are involved in, lens development (Kamachi et al., 1995; Kamachi et al., 1998). One of these, Sox1, has been shown to be essential for complete elongation of LFCs and for expression of γ -crystallins (Nishiguchi et al., 1998). Sox1 was expressed in all lens cells at E14.5-15.5, with a marked increase in differentiating LFCs (Nishiguchi et al., 1998) (Fig. 5G). The same expression pattern was observed in Pax6^{lox/lox};Mlr10 lenses, indicating that Pax6 is not crucial for Sox1 expression (Fig. 5I).

Finally, cMaf is a lens-specific member of the large Maf gene family. cMaf has been shown to be essential for LFC elongation and γ -crystallin expression (Kawauchi et al., 1999; Ring et al., 2000; Yoshida et al., 2001; Yoshida and Yasuda, 2002). In Pax6^{lox/lox};Mlr10

lenses, the expression of cMaf was similar to in controls. cMaf transcripts were detected throughout the lens, with elevated expression at the lens equator (Sakai et al., 1997) (Fig. 5H,J).

Taken together, the apparently normal upregulation of Sox1 and cMaf at the lens equator, as well as the normal distribution of Prox1 protein, demonstrate that despite Pax6 loss, the cells in the transitional zone are able to respond to extracellular signals and activate some differentiation markers. However, even with the activation of these factors, execution of the lens fiber differentiation program requires Pax6.

Pax6 negatively regulates Sox2 in the equatorial zone of the embryonic lens

The Sox2 TF is expressed in the developing lens and has been implicated to function with Pax6 in initiating crystallin expression (Kamachi et al., 1998; Kamachi et al., 2001; Kondoh et al., 2004; Stevanovic et al., 1994; Yang et al., 2004). Furthermore, direct




Fig. 4. Pax6 is not essential for maintaining the expression of α -crystallins or for the downregulation of β / γ -crystallins during LFC differentiation. Control (A-E) and Pax6^{lox/lox};Mlr10 mutant (F-J) E14.5 mouse lenses labeled with antibodies against α A-crystallin (Cryaa, A,F, red), β -crystallins (D,I, green) and γ -crystallins (E,J, green), or subjected to ISH with probes against α A-crystallin (B,G) and α B-crystallin (C,H). Arrows indicate the LE, arrowheads to Pax6⁻ cells accumulating in the posterior lens. co, cornea; re, retina. Scale bar: 100 μm.




Fig. 5. Intrinsic requirement for Pax6 in LFC differentiation is not mediated through Prox1, Sox1 or cMaf. (A-J) Expression patterns in E14.5 (A,B,D,E,G,I) or E15.5 (C,F,H,J), control (A-C,G,H) and *Pax6^{lox/lox};Mlr10* (D-F,I,J) mouse eyes, showing *Prox1* transcript (A,D) and protein (B,E, boxed regions magnified in B',B'',E',E'', red), p57^{Kip2} protein (C,F, red), and *Sox1* (G,I) and *cMaf* (H,J) transcripts. Asterisk in E'' marks a patch of low-level *Prox1* expression. Counterstaining is with DAPI (B-C,E-F, blue). co, cornea; eq, lens equator; LE, lens epithelium; re, retina. Scale bars: 100 μ m.

regulation of *Sox2* by Pax6 has been demonstrated in neural progenitor cells (Wen et al., 2008). The role of *Sox2* and whether it interacts with *Pax6* during later stages of lens development are unknown. We therefore characterized the expression of *Sox2* following *Pax6* loss. We utilized Ap2 α (Tcfap2a), a TF that is expressed in the anterior LE and is essential for early lens development, as a marker for the anterior LE (Pontoriero et al., 2008; West-Mays et al., 1999). Double immunolabeling for Ap2 α and *Sox2* revealed that in the control lens at E14.5, Ap2 α is co-expressed with *Sox2* in the anterior LE, whereas in the transitional zone only *Sox2* was detected (Fig. 6A). In the conditional mutant, Ap2 α was restricted to a small population of the most anterior cells of the LE. By contrast, *Sox2*⁺ cells were detected in a much wider population of cells at the *Pax6^{lox/lox};Mlr10* equator and at a high level of expression, similar to that in the retina. Ectopic cells at the lens posterior were also intensely *Sox2* positive (Fig. 6B). *Sox2* expression was quantified by confocal microscopy. In controls, only a low level of *Sox2* was observed in the anterior LE, the same as in the lens equator and about half of that in the retina (Fig. 6A). Anterior LE cells of the *Pax6^{lox/lox};Mlr10* had expression levels comparable to those of controls, but equatorial and posterior cells showed a 2.2-fold increase in expression ($P=0.0001$), attaining levels greater than in the retina (Fig. 6B,C). Therefore, following *Pax6* ablation, cells of the lens equator fail to differentiate into LFCs, increase at the expense of anterior Ap2 α ⁺ LE, express high levels of *Sox2* and expand into the posterior lens.

The differentiation failure and proliferation of aberrant LE are not mediated through *Sox2*

Sox2 is known to be involved in the determination of stem cell fate and in the proliferation of neural stem cells (Episkopou, 2005). To examine the role of *Sox2* in the lens and to determine whether the significant increase in *Sox2* expression in *Pax6^{lox/lox};Mlr10* lenses mediates the observed differentiation failure, we established the

Sox2^{lox} allele, which includes two loxP sequences flanking the single exon of the murine *Sox2* gene (Fig. 7A). This allele was employed in combination with *Mlr10* to inactivate either *Sox2* alone (*Sox2^{lox/lox};Mlr10*) or *Sox2* together with *Pax6* (*Sox2^{lox/lox};Pax6^{lox/lox};Mlr10*). The *Sox2^{lox/lox};Mlr10* embryos and adult mice did not exhibit any abnormal ocular phenotypes (not shown). This is in agreement with an apparent reduction in *Sox2* expression at E12.5-15.5 (Nishiguchi et al., 1998), suggesting that the low-level expression of *Sox2* in E14.5 lenses is not essential for lens development.

In the *Sox2^{lox/lox};Pax6^{lox/lox};Mlr10* double somatic mutants, both *Pax6* and *Sox2* are deleted exclusively in the lens (Fig. 7). Accordingly, *Sox2* protein was not detected in the *Sox2^{lox/lox};Pax6^{lox/lox};Mlr10* lens, but was preserved in the adjacent optic cup, where Cre is not active (Fig. 7I). Despite the obvious loss of *Sox2*, the ocular phenotype of the double somatic mutant was strikingly similar to that of *Pax6^{lox/lox};Mlr10* mutants. The *Sox2^{lox/lox};Pax6^{lox/lox};Mlr10* lenses were smaller than controls and epithelial cells accumulated posterior to the lens equator (Fig. 7H). The anterior LE, as identified by Ap2 α expression, was reduced in size (Fig. 7I). Moreover, similar to the phenotype of *Pax6^{lox/lox};Mlr10*, in *Sox2^{lox/lox};Pax6^{lox/lox};Mlr10* lenses α A-crystallin protein and transcripts were strongly expressed in the LFCs and weakly in the LE and in the aberrant posterior cells (Fig. 7K and data not shown), whereas β -crystallin was absent from cells of the lens equator and from the aberrant posterior cells (Fig. 7L). Failure of cell cycle exit was also evident in *Sox2^{lox/lox};Pax6^{lox/lox};Mlr10* lenses (Fig. 7J,J'). Finally, some *Sox2^{lox/lox};Pax6^{lox/lox};Mlr10* LE cells underwent apoptosis, as demonstrated by cCas3 immunostaining (Fig. 7M,M'). Therefore, when *Sox2* overexpression is prevented, *Pax6*-null LE cells undergo the same LFC differentiation failure and cell death as observed in lenses that overexpress *Sox2*. Thus, the LFC differentiation failure observed in *Pax6^{lox/lox};Mlr10* mutants is independent of *Sox2*.




Fig. 6. Pax6 downregulates Sox2 in the lens equator. (A,B) Immunofluorescent detection of Sox2 (green) and Ap2α (red) in E14.5 control (A) and Pax6^{lox/lox};Mlr10 (B) mouse lenses. Arrowheads in B indicate elevated expression of Sox2 at the lens equator and in the posterior lens. (A',B') Counterstaining of A,B with DAPI. (C) Quantification of Sox2 protein by confocal image analysis ($n=6$, $**P<0.001$). eq, lens equator; LE, lens epithelium; re, retina. Scale bar: 100 μm.

Ectopic Wnt/β-catenin activity inhibits LFC differentiation

Sox2 is a known target of Wnt signaling in the retina (Van Raay et al., 2005), and members of the Sox family modulate β-catenin activity (Sinner et al., 2007; Sinner et al., 2004). Therefore, we examined a possible connection between loss of Pax6 and canonical Wnt signaling. We first characterized the expression of *Sfrp2*, a secreted inhibitor of Wnt signaling and a target of Pax6 (Kim et al., 2001). In control E14.5 lenses, *Sfrp2* was detected anterior to the lens equator (Fig. 8A) (Chen et al., 2004). By contrast, *Sfrp2* was not detected in Pax6^{lox/lox};Mlr10 lenses (Fig. 8E). Thus, Pax6 regulates *Sfrp2* in the LE, which might play a role in the attenuation of Wnt signaling during LFC differentiation.

Taking this into consideration, we hypothesized that overexpression of β-catenin (*Catnb*; *Ctnnb1*) would result in LFC differentiation failure. To test this hypothesis, we established *Catnb^{lox(ex3)};Mlr10-Cre* gain-of-function mutants. In the *Catnb^{lox(ex3)}* allele, Cre-mediated deletion of exon 3 results in accumulation of β-catenin in the nucleus, enabling expression of its target genes (Harada et al., 1999).

Catnb^{lox(ex3)};Mlr10 adult lenses were significantly smaller than controls (not shown). At E15.5, the morphology of *Catnb^{lox(ex3)};Mlr10* lenses was abnormal, with epithelial cells accumulating at the lens equator and in the posterior lens (Fig. 8F), similar to the Pax6^{lox/lox};Mlr10 phenotype (Fig. 2H-J).

The transcriptional control function of β-catenin, as opposed to its structural role, depends on its cellular localization. In the control, β-catenin was detected primarily in the cell membranes (Fig. 8C), whereas in the *Catnb^{lox(ex3)};Mlr10* lenses it was detected in the cytoplasm and nuclei (Fig. 8G). Nuclear localization was detected by co-immunostaining with an antibody against cyclin D1 (Fig. 8C,G), a plausible target of the canonical Wnt pathway (Shtutman et al., 1999; Tetsu and McCormick, 1999). Similar to in Pax6^{lox/lox};Mlr10, proliferation, as detected by BrdU, was detected in the large mass of small nucleated cells of the equator and posterior lens (Fig. 8H). Apoptotic cells were detected in the *Catnb^{lox(ex3)};Mlr10* lens, but not in controls (Fig. 8I,N).

Pax6 was apparently unaffected by the activation of the Wnt pathway in *Catnb^{lox(ex3)};Mlr10* lenses, as it showed strong expression in the anterior LE and weak expression in the equator and in the aberrant cells of the posterior lens (Fig. 8J,O). In contrast to in Pax6^{lox/lox};Mlr10 lenses, Sox2 was not upregulated at the equator of *Catnb^{lox(ex3)};Mlr10* lenses (Fig. 8P), suggesting Pax6-dependent repression of Sox2 in lens cells. Moreover, it seems that Wnt/β-catenin does not activate Sox2 in the mammalian lens.

The canonical Wnt pathway is inactive during secondary LFC differentiation and is not regulated by Pax6

To directly examine whether Wnt/β-catenin signaling is active in Pax6^{lox/lox};Mlr10 lenses, we employed the *BATlacZ* transgene (Nakaya et al., 2005). In this reporter line, *lacZ* is expressed under control of the *Tcf/Lef* promoter, which is activated by β-catenin. As expected, in *Catnb^{lox(ex3)};Mlr10;BATlacZ* embryos, β-galactosidase activity was detected in most lens cells, especially in the nucleated, undifferentiated cells at the equator and posterior of the lens (Fig. 8Q). In Pax6^{lox/lox};Mlr10;BATlacZ animals, β-galactosidase activity was identical to that of control littermates and was not detected in the lens at E14.5 (Fig. 8L,M). Furthermore, β-catenin remained confined to the cellular membrane and did not enter the nucleus of Pax6^{lox/lox};Mlr10 lenses (Fig. 8R). This indicates that the failure of Pax6-negative cells to differentiate into LFCs is unlikely to be mediated through Wnt/β-catenin transcriptional activity.

DISCUSSION

In this study, we established the first in vivo model in which Pax6 is abolished from a formed embryonic lens, constituting a direct tool for the study of the role of Pax6 during secondary lens fiber differentiation. The findings presented reveal that Pax6 is essential for lens fiber differentiation but is dispensable for maintaining a lens epithelial identity. This role of Pax6 is not mediated by changes in canonical Wnt pathway activity, or by the upregulation of Sox2 observed in Pax6-deficient lenses. Known transcriptional regulators of LFC differentiation – Sox1, cMaf and Prox1 – are not dependent on Pax6 activity, but are, however, insufficient to enable lens fiber differentiation without Pax6. Therefore, Pax6 activity within the lens is crucial for cell cycle exit and for initiation of the lens fiber differentiation program in the mammalian eye.

Robustness of fetal stage LE to haploinsufficiency of Pax6

The vertebrate eye is sensitive to changes in Pax6 dosage: both reduction and elevation result in severe ocular phenotypes (Duncan et al., 2004; Glaser et al., 1994; Glaser et al., 1990; Hogan et al., 1988; Sanyal and Hawkins, 1979; Schedl et al., 1996). We have previously shown that the lens is intrinsically sensitive to Pax6 dosage reduction, as somatic inactivation of one copy of Pax6 in the SE mimics the lens phenotype of Pax6 heterozygotes (Davis-Silberman et al., 2005).




Fig. 7. Arrest of LFC differentiation following Pax6 loss is not mediated by upregulation of Sox2. (A) *Sox2*^{lox} targeting vector and somatic deletion allele. The *neo* selection cassette is flanked by *frt* sites (blue triangles). The single *Sox2* exon is flanked by loxP sites (green triangles). (B–M') Cre-mediated deletion results in the *Sox2*^d allele. *Sox2*^{lox/lox}; *Pax6*^{lox/lox} control (B–G) and *Sox2*^{lox/lox}; *Pax6*^{lox/lox}; *Mr10* E14.5 double somatic mutant (H–M') mouse lenses analyzed by H&E staining (B, H) and antibody labeling for Sox2 and Ap2α (C, I, green and red, respectively), phosphohistone H3 (PH3, red in D, J, J'), αA-crystallin (E, K), β-crystallin (F, L) and cleaved caspase 3 (cCas, red in G, M, M'). Counterstaining was with DAPI (D, F, G, J, L, M). Arrows indicate the lens equator, arrowheads to the aberrant cells in the lens posterior. LE, lens epithelium; re, retina. Scale bar: 100 μm.

In contrast to the phenotype observed in the SE following Pax6 reduction, we observed no phenotypic difference between *Pax6*^{lox/+}; *Mr10* lenses and controls, even in adult mice (1 year old, not shown). Therefore, a diploid dose of *Pax6* is not necessary during the late stages of lens development, in contrast to the sensitivity to *Pax6* reduction during formation of the LP. This confirms previous hypotheses, which attributed the *Pax6* dosage requirement to lens placode formation, based on the analysis of lens development in *Pax6*^{+/-} mutants (van Raamsdonk and Tilghman, 2000) or deletion of the *Pax6* ectoderm enhancer (Dimanlig et al., 2001).

Pax6 is required for cell cycle exit, cell survival and lens fiber differentiation

Pax6 is expressed in both the proliferating anterior LE and in the transitional zone, including non-proliferating cells (Ki67⁻ BrdU⁻; Fig. 3). *Pax6* loss from the whole lens alters cell proliferation in both regions, increasing the proportion of cells in the S phase in the LE and preventing cell cycle exit in the transitional zone (Fig. 3). *Pax6* involvement in cell cycle regulation has been reported in the developing retina (Marquardt et al., 2001; Oron-Karni et al., 2008). During brain development, *Pax6* loss results in a shortened cell cycle during early corticogenesis but a prolonged S phase during later stages (Estivill-Torrus et al., 2002).

Pax6 involvement in cell cycle regulation might be through its direct interactions with cell cycle components, including the retinoblastoma protein (pRb; Rb1), which has been found to be associated with *Pax6* in vitro and in lens extracts (Cvekl et al., 1999). Accordingly, the phenotype of pRb loss-of-function includes cell differentiation arrest, persistent proliferation and reduced survival – a phenotype reminiscent of *Pax6*^{lox/lox}; *Mr10* lenses (Morgenbesser et al., 1994; Pan and Griep, 1994). Other proposed mechanisms include direct association of *Pax6* with the centrosomes or mitotic

chromosomes in proliferating cortical progenitors and cultured cells, respectively (Tamai et al., 2007; Zaccarini et al., 2007). The relevance of the above findings to *Pax6* function in cell cycle regulation in the lens remains to be investigated.

Pax6 is known to bind, activate and repress crystallin gene expression in vitro and in vivo during early stages of development (Cvekl and Duncan, 2007; Cvekl et al., 2004). During the late stages of newt lens regeneration, which emulates normal lens development, *Pax6* has been shown to be needed for LFC differentiation but not for crystallin maintenance (Madhavan et al., 2006). In accordance with this, our results show that removal of *Pax6* does not alter the expression of α-crystallins in the LE, but at the same time precludes the upregulation of crystallin expression observed in differentiating LFCs (Fig. 4). The requirement for *Pax6* for the onset of LFC differentiation can be explained by the recently proposed chromatin remodeling model (Yang et al., 2006), according to which TFs operate in a temporal order on enhancer sequences of the *Cryaa* gene, each TF enabling chromatin remodeling and activity of further TFs. In *Pax6*^{lox/lox}; *Mr10* mutants, *Pax6* might enable basal expression of *Cryaa* and *Cryab* by 'opening' chromatin to transcription prior to mutation onset in LE cells. After the initiation of α-crystallin expression, *Pax6* is dispensable for its maintenance in the LE. In the transitional zone, upregulation of *Cryaa* and the initiation of β- and γ-crystallin activation do require *Pax6*.

Pax6 seems to govern some, but not all, of the processes associated with LFC differentiation. In the *Pax6*^{lox/lox}; *Mr10* lenses, expression of differentiation regulators (*Sox1*, *cMaf* and *Prox1*) and cell cycle inhibitors (*p27*^{Kip1} and *p57*^{Kip2}) is not lost in the transitional zone (Fig. 5). In fact, the transitional zone seems to be expanded, probably due to the continued proliferation of *Pax6*-deficient cells. This expansion might also be occurring at the expense of the anterior LE, as can be seen by the large population of Sox2⁺ cells at the equator and the relatively small population of




Fig. 8. β -catenin overexpression leads to LFC differentiation failure independently of Pax6. (A,E) *Sfrp2* transcripts in control (A) and *Pax6^{lox/lox};Mlr10* (E) E14.5 eyes. (B-D,F-K,N-P) E15.5 control (B-D,I-K) and *Catnb^{lox(ex3)};Mlr10* (F-H,N-P) lenses. (B,F) H&E staining reveals aberrant accumulation of cells at the lens equator of *Catnb^{lox(ex3)};Mlr10* mice (F, arrowhead). (C,G) β -catenin and cyclin D1 (green and red, respectively) detected by antibody labeling. (D,H) BrdU⁺ cells (red) found anterior to the lens equator in the control (D, arrow) accumulate posterior to the lens equator of *Catnb^{lox(ex3)};Mlr10* lenses (H, arrows). (I,N) cCas3 (red) is not detected in controls (I) but is detected in *Catnb^{lox(ex3)};Mlr10* lenses (N, arrows). (J,O) Pax6 protein is detected in the *Catnb^{lox(ex3)};Mlr10* lenses (O), as in the control (J). (K,P) Sox2 is weakly expressed in control LE and in the lens equator of *Catnb^{lox(ex3)};Mlr10* (arrows). (L,M,Q) β -galactosidase activity (β -Gal, blue) in E14.5 lenses of *BATlacZ* (L), *Catnb^{lox(ex3)};Mlr10;BATlacZ* (Q) and *Pax6^{lox/lox};Mlr10;BATlacZ* (M). β -galactosidase activity is detected in the developing eyelid (el, arrow) but not in lenses of *Pax6^{lox/lox};Mlr10;BATlacZ* (M). (R) Antibody labeling against β -catenin (green) and cyclin D1 (red). Counterstaining is with DAPI (blue, D,H,I,N). el, eyelid; eq, lens equator; re, retina; lf, lens fiber. Scale bars: white/black, 100 μ m; gray, 400 μ m.

anterior Ap2 α ⁺ cells (Fig. 6). It appears that cells at the *Pax6*-deficient lens equator are competent to respond to some external cues that trigger the expression of transitional zone markers. However, without *Pax6*, these factors are insufficient to bring about cell cycle exit, or to activate crystallin expression and cellular elongation. Knockout models of *Sox1*, *cMaf* and *Prox1* show that these TFs are directly essential for crystallin accumulation and elongation of LFCs (Kawauchi et al., 1999; Nishiguchi et al., 1998; Ring et al., 2000; Wigle et al., 1999; Yoshida and Yasuda, 2002). In the transitional zone, Pax6 is co-expressed with these factors and has been found to co-operate with cMaf (Sakai et al., 2001; Yoshida et al., 2001). Thus, although Pax6 is not required for the onset of expression of Sox1, cMaf and Prox1, it might function with them to regulate LFC differentiation.

Lens inversion experiments have demonstrated that lens polarity is dependent on the cellular environment (Coulombre and Coulombre, 1963). Since then, numerous growth factor families have been reported to influence LFC differentiation (reviewed by Lovicu and McAvoy, 2005). Most notably, FGFs were shown to initiate LFC differentiation in a concentration-dependent manner (Robinson, 2006). *Mlr10*-Cre-mediated inactivation of three FGF receptors resulted in complete arrest of LFC differentiation at the lens vesicle stage and reduced expression of Prox1, cMaf, p27^{Kip1} and p57^{Kip2} (Zhao et al., 2008). This phenotype was more severe than that of the *Pax6* mutant presented here, which suggests that Pax6 is not absolutely essential for the capacity of cells to respond to FGF signaling, although it might regulate some components of this pathway.

A complex relationship between Pax6 and Sox2: Pax6 inhibits the expression of Sox2 at the lens equator

Pax6 and Sox2 have been shown to form a functional complex that is required for the activation of crystallin genes at the placodal stage (Cvekl et al., 2004; Kamachi et al., 2001; Kondoh et al., 2004; Smith

et al., 2005). In addition, Pax6 has been shown to bind enhancer sequences of *Sox2* and to activate *Sox2* expression in lens cells (Inoue et al., 2007; Lengler et al., 2005) and in neuronal progenitors (Wen et al., 2008), suggesting a positive effect of Pax6 on *Sox2* expression.

We show that during late stages of development, Pax6 ablation results in a dramatic increase in *Sox2* expression in the transitional zone but not in the anterior LE (Fig. 6C). *Sox2* is associated with maintenance of a progenitor phenotype and stem cell characteristics (Graham et al., 2003; Loh et al., 2008; Pan and Thomson, 2007). Therefore, the observed upregulation of *Sox2* might be the result of reversion to a more primal state that lacks the capacity to differentiate. However, by deleting *Sox2* in Pax6-deficient lenses, we demonstrated that the increase in *Sox2* is not the cause of the observed phenotype. The analysis of *Sox2*-deficient lenses suggests that *Sox2* is not required at later stages of lens development (Fig. 7 and not shown). Moreover, when LE cells fail to differentiate because of β -catenin activation, *Sox2* expression does not increase (Fig. 8P), contradicting the notion that *Sox2* upregulation is the default result of differentiation failure in the LE.

The Wnt pathway and LFC differentiation

During lens induction, Wnt signaling in the SE is essential for preventing ectopic lens formation in the surrounding head ectoderm, and overexpression of β -catenin in the SE prevents lens induction and inhibits expression of both *Pax6* and *Sox2* (Miller et al., 2006; Smith et al., 2005; Stump et al., 2003). The involvement of the canonical Wnt pathway in LFC differentiation is still under debate. β -catenin loss-of-function phenotypes have been largely attributed to its structural, rather than transcriptional, role (Kreslova et al., 2007; Smith et al., 2005). Nevertheless, many components of the Wnt signaling pathway are expressed in distinct temporal and spatial patterns throughout lens development (Ang et al., 2004; Chen et al., 2004; Lovicu and McAvoy, 2005). In addition, the Wnt

co-receptor Lrp6 has been shown to delay LFC differentiation (Stump et al., 2003). These findings suggest that canonical Wnt/ β -catenin signaling does play an antagonistic role in LFC differentiation.

Recently, lens-specific β -catenin loss-of-function mutants were established (*Catnb^{lox/lox};Mlr10*). Analysis of these mutants revealed that β -catenin is required for proliferation and differentiation of the LE (Cain et al., 2008). In accordance with this, the constitutive stabilization of β -catenin conducted in the current study resulted in the prevention of cell cycle exit and of LFC differentiation (Fig. 8). The seemingly similar phenotypes of β -catenin gain-of-function and *Pax6*-deficient lenses, together with the downregulation of *Sfrp2* in the latter, led to the hypothesis that the phenotype of the *Pax6^{lox/lox};Mlr10* lens is mediated by alterations in the canonical Wnt/ β -catenin signaling pathway. This hypothesis was tested in this study through the use of the *BATlacZ* transgene (Nakaya et al., 2005). The *lacZ* reporter was activated in the lens of E14.5 β -catenin gain-of-function mutants, enabling detection of canonical Wnt pathway activity in the lens. *lacZ* was not active in control or *Pax6^{lox/lox};Mlr10* lenses. From this, we infer that β -catenin transcriptional control activity does not play a major role in the LE at E14.5. Moreover, it seems that the phenotype of *Pax6^{lox/lox};Mlr10* lenses is not mediated by Wnt/ β -catenin signaling, although *Pax6* involvement in LFC differentiation through the non-canonical Wnt pathways remains to be investigated.

Acknowledgements

We thank Joachim Graw and Lena Remizova for helpful comments on the manuscript; Corinne Lobe and Andreas Nagy for the *Z/AP* reporter line; Terry P. Yamaguchi for the *BATlacZ* reporter line; Peter Gruss for the mouse lines established in his laboratory; and Robin Lovell-Badge, Guillermo Oliver, Cornelia Leimeister, Amir Rattner and Leif Lundh for providing constructs for ISH probes. Research in R.A.-P.'s laboratory is supported by the Israel Science Foundation, the Binational Science Foundation, the AMN foundation, the Glaucoma Research Foundation, the Israeli Ministry of Health and the E. Matilda Ziegler Foundation. The research of M.L.R. is supported by NEI R01EY12995.

Supplementary material

Supplementary material for this article is available at <http://dev.biologists.org/cgi/content/full/136/15/2567/DC1>

References

- Ang, S. J., Stump, R. J., Lovicu, F. J. and McAvoy, J. W. (2004). Spatial and temporal expression of Wnt and Dickkopf genes during murine lens development. *Gene Expr. Patterns* **4**, 289-295.
- Ashery-Padan, R. and Gruss, P. (2001). Pax6 lights-up the way for eye development. *Curr. Opin. Cell Biol.* **13**, 706-714.
- Ashery-Padan, R., Marquardt, T., Zhou, X. and Gruss, P. (2000). Pax6 activity in the lens primordium is required for lens formation and for correct placement of a single retina in the eye. *Genes Dev.* **14**, 2701-2711.
- Bassnett, S. and Beebe, D. C. (1992). Coincident loss of mitochondria and nuclei during lens fiber cell differentiation. *Dev. Dyn.* **194**, 85-93.
- Beebe, D. C. and Piatigorsky, J. (1976). Differential synthesis of crystallin and noncrystallin polypeptides during lens fiber cell differentiation in vitro. *Exp. Eye Res.* **22**, 237-249.
- Beebe, D. C., Compart, P. J., Johnson, M. C., Feagans, D. E. and Feinberg, R. N. (1982). The mechanism of cell elongation during lens fiber cell differentiation. *Dev. Biol.* **92**, 54-59.
- Bhat, S. P. (2001). The ocular lens epithelium. *Biosci. Rep.* **21**, 537-563.
- Brahma, S. K. (1988). Ontogeny of beta B1-crystallin polypeptide during chicken lens development. *Exp. Eye Res.* **47**, 507-510.
- Cain, S., Martinez, G., Kokkinos, M. I., Turner, K., Richardson, R. J., Abud, H. E., Huelsken, J., Robinson, M. L. and de longh, R. U. (2008). Differential requirement for beta-catenin in epithelial and fiber cells during lens development. *Dev. Biol.* **321**, 420-433.
- Chen, Y., Stump, R. J., Lovicu, F. J. and McAvoy, J. W. (2004). Expression of Frizzleds and secreted frizzled-related proteins (Sfrps) during mammalian lens development. *Int. J. Dev. Biol.* **48**, 867-877.
- Chen, Y., Stump, R. J., Lovicu, F. J. and McAvoy, J. W. (2006). A role for Wnt/planar cell polarity signaling during lens fiber cell differentiation? *Semin. Cell Dev. Biol.* **17**, 712-725.
- Coulombre, J. L. and Coulombre, A. J. (1963). Lens Development: Fiber Elongation and Lens Orientation. *Science* **142**, 1489-1490.
- Cvekl, A. and Duncan, M. K. (2007). Genetic and epigenetic mechanisms of gene regulation during lens development. *Prog. Retin. Eye Res.* **26**, 555-597.
- Cvekl, A., Kashanchi, F., Sax, C. M., Brady, J. N. and Piatigorsky, J. (1995). Transcriptional regulation of the mouse alpha A-crystallin gene: activation dependent on a cyclic AMP-responsive element (DE1/CRE) and a Pax-6-binding site. *Mol. Cell. Biol.* **15**, 653-660.
- Cvekl, A., Kashanchi, F., Brady, J. N. and Piatigorsky, J. (1999). Pax-6 interactions with TATA-box-binding protein and retinoblastoma protein. *Invest. Ophthalmol. Vis. Sci.* **40**, 1343-1350.
- Cvekl, A., Yang, Y., Chauhan, B. K. and Cveklova, K. (2004). Regulation of gene expression by Pax6 in ocular cells: a case of tissue-preferred expression of crystallins in lens. *Int. J. Dev. Biol.* **48**, 829-844.
- Davis-Silberman, N., Kalich, T., Oron-Karni, V., Marquardt, T., Kroeber, M., Tamm, E. R. and Ashery-Padan, R. (2005). Genetic dissection of Pax6 dosage requirements in the developing mouse eye. *Hum. Mol. Genet.* **14**, 2265-2276.
- de longh, R. U., Lovicu, F. J., Chamberlain, C. G. and McAvoy, J. W. (1997). Differential expression of fibroblast growth factor receptors during rat lens morphogenesis and growth. *Invest. Ophthalmol. Vis. Sci.* **38**, 1688-1699.
- Dimanlig, P. V., Faber, S. C., Auerbach, W., Makarenkova, H. P. and Lang, R. A. (2001). The upstream ectoderm enhancer in Pax6 has an important role in lens induction. *Development* **128**, 4415-4424.
- Duncan, M. K., Li, X., Ogino, H., Yasuda, K. and Piatigorsky, J. (1996). Developmental regulation of the chicken beta B1-crystallin promoter in transgenic mice. *Mech. Dev.* **57**, 79-89.
- Duncan, M. K., Haynes, J. I., 2nd, Cvekl, A. and Piatigorsky, J. (1998). Dual roles for Pax-6: a transcriptional repressor of lens fiber cell-specific beta-crystallin genes. *Mol. Cell. Biol.* **18**, 5579-5586.
- Duncan, M. K., Cui, W., Oh, D. J. and Tomarev, S. I. (2002). Prox1 is differentially localized during lens development. *Mech. Dev.* **112**, 195-198.
- Duncan, M. K., Xie, L., David, L. L., Robinson, M. L., Taube, J. R., Cui, W. and Reneker, L. W. (2004). Ectopic Pax6 expression disturbs lens fiber cell differentiation. *Invest. Ophthalmol. Vis. Sci.* **45**, 3589-3598.
- Endl, E., Steinbach, P., Knuchel, R. and Hofstadter, F. (1997). Analysis of cell cycle-related Ki-67 and p120 expression by flow cytometric BrdUrd-Hoechst/7AAD and immunolabeling technique. *Cytometry* **29**, 233-241.
- Episkopou, V. (2005). SOX2 functions in adult neural stem cells. *Trends Neurosci.* **28**, 219-221.
- Estivill-Torres, G., Pearson, H., van Heyningen, V., Price, D. J. and Rashbass, P. (2002). Pax6 is required to regulate the cell cycle and the rate of progression from symmetrical to asymmetrical division in mammalian cortical progenitors. *Development* **129**, 455-466.
- Fujiwara, M., Uchida, T., Osumi-Yamashita, N. and Eto, K. (1994). Uchida rat (rSey): a new mutant rat with craniofacial abnormalities resembling those of the mouse Sey mutant. *Differentiation* **57**, 31-38.
- Garcia, C. M., Yu, K., Zhao, H., Ashery-Padan, R., Ornitz, D. M., Robinson, M. L. and Beebe, D. C. (2005). Signaling through FGF receptor-2 is required for lens cell survival and for withdrawal from the cell cycle during lens fiber cell differentiation. *Dev. Dyn.* **233**, 516-527.
- Gehring, W. J. (1996). The master control gene for morphogenesis and evolution of the eye. *Genes Cells* **1**, 11-15.
- Glaser, T., Lane, J. and Housman, D. (1990). A mouse model of the aniridia-Wilms tumor deletion syndrome. *Science* **250**, 823-827.
- Glaser, T., Jepeal, L., Edwards, J. G., Young, S. R., Favor, J. and Maas, R. L. (1994). PAX6 gene dosage effect in a family with congenital cataracts, aniridia, anophthalmia and central nervous system defects. *Nat. Genet.* **7**, 463-471 [erratum appears in *Nat. Genet.* (1994) **8**, 203].
- Gopal-Srivastava, R., Cvekl, A. and Piatigorsky, J. (1996). Pax-6 and alphaB-crystallin/small heat shock protein gene regulation in the murine lens. Interaction with the lens-specific regions, LSR1 and LSR2. *J. Biol. Chem.* **271**, 23029-23036.
- Graham, V., Khudiyakov, J., Ellis, P. and Pevny, L. (2003). SOX2 functions to maintain neural progenitor identity. *Neuron* **39**, 749-765.
- Grainger, R. M., Mannion, J. E., Cook, T. L., Jr and Zygar, C. A. (1997). Defining intermediate stages in cell determination: acquisition of a lens-forming bias in head ectoderm during lens determination. *Dev. Genet.* **20**, 246-257.
- Grindley, J. C., Davidson, D. R. and Hill, R. E. (1995). The role of Pax-6 in eye and nasal development. *Development* **121**, 1433-1442.
- Harada, N., Tamai, Y., Ishikawa, T., Sauer, B., Takaku, K., Oshima, M. and Taketo, M. M. (1999). Intestinal polyposis in mice with a dominant stable mutation of the beta-catenin gene. *EMBO J.* **18**, 5931-5942.
- Hogan, B. L., Hirst, E. M., Horsburgh, G. and Hetherington, C. M. (1988). Small eye (Sey): a mouse model for the genetic analysis of craniofacial abnormalities. *Development* **103 Suppl**, 115-119.

- Inoue, M., Kamachi, Y., Matsunami, H., Imada, K., Uchikawa, M. and Kondoh, H. (2007). PAX6 and SOX2-dependent regulation of the Sox2 enhancer N-3 involved in embryonic visual system development. *Genes Cells* **12**, 1049-1061.
- Jaenicke, R. (1996). Stability and folding of ultrastable proteins: eye lens crystallins and enzymes from thermophiles. *FASEB J.* **10**, 84-92.
- Joyner, A. L. I. (1995). Gene targeting: a practical approach. In *The Practical Approach Series* (ed. B. D. H. D. Rickwood), pp. 234. New York: IRL Press at Oxford University Press.
- Kamachi, Y., Sockanathan, S., Liu, Q., Breitman, M., Lovell-Badge, R. and Kondoh, H. (1995). Involvement of SOX proteins in lens-specific activation of crystallin genes. *EMBO J.* **14**, 3510-3519.
- Kamachi, Y., Uchikawa, M., Collignon, J., Lovell-Badge, R. and Kondoh, H. (1998). Involvement of Sox1, 2 and 3 in the early and subsequent molecular events of lens induction. *Development* **125**, 2521-2532.
- Kamachi, Y., Uchikawa, M., Tanouchi, A., Sekido, R. and Kondoh, H. (2001). Pax6 and SOX2 form a co-DNA-binding partner complex that regulates initiation of lens development. *Genes Dev.* **15**, 1272-1286.
- Kawauchi, S., Takahashi, S., Nakajima, O., Ogino, H., Morita, M., Nishizawa, M., Yasuda, K. and Yamamoto, M. (1999). Regulation of lens fiber cell differentiation by transcription factor c-Maf. *J. Biol. Chem.* **274**, 19254-19260.
- Kim, A. S., Anderson, S. A., Rubenstein, J. L., Lowenstein, D. H. and Pleasure, S. J. (2001). Pax-6 regulates expression of SFRP-2 and Wnt-7b in the developing CNS. *J. Neurosci.* **21**, RC132.
- Kondoh, H., Uchikawa, M. and Kamachi, Y. (2004). Interplay of Pax6 and SOX2 in lens development as a paradigm of genetic switch mechanisms for cell differentiation. *Int. J. Dev. Biol.* **48**, 819-827.
- Kralova, J., Czerny, T., Spanielova, H., Ratajova, V. and Kozmik, Z. (2002). Complex regulatory element within the gammaE- and gammaF-crystallin enhancers mediates Pax6 regulation and is required for induction by retinoic acid. *Gene* **286**, 271-282.
- Kreslova, J., Machon, O., Ruzickova, J., Lachova, J., Wawrousek, E. F., Kemler, R., Krauss, S., Piatigorsky, J. and Kozmik, Z. (2007). Abnormal lens morphogenesis and ectopic lens formation in the absence of beta-catenin function. *Genesis* **45**, 157-168.
- Leimeister, C., Bach, A. and Gessler, M. (1998). Developmental expression patterns of mouse sFRP genes encoding members of the secreted frizzled related protein family. *Mech. Dev.* **75**, 29-42.
- Lengler, J., Bittner, T., Munster, D., Gawad Ael, D. and Graw, J. (2005). Agonistic and antagonistic action of AP2, Msx2, Pax6, Prox1 AND Six3 in the regulation of Sox2 expression. *Ophthalmic Res.* **37**, 301-309.
- Liu, H., Mohamed, O., Dufort, D. and Wallace, V. A. (2003). Characterization of Wnt signaling components and activation of the Wnt canonical pathway in the murine retina. *Dev. Dyn.* **227**, 323-334.
- Lobe, C. G., Koop, K. E., Kreppner, W., Lomeli, H., Gertsenstein, M. and Nagy, A. (1999). Z/AP, a double reporter for cre-mediated recombination. *Dev. Biol.* **208**, 281-292.
- Loh, Y. H., Ng, J. H. and Ng, H. H. (2008). Molecular framework underlying pluripotency. *Cell Cycle* **7**, 885-891.
- Lovicu, F. J. and Robinson, M. L. (2004). *Development of the Ocular Lens*. Cambridge: Cambridge University Press.
- Lovicu, F. J. and McAvoy, J. W. (2005). Growth factor regulation of lens development. *Dev. Biol.* **280**, 1-14.
- Madhavan, M., Haynes, T. L., Frisch, N. C., Call, M. K., Minich, C. M., Tsonis, P. A. and Del Rio-Tsonis, K. (2006). The role of Pax-6 in lens regeneration. *Proc. Natl. Acad. Sci. USA* **103**, 14848-14853.
- Marquardt, T., Ashery-Padan, R., Andrejewski, N., Scardigli, R., Guillemot, F. and Gruss, P. (2001). Pax6 is required for the multipotent state of retinal progenitor cells. *Cell* **105**, 43-55.
- Miller, L. A., Smith, A. N., Taketo, M. M. and Lang, R. A. (2006). Optic cup and facial patterning defects in ocular ectoderm beta-catenin gain-of-function mice. *BMC Dev. Biol.* **6**, 14.
- Morgenbesser, S. D., Williams, B. O., Jacks, T. and DePinho, R. A. (1994). p53-dependent apoptosis produced by Rb-deficiency in the developing mouse lens. *Nature* **371**, 72-74.
- Muta, M., Kamachi, Y., Yoshimoto, A., Higashi, Y. and Kondoh, H. (2002). Distinct roles of SOX2, Pax6 and Maf transcription factors in the regulation of lens-specific delta1-crystallin enhancer. *Genes Cells* **7**, 791-805.
- Nakaya, M. A., Biris, K., Tsukiyama, T., Jaime, S., Rawls, J. A. and Yamaguchi, T. P. (2005). Wnt3a links left-right determination with segmentation and anteroposterior axis elongation. *Development* **132**, 5425-5436.
- Nishiguchi, S., Wood, H., Kondoh, H., Lovell-Badge, R. and Episkopou, V. (1998). Sox1 directly regulates the gamma-crystallin genes and is essential for lens development in mice. *Genes Dev.* **12**, 776-781.
- Ogino, H. and Yasuda, K. (2000). Sequential activation of transcription factors in lens induction. *Dev. Growth Differ.* **42**, 437-448.
- Oliver, G., Mailhos, A., Wehr, R., Copeland, N. G., Jenkins, N. A. and Gruss, P. (1995). Six3, a murine homologue of the sine oculis gene, demarcates the most anterior border of the developing neural plate and is expressed during eye development. *Development* **121**, 4045-4055.
- Oron-Karni, V., Farhy, C., Elgart, M., Marquardt, T., Remizova, L., Yaron, O., Xie, Q., Cvekl, A. and Ashery-Padan, R. (2008). Dual requirement for Pax6 in retinal progenitor cells. *Development* **135**, 4037-4047.
- Pan, G. and Thomson, J. A. (2007). Nanog and transcriptional networks in embryonic stem cell pluripotency. *Cell Res.* **17**, 42-49.
- Pan, H. and Griep, A. E. (1994). Altered cell cycle regulation in the lens of HPV-16 E6 or E7 transgenic mice: implications for tumor suppressor gene function in development. *Genes Dev.* **8**, 1285-1299.
- Pontoriero, G. F., Deschamps, P., Ashery-Padan, R., Wong, R., Yang, Y., Zavadil, J., Cvekl, A., Sullivan, S., Williams, T. and West-Mays, J. A. (2008). Cell autonomous roles for AP-2alpha in lens vesicle separation and maintenance of the lens epithelial cell phenotype. *Dev. Dyn.* **237**, 602-617.
- Quinn, J. C., West, J. D. and Hill, R. E. (1996). Multiple functions for Pax6 in mouse eye and nasal development. *Genes Dev.* **10**, 435-446.
- Rafferty, N. S. and Rafferty, K. A., Jr (1981). Cell population kinetics of the mouse lens epithelium. *J. Cell. Physiol.* **107**, 309-315.
- Ring, B. Z., Cordes, S. P., Overbeek, P. A. and Barsh, G. S. (2000). Regulation of mouse lens fiber cell development and differentiation by the Maf gene. *Development* **127**, 307-317.
- Robinson, M. L. (2006). An essential role for FGF receptor signaling in lens development. *Semin. Cell Dev. Biol.* **17**, 726-740.
- Robinson, M. L. and Overbeek, P. A. (1996). Differential expression of alpha A- and alpha B-crystallin during murine ocular development. *Invest. Ophthalmol. Vis. Sci.* **37**, 2276-2284.
- Rodriguez, C. I., Buchholz, F., Galloway, J., Sequerra, R., Kasper, J., Ayala, R., Stewart, A. F. and Dymecki, S. M. (2000). High-efficiency deleter mice show that FLPe is an alternative to Cre-loxP. *Nat. Genet.* **25**, 139-140.
- Sakai, M., Imaki, J., Yoshida, K., Ogata, A., Matsushima-Hibaya, Y., Kuboki, Y., Nishizawa, M. and Nishi, S. (1997). Rat maf related genes: specific expression in chondrocytes, lens and spinal cord. *Oncogene* **14**, 745-750.
- Sakai, M., Serria, M. S., Ikeda, H., Yoshida, K., Imaki, J. and Nishi, S. (2001). Regulation of c-maf gene expression by Pax6 in cultured cells. *Nucleic Acids Res.* **29**, 1228-1237.
- Sanyal, S. and Hawkins, R. K. (1979). Dysgenetic lens (dyl)-a new gene in the mouse. *Invest. Ophthalmol. Vis. Sci.* **18**, 642-645.
- Schedl, A., Ross, A., Lee, M., Engelkamp, D., Rashbass, P., van Heyningen, V. and Hastie, N. D. (1996). Influence of PAX6 gene dosage on development: overexpression causes severe eye abnormalities. *Cell* **86**, 71-82.
- Sevel, D. and Isaacs, R. (1988). A re-evaluation of corneal development. *Trans. Am. Ophthalmol. Soc.* **86**, 178-207.
- Shtutman, M., Zhurinsky, J., Simcha, I., Albanese, C., D'Amico, M., Pestell, R. and Ben-Ze'ev, A. (1999). The cyclin D1 gene is a target of the beta-catenin/LEF-1 pathway. *Proc. Natl. Acad. Sci. USA* **96**, 5522-5527.
- Sinner, D., Rankin, S., Lee, M. and Zorn, A. M. (2004). Sox17 and beta-catenin cooperate to regulate the transcription of endodermal genes. *Development* **131**, 3069-3080.
- Sinner, D., Kordich, J. J., Spence, J. R., Opoka, R., Rankin, S., Lin, S. C., Jonatan, D., Zorn, A. M. and Wells, J. M. (2007). Sox17 and Sox4 differentially regulate beta-catenin/T-cell factor activity and proliferation of colon carcinoma cells. *Mol. Cell. Biol.* **27**, 7802-7815.
- Smith, A. N., Miller, L. A., Song, N., Taketo, M. M. and Lang, R. A. (2005). The duality of beta-catenin function: a requirement in lens morphogenesis and signaling suppression of lens fate in periocular ectoderm. *Dev. Biol.* **285**, 477-489.
- Stevanovic, M., Zuffardi, O., Collignon, J., Lovell-Badge, R. and Goodfellow, P. (1994). The cDNA sequence and chromosomal location of the human SOX2 gene. *Mamm. Genome* **5**, 640-642.
- Stump, R. J., Ang, S., Chen, Y., von Bahr, T., Lovicu, F. J., Pinson, K., de longh, R. U., Yamaguchi, T. P., Sassoon, D. A. and McAvoy, J. W. (2003). A role for Wnt/beta-catenin signaling in lens epithelial differentiation. *Dev. Biol.* **259**, 48-61.
- Tamai, H., Shinohara, H., Miyata, T., Saito, K., Nishizawa, Y., Nomura, T. and Osumi, N. (2007). Pax6 transcription factor is required for the interkinetic nuclear movement of neuroepithelial cells. *Genes Cells* **12**, 983-996.
- Tetsu, O. and McCormick, F. (1999). Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature* **398**, 422-426.
- van Raamsdonk, C. D. and Tilghman, S. M. (2000). Dosage requirement and allelic expression of PAX6 during lens placode formation. *Development* **127**, 5439-5448.
- Van Raay, T. J., Moore, K. B., Iordanova, I., Steele, M., Jamrich, M., Harris, W. A. and Vetter, M. L. (2005). Frizzled 5 signaling governs the neural potential of progenitors in the developing *Xenopus* retina. *Neuron* **46**, 23-36.
- Walther, C., Guenet, J. L., Simon, D., Deutsch, U., Jostes, B., Goulding, M. D., Plachov, D., Balling, R. and Gruss, P. (1991). Pax: a murine multigene family of paired box-containing genes. *Genomics* **11**, 424-434.
- Wawrousek, E. F., Chepelinsky, A. B., McDermott, J. B. and Piatigorsky, J. (1990). Regulation of the murine alpha A-crystallin promoter in transgenic mice. *Dev. Biol.* **137**, 68-76.

- Wen, J., Hu, Q., Li, M., Wang, S., Zhang, L., Chen, Y. and Li, L. (2008). Pax6 directly modulate Sox2 expression in the neural progenitor cells. *NeuroReport* **19**, 413-417.
- West-Mays, J. A., Zhang, J., Nottoli, T., Hagopian-Donaldson, S., Libby, D., Strissel, K. J. and Williams, T. (1999). AP-2alpha transcription factor is required for early morphogenesis of the lens vesicle. *Dev. Biol.* **206**, 46-62.
- Wigle, J. T., Chowdhury, K., Gruss, P. and Oliver, G. (1999). Prox1 function is crucial for mouse lens-fibre elongation. *Nat. Genet.* **21**, 318-322.
- Yang, Y. and Cvekl, A. (2005). Tissue-specific regulation of the mouse alphaA-crystallin gene in lens via recruitment of Pax6 and c-Maf to its promoter. *J. Mol. Biol.* **351**, 453-469.
- Yang, Y., Chauhan, B. K., Cvekl, K. and Cvekl, A. (2004). Transcriptional regulation of mouse alphaB- and gammaF-crystallin genes in lens: opposite promoter-specific interactions between Pax6 and large Maf transcription factors. *J. Mol. Biol.* **344**, 351-368.
- Yang, Y., Stopka, T., Golestaneh, N., Wang, Y., Wu, K., Li, A., Chauhan, B. K., Gao, C. Y., Cvekl, K., Duncan, M. K. et al. (2006). Regulation of alphaA-crystallin via Pax6, c-Maf, CREB and a broad domain of lens-specific chromatin. *EMBO J.* **25**, 2107-2118.
- Yaron, O., Farhy, C., Marquardt, T., Applebury, M. and Ashery-Padan, R. (2006). Notch1 functions to suppress cone-photoreceptor fate specification in the developing mouse retina. *Development* **133**, 1367-1378.
- Yoshida, K., Kim, J. I., Imaki, J., Hiromi, I., Nishi, S., Matsuda, H., Harada, T., Harada, C., Ohno, S. and Sakai, M. (2001). Proliferation in the posterior region of the lens of c-maf^{-/-} mice. *Curr. Eye Res.* **23**, 116-119.
- Yoshida, T. and Yasuda, K. (2002). Characterization of the chicken L-Maf, MafB and c-Maf in crystallin gene regulation and lens differentiation. *Genes Cells* **7**, 693-706.
- Zaccarini, R., Cordelieres, F. P., Martin, P. and Saule, S. (2007). Pax6p46 binds chromosomes in the pericentromeric region and induces a mitosis defect when overexpressed. *Invest. Ophthalmol. Vis. Sci.* **48**, 5408-5419.
- Zhao, H., Yang, Y., Rizo, C. M., Overbeek, P. A. and Robinson, M. L. (2004). Insertion of a Pax6 consensus binding site into the alphaA-crystallin promoter acts as a lens epithelial cell enhancer in transgenic mice. *Invest. Ophthalmol. Vis. Sci.* **45**, 1930-1939.
- Zhao, H., Yang, T., Madakashira, B. P., Thiels, C. A., Bechtel, C. A., Garcia, C. M., Zhang, H., Yu, K., Ornitz, D. M., Beebe, D. C. et al. (2008). Fibroblast growth factor receptor signaling is essential for lens fiber cell differentiation. *Dev. Biol.* **318**, 276-288.