

Development 138, 1247-1257 (2011) doi:10.1242/dev.057646  
© 2011. Published by The Company of Biologists Ltd

# Canonical Wnt9b signaling balances progenitor cell expansion and differentiation during kidney development

Courtney M. Karner<sup>1</sup>, Amrita Das<sup>1</sup>, Zhendong Ma<sup>1</sup>, Michelle Self<sup>2</sup>, Chuo Chen<sup>3</sup>, Lawrence Lum<sup>4</sup>, Guillermo Oliver<sup>2</sup> and Thomas J. Carroll<sup>1,\*</sup>

## SUMMARY

The mammalian kidney is composed of thousands of individual epithelial tubules known as nephrons. Deficits in nephron number are associated with myriad diseases ranging from complete organ failure to congenital hypertension. A balance between differentiation and maintenance of a mesenchymal progenitor cell population determines the final number of nephrons. How this balance is struck is poorly understood. Previous studies have suggested that Wnt9b/ $\beta$ -catenin signaling induced differentiation (mesenchymal-to-epithelial transition) in a subset of the progenitors but needed to be repressed in the remaining progenitors to keep them in the undifferentiated state. Here, we report that Wnt9b/ $\beta$ -catenin signaling is active in the progenitors and is required for their renewal/proliferation. Using a combination of approaches, we have revealed a mechanism through which cells receiving the same Wnt9b/ $\beta$ -catenin signal can respond in distinct ways (proliferate versus differentiate) depending on the cellular environment in which the signal is received. Interpretation of the signal is dependent, at least in part, on the activity of the transcription factor Six2. Six2-positive cells that receive the Wnt9b signal are maintained as progenitors whereas cells with reduced levels of Six2 are induced to differentiate by Wnt9b. Using this simple mechanism, the kidney is able to balance progenitor cell expansion and differentiation insuring proper nephron endowment. These findings provide novel insights into the molecular mechanisms that regulate progenitor cell differentiation during normal and pathological conditions.

**KEY WORDS:** Wnt, Progenitor cells, Kidney, Six2, Mouse

## INTRODUCTION

During organogenesis, a balance must be struck between progenitor cell proliferation/renewal and differentiation to ensure proper organ size. The precarious nature of this equilibrium is particularly evident in the developing metanephric kidney. During development, signals from the ureteric bud epithelium stimulate the survival, proliferation and differentiation of an adjacent population of progenitor cells known as the metanephric mesenchyme. In addition, a bud-derived signal(s) causes a subset of the progenitors to differentiate into an epithelial structure known as a renal vesicle. The renal vesicle will undergo significant morphogenesis to form a nephron, a vascularized tubule that maintains blood chemistry. As a result of these inductive interactions, each mouse kidney will form up to 20,000 nephrons [ $\sim$ 1,000,000 in an adult human (Hoy et al., 2003; Keller et al., 2003; Nyengaard and Bendtsen, 1992)] all derived from an initial progenitor population of approximately 12,000 cells (Kobayashi et al., 2008).

An essential step in kidney development is establishment of a balance between the expansion and differentiation of the nephron progenitor cell population. Tipping this balance in favor of one or the other results in a reduction in nephron endowment and can have dire consequences, including the formation of progenitor cell

cancers (Wilms' tumors), renal hypoplasia, chronic hypertension and kidney failure (Keller et al., 2003; Koesters et al., 2003; Koesters et al., 1999; Li et al., 2004). Although several factors regulating progenitor cell expansion and differentiation have been identified (Barasch et al., 1999; Blank et al., 2009; Carroll et al., 2005; Dudley et al., 1995; Grieshammer et al., 2005; Perantoni et al., 2005; Schmidt-Ott et al., 2007; Stark et al., 1994), it remains unclear how the balance between these two events is controlled.

Nephron induction requires the sequential activity of two Wnts: Wnt9b and Wnt4 (Carroll et al., 2005; Stark et al., 1994). Wnt9b is secreted from the ureteric bud and induces a subset of the renal progenitor cells to aggregate and express Wnt4. Wnt4, by signaling through the canonical Wnt/ $\beta$ -catenin pathway, further induces these pre-tubular aggregates (PTAs) to transition into epithelial structures known as renal vesicles (RVs) (Park et al., 2007). Although the molecular nature of the Wnt4 signal has been determined, the pathway used by Wnt9b is still uncertain, although it has been suggested that it also signals through  $\beta$ -catenin (Park et al., 2007).

*Wnt4*, and other characterized Wnt9b targets such as *Pax8* and *Fgf8*, are expressed in only a small proportion of the progenitor cells (those undergoing differentiation), while the remaining 'uninduced' progenitors do not express these genes. This observation, and others, has led to a model suggesting that the Wnt9b signal is not active in the renewing progenitor cells, which keeps them from differentiating. How Wnt9b signaling is repressed in these cells is unclear, although one hypothesis, that has some experimental support, suggests that the transcription factor Six2 blocks the ability of cells to receive the Wnt9b signal. *Six2* is expressed in the renewing progenitors and its loss results in the precocious differentiation of the entire progenitor population in a Wnt9b-dependent manner (Kobayashi et al., 2008; Self et al., 2006). How Six2 affects Wnt9b activity is unclear.

<sup>1</sup>Department of Internal Medicine (Nephrology) and Molecular Biology, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard Dallas, TX 75390, USA. <sup>2</sup>Department of Genetics and Tumor Cell Biology, St Jude Children's Research Hospital, 262 Danny Thomas Place, Memphis, TN 38105, USA. <sup>3</sup>Department of Chemistry, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard Dallas, TX 75390, USA. <sup>4</sup>Department of Cell Biology, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, TX 75390, USA.

\*Author for correspondence (thomas.carroll@utsouthwestern.edu)

In this study, we have identified several novel targets of Wnt9b. Our data suggest that, contrary to current models, Wnt9b/ $\beta$ -catenin signaling is active in both the differentiating and renewing progenitor cells. We provide evidence that, rather than inhibiting Wnt9b, Six2 appears to cooperate with it to promote progenitor cell proliferation. Based on our results, we propose a model whereby Wnt9b is the key factor that regulates the balance between progenitor cell proliferation and differentiation. In the renewing progenitor cells, Wnt9b/ $\beta$ -catenin cooperates with Six2 to elicit progenitor cell expansion. In cells where Six2 activity is low or absent, Wnt9b/ $\beta$ -catenin promotes differentiation. These results modify our current understanding of the molecular mechanisms that regulate progenitor cell expansion and differentiation during normal and abnormal kidney development.

## MATERIALS AND METHODS

### Mice

All mouse alleles (*Wnt9b*<sup>-</sup>, *Wnt9b*<sup>neo</sup>, *Wnt9b*<sup>lox</sup>, *Six2*<sup>-</sup>, *Wnt4*<sup>-</sup>, *Rarb2Cre*; *KspCre*, *RosaYFP*, *catnb*<sup>exon3lox</sup> and *catnb*<sup>lox</sup>) are as previously described (Brault et al., 2001; Carroll et al., 2005; Harada et al., 1999; Karner et al., 2009; Kobayashi et al., 2005; Self et al., 2006; Shao et al., 2002; Stark et al., 1994).

### Ex vivo organ culture

Organ culture, mesenchymal isolation and inhibitor of Wnt (IW) or lithium treatment were as previously described (Carroll et al., 2005; Karner et al., 2010). Briefly, E11.5 kidneys or isolated mesenchyme were grown on transwell filters at the air/media interface for 24, 48 or 72 hours. The media was supplemented with LiCl (15 mM), IWR1 (100  $\mu$ M), IWP2 (5  $\mu$ M) or DMSO as a negative control. The media was replaced with fresh media containing the above listed compounds every 12 hours. All treatments were repeated at least three times with a minimum of six individual kidneys from six distinct embryos assayed per replicate.

### Heterochronic recombination

Metanephric mesenchyme from either E11.5 wild-type or E13.5 *Wnt9b*<sup>-</sup> animals was isolated as previously described (Carroll et al., 2005). Once isolated, both wild-type and *Wnt9b*<sup>-</sup> metanephric mesenchyme were individually recombined with the E11.5 wild-type ureteric bud and cultured for 48 hours. Control mesenchymes were recombined with E13.5 *Wnt9b*<sup>-</sup> ureteric bud.

### Microarray analysis

Total mRNA was extracted from 30-35 wild-type or *Wnt9b*<sup>-</sup> E11.5 mesenchymes to provide 1  $\mu$ g total RNA. This combined mRNA constitutes a single replicate. Total RNA was hybridized to the Affymetrix GeneChip Mouse Genome 430 2.0 Array by the UT Southwestern Genomics & Microarray Core facility. This analysis was repeated for five total replicates. The results of these replicates were analyzed using the Gene Set Analysis Toolkit (<http://bioinfo.vanderbilt.edu/webgestalt>) (Zhang et al., 2005).

### In situ hybridization

Both whole-mount and section in situ hybridization was performed as previously described (Carroll et al., 2005; Self et al., 2006). Tissue was hybridized with antisense probes for the following genes: *C1qdc2* (*Fam132a* – Mouse Genome Informatics), *Cited1*, *Pla2g7*, *Tafa5* (*Fam19a5* – Mouse Genome Informatics), *Uncx4.1* (*Uncx* – Mouse Genome Informatics), *Rspo1*, *Pax8*, *Pax2*, *Six2*, *Gdnf* and *Eya1*. Restriction enzymes and polymerase have been previously described, except for *C1qdc2* (Accession BC026939), *Cited1* (Accession BC052030), *Pla2g7* (Accession BC010726), *Tafa5* (Accession BC015306), *Uncx4.1* (Accession AJ001116), *Rspo1* (Accession BQ933737) and *Eya1* (Accession BC060260), that were cut with *EcoRI* and transcribed with T3 Polymerase (*Cited1* and *Eya1*), *EcoRI* and T7 (*C1qdc2*, *Tafa5* and *Rspo1*), *EcoRV* and T7 (*Pla2g7*) or *XhoI* and T7 (*Uncx4.1*).

### Immunohistochemistry

Tissue fixation, staining and image capture were as previously described (Karner et al., 2009). Tissue was incubated with the following antibodies: anti-phospho-histone H3 (Ser10) (1:500, Sigma, St Louis, MO, USA), anti-Six2 (rabbit; 1:500, ProteinTech, Chicago, IL, USA), anti-Pax2 (rabbit; 1:500, Covance, Princeton, NJ, USA) and anti-DBA (Biotinylated; 1:500, VectorLabs, Burlingame, CA, USA), anti-Cited 1 (rabbit; 1:500, NeoMarkers, Fremont, CA, USA) and anti-amphiphysin (rabbit; 1:500, ProteinTech). Nuclei were counterstained using either Topro-3 (1:1000, Invitrogen, Carlsbad, CA, USA) or Sytox Green (1:5000, Invitrogen). Cell death was quantitated using LysoTracker (1:200, Invitrogen) as previously described (Zucker et al., 1999). Results shown are representative examples from one of at least three different stainings of three different kidneys.

### Chromatin immunoprecipitation

Approximately 20-25 E15.5 mouse kidneys were isolated, homogenized and crosslinked in 1% formaldehyde. Crosslinked tissues were homogenized into a single-cell suspension. Nuclei from crosslinked cells were resuspended in Tris-EDTA buffer and sonicated. The soluble chromatin was transferred into radioimmunoprecipitation assay (RIPA) buffer and precleared. Immunoprecipitation was performed with 5  $\mu$ g of rabbit anti- $\beta$ -catenin (Santa Cruz, sc-7199) or isotype-specific IgG as a negative control (Schmidt-Ott et al., 2007), and the immune complexes were absorbed with protein A/G beads (Pierce) and blocked with bovine serum albumin. Purified genomic DNA was amplified using promoter-specific primers and visualized on a 2% agarose gel. One percent of the input DNA was amplified to normalize results.

To quantify the fold enrichment, the chromatin from two separate pull-downs was pooled and the relative amplification values were identified using an iCycler (Bio-Rad) real-time PCR detection system. The amplification was determined by normalizing the expression levels of the  $\beta$ -catenin pull-down samples to the input. The relative fold change of expression was further calculated with respect to the amplification in the IgG pull-down samples, which were arbitrarily assigned a value of 1 for each primer set. A 150 bp fragment with no consensus  $\beta$ -catenin-binding sites was amplified as the non-specific internal control site. The data provided are the averages of the fold change values relative to the IgG control from two independent runs (three replicates per run) and the error bars represent the standard deviation. As the IgG controls are always assigned a value of 1, no standard deviation is assigned.

### Primer sequences for ChIP

Primer sequences were as follows: C1qdc2 site 1 (forward) 5'-TGCTCTCCATTTCAGGTTG-3' and (reverse) 5'-GGTGACCCATTCATTGCG-3'. C1qdc2 site 2 (forward) 5'-CGTCTCATTGCTGTGGC-3' and (reverse) 5'-TGGTCATAATCTCCCTCCGC-3'. C1qdc2 site 3 (forward) 5'-GCAGGCAGGCTGAGAAAATCAC-3' and (reverse) 5'-AAATCCTCCAAACAAGACCC-3'. Tafa5 site 1 (forward) 5'-TTTGAATCCCGAAATGCC-3' and (reverse) 5'-TGAAGGAGAA-GCAGTTTGGTTACC-3'. Tafa5 site 2 (forward) 5'-TGTCTCCTTCACTAGCATCGTC-3' and (reverse) 5'-TCTCACAGAGCAC-CAAGCATCC-3'. Tafa5 site 3 (forward) 5'-GCTGAAGAGCAGAT-GAAGTCGG-3' and (reverse) 5'-CCAAGGAGAAATGAGAAATC-CAGAGC-3'. Pla2g7 site 1 (forward) 5'-GAGAAGGAAGGATTC-CACAGGTC-3' and (reverse) 5'-TGAGTCTGAGATTCATCACAC-CCAG-3'. Pla2g7 site 2 (forward) 5'-TTGAGGATCCCTGTGAGGTG-TG and (reverse) 5'-TCCCAAGGACCCCATCTAAG.

### Statistics

All statistical analysis was performed using Student's *t*-test.

## RESULTS

### Identification of novel Wnt9b target genes

We previously showed that Wnt9b is necessary and sufficient for the induction of renal vesicles in the metanephric mesenchyme (Carroll et al., 2005). To identify novel molecular targets of Wnt9b, microarray analysis was performed comparing E11.5 wild-type and

Wnt9b<sup>-/-</sup> mesenchymes. Analysis of this data resulted in the identification of 30 putative Wnt9b targets that were downregulated at least twofold in the absence of Wnt9b function (Table 1).

The expression of the Wnt9b target genes was validated in wild-type E11.5 kidneys using in situ hybridization. We generated riboprobes for 28 of the 30 target genes identified. Twenty-seven

**Table 1. Wnt9b target genes**

Gene description	Gene name	Fold change	Class	Expression in Wnt9b <sup>-/-</sup>	Expression in Wnt4 <sup>-/-</sup>	LiCl induction	Expression in IWR1 treated explants	Expression in IWP2 treated explants	Lef/Tcf sites
C1q domain containing 2	<i>C1qdc2</i>	0.481	PTA/I	---	+/-	Yes	---	---	Yes*
Cadherin 4	<i>Cdh4</i>	0.362, 0.328	PTA/I	---	+/-	ND	---	---	Yes†
Fibroblast growth factor 9	<i>Fgf9</i>	0.399	PTA/I	---	---	ND	ND	ND	Yes†
Glutathione peroxidase 6	<i>Gpx6</i>	0.203	PTA/I	---	ND	ND	ND	ND	Yes <sup>§</sup>
Lymphoid enhancer binding factor 1	<i>Lef1</i>	0.44	PTA/I	---	---	Yes	---	ND	Yes†
Musashi homolog 2	<i>Msi2</i>	0.486	PTA/I	-	ND	ND	ND	ND	Yes <sup>§</sup>
Paired box gene 8	<i>Pax8</i>	0.49	PTA/I	---	+/-	Yes	---	---	Yes††
Phospholipase C-like 3	<i>Plcl3</i>	0.496	PTA/I	---	---	ND	ND	ND	Yes†
Riken cDNA 0610010D24	<i>Daple</i>	0.494	PTA/I	---	---	Yes	ND	ND	Yes <sup>§</sup>
Solute carrier family 45, member 3	<i>Slc45a3</i>	0.399	PTA/I	---	ND	ND	ND	ND	Yes**
Sorbin and SH3 domain containing 2	<i>Sorbs2</i>	0.434	PTA/I	---	---	ND	ND	ND	Yes†
Winless-related MMTV integration site 4	<i>Wnt4</i>	0.38, 0.347	PTA/I	---	---	Yes	---	---	Yes†
Amphiphysin	<i>Amph</i>	0.492	Prog/RV/II	--	ND	Yes	ND	ND	None
Unc4.1 homeobox	<i>Uncx4.1</i>	0.36	Prog/RV/II	--	+/-	ND	---	---	Yes**
BTB (POZ) domain containing 11	<i>Btbd11</i>	0.481, 0.257, 0.209	Prog/II	---	+/-	Yes	-	---	Yes†
Cbp/p300-interacting transactivator with Glu/Asp-rich carboxy-terminal domain 1	<i>Cited1</i>	0.453	Prog/II	---	+/-	Yes	---	---	None
Chemokine (C-X-C motif) receptor 4	<i>Cxcr4</i>	0.248	Prog/II	---	---	Yes	ND	ND	Yes†
Claudin 9	<i>Cldn9</i>	0.47	Prog/II	---	+/-	Yes	ND	ND	ND
Crystallin, mu	<i>Crym</i>	0.46	Prog/II	--	+/-	ND	ND	ND	Yes <sup>§</sup>
Ets variant gene 5	<i>Etv5</i>	0.496, 0.476	Prog/II	--	ND	ND	ND	ND	Yes†
Expressed sequence AW049604	<i>Tafa5</i>	0.448	Prog/II	---	+/-	Yes	---	---	Yes†
Glial cell line derived neurotrophic factor	<i>Gdnf</i>	0.495	Prog/II	-	ND	ND	--	--	Yes†
Integrin alpha 8	<i>Itga8</i>	0.347	Prog/II	-	ND	ND	ND	ND	ND
Phospholipase A2, group VII	<i>Pla2g7</i>	0.441	Prog/RV/II	---	+/-	Yes	---	---	Yes†
Riken cDNA 2310045A20	<i>Rik2310045A20</i>	0.305	Prog/II	--	+/-	Yes	---	---	Yes†
Riken cDNA 5930427L02	<i>Rik5930427L02</i>	0.411	Prog/II	---	--	Yes	ND	ND	ND
Solute carrier family 12, member 2	<i>Slc12a2</i>	0.479	Prog/II	---	ND	ND	ND	ND	Yes <sup>§</sup>
Transferrin receptor	<i>Tfrc</i>	0.483	Prog/RV/II	--	ND	ND	ND	ND	Yes*
Expressed sequence AL022943	<i>AL022943</i>	0.44	ND	ND	ND	ND	ND	ND	ND
Riken cDNA 9430071P14	<i>Rik9430071P14</i>	0.418	ND	ND	ND	ND	ND	ND	ND

Expression domains of genes that were reduced by half or more were determined by in situ hybridization. PTA, Prog and ND indicates genes expressed in the pretubular aggregates, mesenchymal progenitors or whose expression patterns were not determined, respectively. Qualitative comparison of in situ results in wild type, Wnt9b or Wnt4 mutants indicated that some genes were undetectable (---), significantly reduced but still detectable (--), minimally but noticeably reduced (-) or not changed (+/-).

Using rVista, Lef/TCF sites were found to be conserved through the following: \*mouse (M), rat (R) and human (H); †M, R, H and Dog (D); ††M, R, H, D, Chimp (C) and Rhesus monkey (R); †††M, H, D; †††M, R, C, H; \*\*M, R, C, R, H; †††Schmidt-Ott et al., 2007.

out of 28 showed expression in the metanephric mesenchyme at E11.5 (Table 1). Based on spatial differences in the observed expression patterns, the genes were grouped into two distinct classes. Class I targets represented genes expressed in the differentiating pre-tubular aggregates (PTA, see Table 1). This class included the previously identified Wnt9b targets *Pax8* and *Wnt4* (Carroll et al., 2005), as well as several newly identified targets including cadherin 4 (*Cdh4*), *Lef1* and *C1qdc2* (Fig. 1A; Table 1). Class II targets were characterized by broader expression in the metanephric mesenchyme. This class of genes included *Cited1*, phospholipase A2 group 7 (*Pla2g7*), amphiphysin (*Amph*) expressed sequence AW049604 (*Tafa5*) and *Uncx4.1* (Fig. 1D,G,J,M; Table 1).

In situ hybridization and immunofluorescence on adjacent sections of E11.5 and P1 wild-type kidneys revealed that the class II target genes had significant overlap in expression with the bona fide progenitor cell marker *Six2*. In fact, one Class II target, *Cited1*, has previously been shown to be a marker of the self-renewing mesenchymal progenitor cells (Boyle et al., 2008). Although several of the Class II targets appeared to be specifically expressed in the progenitor cells, others had patterns that were more expansive and included the PTA domain and later structures. By contrast, Class I targets showed little if any expression in the progenitor cells (see Fig. S1A in the supplementary material and data not shown).


To verify that the identified genes were indeed targets of Wnt9b, we examined their expression in Wnt9b mutants using in situ hybridization. Irrespective of their expression domain, the expression of most genes examined was noticeably reduced or undetectable in Wnt9b mutant mesenchyme at E11.5 (Fig. 1; Table 1 and data not shown).

The expression of Wnt9b target genes in the progenitor population was unexpected. Previous studies have suggested that Wnt9b was only active in the subpopulation of mesenchymal progenitor cells that was induced to differentiate and blocked in the remaining progenitors to allow expansion (Carroll et al., 2005; Kobayashi et al., 2008; Self et al., 2006). We therefore investigated the molecular basis of the progenitor signaling.

### **β-Catenin signaling induces the expression of Wnt9b target genes**


Wnt signals can be transduced down β-catenin-dependent or -independent pathways. Although previous studies have suggested that Wnt9b signaled through β-catenin to induce tubule differentiation, this has not been directly demonstrated (Park et al., 2007). The identification of cell type-specific Wnt9b target genes allows us to address more precisely Wnt9b pathway usage in both the PTAs and the progenitor cells.

To determine which pathway was responsible for the activation of Class I and II target genes, we used ex vivo organ culture. E11.5 metanephric mesenchyme was isolated and cultured in the presence or absence of the *Gsk-3beta* antagonist LiCl (Klein and Melton, 1996), which results in stabilization of β-catenin. Mesenchymes were cultured for 48 hours in 15 mM LiCl, a treatment that effectively induces tubulogenesis in cultured kidneys (Davies and Garrod, 1995), and assayed for expression of Wnt9b targets. As expected, LiCl treatment was sufficient to induce the expression of the Class I genes *C1qdc2* and *Pax8*, whereas DMEM alone was not (Fig. 2A,B; Table 1). Next, we evaluated the expression of Class II genes. Mesenchymes cultured with LiCl expressed every Wnt9b progenitor target tested, including *Cited1*, *Pla2g7* and *Tafa5*, whereas mesenchyme cultured in DMEM alone exhibited no



**Fig. 1. Identification of novel Wnt9b target genes in kidney progenitor cells.**

(A-O) In situ hybridization evaluating the expression of *C1qdc2* (A-C), *Cited1* (D-F), *Pla2g7* (G-I), *Tafa5* (J-L) or *Uncx4.1* (M-O) in wild-type (A,D,G,J,M), *Wnt9b*<sup>-/-</sup> (B,E,H,K,N) or *Wnt4*<sup>-/-</sup> (C,F,I,L,O) animals at E11.5. In all images, the pretubular aggregates are located on the left, whereas the progenitor cells are on the right. The ureteric bud tips are outlined.



**Fig. 2.  $\beta$ -Catenin signaling is sufficient and necessary for Wnt9b progenitor target gene expression.** (A-R) In situ hybridization evaluating the expression of *C1qdc2* (A,B,G-I), *Pla2g7* (C,D,J-L) and *Tafa5* (E,F,M-O) and *Uncx4.1* (P-R). A-F represent E11.5 metanephric mesenchyme cultured for 48 hours in media alone (A,C,E) or in media containing 15 mM LiCl (B,D,F). (G-R) Embryonic kidneys grown in the presence of DMSO (G,J,M,P), 5  $\mu$ M IWP2 (H,K,N,Q) or 100  $\mu$ M IWR1 (I,L,O,R).

expression (Fig. 2C-F; Table 1). These data show that activation of  $\beta$ -catenin is sufficient to stimulate expression of both Class I and II Wnt9b target genes.

### $\beta$ -Catenin activity is necessary for expression of Wnt9b targets

The studies listed above suggest that  $\beta$ -catenin activation is sufficient to activate expression of Class I and II targets. We next sought to determine whether it was necessary. Recently identified small molecule inhibitors of the Wnt pathway have been used to study the role of Wnt signaling in kidney organ culture (Chen et al., 2009; Huang et al., 2009; Karner et al., 2010). To test pathway requirements in the activation of Wnt9b targets, we used two individual inhibitors, IWP2 and IWR1, that function at distinct points in the Wnt pathway. IWP2 inhibits porcupine, a molecule necessary for Wnt ligand secretion (Kadowaki et al., 1996). Application of this molecule blocks both canonical and non-canonical Wnt signaling. IWR1 specifically inhibits the canonical/ $\beta$ -catenin pathway through inhibition of tankyrase 1 and 2 (Chen et al., 2009; Huang et al., 2009). Both of these factors act specifically and effectively in the context of kidney organ culture (Karner et al., 2010).

E11.5 wild-type kidneys were cultured for 24, 48 or 96 hours in the presence of either DMSO or IWP2 and assayed for the expression of Wnt9b targets. As expected, the class I targets *C1qdc2* and *Pax8*, and the class II targets *Cited1*, *Pla2g7*, *Tafa5* and *Uncx4.1* were all expressed in the induced mesenchyme of kidneys cultured in DMSO (Fig. 2G,J,M,P; Table 1). However, inhibition of Wnt ligand secretion using IWP2 resulted in a significant reduction of all Wnt9b target genes examined (Fig. 2, compare G with H, J with K, M with N and P with Q; data not shown). This was not a delay in kidney development as markers were still undetectable after 96 hours of culture (Karner et al., 2010). These data validate the efficacy of this approach and set a baseline for further comparison.

The data above verify that Wnt production is necessary for the expression of Wnt9b target genes. To test whether the loss of the target genes can be specifically attributed to inhibition of the canonical Wnt pathway, we cultured kidneys in the presence of IWR1. IWR1 administration leads to the complete inhibition of the Class I Wnt9b target gene *Pax8* within 16 hours (Karner et al., 2010) (data not shown). Similarly, IWR1 co-culture inhibited the expression of other class I genes, including *C1qdc2* and *Cdh4*, and all Class II target genes tested, including *Cited1*, *Pla2g7*, *Tafa5* and *Uncx4.1* (Fig. 2, compare G with I, J with L, M with O and P with R; Table 1). The kinetics of gene loss after treatment with IWR1 were indistinguishable for Class I and II targets. Together, these data suggest that  $\beta$ -catenin-dependent Wnt signaling is necessary for expression of both Class I and II Wnt9b target genes.

### Wnt9b/ $\beta$ -catenin cell autonomously activates progenitor target genes

Although the data presented above indicate that  $\beta$ -catenin is required for the expression of Wnt9b target genes in the progenitors, treatment of isolated mesenchymes with LiCl or the IW inhibitors does not provide the cell type specificity to determine the precise cellular targets of pathway activation. These compounds could be affecting Wnt4 activity in the PTAs, which might secondarily affect expression of target genes in the progenitors.


To test the possibility that a signal(s) from the PTAs regulates expression of genes in the progenitor cells, we evaluated the expression of Class I and II Wnt9b targets in E11.5 *Wnt4*<sup>-/-</sup> kidneys (Stark et al., 1994). Not surprisingly, 6/12 class I targets (PTA markers) were completely absent from *Wnt4*<sup>-/-</sup> metanephric mesenchyme (MM) (Table 1), suggesting that they were indirect targets of Wnt9b identified due to the loss of Wnt4 activity in *Wnt9b* mutants. However, the majority of Wnt9b target genes, including the Class I targets *C1qdc2* and *Cadh4*, and the Class II targets *Cited1*, *Pla2G7*, *Tafa5* and *Uncx4.1*, were still expressed in *Wnt4*<sup>-/-</sup> MM at E11.5 (Fig. 1, compare A with C, D with F, G with I, J with L, M with O; Table 1). In fact, one class II target, *Cited1*, appeared to be expanded into what would normally be the PTA domain in *Wnt4* mutants. This expansion most probably reflects a failure to induce differentiation in the mutant progenitor cells. *Cited1*, and all other class II targets examined, are still expressed through at least E13.5 (Table 1), even though all Class I targets examined were lost from *Wnt4* mutants by E12.5.

To provide further evidence that  $\beta$ -catenin signaling was required in the progenitors, we determined whether activation of  $\beta$ -catenin in these cells was sufficient to rescue the expression of progenitor targets in a *Wnt9b* mutant. To accomplish this, we expressed an activated allele of  $\beta$ -catenin [*catnb1*<sup>exon3lox</sup> (Harada et al., 1999)] in the progenitor cells of *Wnt9b* mutants using the *Rar $\beta$ 2Cre* transgene (Kobayashi et al., 2005). *Rar $\beta$ 2Cre* is active throughout the


nephrogenic mesenchyme (including the mesonephric and metanephric mesenchyme) from E10.5 through birth as well as in a subset of the cortical stroma at E12.5 (Kobayashi et al., 2005). Activation of  $\beta$ -catenin with *Rarb2Cre* in otherwise wild-type animals resulted in an anterior expansion of all evaluated genes, including *C1qdc2*, *Tafa5* and *Pla2g7*, relative to wild-type littermates (Fig. 3A,B,E,F,I,J). This expansion is probably reflective of the expression of *Rarb2cre* in the mesonephric mesenchyme. Consistent with this hypothesis, we also observed ectopic expression of *C1qdc2*, *Tafa5* and *Pla2g7* in the mesonephric tubules in *Rarb2Cre;catnb1<sup>exon3floxed</sup>* animals (data not shown).

To test whether activation of  $\beta$ -catenin was sufficient to drive expression of *Wnt9b* target genes in the absence of *Wnt9b*, we generated *Rarb2Cre;catnb<sup>exon3floxed</sup>;Wnt9b<sup>-/-</sup>* mice. Activation of  $\beta$ -catenin in the *Rarb2Cre* expression domain of *Wnt9b* mutants was sufficient to rescue the expression of both the class I target gene *C1qdc2* (compare Fig. 3C with 3D) and the class II targets *Tafa5* and *Pla2g7* (compare Fig. 3G with 3H and 3K with 3L).

We next asked whether  $\beta$ -catenin activity was required in the progenitors for the expression of *Wnt9b* targets using a floxed null allele of  $\beta$ -catenin (Brault et al., 2001) and the *Rarb2Cre* strain (Kobayashi et al., 2005). We found that most *Rarb2Cre; $\beta$ -catenin<sup>-floxed</sup>* pups (8/10) had severely hypoplastic kidneys and died within 2 days of birth (not shown). At E12.5, *Rarb2Cre;Catnb<sup>-floxed</sup>* kidneys appeared normal or were slightly smaller than wild type (Fig. 4A-D). Both *Pla2g7* and *Tafa5* were significantly down regulated in the MM of *Rarb2Cre;Catnb<sup>-floxed</sup>* animals at E12.5, with some areas showing a complete loss of expression (arrows in Fig. 4C,D). Immunofluorescence with a  $\beta$ -catenin antibody revealed mosaic expression of  $\beta$ -catenin protein in mutants at E12.5, indicating that recombination with this line was inefficient (data not shown). By E13.5, we were able to find large domains of progenitors (as marked by *Six2* protein) that lacked detectable levels of  $\beta$ -catenin protein (Fig. 4E-H). Although mutant cells continued to express markers of the progenitor domains (such as *Six2*), they showed greatly reduced protein levels for two Class II



**Fig. 3.  $\beta$ -Catenin signaling regulates *Wnt9b* progenitor target gene expression in vivo.** (A-L) In situ hybridization evaluating the expression of *C1qdc2* (A-D), *Tafa5* (E-H) and *Pla2g7* (I-L) in wild-type (A,E,I), *Rarb2Cre;Catnb<sup>ex3floxed</sup>* (B,F,J), *Wnt9b<sup>-/-</sup>* (C,G,K) and *Rarb2Cre;Catnb<sup>ex3floxed</sup>;Wnt9b<sup>-/-</sup>* (D,H,L) animals at E11.5. In all images, the Wolffian duct is located towards the left while the progenitor cells are towards the right. The ureteric bud tips are outlined. Direct genetic stabilization of  $\beta$ -catenin in the mesenchymal progenitors is sufficient to induce expression of *Wnt9b* targets, even in the absence of *Wnt9b*. (M-O) Quantification of ChIP with  $\beta$ -catenin antibody on E15.5 kidneys. Enriched enhancer sequences containing *Lef/TCF* sites are indicated by red ovals. Non-enriched sites are indicated by blue ovals. The graph shows fold increase relative to anti-IgG control. Data are mean  $\pm$  s.e.m.  $\beta$ -Catenin protein is associated with conserved *Lef/TCF* binding sites in the enhancers of at least three of the *Wnt9b* target genes.



**Fig. 4.  $\beta$ -Catenin is required cell-autonomously for the expression of progenitor markers.** (A–J) Expression analysis of Wnt9b target genes in wild-type (A, B, E–G) or *Rarb2Cre;catnb<sup>flox/-</sup>* mutant (C, D, H–J) E12.5 kidneys. A and C are images of whole-mount kidneys hybridized with antisense probes to *Tafa5* (A, C) or *Pla2g7* (B, D). E–J are images of sections stained with antibodies to  $\beta$ -catenin (red), *Six2* (green in E, H), *Cited1* (green in F, I) and amphiphysin (green in G, J). All sections were counterstained with the nuclear marker To-Pro 3 (blue). Arrows indicate progenitor cells. Removal of  $\beta$ -catenin from the progenitors results in cell-autonomous loss of Wnt9b target genes.

Wnt9b targets, amphiphysin and *Cited1* (Fig. 4I, J). In some cases, we were able to find small clusters of wild-type cells expressing the Class II targets that were surrounded by mutant cells, suggesting that loss of expression was cell-autonomous (data not shown). The loss and gain of expression studies suggest that  $\beta$ -catenin activity in the progenitors is necessary and sufficient for the expression of Class II Wnt9b target genes in the progenitors.

### **$\beta$ -Catenin is associated with progenitor gene promoters**

Collectively, our data supports the hypothesis that Wnt9b signals to both the PTAs and progenitor cells via  $\beta$ -catenin. However, the data still do not indicate whether the Wnt9b target genes identified here are direct targets of  $\beta$ -catenin. To identify prospective  $\beta$ -catenin regulatory elements, we scanned the genomic loci of Wnt9b targets using the genome alignment tool in the ECR

browser (<http://ecrbrowser.dcode.org>) looking for consensus Lef/Tcf-binding sites [5'-(A/T)(A/T)CAAAG-3']. Of the genes evaluated, most (23/25) contained multiple consensus Lef/Tcf-binding sites within 50 kb of the transcriptional start (Table 1).


Previous studies showed that  $\beta$ -catenin was associated with the promoter of at least one of our Class I targets, *Pax8* (Schmidt-Ott et al., 2007). Examination of another Class I target, *C1qdc2*, revealed that it contained three conserved Lef/Tcf-binding sites. Two of the sites were located in the first intron, while a third site was located downstream of the 3' UTR (Fig. 3M). To determine whether  $\beta$ -catenin was physically associated with these sites, we performed chromatin immunoprecipitation (ChIP) with an antibody to  $\beta$ -catenin and real time-PCR with primers flanking each of these sites. ChIP against  $\beta$ -catenin significantly enriched for the DNA surrounding Lef/Tcf-binding sites 1 and 2 in *C1qdc2* over a control (IgG) IP (6.3±0.8 and 2.8±0.9 fold, respectively) (Fig. 3M). No enrichment was observed for site 3 (not shown).

We next evaluated the genomic loci of the class II targets *Tafa5* and *Pla2g7*. *Tafa5* contained three conserved Lef/Tcf sites. The first site was located ~20 kb upstream of the first exon, while sites 2 and 3 were located in the second intron. Again, using site-specific primers, we found that the second site was not enriched in the  $\beta$ -catenin precipitated chromatin (relative to IgG control), whereas the first and third sites were (3.1±0.7 and 6.1±1.6 fold, respectively) (Fig. 3N). *Pla2g7* contained two conserved Lef/Tcf sites. The first site was located upstream of the 5' UTR, whereas the second site was located in the first intron. Both site 1 and 2 were enriched by  $\beta$ -catenin precipitation (5.5±0.4 and 5.1±0.2 fold relative to control, respectively), while a randomly chosen site upstream of the 5' UTR that did not contain a consensus site was not enriched (Fig. 3O). These data indicate that  $\beta$ -catenin is physically associated with DNA surrounding the loci of both classes of Wnt9b target genes. Based on these studies, as well as the genetic and ex vivo culture data, we propose that Wnt9b signals through  $\beta$ -catenin to both the pre-tubular aggregates and the progenitor cells.

### **Wnt9b and *Six2* act cooperatively to regulate expression of target genes**

Previous studies suggested that *Six2* inhibited Wnt9b and  $\beta$ -catenin signaling within the progenitor cells (Kobayashi et al., 2008). Loss of *Six2* resulted in the expansion of several Wnt9b target genes including *Pax8*, *Sfrp2* and *Wnt4*. However, our data suggest that Wnt9b actively signals to the *Six2*-expressing cells. To gain insight into these seemingly contradictory results, we determined what effect loss of *Six2* had on our newly identified Wnt9b target genes. As expected, in situ hybridization revealed that, similar to previous observations (Kobayashi et al., 2008; Self et al., 2006), several Class I targets of Wnt9b, including *Pax8*, *Wnt4* and *C1qdc2*, were expanded into the progenitor domain in *Six2<sup>-/-</sup>* mesenchyme at E11.5 [see Fig. 5A, B, Kobayashi et al. (Kobayashi et al., 2008) and Self et al. (Self et al., 2006)].

We next evaluated the expression of Class II targets in *Six2<sup>-/-</sup>* mutants. Somewhat surprisingly, we found that the Wnt9b targets *Cited1*, *Pla2g7* and *Tafa5* were absent from *Six2<sup>-/-</sup>* mesenchyme at E11.5 (Fig. 5C–H). Importantly, the loss of Wnt9b targets did not reflect a loss of the progenitor domain as several Wnt9b-independent progenitor markers, including *Eya1* and *Pax2*, were still present in both *Wnt9b* and *Six2* mutants at this stage (Fig. 5I–K) (Carroll et al., 2005; Kobayashi et al., 2008; Self et al., 2006). Furthermore, *Tafa5* mRNA is absent from the mesenchyme of both *Wnt9b<sup>-/-</sup>* and *Six2<sup>-/-</sup>* embryos at E10.5 (see Fig. S2 in the



**Fig. 5. Wnt9b and Six2 cooperate to regulate target gene expression.** (A-P) In situ hybridization evaluating the expression of *C1qdc2* (A,B), *Cited1* (C,D), *Pla2g7* (E,F), *Tafa5* (G,H), *Eya1* (I-L) and *Gdnf* (M-P) in E11.5 wild-type (A,C,E,G,I,M), *Six2*<sup>-/-</sup> (B,D,F,H,K,O), *Wnt9b*<sup>-/-</sup> (J,N) and *Wnt9b*<sup>-/-</sup> *Six2*<sup>-/-</sup> (L,P) animals at E11.5. In all images, the Wolffian duct is located towards the left while the progenitor cells are towards the right. The ureteric bud tips are outlined. Six2 is necessary for the expression of Class II Wnt9b target genes (D,F,H) but represses Class I targets (B). Wnt9b and Six2 genetically interact to regulate expression of Class II genes (M-P), and the loss of Class II gene expression in Wnt9b and Six2 mutants is not caused by loss of the progenitor population (I-L).

supplementary material), a time point prior to the manifestation of morphological defects in either mutant (Carroll et al., 2005; Self et al., 2006). These results suggest that Wnt9b and Six2 are both required for the expression of Wnt9b target genes in the progenitor population.

In both *Wnt9b* and *Six2* mutants, the initial branching of the ureteric bud to form a T shape occurs normally (Carroll et al., 2005; Self et al., 2006). By contrast, E11.5 *Wnt9b*<sup>-/-</sup>; *Six2*<sup>-/-</sup> ureteric buds failed to branch within the metanephric mesenchyme (Kobayashi et al., 2008) (Fig. 5P). Branching of the ureteric bud is mediated by several factors expressed in the progenitor cells, including glial-derived neurotrophic factor (*Gdnf*). Although *Gdnf* mRNA is expressed in the mesenchyme of *Wnt9b*<sup>-/-</sup> and *Six2*<sup>-/-</sup> embryos at E11.5, the domain of expression is slightly reduced in both mutants relative to wild type (Fig. 5M-O). To determine

whether the reduced branching phenotype was caused by a further decrease in *Gdnf* levels, we examined its expression in *Wnt9b*<sup>-/-</sup>; *Six2*<sup>-/-</sup> kidneys. *Gdnf* expression was completely absent from the mesenchyme of *Wnt9b*<sup>-/-</sup>; *Six2*<sup>-/-</sup> animals at E11.5 (Fig. 5P). To ensure that the loss of *Gdnf* was not due to a loss of progenitor cells, we evaluated the expression of an additional progenitor marker, *Eya1*, the expression of which is independent of Wnt9b and Six2 activity. *Eya1* was expressed in the mesenchyme of *Wnt9b*<sup>-/-</sup>; *Six2*<sup>-/-</sup> animals at E11.5 (Fig. 5L). These data indicate that Six2 and Wnt9b act cooperatively within the mesenchymal progenitor cells to activate expression of target genes.

### Wnt9b regulates progenitor cell proliferation

The data presented above suggest that canonical Wnt9b signaling is active in the progenitor cells. Previous studies have demonstrated roles for Wnt signaling in progenitor cell specification, survival/maintenance and renewal/proliferation (Chenn and Walsh, 2002; Reya et al., 2003; Zechner et al., 2003). As mentioned *Six2*, *Eya1*, *Gdnf* and *Pax2* are all expressed in Wnt9b mutant mesenchyme at E11.5. *Pax2* and *Six2* are still expressed in Wnt9b mutants at E13.5, although relative to wild type, the domain of expression is greatly reduced (see Fig. S3 in the supplementary material). Furthermore, heterochronic recombination and culture of the E13.5 mutant mesenchyme with wild-type E11.5 ureteric bud results in activation of class II targets *Cited1* and Class I targets *Pax8* and *C1qdc2* (see Fig. S3 in the supplementary material and data not shown). Together, these results indicate that the Wnt9b mutant progenitors are properly specified and retain competence to respond to the inductive signal.


The failure of the progenitor population to expand in Wnt9b mutants suggested that Wnt9b might regulate apoptosis or cell proliferation of the mesenchyme. Using Lyso-tracker, we were unable to detect differences in the rate of apoptosis between wild-type and Wnt9b-mutant mesenchymal cells at E11.5 (see Fig. S4 in the supplementary material). By contrast, wild-type E11.5 mesenchymal cells had a rate of proliferation approximately five times greater than those of Wnt9b mutants [2.48% versus 0.48% of cells positive for phospho-histone H3 staining;  $P=0.007$ ,  $t$ -test,  $n=3$  animals and 2376 or 2122 total cells for wild type and Wnt9b null, respectively (see Fig. S4 in the supplementary material)].

To ensure the effect on proliferation was a direct result of loss of Wnt9b and not a secondary consequence of failure to induce renal vesicles, we also evaluated proliferation in *Wnt4*<sup>-/-</sup> mesenchyme. Although loss of Wnt4 also resulted in a significant decrease in proliferation (3.12% and 1.81% in wild type and null, respectively;  $P=0.006$ ,  $t$ -test,  $n=3$  animals and 2922 or 2076 total cells for wild type and Wnt4 null), the rate of proliferation in Wnt4 mutants was much higher than observed in Wnt9b mutants. These data suggest that Wnt9b regulates proliferation/renewal of the mesenchymal progenitor cells.

### Ablation of Wnt9b results in premature exhaustion of the progenitor cells

Wnt9b is expressed in mouse ureteric bud/collecting ducts through the embryonic period and into adult stages (Karner et al., 2009). However, despite continued presence of active Wnt9b ligand, renal tubule formation ceases by postnatal (P) day 5 (Hartman et al., 2007). For most of the developmental period, progenitor cell renewal outpaces differentiation. However, around P3, the rate of progenitor cell expansion decreases, and by P5, the progenitor





**Fig. 6. Removal of Wnt9b results in a premature loss of progenitor cells.** (A-P) In situ hybridization evaluating the expression of *Six2* (A,B,I,J), *Pla2g7* (C,D,K,L), *Tafa5* (E,F,M,N) and *Uncx4.1* (G,H,O,P) in wild-type (A,C,E,G,I,K,M,O) or *KspCre;Wnt9b<sup>-/-</sup>* (B,D,F,H,J,L,N,P) kidneys at E17.5 (A-H) or P1 (I-P). Wnt9b is necessary for the maintenance of the progenitor population at later stages.

population is exhausted after a wave of differentiation (Hartman et al., 2007). A reasonable hypothesis based on the data presented in this study is that a loss in ability of the progenitor cells to respond to Wnt9b may play a causal role in the failure of progenitor cells to self-renew postnatally.

To test this hypothesis, we compared the expression of Wnt9b target genes to the progenitor expression of *Six2* in wild-type mice from P3 to P4. As expected, *Six2* mRNA and protein were present in P3 kidneys, although the progenitor expression is reduced and, in some cases, *Six2* appeared to be only expressed in the PTAs (see Fig. S5A in the supplementary material). This most probably represents the period where the last progenitors are being converted to tubules. *Six2* mRNA and protein were completely undetectable by P4 (see Fig. S5B in the supplementary material). The Wnt9b target genes *Pla2g7* and *Cited1* were both present in the progenitor cells at P3 but completely absent (for *Cited1*) or present only in epithelial structures (for *Pla2g7*) at P4, indicating that, although Wnt9b was still expressed, it no longer actively signaled to the progenitor population (see Fig. S5C-F in the supplementary material). Importantly, the class I targets *Clqdc2* and *Pax8* were still expressed at P4, a full day after *Cited1* was lost, indicating that Wnt9b is still active and inducing tubule formation (see Fig. S5H in the supplementary material and not shown). These data suggest that a loss of Wnt9b signaling to the progenitor cells normally coincides with or immediately follows loss of *Six2* expression in the progenitors.

We have previously shown that removal of Wnt9b from the kidney between E15.5 and E17.5 resulted in significantly smaller kidneys than in wild type (Karner et al., 2009). We originally attributed this hypoplastic phenotype to a deficit in tubule induction; however, our new findings suggest this phenotype may be equally attributable to a deficit in progenitor cell expansion. To test this hypothesis, we assessed progenitor cell maintenance/expansion in *KspCre;Wnt9b<sup>-/-</sup>* mice. Prior to E15.5, KspCre is active only in the distal collecting ducts and has no apparent effect on the expression of Wnt9b target genes (Karner et al., 2009) (data not shown). At some time between E15.5 and E17.5, KspCre activity expands to include the ureteric bud tips (adjacent to

the progenitors). We therefore examined the expression of Wnt9b-dependent and -independent progenitor markers in *KspCre;Wnt9b<sup>-/-</sup>* kidneys between E15.5 and P1. *Pax2*, *Six2* and the Wnt9b target genes *Pla2g7*, *Tafa5* and *Uncx4.1* were all present and expressed at comparable levels with wild-type kidneys at E15.5 (data not shown). At E17.5, *Pax2* and *Six2* levels were reduced in the progenitor cells of *KspCre;Wnt9b<sup>-/-</sup>* animals (Fig. 6A,B; data not shown), whereas *Pla2g7* and *Uncx4.1* were completely lost (Fig. 6C,D,G,H). Levels of *Tafa5* were highly reduced at E17.5 and completely absent at P1 (Fig. 6E,F,M,N). The loss of expression for these genes was specific to progenitor cells as *Pla2g7* expression in the medullary stroma was maintained in mutants (not shown). By P1, the progenitor population (as indicated by *Six2* expression) appeared to be completely lost (Fig. 6I-P).

We next tested the effects of late Wnt9b ablation on progenitor cell proliferation. At E17.5, there was a significant reduction in cell proliferation rates in Wnt9b mutants (2.44% and 1.75% for wild type and *KspCre;Wnt9b<sup>-/-</sup>*, respectively;  $P=0.027$ ). However, as the progenitor cells in Wnt9b mutants are not being converted into new tubules (as determined by *Clqdc2* expression) decreases in proliferation alone cannot explain the complete absence of the progenitor population by P1. We also evaluated the rate of apoptosis in *KspCre;Wnt9b<sup>-/-</sup>* kidneys at E17.5. Somewhat surprisingly, this analysis revealed a significant increase in cell death upon loss of Wnt9b (1.62% and 26.83% apoptosis in wild-type and *KspCre;Wnt9b<sup>-/-</sup>* animals, respectively,  $P=0.0039$ ,  $t$ -test), even though apoptosis was not affected by Wnt9b loss at earlier stages. These data suggest that Wnt9b plays an additional (most probably secondary) role in survival of the progenitor population.

## DISCUSSION

In this study, we present data suggesting that Wnt9b is actively signaling to the kidney progenitor cells where it cooperates with *Six2* to mediate expansion and/or self-renewal. Multiple pieces of data suggest that the role for Wnt9b in the progenitors is direct and through the canonical/ $\beta$ -catenin-dependent pathway. First, we show

that canonical activity is required for normal expression of progenitor targets. Second, we show that activation of the canonical pathway is sufficient to induce expression of these genes, even in the absence of Wnt9b. That this activation is not mimicking signaling downstream of Wnt4 (and tubule differentiation) is supported by the observation that progenitor targets are expressed normally in *Wnt4* mutants and are lost prior to the cessation of renal vesicle formation (and Wnt4 loss) in wild-type post-natal kidneys. Finally, several Wnt9b progenitor target genes (11/13) contain highly conserved consensus Lef/Tcf-binding sites within their loci and several of these elements are associated with  $\beta$ -catenin in vivo.

Our data suggest that Wnt9b signals through  $\beta$ -catenin to both cell populations with distinct cellular and molecular results. In the progenitors, it promotes renewal/proliferation, whereas in the PTAs it promotes differentiation. Reiterative use of  $\beta$ -catenin in phases of stem cell activation and differentiation has been shown in other systems, most notably the hair follicle stem cells (Lowry et al., 2005). How does Wnt9b induce disparate responses (self-renewal versus differentiation) in different cell types using the same signal transduction cascade? Our data suggest that, in the kidney, *Six2* is an important player in this decision.

A simple model that explains all of the genetic and molecular analysis in this and previous studies is that a combination of  $\beta$ -catenin and *Six2* results in progenitor renewal, while  $\beta$ -catenin alone (or in combination with another factor) results in differentiation. Cells that express *Six2* and receive a Wnt9b signal are induced to proliferate and to maintain the progenitor pool, while cells that receive the Wnt9b/ $\beta$ -catenin signal but do not have active *Six2* (and/or do express another, unknown factor) are induced to differentiate.

The molecular nature of the interaction between  $\beta$ -catenin and *Six2* is unclear. It is plausible that *Six2* regulates the expression of a separate set of progenitor specific factors that interacts with and/or alters the response to Wnt9b/ $\beta$ -catenin, perhaps amplifying or dampening the signal, as has been suggested in the hair follicles. It is also possible that *Six2* directly (or indirectly) interacts with  $\beta$ -catenin to drive expression of individual genes. Determining which, if any, of the above scenarios exists within the kidney mesenchyme will certainly enhance our understanding of progenitor cell differentiation during normal development, as well as in pathological states.

It is important to note that, although levels are highest in the progenitors, *Six2* also shows low level expression in the PTAs. It is possible that *Six2* cooperates with  $\beta$ -catenin in both cell types to activate expression of target genes. This is consistent with our finding that many of the Wnt9b targets are expressed in both the progenitors and the PTAs, and are completely lost in *Six2* mutants. However, some, such as *Cited1*, are expressed only in the progenitors whereas others, such as *C1qdc2*, are expressed only in the PTAs (and are expanded in *Six2* mutants). These observations suggest that there are additional cell type-specific transcriptional regulators that cooperate with *Six2* and  $\beta$ -catenin to drive expression in the various cell types.

In summary, we have identified a novel mechanism by which renal progenitor cells respond to the same Wnt9b signal with distinct results: self-renewal versus differentiation. By using this relatively simple mechanism, the kidney is able to balance tubule induction and progenitor cell maintenance, insuring that the proper numbers of nephrons form. These findings are highly relevant to human diseases that affect kidney progenitors, such as aplasia, hypoplasia, decreased nephron endowment and renal progenitor

tumors. Furthermore, they provide novel insights into the molecular mechanisms that regulate kidney progenitor cell differentiation and will enhance future attempts to repair or replace missing or damaged organs.

#### Acknowledgements

We thank members of the Carroll, Cleaver and Igarashi labs for their comments on this work. This work was supported by grants from the American Heart Association (0730236N), the NIH (R01DK080004) and the Susan Irene Simons Kidney Cancer Foundation to T.C.; by grants from the NIGMS (R01GM079554) and Welch Foundation (Welch I-1596) to C.C.; and by the NIH (R01GM076398) and Welch Foundation to L.L. Deposited in PMC for release after 12 months.

#### Competing interests statement

The authors declare no competing financial interests.

#### Supplementary material

Supplementary material for this article is available at <http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.057646/-/DC1>

#### References

- Barasch, J., Yang, J., Ware, C. B., Taga, T., Yoshida, K., Erdjument-Bromage, H., Tempst, P., Parravicini, E., Malach, S., Aranoff, T. et al. (1999). Mesenchymal to epithelial conversion in rat metanephros is induced by LIF. *Cell* **99**, 377-386.
- Blank, U., Brown, A., Adams, D. C., Karolak, M. J. and Oxburgh, L. (2009). BMP7 promotes proliferation of nephron progenitor cells via a JNK-dependent mechanism. *Development* **136**, 3557-3566.
- Boyle, S., Misfeldt, A., Chandler, K. J., Deal, K. K., Southard-Smith, E. M., Mortlock, D. P., Baldwin, H. S. and de Caestecker, M. (2008). Fate mapping using *Cited1-CreERT2* mice demonstrates that the cap mesenchyme contains self-renewing progenitor cells and gives rise exclusively to nephronic epithelia. *Dev. Biol.* **313**, 234-245.
- Braut, V., Moore, R., Kutsch, S., Ishibashi, M., Rowitch, D. H., McMahon, A. P., Sommer, L., Boussadia, O. and Kemler, R. (2001). Inactivation of the beta-catenin gene by Wnt1-Cre-mediated deletion results in dramatic brain malformation and failure of craniofacial development. *Development* **128**, 1253-1264.
- Carroll, T. J., Park, J. S., Hayashi, S., Majumdar, A. and McMahon, A. P. (2005). Wnt9b plays a central role in the regulation of mesenchymal to epithelial transitions underlying organogenesis of the mammalian urogenital system. *Dev. Cell* **9**, 283-292.
- Chen, B., Dodge, M. E., Tang, W., Lu, J., Ma, Z., Fan, C. W., Wei, S., Hao, W., Kilgore, J., Williams, N. S. et al. (2009). Small molecule-mediated disruption of Wnt-dependent signaling in tissue regeneration and cancer. *Nat. Chem. Biol.* **5**, 100-107.
- Chenn, A. and Walsh, C. A. (2002). Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. *Science* **297**, 365-369.
- Davies, J. A. and Garrod, D. R. (1995). Induction of early stages of kidney tubule differentiation by lithium ions. *Dev. Biol.* **167**, 50-60.
- Dudley, A. T., Lyons, K. M. and Robertson, E. J. (1995). A requirement for bone morphogenetic protein-7 during development of the mammalian kidney and eye. *Genes Dev.* **9**, 2795-2807.
- Grieshammer, U., Cebrian, C., Ilagan, R., Meyers, E., Herzlinger, D. and Martin, G. R. (2005). FGF8 is required for cell survival at distinct stages of nephrogenesis and for regulation of gene expression in nascent nephrons. *Development* **132**, 3847-3857.
- Harada, N., Tamai, Y., Ishikawa, T., Sauer, B., Takaku, K., Oshima, M. and Taketo, M. M. (1999). Intestinal polyposis in mice with a dominant stable mutation of the beta-catenin gene. *EMBO J.* **18**, 5931-5942.
- Hartman, H. A., Lai, H. L. and Patterson, L. T. (2007). Cessation of renal morphogenesis in mice. *Dev. Biol.* **310**, 379-387.
- Hoy, W. E., Douglas-Denton, R. N., Hughson, M. D., Cass, A., Johnson, K. and Bertram, J. F. (2003). A stereological study of glomerular number and volume: preliminary findings in a multiracial study of kidneys at autopsy. *Kidney Int.* **63**, S31-S37.
- Huang, S. M., Mishina, Y. M., Liu, S., Cheung, A., Stegmeier, F., Michaud, G. A., Charlat, O., Wiellette, E., Zhang, Y., Wiessner, S. et al. (2009). Tankyrase inhibition stabilizes axin and antagonizes Wnt signalling. *Nature* **461**, 614-620.
- Kadowaki, T., Wilder, E., Klingensmith, J., Zachary, K. and Perrimon, N. (1996). The segment polarity gene *porcupine* encodes a putative multitransmembrane protein involved in Wingless processing. *Genes Dev.* **10**, 3116-3128.
- Karner, C. M., Chirumamilla, R., Aoki, S., Igarashi, P., Wallingford, J. B. and Carroll, T. J. (2009). Wnt9b signaling regulates planar cell polarity and kidney tubule morphogenesis. *Nat. Genet.* **41**, 793-799.

- Karner, C. M., Merkel, C. E., Dodge, M., Ma, Z., Lu, J., Chen, C., Lum, L. and Carroll, T. J. (2010). Tankyrase is necessary for canonical Wnt signaling during kidney development. *Dev. Dyn.* **239**, 2014-2023.
- Keller, G., Zimmer, G., Mall, G., Ritz, E. and Amann, K. (2003). Nephron number in patients with primary hypertension. *N. Engl. J. Med.* **348**, 101-108.
- Klein, P. S. and Melton, D. A. (1996). A molecular mechanism for the effect of lithium on development. *Proc. Natl. Acad. Sci. USA* **93**, 8455-8459.
- Kobayashi, A., Kwan, K. M., Carroll, T. J., McMahon, A. P., Mendelsohn, C. L. and Behringer, R. R. (2005). Distinct and sequential tissue-specific activities of the LIM-class homeobox gene *Lim1* for tubular morphogenesis during kidney development. *Development* **132**, 2809-2823.
- Kobayashi, A., Valerius, M. T., Mugford, J. W., Carroll, T. J., Self, M., Oliver, G. and McMahon, A. P. (2008). *Six2* defines and regulates a multipotent self-renewing nephron progenitor population throughout mammalian kidney development. *Cell Stem Cell* **3**, 169-181.
- Koesters, R., Ridder, R., Kopp-Schneider, A., Betts, D., Adams, V., Niggli, F., Briner, J. and von Knebel Doeberitz, M. (1999). Mutational activation of the beta-catenin proto-oncogene is a common event in the development of Wilms' tumors. *Cancer Res.* **59**, 3880-3882.
- Koesters, R., Niggli, F., von Knebel Doeberitz, M. and Stallmach, T. (2003). Nuclear accumulation of beta-catenin protein in Wilms' tumours. *J. Pathol.* **199**, 68-76.
- Li, C. M., Kim, C. E., Margolin, A. A., Guo, M., Zhu, J., Mason, J. M., Hensle, T. W., Murty, V. V., Grundy, P. E., Fearon, E. R. et al. (2004). CTNNB1 mutations and overexpression of Wnt/beta-catenin target genes in WT1-mutant Wilms' tumors. *Am. J. Pathol.* **165**, 1943-1953.
- Lowry, W. E., Blanpain, C., Nowak, J. A., Guasch, G., Lewis, L. and Fuchs, E. (2005). Defining the impact of beta-catenin/Tcf transactivation on epithelial stem cells. *Genes Dev.* **19**, 1596-1611.
- Nyengaard, J. R. and Bendtsen, T. F. (1992). Glomerular number and size in relation to age, kidney weight, and body surface in normal man. *Anat. Rec.* **232**, 194-201.
- Park, J. S., Valerius, M. T. and McMahon, A. P. (2007). Wnt/beta-catenin signaling regulates nephron induction during mouse kidney development. *Development* **134**, 2533-2539.
- Perantoni, A. O., Timofeeva, O., Naillat, F., Richman, C., Pajni-Underwood, S., Wilson, C., Vainio, S., Dove, L. F. and Lewandoski, M. (2005). Inactivation of *FGF8* in early mesoderm reveals an essential role in kidney development. *Development* **132**, 3859-3871.
- Reya, T., Duncan, A. W., Ailles, L., Domen, J., Scherer, D. C., Willert, K., Hintz, L., Nusse, R. and Weissman, I. L. (2003). A role for Wnt signalling in self-renewal of haematopoietic stem cells. *Nature* **423**, 409-414.
- Schmidt-Ott, K. M., Masckauchan, T. N., Chen, X., Hirsh, B. J., Sarkar, A., Yang, J., Paragas, N., Wallace, V. A., Dufort, D., Pavlidis, P. et al. (2007). {beta}-catenin/TCF/Lef controls a differentiation-associated transcriptional program in renal epithelial progenitors. *Development* **134**, 3177-3190.
- Self, M., Lagutin, O. V., Bowling, B., Hendrix, J., Cai, Y., Dressler, G. R. and Oliver, G. (2006). *Six2* is required for suppression of nephrogenesis and progenitor renewal in the developing kidney. *EMBO J.* **25**, 5214-5228.
- Shao, X., Somlo, S. and Igarashi, P. (2002). Epithelial-specific *Cre/lox* recombination in the developing kidney and genitourinary tract. *J. Am. Soc. Nephrol.* **13**, 1837-1846.
- Stark, K., Vainio, S., Vassileva, G. and McMahon, A. P. (1994). Epithelial transformation of metanephric mesenchyme in the developing kidney regulated by Wnt-4. *Nature* **372**, 679-683.
- Zechner, D., Fujita, Y., Hulsken, J., Muller, T., Walther, I., Taketo, M. M., Crenshaw, E. B., 3rd, Birchmeier, W. and Birchmeier, C. (2003). beta-Catenin signals regulate cell growth and the balance between progenitor cell expansion and differentiation in the nervous system. *Dev. Biol.* **258**, 406-418.
- Zhang, B., Kirov, S. and Snoddy, J. (2005). WebGestalt: an integrated system for exploring gene sets in various biological contexts. *Nucleic Acids Res.* **33**, W741-W748.
- Zucker, R. M., Hunter, E. S., 3rd and Rogers, J. M. (1999). Apoptosis and morphology in mouse embryos by confocal laser scanning microscopy. *Methods* **18**, 473-480.