

CORRECTION

Doublecortin marks a new population of transiently amplifying muscle progenitor cells and is required for myofiber maturation during skeletal muscle regeneration

Ryo Ogawa, Yuran Ma, Masahiko Yamaguchi, Takahito Ito, Yoko Watanabe, Takuji Ohtani, Satoshi Murakami, Shizuka Uchida, Piera De Gaspari, Akiyoshi Uezumi, Miki Nakamura, Yuko Miyagoe-Suzuki, Kazutake Tsujikawa, Naohiro Hashimoto, Thomas Braun, Teruyuki Tanaka, Shin'ichi Takeda, Hiroshi Yamamoto and So-ichiro Fukada

There was an error published in *Development* 142, 51-61.

In the Discussion (p. 57), it was stated that: Pallafacchina et al. reported that Dcx is expressed in neonatal satellite cells but not in adult satellite cells (Pallafacchina et al., 2010). This should have stated: Pallafacchina et al. reported that Dcx is expressed in satellite cells from one-week-old mice and adult mice depleted for dystrophin, where satellite cells are reactivated, but not (or at only very low levels) in resting satellite cells in the wild-type adult or in activated myoblasts in culture (Pallafacchina et al., 2010).

The authors apologise to readers for this mistake.