

Binding of zona pellucida proteins to a boar sperm polypeptide of M_r 53 000 and identification of zona moieties involved

C. R. BROWN and R. JONES

AFRC Institute of Animal Physiology & Genetics Research, Animal Research Station, 307 Huntingdon Road, Cambridge CB3 0JQ, UK

Present address: AFRC Institute of Animal Physiology & Genetics Research, Department of Molecular Embryology, Babraham, Cambridge CB2 4AT, UK

Summary

Experiments have been carried out to identify proteins on boar spermatozoa that bind to components of the zona pellucida. Polypeptides in sodium deoxycholate extracts of boar spermatozoa and in whole seminal plasma have been separated by sodium dodecyl sulphate–polyacrylamide gel electrophoresis, transferred onto nitrocellulose sheet by electroblotting and probed with ^{125}I -labelled heat-solubilized zona pellucida from pig oocytes or ovulated eggs. Zona proteins bound avidly and consistently to a polypeptide of M_r 53 000 on blots of capacitated and noncapacitated sperm and weakly to polypeptides of M_r 67 000, 38 000 and 18 000. On blots of seminal plasma the ^{125}I -

labelled probes bound to two polypeptides of M_r 65 000 and 19–24 000. Identification of the zona proteins that were binding to the aforementioned proteins on blots showed that all the major zona pellucida glycoproteins were involved, including those acquired from oviduct secretions. Binding of ^{125}I -ovulated zona pellucida to the polypeptide of M_r 53 000 also occurred in extracts of testicular and epididymal boar spermatozoa. The results are discussed in relation to sperm–egg recognition in the pig.

Key words: porcine, zona pellucida, sperm, polypeptide, boar.

Introduction

A critical stage in mammalian fertilization is the recognition and binding of spermatozoa to the egg in the conducive environment of the oestrus oviduct. Once this propensity to unite has been acquired it seems logical to assume that the initial attachment of gametes is achieved *via* the interaction of adhesion molecules or receptors at their limiting surfaces. In the female gamete such molecules probably reside at the surface of the zona pellucida, the relatively thick glycoprotein matrix that encompasses the egg, and in sperm they are likely to be associated with the membranes overlying the head. The process of recognition and initial attachment of sperm to the egg is not entirely species specific unlike consolidated binding and sperm penetration through the zona pellucida (Gwatkin, 1977, review). While these broad concepts are widely accepted very little is known about the identity of such recognition molecules or their eventual interaction, although recently some progress has been made in this direction. For instance, in the

mouse (Bleil & Wassarman, 1983) and pig (Sacco, Subramanian & Yurewicz, 1984) pre-exposure of spermatozoa to a component isolated from the zona pellucida has been shown to prevent sperm–egg recognition. There is also good evidence in the mouse that initial binding of spermatozoa to the zona pellucida is promoted by a sperm surface galactosyltransferase with a lectin-like affinity for *N*-acetylglucosamine residues on zona glycoproteins (Shur & Hall, 1982). In the guinea pig a role in sperm binding to the egg has been proposed for a mobile sperm surface antigen which transfers from its position over the posterior head of intact sperm to the inner acrosomal membrane of the acrosome-reacted sperm, presumably there to participate in consolidation of binding (Primakoff, Hyatt & Myles, 1985). Other putative sperm–zona pellucida binding molecules have been identified in sperm of various species by means of the novel approach of probing Western blots of sperm proteins with labelled zona proteins (Sullivan & Bleau, 1985; O’Rand, Matthews, Welch & Fisher, 1985). However, in some cases, e.g. in the pig, zonae

pellucidae invariably have been prepared on a large scale from ovarian oocytes and as such have not been exposed to the milieu of the oestrus oviduct. This may be significant in terms of identification of recognition molecules because it has recently been shown that the oestrus oviduct of the pig contains two glycoproteins which bind firmly to the zona pellucida of the freshly ovulated egg and which remain bound until at least the 2- to 4-cell stage (Brown & Cheng, 1986).

In view of the above points we have attempted to identify molecules on boar spermatozoa that have the ability to recognize zona pellucida moieties. In this paper we compare zonae pellucidae from pig oocytes with those from ovulated eggs in their respective ability to recognize and bind to boar sperm proteins on Western blots and we also identify the zona pellucida moieties involved in recognition.

Materials and methods

Chemicals

Chemicals for electrophoresis and blotting procedures were obtained from Sigma Chemical Co. (London, UK) or from BDH (Poole, Dorset), and the reagent *N*-succinimidyl 3-(4-hydroxy, 5-[¹²⁵I]iodophenyl) propionate and ¹⁴C-methylated molecular weight (M_r) markers from Amersham International, Amersham, Bucks, UK. Sodium deoxycholate (specially purified) was supplied by Fisons Scientific, Loughborough, Leics, UK, and pregnant mare's serum gonadotrophin (PMSG) and human chorionic gonadotrophin (hCG) by Intervet Laboratories Ltd, Cambridge, UK.

Collection of eggs and oocytes

Eggs were recovered surgically from 8- to 10-month-old crossbred (Large White and Landrace) gilts which had been injected on day 16 of the oestrus cycle with 1200 i.u. PMSG followed by 50 i.u. hCG on day 19.5. At 36 h post hCG oviducts were flushed with phosphate-buffered saline (PBS) containing 4 mg ml⁻¹ bovine serum albumin (BSA) (Brown & Cheng, 1986) and recovered eggs were washed thoroughly in PBS with polyvinyl alcohol (0.2 mg ml⁻¹) in place of BSA. Oocytes were aspirated from the follicles of ovaries obtained fresh from a slaughterhouse.

Iodination and solubilization of zonae pellucidae

The zona pellucida of eggs and oocytes was freed of corona radiata cells by gentle pipetting and the investment carefully removed with the aid of a Leitz micromanipulator. To groups of 25–50 isolated zonae, in 10 μ l PBS, was added 2.5 μ Ci *N*-succinimidyl 3-(4-hydroxy, 5-[¹²⁵I]iodophenyl) propionate in 20 μ l 0.1 M-borate buffer pH 8.5 (Bolton & Hunter, 1973). After 15 min at 25°C the zonae were washed thoroughly by centrifugation through five changes of 1 ml PBS (5 min at 1000g). They were then solubilized by heating for 2 h at 70°C in 0.4 ml 0.03 M-sodium carbonate buffer pH 9.2 and stored frozen at -20°C.

Collection and extraction of spermatozoa

To obtain noncapacitated boar spermatozoa 5 ml of sperm-rich fraction of an ejaculate was centrifuged for 15 min at 650g (25°C) and the loosely sedimented cells washed twice with 5 ml PBS. Sperm were capacitated by first washing in 0.9% (w/v) NaCl containing 1 mg ml⁻¹ BSA followed by 4–6 h incubation at room temperature in supplemented Medium 199 (Cheng, 1985). Seminal plasma was cleared of residual spermatozoa and cellular debris by centrifugation at 10000g for 10 min before use. Cauda epididymal spermatozoa were obtained by retrograde flushing of the lumen of the cauda epididymidis with PBS and testicular spermatozoa by puncturing the efferent ductules. They were separated from epididymal and testicular fluids in the same way as ejaculated spermatozoa from seminal plasma. Spermatozoa were collected from the caput epididymidis by slicing the tissue in PBS. After incubating for 30 min supernatant fluids were drained, filtered through four layers of gauze and centrifuged for 30 s at 600g. Supernatants contained spermatozoa with negligible contamination from tissue fragments. Sperm membrane proteins were solubilized by incubating the cells in 0.4% w/v sodium deoxycholate (DOC) in PBS for 30 min at 4°C, and extracts recovered by centrifugation at 10000g for 15 min.

Extraction of other tissues

Fresh tissue (0.5 g) from porcine liver, kidney, spleen, heart and lung was homogenized for 0.5 min at 4°C in 5 ml 0.4% w/v DOC in 20 mM-Tris-HCl pH 8.4. Homogenates were incubated for 30 min at 4°C and extracts recovered by centrifugation at 10000g for 30 min.

Procedure for probing proteins from sperm and seminal plasma on Western blots with ¹²⁵I-zonae pellucidae glycoproteins

Polypeptides in seminal plasma and in DOC extracts of spermatozoa were separated on nonreducing sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) employing a 7.7–14% w/v gradient of polyacrylamide (Laemmli, 1970). Separated proteins were visualized either with Coomassie Brilliant Blue R250 (0.05% w/v) or stained for carbohydrate with the periodic acid Schiff reaction (PAS) (Fairbanks, Steck & Wallach, 1971). Alternatively, proteins were transferred onto nitrocellulose paper (0.45 μ m; Schleicher & Schull Ltd) using the electroblotting method of Towbin (1979). Blots were washed three times in PBS (10 min with shaking) blocked for 1 h at 25°C with 5% w/v BSA in PBS and then probed with ¹²⁵I-zonae pellucidae (approx. 1–2 \times 10⁶ cts min⁻¹ in 0.4 ml) for 3 h at 25°C. The blots were washed three times in PBS which, in some experiments, included Nonidet P-40 (0.5% w/v), and dried. Labelled proteins were detected by autoradiography using preflashed Kodak X-Omat S film.

Several technical problems arose during initial experiments with the above protocol. One was the random appearance on autoradiographs of 'hot' spots, e.g. see Fig. 2. These small areas of intense radioactivity were caused by the adherence of tiny particles of incompletely solubilized ¹²⁵I-labelled zonae to the nitrocellulose. This problem could be circumvented by centrifuging the melted

zonae at 10 000 g for 5 min before probing the blot. Preliminary experiments showed that despite incomplete dissolution zona proteins do not behave preferentially in their solubility properties during heating at 70°C (results not shown). A second difficulty was the requirement for a 'preblot' with the freshly solubilized ^{125}I -labelled zona proteins. It was found that the first time the probe was used background radioactivity on blots was unacceptably high, despite effective blocking of the nitrocellulose with 5% w/v BSA. O'Rand *et al.* (1985) have reported the same phenomenon and also employed a preblot. However, the second and third times the probe was used background levels became weak and it was possible to detect specific binding. Again preliminary experiments showed that the preblot did not remove preferentially any particular class of zona proteins (results not shown).

Extraction and electrophoresis of ^{125}I -zonae pellucidae recognized on blots

It is known that depending upon whether or not the egg has been ovulated the pig zona pellucida may consist of up to three major glycoproteins (Brown & Cheng, 1986) and in order to examine if one or all of these glycoproteins are involved in binding to sperm and seminal plasma polypeptides, relevant areas of the probed Western blot were cut out and extracted for 5 min at 100°C in 50 μl 62.5 mM-Tris-HCl pH 7.4 containing 2% w/v SDS. SDS-PAGE was carried out as described above.

Results

Electrophoretic analysis of solubilized zonae pellucidae

In preliminary experiments we investigated if the solubilization process had any effect on the macromolecular constituents of the investment. A comparison was made of the profile of ^{125}I -labelled polypeptides, separated on nonreducing SDS-PAGE, of zonae removed manually from freshly ovulated eggs and either dissolved directly in SDS or solubilized by heating as described in Materials and methods. There was no difference between the two procedures as each profile of proteins was identical to that shown in Fig. 1j. Similarly, solubilizing zonae in acid Tyrode's solution pH 2.9 or PBS pH 2.5 had no significant effect on the profile of polypeptides on SDS-PAGE compared to those dissolved directly in SDS.

Identification of molecules in sperm and seminal plasma that recognize ^{125}I -zonae pellucidae glycoproteins

The range of polypeptides present in sodium deoxycholate extracts of ejaculated sperm and in seminal plasma after staining with Coomassie blue is shown in Fig. 1a,b. Sperm extracts separated into a relatively complex profile of proteins with M_r between 14 000 to 100 000 whereas seminal plasma contained fewer

proteins but with a major group between M_r 14 000 and 25 000. Those proteins that stain with the PAS procedure to detect carbohydrate are indicated by an asterisk; in some cases the reaction was quite strong although, as can be seen from Fig. 1, the Coomassie blue stain appeared weak.

When Western blots of parallel gels of those shown in Fig. 1a,b were probed with ^{125}I -zonae from follicular oocytes (^{125}I -FZP) consistent binding was observed to a sperm component of M_r 53 000 (Fig. 1c). Weaker affinity binding was found over a polypeptide M_r 18 000 while two others, M_r 67 000 and 38 000, varied in the amount of probe they bound, the maximum being seen in track c. In seminal plasma (track d) there was strong binding of ^{125}I -FZP over an area ranging from M_r 19 000 to 24 000 and weak binding towards a polypeptide of M_r 65 000.

Probing Western blots of sperm proteins with zonae from ovulated eggs (^{125}I -OZP) gave essentially the same profile of polypeptide recognition as did probing with follicular zonae (Fig. 1e) although in general there was an increased avidity of the probe, particularly for the sperm polypeptide of M_r 53 000. We detected no difference in binding either qualitatively or quantitatively between capacitated and non-capacitated sperm (tracks e and f).

On blots of seminal plasma proteins the ^{125}I -OZP probe bound to a polypeptide doublet M_r 65 000 and an area of polypeptides M_r 19 000–24 000 (tracks d and g).

Washing blots probed with either ^{125}I -FZP or ^{125}I -OZP in PBS containing 0.5% w/v NP-40 reduced considerably background labelling as well as the amount of radiolabel associated with the M_r 67 000, 38 000 and 18 000 polypeptides on spermatozoa. However, the amount of label associated with the sperm M_r 53 000 polypeptide remained undiminished after washing four times with PBS/NP-40.

Identification of zona moieties recognizing sperm and seminal plasma polypeptides

When those areas of the blots occupied by the polypeptides of M_r 18 000, 19 000–24 000, 53 000 and 65 000 were cut out and the bound ^{125}I -zona proteins extracted as described in the Materials and methods section it was possible to identify which of the zona moieties were involved in recognizing the sperm proteins. As shown in Fig. 1 there was a marked difference in the polypeptide profiles depending upon which ^{125}I -zona probe was used. In all cases where ^{125}I -FZP was employed a single polypeptide of M_r 79 000 was resolved with a broader area of label spread directly below it, whereas when ^{125}I -OZP was used each extract separated into three polypeptides of M_r 250 000, 90 000 and 79 000. Again there was a broader area of label below with a lower molecular




Fig. 1. SDS-PAGE of sodium deoxycholate extracts of ejaculated sperm (a) and of seminal plasma (b) stained with Coomassie brilliant blue R250; tracks c and d are similar separations transferred to nitrocellulose and probed with ^{125}I -FZP and f and g represent the same but probed with ^{125}I -OZP. For track e sperm were capacitated before extraction. The electrophoretograms in the lower half of the figure are nonreduced SDS-PAGE separations of zona components extracted from areas of the blot as indicated by their M_r ; tracks h and j represent an individual zona from an oocyte and from an ovulated egg, respectively.

weight limit of approximately 55 000. This difference is shown in Fig. 1 together with polypeptide profiles of the zona from an oocyte and an ovulated egg and it can be seen that the former matches extracts of polypeptides probed with follicular zonae and the latter those probed with ovulated zonae.

Recognition of ^{125}I -zona pellucida by polypeptides of immature and mature spermatozoa

The M_r 53 000 polypeptide was present in DOC extracts of testicular, epididymal and ejaculated sperm as revealed by staining with Coomassie blue and in each case it was recognized by ^{125}I -OZP glycoproteins

after blotting (Fig. 2). It was noticeable that more ^{125}I -OZP bound to the M_r 53 000 polypeptide from cauda epididymal and ejaculated sperm than from testicular and proximal caput epididymal sperm. There was also weak recognition of a polypeptide of approximate M_r 67–68 000 in testicular and proximal caput sperm extracts and of an approximate M_r 38 000 polypeptide in epididymal and ejaculated, but not testicular, sperm extracts.

Tissue specificity of the M_r 53 000 polypeptide

When blots of DOC extracts of porcine liver, kidney, spleen, heart and lung polypeptides separated on

SDS-PAGE were probed with ^{125}I -OZP there was no binding of label to an area of the blot corresponding to M_r 53 000 (Fig. 3).




Fig. 2. SDS-PAGE of sodium deoxycholate extracts of testicular (a), caput (b) and cauda (c) epididymal and ejaculated (d) spermatozoa blotted and probed with ^{125}I -OZP.




Fig. 3. SDS-PAGE of sodium deoxycholate extracts of porcine spermatozoa (a), liver (b), kidney (c), spleen (d), heart (e) and lung (f) blotted and probed with ^{125}I -OZP. Each track received approximately 150 μg protein.

Discussion

A major conclusion from the work described in this paper is that glycoproteins from the zona pellucida of pig eggs have a strong avidity for a protein of M_r 53 000 present in DOC extracts of boar spermatozoa. This protein is recognized by zona glycoproteins from follicular as well as ovulated eggs. It is also present in detergent extracts of spermatozoa from the testis and various levels of the epididymis so it is therefore unlikely that the protein is acquired during sperm maturation in the epididymis. Another aspect of these results is the specificity and strong avidity of the zona glycoproteins for the sperm M_r 53 000 component which, due to its absence from several porcine tissues, is apparently not a ubiquitous protein.

Other sperm polypeptides recognized by the zona glycoproteins were those of M_r 67 000, 38 000 and 18 000 although binding to these was more variable and with lower affinity. O'Rand *et al.* (1985), using the same species and experimental approach, reported major binding of zona proteins to sperm polypeptides of M_r 16 000 and 18 000 with less binding to other species of M_r between 63 000 and 88 000, and 45 000 and 53 000. Our results differ mainly in that the polypeptide M_r 53 000 is consistently the major species recognized in our experiments although we did detect a small amount of binding at M_r 18 000. The only major area of recognition that we saw corresponding to an M_r value of 18–19 000 was in seminal plasma. Possible reasons why the results of O'Rand *et al.* are different from those reported here could be that they adopted a different labelling protocol (solubilized zonae were labelled with ^{125}I -iodogen) or that their proteins were prepared by extracting spermatozoa directly with SDS. It is possible that the polypeptide M_r 67 000 represents one of the minor bands occasionally seen by O'Rand *et al.* and it is interesting that we also found this to be a spasmodic and in our case low-affinity recognition. Using the same approach in the hamster, Sullivan & Bleau (1985) identified a sperm plasma membrane polypeptide of M_r 26 400. This is thought to be a glycoprotein because of its ability to bind ^{125}I -concanavalin A. However, we were unable to detect carbohydrate on the M_r 53 000 polypeptide whereas several others in sperm and seminal plasma were quite strongly PAS-positive.

It is interesting that zona glycoproteins bind to polypeptides in seminal plasma as well as in spermatozoa. Both zona probes (^{125}I -FZP and ^{125}I -OZP) recognized several polypeptides with an M_r value ranging from 19 000 to 24 000 and also, though again to a variable extent, a polypeptide doublet of M_r 65 000. Peterson, Russell & Hunt (1984) have

raised the possibility that boar seminal plasma proteins derived from the seminal vesicle secretion (Hunt, Russell, Peterson & Strack, 1983) can bind to the porcine zona pellucida. Hunt *et al.* have shown that relatively high levels of seminal plasma proteins bind readily to boar spermatozoa and that a group of M_r 16–19 000 is represented in plasma membrane preparations. These are present in seminal vesicle secretion and thus could be responsible for the binding of the zona proteins to blots of seminal plasma components, particularly the lower molecular weight group. Since there is likely to be less of these proteins on sperm after washing it is possible that the sperm polypeptide of M_r 18 000 originates from the seminal plasma.

An alternative approach used to identify molecules involved in sperm–egg recognition utilizes purified or enriched components of either the zona pellucida or sperm membrane to block sperm–zona binding *in vitro*. For instance in the pig a zona component of M_r 85 000 is effective in this respect (Sacco *et al.* 1984) as are enriched sperm membrane preparations (Peterson, Henry, Hunt, Saxena & Russell, 1985). Although the experimental approach that we have used is arguably less physiological than the above it has the advantage of testing the complete profile of zona glycoproteins against a wide range of sperm polypeptides, thus increasing the chance of identifying potentially interesting molecules. Also in the pig a similarly comprehensive approach has been adopted by Peterson *et al.* (1985). They incubated solubilized sperm plasma membrane preparations with isolated zonae and like us found that several proteins with molecular weights ranging from approximately 35 000 to 100 000 showed affinity for the zona.

The mouse zona pellucida consists of three major glycoproteins and one of these, M_r 83 000 (ZP3), has sperm-binding properties (Bleil & Wassarman, 1980). Similarly in the pig sperm receptor activity has been attributed to a glycoprotein of M_r 58 000 (Sacco *et al.* 1984). However, in our experiments recognition of sperm polypeptides was not restricted to a particular zona component. Analysis of ^{125}I -zona probe that bound to sperm polypeptides clearly showed that all zona moieties were involved. Although at present we are unable to elaborate on this common binding property of zona glycoproteins the results do suggest that a significant structural homology between them exists. The zona pellucida of the oocyte is heavily glycosylated (Dunbar, Wardrip & Hedrick, 1980) and both oviduct glycoproteins react strongly with PAS (Brown & Cheng, 1986) so it is possible that the sperm polypeptide M_r 53 000 is recognizing common sugar sequences. This would not be surprising for evidence is accruing that carbohydrates are critically involved in the initial events of gamete fusion (Lopez,

Bayna, Litoff, Shaper, Shaper & Shur, 1985; Florman & Wassarman, 1985). Further experiments are in progress to investigate the nature of the binding we have observed and its possible significance in relation to sperm–egg recognition processes.

We are grateful to Dr S. J. Gaunt for helpful discussions, Mr K. I. von Gloß for technical assistance and Mr K. T. Elsome for supplying pig eggs.

References

- BLEIL, J. D. & WASSARMAN, P. M. (1980). Mammalian sperm-egg interaction: identification of a glycoprotein in mouse egg zonae pellucidae possessing receptor activity for sperm. *Cell* **20**, 873–882.
- BOLTON, A. E. & HUNTER, W. M. (1973). The labelling of proteins to high specific activities by conjugation to a ^{125}I -containing acylating agent. *Biochem. J.* **133**, 529–539.
- BROWN, C. R. & CHENG, W. K. T. (1986). Changes in composition of the porcine zona pellucida during development of the oocyte to the 2- to 4-cell embryo. *J. Embryol. exp. Morph.* **92**, 183–191.
- CHENG, W. K. T. (1985). *In vitro* fertilization of farm animal oocytes. PhD thesis. Council for National Academic Awards. Institute of Animal Physiology, Cambridge, UK.
- DUNBAR, B. S., WARDRIP, N. J. & HEDRICK, J. L. (1980). Isolation, physicochemical properties, and macromolecular composition of zona pellucida from porcine oocytes. *Biochemistry, N.Y.* **19**, 356–364.
- FAIRBANKS, G., STECK, T. L. & WALLACH, D. F. H. (1971). Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochemistry, N.Y.* **10**, 2606–2617.
- FLORMAN, M. H. & WASSARMAN, P. M. (1985). O-Linked oligosaccharides of mouse egg ZP3 account for its sperm receptor activity. *Cell* **41**, 313–324.
- GWATKIN, R. B. L. (1977). *Fertilization Mechanisms in Man and Mammals*. New York: Plenum Press.
- HUNT, W. P., RUSSELL, L. D., PETERSON, R. N. & STRACK, L. E. (1983). Association of polypeptides from the boar reproductive tract fluids with boar sperm plasma membranes. *Fed. Proc.* **42**(3), 615.
- LAEMMLI, U. K. (1970). Cleavage of structural proteins during assembly of the head of bacteriophage T₄. *Nature, Lond.* **227**, 680–685.
- LOPEZ, L. C., BAYNA, E. M., LITOFF, D., SHAPER, N. L., SHAPER, J. H. & SHUR, B. D. (1985). Receptor function of mouse sperm surface galactosyltransferase during fertilization. *J. Cell Biol.* **101**, 1501–1510.
- O'RAND, M. G., MATTHEWS, J. E., WELCH, J. E. & FISHER, S. J. (1985). Identification of zona binding proteins of rabbit, pig, human and mouse spermatozoa on nitrocellulose blots. *J. exp. Zool.* **235**, 423–428.
- PETERSON, R. N., RUSSELL, L. D. & HUNT, W. P. (1984). Evidence for specific binding of uncapacitated boar spermatozoa to porcine zonae pellucidae *in vitro*. *J. exp. Zool.* **231**, 137–147.

- PETERSON, R. N., HENRY, L., HUNT, W., SAXENA, N. & RUSSELL, L. D. (1985). Further characterization of boar sperm plasma membrane proteins with affinity for the porcine zona pellucida. *Gamete Res.* **12**, 91–100.
- PRIMAKOFF, P., HYATT, H. & MYLES, D. G. (1985). A role for the migrating sperm surface antigen PH-20 in guinea pig sperm binding to the egg zona pellucida. *J. Cell Biol.* **101**, 2239–2244.
- SACCO, A. G., SUBRAMANIAN, M. G. & YUREWICZ, E. C. (1984). Association of sperm receptor activity with a purified pig zona antigen (PPZA). *J. Reprod. Immunol.* **6**, 89–103.
- SHUR, B. D. & HALL, N. J. (1982). A role for mouse sperm surface galactosyltransferase in sperm binding to the egg zona pellucida. *J. Cell Biol.* **95**, 574–579.
- SULLIVAN, R. & BLEAU, G. (1985). Interaction of isolated components from mammalian sperm and egg. *Gamete Res.* **12**, 101–116.
- TOWBIN, H., STAEBELIN, T. T. & GORDON, J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. natn. Acad. Sci. U.S.A.* **76**, 4350–4354.