

RESEARCH ARTICLE

Increased host ATP efflux and its conversion to extracellular adenosine is crucial for establishing *Leishmania* infection

Moumita Basu¹, Purnima Gupta², Ananya Dutta³, Kuladip Jana³ and Anindita Ukil^{1,*}

ABSTRACT

Intracellular survival of *Leishmania donovani* demands rapid production of host ATP for its sustenance. However, a gradual decrease in intracellular ATP in spite of increased glycolysis suggests ATP efflux during infection. Accordingly, upon infection, we show here that ATP is exported and the major exporter was pannexin-1, leading to raised extracellular ATP levels. Extracellular ATP shows a gradual decrease after the initial increase, and analysis of cell surface ATP-degrading enzymes revealed induction of the ectonucleotidases CD39 and CD73. Ectonucleotidase-mediated ATP degradation leads to increased extracellular adenosine (eADO), and inhibition of CD39 and CD73 in infected cells decreased adenosine concentration and parasite survival, documenting the importance of adenosine in infection. Inhibiting adenosine uptake by cells did not affect parasite survival, suggesting that eADO exerts its effect through receptor-mediated signalling. We also show that *Leishmania* induces the expression of adenosine receptors A_{2A}R and A_{2B}R, both of which are important for anti-inflammatory responses. Treating infected BALB/c mice with CD39 and CD73 inhibitors resulted in decreased parasite burden and increased host-favourable cytokine production. Collectively, these observations indicate that infection-induced ATP is exported, and after conversion into adenosine, propagates infection via receptor-mediated signalling.

KEY WORDS: ATP, Ectonucleotidase, Adenosine receptor, *Leishmania donovani*, Macrophage

INTRODUCTION


Leishmania donovani, the causative agent of fatal visceral leishmaniasis, is an obligate intracellular parasite inoculated into mammals by the bite of its vector, the sand fly. Once transmitted to the host, the infective promastigote forms are rapidly phagocytosed by macrophages (de Menezes et al., 2016) and the parasite, along with combating the antimicrobial defence arsenals of the macrophages, has to survive within the harsh environment of the phagolysosomes (Cunningham, 2002). In order to succeed, *Leishmania* requires high energy levels; thus, the energy needs of both the parasite for its survival and of the host cell to resist infection, have drawn attention (Schaible and Kaufmann, 2007). As

the intracellular parasite and the host cell utilise the same host energy sources, knowledge about the source of cellular energy and the pathway in which it is utilised within infected macrophage seems extremely important to address the molecular signalling of infection.

Adenosine 5'-triphosphate (ATP), the universal energy currency of a cell, fuels various vital processes necessary for maintenance of a cell and also acts as an intracellular and extracellular messenger (Corriden and Insel, 2010; Trautmann, 2009). The main source of ATP is the catabolic part of carbohydrate metabolism where glycolysis is followed by the tricarboxylic acid (TCA) cycle and oxidative phosphorylation (O'Neill and Pearce, 2016). However, during certain infections or inflammatory conditions, the pyruvate generated via glycolysis gets converted into lactate and leads to lesser but faster ATP production (Kelly and O'Neill, 2015). ATP generated within a cell, apart from acting as a source of energy, may act as a signalling molecule both intracellularly and even extracellularly (Corriden and Insel, 2010; Trautmann, 2009). Various stimuli such as stress, hypoxia, inflammation and reactive oxygen species (ROS) are found to induce ATP release from cells with the help of various ATP release channels such as the maxi-anion channel, the gap junction hemichannel, the P2X7 receptor and even via vesicular exocytosis in a non-lytic mechanism (Kojima et al., 2017). Released extracellular ATP (eATP) often act as a signalling molecule and regulates various cellular processes (Corriden and Insel, 2010). Initially, it was found to play an important role in modulating the nervous and vascular systems (Housley et al., 2009; North and Verkhatsky, 2006; Schwiebert et al., 2002). Recent studies have shown that eATP may inhibit infection by intracellular bacterial pathogens (Coutinho-Silva et al., 2003; Lammass et al., 1997) and also regulate chemotaxis of immune cells (Chen et al., 2006; Kronlage et al., 2010). However, the concentration of eATP is critically maintained as a high eATP level acts as a 'danger signal' and can promote inflammation through the P2X7-receptor-dependent activation of the NLRP3 inflammasome, leading to production of pro-inflammatory cytokine IL-1 β (Cassel and Sutterwala, 2010). The ATP level in the extracellular environment is usually maintained with the help of various ATP hydrolysing enzymes such as ecto-apyrase, pyrophosphatases, phosphodiesterases and ecto-adenosine triphosphatases (eATPase), amongst which the prevalent one is CD39 (also known as ENTPD1), a member of the ectonucleotide diphosphohydrolase (e-NTDPase) family. CD39 dephosphorylates ATP to ADP and AMP (Antonioli et al., 2013). AMP thus generated is further converted to adenosine (ADO) by the ecto-5'-nucleotidase CD73 (also known as NT5E). These enzymes are found to be expressed on the surface of macrophages, endothelial cells and various subsets of lymphocytes (Antonioli et al., 2013). Therefore, these enzymes are not only important for maintaining intermediate metabolite concentration but also play a significant role in modulating cellular immune responses. CD73 expression on tissues and T helper cells was found to promote

¹Department of Biochemistry, University of Calcutta, Kolkata 700019, West Bengal, India. ²Infections and Cancer Biology Group, International Agency for Research on Cancer, 69372, Lyon Cedex 08, France. ³Division of Molecular Medicine, Bose Institute, P1/12 Calcutta Improvement Trust Scheme, VIIM, Kolkata, 700054, West Bengal, India.

*Author for correspondence (aubicc@caluniv.ac.in)

 M.B., 0000-0001-6691-5279; K.J., 0000-0003-4748-253X; A.U., 0000-0002-1102-9437

colonisation and thus prolonged murine salmonellosis infection (Alam et al., 2014). Infection of dendritic cells with different species of *Leishmania* has also been shown to upregulate the expression and activity of CD39 and CD73 (Figueiredo et al., 2012), although the consequences have not been studied in detail.

Extracellular adenosine (eADO) generated by CD39 and CD73 is further acted upon by adenosine deaminase (ADA), which catalyses the deamination of adenosine to inosine. This is the only enzyme known to regulate the adenosine concentration, and absence of ADA has been associated with increased eADO level, leading to anti-inflammatory responses (Cronstein, 1994; Haskó and Cronstein, 2013). Adenosine is an immunomodulator generated at sites during tissue injury, inflammation and ischemia, and mainly elicits its action on cells through one of its four G-protein-coupled receptors: A₁, A_{2A}, A_{2B} and A₃ (Fredholm et al., 2001; Olah and Stiles, 1995). Adenosine was found to inhibit superoxide production, adhesion of cells to the endothelial barrier (Bouma et al., 1996), and proliferation and expansion of T-cells through the A_{2A} receptor (Hoskin et al., 2008; Huang et al., 1997). A₁R on the other hand, has been found to transduce pro-inflammatory signalling of adenosine (Haskó and Cronstein, 2004). Thus, the effect of adenosine on a given cell depends on the adenosine receptor subtype present and the signal it transduces.

The present study was initiated to address how the infective parasite and the infected macrophage cope with the demand for rapid requirement of ATP. Characterisation of infected host cell metabolic profiling and possible ATP efflux encouraged us to determine the fate of the extracellular ATP produced. Further investigation revealed that ATP is exported in the extracellular environment via a specific channel and gets hydrolysed to adenosine by cell surface ectonucleotidases. This not only enables the parasite to escape macrophage immune activation, but creates an anti-inflammatory cellular environment necessary for intra-macrophage survival of *Leishmania* through receptor-mediated signalling.

RESULTS

Host ATP generation during *L. donovani* infection

Successful survival and replication of intracellular pathogens require substantial amount of energy, i.e. ATP production. To quantify the energy status of an infected cell, we measured the intracellular ATP (iATP) concentration, which is expected to be increased following *L. donovani* infection. However, the iATP level showed a decreasing pattern up to 8 h post-infection in RAW macrophages (Fig. 1A). This contradictory observation led us to check the level of ATP in peritoneal macrophages, which also depicted a similar profile (Fig. 1B). Therefore, to investigate the metabolic status of RAW 264.7 macrophage cells during the early stages of *L. donovani* infection (0–8 h), the extracellular acidification rate (ECAR) and the mitochondrial oxygen consumption rate (OCR) were measured using live-cell extracellular flux analysis. ECAR values were found to be higher in infected cells (3.6-fold increase compared with uninfected cells at 3 h post-infection, $P<0.001$) indicating higher glycolytic activity (Fig. 1C). In contrast, the OCR values at 3 h post-infection showed a significant reduction (29.5% decrease compared with uninfected control, $P<0.01$), suggesting a lower mitochondrial respiration rate (Fig. 1C). A similar profile was obtained for peritoneal macrophages (Fig. 1D). Expression of the three enzymes involved in the irreversible steps of glycolysis, namely hexokinase 1 (Hk1), phosphofructokinase (Pfk) and pyruvate kinase M2 (Pkm2) also showed significant transcriptional upregulation during infection (3.3-, 4.2- and 5.1-fold, respectively, compared with uninfected control at 4 h post-infection, $P<0.001$) (Fig. 1E). In order to determine whether this upregulation of glycolysis is induced by live

L. donovani, the transcript level of the enzymes was also assessed with paraformaldehyde fixed (PFA Fx) *L. donovani*, which showed almost unaltered levels of all three enzymes 4 h post-infection (Fig. 1F). Reduced mitochondrial respiration (Fig. 1C,D), led us to check whether pyruvate generated in infected macrophage cells from increased glycolysis is converted to lactate instead of entering the TCA cycle. Extracellular lactate concentration was found to be gradually increased with a maximum (3.5-fold increase, $P<0.001$) at 8 h post-infection (Fig. 1G), which corroborated well with our earlier observations. However, PFA-fixed *L. donovani* did not show any significant change in lactate level (Fig. 1G). To find out whether this early upregulation of glycolysis has an effect on intra-macrophage survival of *L. donovani*, macrophages were pre-treated with a graded concentration of 2-deoxyglucose (2-DG), an inhibitor of glycolysis, for 4 h followed by infection with promastigotes for 24 h. Treatment with 2-DG did not affect the viability of RAW 264.7 cells but the number of intracellular parasites was found to be significantly reduced at 10 mM concentration of 2-DG (59.9% decrease compared with untreated infected cells, $P<0.01$) (Fig. 1H). However, treatment of both promastigotes and axenic amastigotes with 10 mM 2-DG did not substantially affect parasite viability (Fig. S1B). Taken together, these results indicate that *L. donovani* infection causes a metabolic shift towards glycolysis during the initial hours of infection, which is essential for parasite survival within macrophages. However, the ATP generated via glycolysis is either utilised by the cell or effluxes to the extracellular environment.

ATP efflux during infection promotes parasite survival

Since ATP concentration showed a decreasing pattern (Fig. 1A,B) in spite of a higher glycolytic rate in infected macrophages, to understand the fate of glycolysis generated ATP, we measured the extracellular ATP concentration of infected cells, as macrophages are capable of releasing ATP (Dosch et al., 2018). Accordingly, it was found that at 2 h post-infection, the ATP concentration was significantly higher in the cell culture supernatant in comparison to uninfected cells (2.9- and 2.6-fold increase in RAW and peritoneal macrophages, respectively, $P<0.001$), indicating release of intracellular ATP during infection (Fig. 2A). Pre-treatment of infected macrophages with 2-DG, reduced the eATP concentration implying that the infection-induced ATP release was glycolysis dependent (Fig. 2A). Next, to evaluate the mechanism of ATP release, infected RAW cells were pre-treated with 100 μ M amounts of inhibitors of various ATP release channels [¹⁰Panx for pannexin-1 (also known as PANX1; Pelegrin and Surprenant, 2006), carbenoxolone (CBX) for pannexin and connexin-related channels (Chekeni et al., 2010), flufenamic acid (FFA) for anion transporter (Guinamard et al., 2013) and gadolinium chloride (GdCl₃) for maxi-anion channel (Zhao et al., 2017)], then the extracellular ATP level was measured at 2 h post-infection. Of all the inhibitors, ¹⁰Panx and CBX treatment led to significant reduction of extracellular ATP level (Fig. 2B). However, the extent of inhibition was much less (36.2% decrease, $P<0.01$) in case of CBX in comparison to ¹⁰Panx, which showed a 56.3% decrease ($P<0.001$) (Fig. 2B). Higher doses of ¹⁰Panx and CBX did not lead to any further significant decrease in the eATP concentration, as revealed by a dose titration experiment (Fig. 2C). Maximum significant inhibition was obtained at a dose of 100 μ M for both ¹⁰Panx and CBX (Fig. 2C), which agreed with the administered dose used in our inhibition study (Fig. 2B). Since inhibition of ¹⁰Panx exerted maximum effect on ATP release, we next assessed the expression of pannexin-1 in infected macrophages at both the mRNA and protein level. Expression of pannexin-1 was found to be significantly higher in both cases (4.3- and 6.0-fold




Fig. 1. ATP production during *Leishmania donovani* infection. (A,B) Macrophages were infected with *L. donovani* promastigotes (*L.d.*) for the indicated time points (0–8 h) and intracellular ATP (iATP) level was estimated using a luminescent kit for the control and infected samples of RAW 264.7 (A) and peritoneal macrophages (PM ϕ) (B). (C,D) RAW 264.7 cells (8×10^4) (C) and peritoneal macrophages (2×10^5) (D) were infected with *L. donovani* promastigotes with a parasite/macrophage ratio of 10:1 for 1 and 3 h. At both time points, ECAR (mpH/min/ μ g protein) and OCR (pmol/min/ μ g protein) were measured in real time under basal conditions. (E) mRNA level expressions of hexokinase 1 (Hk1), phosphofructokinase (Pfk) and pyruvate kinase M2 (Pkm2) were assessed in *L. donovani*-infected RAW cells for the indicated time points using β -actin as an endogenous control by RT-PCR. (F) RAW cells were infected with live or paraformaldehyde-fixed (PFA Fx) *L. donovani* promastigotes for 4 h and expression of Hk1, Pfk and Pkm2 were assessed at mRNA level. (G) Extracellular lactate concentration in the cell culture supernatant of RAW cells following *L. donovani* infection for varying time periods was measured by colorimetric assay. (H) RAW cells were infected with *L. donovani* alone or in the presence of increasing concentrations of 2-deoxyglucose (2-DG; 0.1, 1, 10 and 20 mM). Percentage cell viability was measured by MTT assay at 4 h and number of intracellular parasites per 100 macrophages was measured by DAPI staining at 24 h post-infection. Bands were quantified by densitometry and are shown as bar graphs. Results are representative of three independent experiments and error bars are expressed as mean \pm s.d. $n=3$; ns, not significant, ** $P<0.01$, *** $P<0.001$ (Student's *t*-test).

induction for mRNA and protein level at 2 h post-infection, respectively, compared with uninfected control, $P<0.001$) up to 6 h post-infection (Fig. 2D,E). Inhibition of export of ATP from infected macrophages using $^{10}\text{Panx}$ further led to a significant reduction in the number of intracellular parasites (56.9% decrease over infected control, $P<0.001$) (Fig. 2F). Infected cells when supplemented with 100 μM ATP showed a marked reversal in the inhibition of intracellular parasite count even upon treatment with $^{10}\text{Panx}$ (1.6-fold increase over infected cells pre-treated with $^{10}\text{Panx}$, $P<0.01$) (Fig. 2F) thereby suggesting the relevance of ATP export during infection. The cell viability of the promastigotes and axenic

amastigotes in the presence of 100 μM $^{10}\text{Panx}$ was found to be almost unaltered (Fig. S1B). These observations suggest that the ATP generated by the increased rate of glycolysis during *L. donovani* infection gets exported outside cell, mostly through pannexin-1, and this export seems to be beneficial for intracellular parasite survival.

Role of macrophage ectonucleotidases in the hydrolysis of eATP

Excess eATP is also detrimental for cell viability as it may lead to inflammatory signals and even to apoptosis (Cauwels et al., 2014). Therefore, the time kinetics of eATP level was investigated in




Fig. 2. ATP efflux during infection promotes parasite survival. (A) Extracellular ATP (eATP) concentration in the cell culture supernatant was quantified in both RAW and PM ϕ cells infected with *L. donovani* promastigotes (*L.d.*) for 2 h and pre-treated or not with 10 mM 2-DG for 4 h. (B) eATP concentration was measured in the cell culture supernatant of 2-h-infected RAW cells which were pre-incubated for 30 min with 100 μ M each of 10 Panx or carbenoxolone (CBX) or flufenamic acid (FFA) or gadolinium chloride (GdCl $_3$). (C) eATP was measured in the culture supernatant of 2-h-infected RAW cells, pre-incubated for 30 min with increasing concentrations of 10 Panx (50, 100 and 200 μ M) or carbenoxolone (CBX; 10, 50 and 100 μ M). (D,E) RAW cells were infected with *L. donovani* promastigotes for the indicated time periods (0–6 h) and the expression of pannexin-1 was measured at the mRNA level by RT-PCR (D) and at protein level by immunoblotting (E). (F) Macrophages were infected with *L. donovani* promastigotes for 24 h in the presence of 10 Panx (100 μ M) or both ATP (100 μ M) and 10 Panx (100 μ M) and the number of parasites per 100 macrophages were analysed by DAPI staining. Bands were quantified by densitometry and are shown as bar graphs. Results are representative of three independent experiments, and error bars are expressed as mean \pm s.d. $n=3$; ns, not significant, ** $P<0.01$, *** $P<0.001$ (Student's *t*-test).

infected peritoneal macrophages during the early hours of infection (Fig. 3A). The eATP level started decreasing from 2 h post-infection and was markedly reduced at 8 h post-infection when infected with live promastigotes (Fig. 3A). Unlike live *Leishmania*, PFA-fixed *L. donovani* significantly increased the extracellular ATP concentration even during the later stages of infection (6 and 8 h) (Fig. 3A). In order to counter the detrimental effect of eATP, cells normally utilise various ATP hydrolysing enzymes to control the levels of ATP (Antonioli et al., 2013). We therefore evaluated expression of the main ATP hydrolysing enzymes of macrophage cells: CD39, CD73 and adenosine deaminase (ADA). Time kinetics revealed increased expression of CD39 and CD73 in both RAW and peritoneal macrophages as observed up to 24 h post infection (2.8- and 3.9-fold in RAW and 2.6- and 3.8-fold in peritoneal macrophages, respectively, compared with uninfected control at 6 h post-infection, $P<0.001$) (Fig. 3B,C). However, expression of ADA was found to be very low throughout the time course in both cell types (Fig. 3B,C). The expression kinetics of macrophage-associated ectonucleotidases were also assessed following infection

with PFA-fixed *L. donovani*, which showed almost undetectable expression of all the three enzymes CD39, CD73 and ADA (Fig. 3D). Increased expression of CD39 and CD73 prompted us to evaluate their ectonucleotidase activity. To this end, infected peritoneal macrophages were incubated in the presence of ATP (to evaluate CD39 activity) or AMP (to evaluate CD73 activity) and the ectonucleotidase activity was assessed by measuring inorganic phosphate (P $_i$) release (Fig. 3E,F). A time-dependent increase in hydrolysis was found for both ATP and AMP, and pre-treatment of cells with specific inhibitors of CD39 (POM-1) and CD73 (APCP) reversed this hydrolysis, thereby documenting specific involvement of CD39 and CD73 in hydrolysing extracellular ATP. Low expression of ADA indicated that extracellular adenosine produced during nucleotide hydrolysis might not be processed further and extracellular adenosine level indeed showed an increasing pattern with maximum level 8 h post infection (3.4-fold increase in comparison to uninfected cells, $P<0.001$) (Fig. 3G). On the other hand, infection with PFA-fixed *L. donovani* failed to increase the eADO level as opposed to infection with live parasites




Fig. 3. See next page for legend.

(Fig. 3G), thereby suggesting specific contribution of live parasites in converting host ATP to eADO. In order to determine whether generated adenosine aids in parasite survival, infected macrophages were pre-incubated with increasing concentrations of POM-1 and APCP. It was found that there was a significant decrease in the number of intracellular parasites upon treatment with either POM-1

(100 μM) or APCP (100 μM) (54.5% and 58.6% reduction, respectively, compared with infected control, $P < 0.001$) (Fig. 3H). However, neither POM-1 nor APCP had any effect on the viability of the macrophages or promastigotes or axenic amastigotes at the above concentrations (Fig. S1A,B). Infected cells were also treated with POM-1 and APCP in the presence or absence of adenosine

Fig. 3. Role of macrophage ectonucleotidases in the hydrolysis of ATP.

(A) Peritoneal macrophages (PM ϕ) were infected with live or PFA-fixed *L. donovani* promastigotes for the indicated time periods and the eATP concentration in the culture supernatant was measured. (B,C) Cells were infected with live *L. donovani* promastigotes for indicated time periods (0–24 h) and the protein level expression of CD39, CD73 and ADA measured by immunoblotting in RAW (B) and PM ϕ (C) cells. (D) Peritoneal macrophages were infected with PFA-fixed *L. donovani* promastigotes for the indicated time periods and the protein level expression of CD39, CD73 and ADA measured by immunoblotting. β -actin was used as an endogenous control. (E) PM ϕ were infected with *L. donovani* promastigotes alone or in the presence of increasing concentrations of POM-1 (50, 100 and 200 μ M). Ectonucleotidase activity was then measured for CD39 by the amount of P_i released using ATP as substrate. (F) PM ϕ were infected with *L. donovani* promastigotes alone or in the presence of increasing concentrations of APCP (50, 100 and 200 μ M). Ectonucleotidase activity was then measured for CD73 using AMP as substrate. (G) PM ϕ were infected with live or PFA-fixed *L. donovani* for the indicated time periods and extracellular adenosine concentration (eADO) was measured in culture supernatant. (H) PM ϕ were infected with *L. donovani* for 24 h in the presence of increasing concentrations of POM-1 (50, 100 and 200 μ M) or APCP (50, 100 and 200 μ M) and the number of parasites per 100 macrophages were analysed by DAPI staining. (I) Number of parasites per 100 macrophages were measured in 24-h-infected PM ϕ in the presence of POM-1 (100 μ M) and APCP (100 μ M) alone or in combination or POM-1+APCP+ADO (100 μ M) or ATP- γ s (100 μ M). Bands were quantified by densitometry and are shown as bar graphs. Results are representative of three independent experiments and error bars are expressed as mean \pm s.d. $n=3$; ns, not significant, ** $P<0.01$, *** $P<0.001$ (Student's *t*-test).

(ADO) or a non-hydrolysable ATP analogue (ATP- γ s) (Fig. 3I). The extent of inhibition in the presence of POM-1 and APCP in combination was found to be 62.4%, which is not a very significant change in comparison to pre-treatment with either POM-1 or APCP alone. Treatment of cells with ATP- γ s (100 μ M) also resulted in marked reduction in the number of intracellular parasites (39.5% reduction compared with infected control, $P<0.001$) (Fig. 3I). However, the decrease in intracellular parasite count in POM-1- and APCP-treated cells was found to be markedly reversed upon treatment with adenosine (100 μ M) (2.2-fold increase in comparison to parasite count in POM-1- and APCP-treated cells, $P<0.001$) (Fig. 3I). All these observations suggest that extracellular ATP generated during infection is degraded by the cell surface ectonucleotidases to produce adenosine to propagate infection.

Extracellular adenosine exerts its effect via A_{2A} and A_{2B} receptors

The fact that increased extracellular adenosine concentration facilitated parasite survival prompted us to determine its role on infection. Adenosine generated by the action of ectonucleotidases may either be taken up by the host cell through nucleoside transporters (Khan et al., 2007) or may exert its effect through signalling via specific cell surface receptors (Haskó and Cronstein, 2004). However, pre-treatment of cells with various concentrations of the adenosine uptake blocker dipyrindamole (Di) did not have any effect on the intracellular parasite count (Fig. 4A) suggesting thereby that adenosine exerts its effect via specific adenosine receptor-mediated signalling. Adenosine primarily acts through one of the subtypes of four G-protein-coupled receptors A₁, A_{2A}, A_{2B} and A₃ to establish an anti-inflammatory environment (Fredholm et al., 2001; Olah and Stiles, 1995). Infection increased expression of both A_{2A} receptor (A_{2A}R) and A_{2B} receptor (A_{2B}R) (2.2- and 3.0-fold, respectively, compared with control, $P<0.001$), whereas almost undetectable expression was obtained for A₁R and A₃R (Fig. 4B). Time kinetics studies further showed persistence of enhanced expression of A_{2A}R and A_{2B}R at both the mRNA and protein level (2.9- and 3.4-fold, respectively, at 8 h post-infection

in comparison to uninfected cells at mRNA level and 2.9- and 4.5-fold, respectively, at protein level, $P<0.001$) (Fig. 4C,D). The observation was further verified by fluorescence microscopy using Texas Red-conjugated anti-A_{2A}R antibody and FITC labelled anti-A_{2B}R antibody, which showed significantly enhanced expression of both A_{2A}R and A_{2B}R (Fig. 4E). Next, to determine the role of adenosine receptor-mediated signalling, infected macrophages were pre-treated with specific inhibitors of A_{2A}R and A_{2B}R, ZM 241385 and MRS 1754, respectively, and parasite survival was determined by measuring intracellular parasite count. Individual treatment with either ZM 241385 (1 μ M) (Kreckler et al., 2006) or MRS 1754 (1 μ M) (Kreckler et al., 2006) led to moderate decrease in intracellular parasite count (48.1% and 54.2%, respectively, compared with the control, $P<0.001$) (Fig. 4F). However, a marked reduction in intracellular parasite count was obtained when both the inhibitors were used in combination (61.5% reduction compared with control infected, $P<0.001$) suggesting synergistic inhibition (Fig. 4F). None of the inhibitors showed any significant effect on macrophage or parasite survival in promastigote or axenic amastigote culture (Fig. S1A,B). Furthermore, this inhibition was obtained despite the presence of adequate concentration of adenosine in extracellular environment (Fig. 4F, inset) confirming that adenosine-mediated effects were exerted by specific receptors. To investigate whether inhibitor treatment modulates the expression profile of A_{2A}R and A_{2B}R, control and infected macrophages were pre-treated with either ZM 241385 or MRS 1754, and expression of A_{2A}R and A_{2B}R was determined by western blot analysis. The expression profile of both receptors remained unaltered upon treatment with specific inhibitors (Fig. 4G), confirming that inhibitors modulate only the receptor-mediated signalling, not their expression level. Since adenosine is an endogenous regulator of inflammation (Haskó and Cronstein, 2013), we investigated whether this upregulated expression of A_{2A}R and A_{2B}R following infection had any role in modulation of cytokine profile, by measuring levels of cytokines and chemokines in infected cells pre-treated with ZM 241385 and MRS 1754. LPS-induced production of IL-12 and TNF- α , which was significantly decreased in *L. donovani*-infected cells, was found to be increased upon treatment with ZM241385 (1 μ M) (4.1- and 3.0-fold increase for TNF- α and IL-12, respectively, compared with infected cells, $P<0.001$ and $P<0.01$, respectively) (Fig. 4H). However, pre-treatment with the A_{2B}R-specific antagonist MRS1754 (1 μ M) did not affect pro-inflammatory cytokine production (Fig. 4H). On the other hand, the infection-induced increased IL-10 and TGF- β production was found to be reduced by treatment with MRS 1754 (71.8% and 44.1% reduction in comparison to infected cells, $P<0.001$) and did not show any significant change when infected cells were pre-treated with ZM 241385 (Fig. 4I). Pre-treatment of infected cells with adenosine (100 μ M) marginally increased the levels of IL-10 and TGF- β (Fig. 4I). Like cytokines, inflammatory chemokines CCL3 and CCL5 production were also upregulated over infected control on a very small scale upon treatment with ZM (Fig. 4J). However, MRS1754 did not seem to have any effect on either CCL3 or CCL5 modulation. All these observations suggest that extracellular adenosine produced during infection exerts its effect via the receptors A_{2A}R and A_{2B}R and maintains an intracellular anti-inflammatory milieu necessary for parasite survival.

Effect of the ectonucleotidase inhibitors POM-1 and APCP on infection in BALB/c mice

To evaluate the importance of extracellular ATP degradation on parasite survival in visceral leishmaniasis, *L. donovani*-infected




Fig. 4. Extracellular adenosine exerts effect via $A_{2A}R$ and $A_{2B}R$. (A) Number of parasites per 100 cells was measured by DAPI staining in 24-h-infected PM ϕ in the presence of increasing concentrations of dipyrindamole (5, 10 and 20 μ M). (B) mRNA level expression of the adenosine receptors $A_{1}R$, $A_{2A}R$, $A_{2B}R$ and $A_{3}R$ was assessed by RT-PCR in uninfected and 8-h *L. donovani*-infected RAW cells. (C,D) RAW cells were infected with *L. donovani* promastigotes (*L.d.*) for the indicated time periods (0–24 h) and the expression of $A_{2A}R$ and $A_{2B}R$ were measured at the mRNA level by RT-PCR (C) and at protein level by immunoblotting (D). (E) RAW cells were infected with promastigotes for 8 h and then stained with anti- $A_{2A}R$ and anti- $A_{2B}R$ monoclonal antibodies followed by Texas Red-conjugated anti- $A_{2A}R$ secondary antibody and FITC-conjugated anti- $A_{2B}R$ secondary antibody. Nuclei were stained with DAPI and the cells were analysed under a confocal microscope. The intensity of staining for both antibodies were measured in each cell using ImageJ software and expressed as mean fluorescence intensity. (F) PM ϕ were pre-treated either with ZM241385 (1 μ M) or MRS1754 (1 μ M) or in combination for 30 min followed by infection with *L. donovani* for 24 h. The number of parasites per 100 macrophages were evaluated by DAPI staining. (G) Infected RAW 264.7 cells were pre-treated either with ZM241385 or MRS1754 and the expression of $A_{2A}R$ and $A_{2B}R$ were measured at protein level by immunoblotting. (H,I,J) Macrophages were treated with either LPS (100 ng/ml) or ZM241385 (1 μ M) or MRS1754 (1 μ M) or both ZM241385 and MRS1754 for 30 min followed by infection with *L. donovani* for 24 h. Culture supernatants were assayed for TNF- α and IL-12 (H) and IL-10 and TGF- β (I) and CCL-3 and CCL-5 (J) using ELISA. Bands were quantified by densitometry and are shown as bar graphs. Results are representative of three independent experiments and error bars are expressed as mean \pm s.d. $n=3$; ns, not significant, ** $P<0.01$, *** $P<0.001$ (Student's *t*-test).

mice were administered intra-peritoneally (i.p.) with either the CD39-specific inhibitor POM-1 or the CD73-specific inhibitor APCP. Dose-titration experiments were carried out to assess the efficacy of the inhibitors with a dose range of 2.5–20 mg/kg body weight/day given up to 4 weeks at 3 day intervals starting at 11 days

post-infection. The infection was allowed to proceed for 4 weeks, after which the animals were sacrificed, and the anti-leishmanial potency was assessed in terms of parasite burden in liver and spleen. During the experimental period, all the animals remained healthy. Maximum suppression of organ parasite burden was obtained




Fig. 5. Effect of POM-1 and APCP on infection in mice. (A,B) Various doses of POM-1 and APCP ranging from 2.5 to 20 mg/kg body weight/day were given i.p. up to 4 weeks at every 3 days interval starting at 11 days post-infection. The parasite burdens were then determined at 4 weeks post-infection in liver (A) and spleen (B) for both the inhibitors. (C) The course of visceral infection was followed in BALB/c mice that had received i.p. injections of POM-1 (10 mg/kg body weight/day) and APCP (10 mg/kg body weight/day) for 4 weeks. (D,E) Parasite burden expressed as LDU in liver (D) and spleen (E). (F) Representative microscopic images of H&E-stained liver sections of infected (4 week) and POM-1- and APCP-treated BALB/c mice. Original magnification $\times 40$. (G) Total number of hepatic granulomas (immature and mature) in the H&E-stained liver sections of infected (4 week) and infected+POM-1- or APCP-treated BALB/c mice were determined in 50 consecutive microscopic fields. (H–J) Splenocytes were isolated from control, infected and infected plus POM-1 (10 mg/kg body weight/day) or APCP (10 mg/kg body weight/day)-treated mice at 4 weeks post-infection. Splenocyte cell lysates were subjected to measurement of protein level expression of CD39 and CD73 by immunoblotting (H), and the levels of TNF- α , IL-12 and IFN- γ (I) and IL-10, TGF- β and IL-4 (J) by ELISA. In a set of splenocytes isolated from inhibitor-treated mice, the cells were treated with adenosine (100 μ M) for measurement of the cytokines (I,J). Bands were quantified by densitometry and are shown as bar graphs. Results are representative of three independent experiments and error bars are expressed as mean \pm s.d. $n=5$; ns, not significant, ** $P<0.01$, *** $P<0.001$ (Student's *t*-test).

at a dose of 10 mg/kg/day (Fig. 5A,B). Next, to find out the effectiveness of POM-1 and APCP on the progression of visceral leishmaniasis, infected BALB/c mice were injected with POM-1

and APCP at a dose of 10 mg/kg body weight/day up to 4 weeks at 3 day intervals after 11 days of infection (Fig. 5C) and the liver and spleen parasite burden (LDU) was determined. In both POM-1- and

APCP-treated mice, the parasite burden was found to be markedly lowered (65.1% and 71.4% in liver and 68.6% and 74.8%, respectively, in spleen at 4 weeks post-infection compared with infected control, $P < 0.001$) (Fig. 5D,E). Haematoxylin and Eosin (H&E) staining of sections of liver from both POM-1- and APCP-treated infected mice at 4 weeks post-infection revealed well-defined zones of granuloma formation, a hallmark of macrophage infiltration, compared with untreated infected mice (Fig. 5F). Although POM-1 or APCP treatment did not cause any visible health issues, the tissue architecture was found to be a little disarrayed even when administered without infection (Fig. 5I,C). The number of hepatic mature granulomas was found to be significantly increased in sections from both POM-1- and APCP-treated infected mice at 4 weeks post-infection in comparison to untreated infected mice (Fig. 5G). The protein expression of CD39 and CD73 was also determined in splenocytes isolated from control, infected and POM-1 and APCP administered mice after 4 weeks of infection (Fig. 5H). Although expression of CD39 and CD73 was significantly increased in splenocytes of infected mice at 4 weeks, treatment with the inhibitors did not show any change in the expression of CD39 and CD73. This is not surprising as POM-1 and APCP inhibit the ectonucleotidase activity rather than the expression level of CD39 and CD73. The levels of pro-inflammatory cytokines TNF- α and IL-12 were next analysed in the supernatant of splenocytes isolated from control, infected and POM-1- and APCP-treated infected mice at 4 weeks post-infection. In the case of POM-1 and APCP treatment, a marked increase in the level of TNF- α and IL-12 was observed at 4 weeks post-infection compared with infected controls (774.6 \pm 39.8 pg/ml TNF- α and 974.4 \pm 22.9 pg/ml IL-12 for POM-1 and 769.4 \pm 22.1 pg/ml TNF- α and 990.4 \pm 30.9 pg/ml IL-12 for APCP compared with 102.6 \pm 21 pg/ml TNF- α and 101.4 \pm 24.8 pg/ml IL-12 for infected control, respectively, $P < 0.001$) (Fig. 5I). In contrast, both POM-1 and APCP treatment led to marked decrease in the level of IL-10 and TGF- β evaluated at 4 weeks post-infection (403.2 \pm 35.3 pg/ml IL-10 and 439.1 \pm 52.1 pg/ml TGF- β for POM-1 and 408.4 \pm 22.8 pg/ml IL-10 and 434.4 \pm 50.7 pg/ml TGF- β for APCP compared with 1333.2 \pm 48.2 pg/ml IL-10 and 748.2 \pm 56.8 pg/ml TGF- β for infected control, respectively, $P < 0.001$) (Fig. 5J). Apart from regulating pro- and anti-inflammatory cytokines, POM-1 and APCP significantly modulated the Th1 cytokine IFN- γ and the Th2 cytokine IL-4. Both POM-1- and APCP-treated infected mice showed increased levels of IFN- γ (679.3 \pm 20.0 pg/ml for POM-1 and 597.7 \pm 27.5 pg/ml for APCP compared with 99.7 \pm 24.3 pg/ml for infected control, $P < 0.001$) at 4 weeks post infection (Fig. 5I). Regarding IL-4, a decrease in concentration was observed in inhibitor-treated infected mice (167.0 \pm 25.9 pg/ml for POM-1 and 128.3 \pm 30.5 pg/ml for APCP compared with 406.7 \pm 27.1 pg/ml for infected control, $P < 0.001$) (Fig. 5J). While measuring the cytokines in splenocytes isolated from infected inhibitor-treated mice, in one set of experiments, adenosine was added in the culture medium. The presence of adenosine partially reversed the inhibitory effect of POM-1 and APCP on cytokine production, thereby suggesting the importance of adenosine-receptor mediated signalling in infection (Fig. 5I,J). All these results validate our *in vitro* observations and further suggest the potential of CD39 and CD73 as chemotherapeutic targets in visceral leishmaniasis.

DISCUSSION

Immediately upon entry inside the macrophages, *Leishmania* not only has to fight against the antimicrobial defence system of the host, but also has to adjust to the harsh environment inside phagolysosomes

(Cunningham, 2002; Liu and Uzonna, 2012). This process calls for a rapid supply of energy, i.e. ATP, within the infected host cell. The present study was performed to delineate the source and fate of ATP of infected macrophages; which apart from being an energy currency, may also act as an important intracellular and extracellular signalling molecule (Corriden and Insel, 2010; Trautmann, 2009). In the study, we documented that upon *L. donovani* infection, there is a diminishing pattern of intracellular ATP during early hours of infection even though there was an increased rate of host cell glycolysis (Warburg effect) suggesting that there is an upregulated ATP concentration in infected cells. It was found that the parasite effluxes excess ATP to the extracellular environment via ion exporter pannexin-1. An increase in extracellular ATP may lead to inflammasome-mediated host cell immune activation (Cassel and Sutterwala, 2010) and to escape this, the parasite activates the host macrophage's ATP-degrading enzymes CD39 and CD73 for converting extracellular ATP (eATP) to adenosine. Extracellular adenosine signals through adenosine receptors A_{2A}R and A_{2B}R and provides an anti-inflammatory milieu essential for parasite survival (Fig. 6). To our knowledge, this is the first comprehensive report to demonstrate that ATP produced through altered host cell metabolism not only serves as an energy provider, but it also leads to generation of extracellular adenosine as an immune signalling molecule to facilitate infection (Fig. 6).

ATP is the primary energy currency of a cell, and according to our hypothesis, the iATP level of infected macrophages should increase. However, quantification of the iATP level significantly declined over time and this compelled us to evaluate the source of ATP, i.e. the metabolic status of *Leishmania*-infected macrophages during early infection. Contradictory to the iATP level, infection resulted in a heightened rate of glycolysis followed by lactate production. Intracellular pathogens are at times known to exploit host metabolic resources for their own benefit (Abu Kwaik and Bumann, 2013). Alteration of carbohydrate metabolism, the major ATP-generating pathway of the cell has also been reported (Bravo-Santano et al., 2018; Czyn et al., 2017). Pyruvate produced via glycolysis usually enters into the TCA cycle, but under certain circumstances gets converted into lactate. Although this is energetically less efficient compared with oxidative phosphorylation, it provides energy at a much faster rate. Intracellular infection by various pathogens like *Plasmodium berghei*, *Mycobacterium tuberculosis* and *Toxoplasma gondii* were also found to induce glycolysis and favour the intracellular persistence of the parasites (Li et al., 2008; Nelson et al., 2008; Shi et al., 2015). Metabolic profiling data of dengue virus-infected human cells have also shown that upregulation of glycolysis favours virus replication (Fontaine et al., 2015). Transcriptomic analysis of *L. major*-infected BMDM cells and extracellular flux analysis of *L. infantum*-infected cells have also shown an early upregulation of glycolysis following infection (Moreira et al., 2015; Rabhi et al., 2012).

However, the significant decrease in iATP level despite a higher glycolytic rate needed an explanation. Although ATP is a chemical source of cellular energy, it also plays an important role as a signalling molecule. An increase in cellular ATP concentration may lead to induction of apoptosis, and inhibition of macrophage apoptosis has been considered to be a survival strategy of the parasite (Saha et al., 2018). One way out for *Leishmania* to get rid of excess ATP is via its efflux from the host cell. Extracellular ATP level in infected macrophages indeed showed an increase over control. Inhibition studies in infected macrophages documented that amongst ATP release channel proteins (Kojima et al., 2017), pannexin-1 was mainly involved in the export of ATP and this export is necessary for parasite survival. A similar observation of ATP efflux was shown




Fig. 6. *L. donovani* channels ATP of host macrophages for establishment of infection. *L. donovani* upon infecting host macrophage cell leads to upregulation of glycolysis and lactate production. ATP generated from the upregulated glycolysis, however, is exported out of the cell through pannexin-1 channel proteins. The increased extracellular ATP (eATP) thus produced gets degraded by the cell surface ATP-hydrolyzing ectonucleotidases CD39 and CD73, leading to generation of adenosine. A low level of adenosine deaminase (ADA), which deaminates adenosine (ADO), leads to sustained level of extracellular adenosine (eADO) resulting in an anti-inflammatory environment via signalling through the adenosine receptors A_{2A}R and A_{2B}R. A_{2A}R signalling leads to downregulation of pro-inflammatory cytokines TNF- α and IL-12 whereas A_{2B}R signalling leads to upregulation of anti-inflammatory cytokines IL-10 and TGF- β . Thus, *Leishmania* utilizes host cell ATP in favour of its own survival.

with *P. falciparum* infection of human erythrocytes, where ATP is released via pannexin-1 channels (Alvarez et al., 2014). A significant increase of ATP in the cell supernatant was also found in vesicular stomatitis virus-infected cells and mice, where the increase took place in an exocytosis- and pannexin channel-dependent manner, leading to a reduction in virus replication (Zhang et al., 2017).

ATP in the extracellular environment (eATP) is kept under tight regulation as this molecule primarily acts as a 'danger signal' leading to activation of an inflammatory state. eATP is known to induce IL-12 production (Langston et al., 2003; Schnurr et al., 2000) and even lead to elimination of intracellular pathogens (Junger, 2011; Langston et al., 2003; Piccini et al., 2008). *L. amazonensis* has been shown to be sensitive in response to eATP possibly because of induction of host cell apoptosis (Chaves et al., 2009). However, in the present study, eATP showed a gradual decrease over time and coincided well with increased expression of two major ATP degrading enzymes, CD39 and CD73. Infection with *L. brazillensis*, *L. amazonensis* and *L. major* has also been shown to increase the expression and activity of CD39 and CD73 in dendritic cells (Figueiredo et al., 2012). However, a study involving PBMCs isolated from patients of visceral leishmaniasis has shown decreased frequency of CD73⁺ T-cells in comparison to healthy controls (Rai et al., 2011). This could be because similar stimulus is capable of differential induction of the same enzymes in different cell types but more investigation is needed to be carried out to confirm this. Although *Leishmania* upregulated the expression of both CD39 and CD73, the expression of ADA, which irreversibly deaminates adenosine, the final product of the reaction catalysed by CD73, remained almost undetectable. An increased level of extracellular adenosine during infection further validated the observation. Moreover, decreased parasite survival in

the CD39 and CD73 inhibited system was found to be reversed by adenosine added externally, which suggests that the parasite needs adenosine and deliberately keeps the expression of ADA at a low level. A similar finding was reported in *L. brazillensis* infection, where the presence of adenosine increased the severity of the disease (de Almeida Marques-da-Silva et al., 2008). It is worth mentioning that expression of CD26, which helps ADA to anchor on the surface of cells, has been found to be impaired in visceral leishmaniasis (Rai et al., 2012). The parasite *Leishmania* itself is also found to express surface ectonucleotidases to convert tri-nucleotides like ATP to adenosine (Berrêdo-Pinho et al., 2001; Meyer-Fernandes et al., 1997; Vasconcellos Rde et al., 2014). It has been shown that increased ectonucleotidase activity on the surface of these parasites is directly correlated with increased infectivity and virulence levels of the parasite (de Almeida Marques-da-Silva et al., 2008; Maioli et al., 2004; Peres et al., 2017). *L. amazonensis* has been found to have the highest activity of ectonucleotidases, leading to a higher concentration of adenosine (de Souza et al., 2010; Leite et al., 2012). *Leishmania* promastigotes having higher ectonucleotidase activity were also found to have greater inhibitory effect on immune activation of dendritic and Th cells (de Almeida Marques-da-Silva et al., 2008; Ribeiro et al., 2017). However, following engulfment by macrophages, *Leishmania* will be present within the phagolysosomes (Cunningham, 2002) and *Leishmania* ectonucleotidases probably will not have any effect on the macrophage's extracellular ATP concentration. Adenosine is a well-known anti-inflammatory molecule and has been found to favour the intracellular growth of many pathogens. Pathogenesis of *Toxoplasma*, *Salmonella* and *Helicobacter* has also been found to be dependent on the presence of host adenosine (Alam et al., 2009; Costales et al.,

2018; Mahamed et al., 2015). Adenosine generated outside a cell may exert its effect either by entering the cell through nucleoside transporter channels or through adenosine-receptor-mediated signalling pathway (Haskó and Cronstein, 2004; Khan et al., 2007). Both instances were found to be associated with creating an anti-inflammatory milieu inside the cell (Haskó and Cronstein, 2004), and justified the action of *Leishmania* to maintain a high adenosine level in the extracellular environment. Treatment with the adenosine uptake blocker dipyridamole, did not have any effect on the intracellular survival of parasites and indicated the involvement of adenosine receptors in *L. donovani* infection. Accordingly, we found that among the four adenosine receptor subtypes A₁, A_{2A}, A_{2B} and A₃, infection led to specific upregulation of A_{2A} and A_{2B}. Inhibition of signal transduction via these receptors showed upregulation of pro-inflammatory cytokines and downregulation of anti-inflammatory cytokines, ultimately suppressing the intracellular persistence of *Leishmania*. Few studies have shown an individual contribution of adenosine receptors in infection (Figueiredo et al., 2017; Lima et al., 2017; Vijayamahantesh et al., 2016); however, the present study is perhaps the first to report the crucial role played by both A_{2A}R and A_{2B}R in the pathogenesis caused by *L. donovani*.

In summary, the present study highlights how a parasite channel host ATP to the extracellular milieu for favouring its survival. Adenosine generated as a result of the process is crucial, as it inhibits inflammasome activation and activates parasite-favourable cytokine production. The importance of adenosine was also verified *in vivo*, where infected mice were treated with either the CD39 inhibitor, POM-1 or the CD73 inhibitor, APCP. Treatment with these inhibitors resulted in a significantly decreased liver and spleen parasite burden associated with a host-favourable pro-inflammatory environment. This study therefore not only unveiled the role of ATP in infection but identified ectonucleotidases as a potential therapeutic target in visceral leishmaniasis.

MATERIALS AND METHODS

Cells and parasites

L. donovani promastigotes of strain MHOM/IN/1983/AG83 were maintained in Medium 199 (M199, Invitrogen) supplemented with Hanks salt containing HEPES (12 mM), L-glutamine (20 mM), 10% heat inactivated FBS, 50 U/ml penicillin and 50 µg/ml streptomycin (Invitrogen) at 22°C. *L. donovani* axenic amastigotes were cultured as described previously (Ghosh et al., 2013). The murine macrophage cell line RAW 264.7 (National Repository for Cell lines/Hybridomas, India) was maintained at 37°C, 5% CO₂ in DMEM (Invitrogen) supplemented with 10% FBS, penicillin (100 U/ml) and streptomycin (100 µg/ml). Peritoneal macrophages (PM ϕ) were isolated from peritoneum of 6- to 8-week-old female BALB/c mice as described earlier (Gross et al., 2003) and cultured at 37°C with 5% CO₂ in DMEM supplemented with 10% heat-inactivated FBS, 100 µg/ml streptomycin and 100 U/ml penicillin. All the cells were checked for contamination before starting experiments.

Reagents and antibodies

Adenosine (A9251), adenosine 5'-monophosphate sodium salt (AMP, A1752), adenosine 5'-(α , β -methylene) diphosphate (APCP, M3763), adenosine 5'-triphosphate disodium salt hydrate (ATP, A6419), Adenosine 5'-[γ -thio]triphosphate tetralithium salt (ATP- γ S, A1388), carboxolone disodium salt (C4790), dipyridamole (D9766), D-(+)-glucose solution (G8769), 2-deoxyglucose (2-DG, D6134), gadolinium(III) chloride hexahydrate (G7532), L-glutamine (G8540), oligomycin (O4876) and sodium pyruvate solution (S8636) were obtained from Sigma-Aldrich (St Louis, MO, USA). Reagents such as ZM 241385 (ab120218) and MRS 1754 (ab120397) were purchased from Abcam (Cambridge, UK). POM-1 (21160) and ¹⁰Panx (3348) were obtained from Cayman Chemical (Ann Arbor, Michigan, USA) and Tocris (Bristol, UK), respectively.

The antibodies against pannexin-1 (Chen et al., 2019) (ab139715, 1:1000; Lot# GR3241457-3), CD39 (Shah et al., 2014) [ab108248, EPR3678(2), 1:1000; Lot# GR60408-12], CD73 (Xu et al., 2018a) (ab175396, 1:1000; Lot# GR318955-11) and ADA (ab175310, 1:1000; Lot# GR3181658-9) were purchased from Abcam (Cambridge, UK). The antibodies against adenosine A_{2A} receptor (Haschemi et al., 2007) (NBP1-39474, 7F6-G5-A2, 1:1000 for WB, 1:100 for microscopy; Lot# B-1) and Adenosine A_{2B} receptor (Xu et al., 2018b) (NBP2-41312, 1:2000 for WB, 1:100 for microscopy; Lot# 811-1802) were purchased from Novus Biologicals (Centennial, Colorado, USA). Mouse monoclonal β -actin antibody (A2228, 1:5000; Lot# 066M4860V), alkaline phosphatase-conjugated anti-mouse (A3562, 1:10,000; Lot # SLBV4379) and anti-rabbit (A3687, 1:10,000; Lot# SLBV1976) secondary antibodies were obtained from Sigma-Aldrich (St Louis, MO, USA). Goat polyclonal secondary antibody to Mouse IgG - H&L (TR) (ab6787, 1:200; Lot# GR3207552-3) and goat anti-Rabbit IgG H&L (FITC) (ab6717, 1:200; Lot# 3237804-2) were obtained from Abcam (Cambridge, UK).

Glycolysis and oxidative phosphorylation assay

Cellular glycolytic rate and oxidative phosphorylation levels were assessed using the Seahorse Bioscience Extracellular Flux Analyzer (XF24, Agilent Technologies, Santa Clara, California, USA) by measuring the extracellular acidification rate (ECAR) (indicative of glycolysis) and oxygen consumption rate (OCR) (indicative of respiration). RAW 264.7 cells (8×10^4) and peritoneal macrophages (2×10^5) were seeded in wells of XF24-cell culture microplates and incubated overnight. The monolayer of cells were infected with *L. donovani* parasites in a 1:10 ratio for the indicated time points. Before assessment, the cells were washed with XF assay medium, i.e. XF base medium supplemented with 4 mM glutamine for glycolysis stress test kit assay and for the mito stress test kit assay, the base medium was supplemented with 4 mM glutamine, 1 mM pyruvate and 5.5 mM glucose. The cells were then incubated with 525 µl of the respective media for 1 h at 37°C in non-CO₂ incubator. ECAR and OCR were then measured as recommended by the manufacturer. After completion of measurements, the cells were lysed in the plates using 25 µl RIPA lysis buffer (Sigma; St Louis, MO, USA) supplemented with 3 mM PMSF and 3 mM protease inhibitor mixture (Calbiochem; San Diego, CA, USA). Protein concentrations in the clear supernatant were determined using the Bradford assay and used to normalise the data which were analysed by Wave Desktop software (Wave 2.6.0) which is freely available for download.

Lactate measurement

Lactate in the cell culture medium was assessed using a colorimetric L-Lactate Assay Kit (ab65331, Abcam) according to the manufacturer's instructions. After the indicated time points, the cell culture supernatant was collected and deproteinised using the perchloric acid/KOH method. 50 µl of the deproteinised samples were mixed with 50 µl of reaction mixture and incubated at room temperature for 30 min in the dark. The samples were analysed by measuring absorbance at 450 nm.

ATP measurement

Intracellular and extracellular ATP levels were quantified using the luminescent ATP Determination Kit (A22066, Molecular Probes; Eugene, Oregon, USA). For intracellular ATP, cells (3×10^6) were infected with *L. donovani* promastigotes for varying periods of time. Cells were then washed with PBS and lysed in 200 µl Glo-lysis buffer (Promega; Madison, Wisconsin, USA). Samples were boiled; centrifuged and 500 ng protein was used to measure ATP content using the kit. For extracellular ATP, cells (5×10^4 /well) in 500 µl of fresh serum-free DMEM in 24-well plates were infected with live or PFA-fixed *L. donovani* promastigotes for varying time periods. At specific time points, supernatants were collected and used to quantify ATP.

Adenosine measurement

Adenosine in the cell culture supernatant after the indicated time points were quantified using the Adenosine Assay Kit (MET-5090, Cell Biolabs; San Diego, California, USA) according to the manufacturer's instructions.

The macrophage cells were incubated in serum-free DMEM and infected with live or PFA-fixed *L. donovani* for the specified time periods and then the cell culture medium was collected and centrifuged to obtain the clear supernatant. 50 μ l of the sample was used to quantify adenosine.

Ectonucleotidase activity

Ectonucleotidase activity of *L. donovani*-infected macrophages was determined using the Malachite Green colorimetric assay, which measures the amount of liberated inorganic phosphate (P_i) (Lanzetta et al., 1979). 2×10^5 macrophage cells (post-infection and treatment) were washed with reaction buffer (1 mM $CaCl_2$, 145 mM NaCl, 5 mM KCl, 10 mM HEPES and 1 mM $MgCl_2$, pH 7.4). The reaction was initiated by adding 1 mM ATP for CD39 or 1 mM AMP for CD73 to the culture dish and stopped by addition of the cell supernatant to a vial containing 1 ml of cold 25% activated charcoal in 0.1 M HCl. The vials were then centrifuged at 1500 g for 10 min and 50 μ l of each of the centrifuged samples were quantified for released inorganic P_i using different concentration of P_i as standard. Nonspecific P_i release was determined by incubating cells in the absence of substrate ATP or AMP, which were added after stopping the reaction.

Immunoblotting

Cells (RAW 264.7 and PM ϕ) were harvested and lysed using ice-cold lysis buffer (Cell Signaling Technology; Danvers, Massachusetts, USA) supplemented with 3 mM PMSF (Sigma) and 3 mM protease inhibitor mixture (Calbiochem; San Diego, CA, USA). Protein concentration in the clear supernatant was determined using the Bradford assay. From each sample, an equal amount of protein (50 μ g or 20 μ g) was resolved by 10% SDS-PAGE and transferred to nitrocellulose membrane, followed by blocking the membrane with 5% BSA and overnight incubation with specific primary antibody in the desired dilution. After washing with TBS, membranes were incubated with alkaline phosphatase-conjugated secondary antibody and detected by hydrolysis of 5-bromo-4-chloro-3-indolylphosphate (BCIP) chromogenic substrate. β -actin was used as a loading control. Band intensities were quantified by using ImageJ.

Cytotoxicity assay

Cytotoxic or cytostatic effects were observed for the administered doses of the reagents by examination of cellular morphology and cell viability of RAW 264.7, peritoneal macrophages, promastigotes and axenic amastigotes by MTT assay kit (Roche Applied Science; Penzberg, Germany).

Assessment of intracellular infection

In *in vitro* infection experiments, RAW 264.7 cells and peritoneal macrophage cells were infected with stationary phase *L. donovani* promastigotes at a 10:1 parasite/macrophage ratio (Kar et al., 2010). Infection was allowed to proceed for 4 h, the non-internalised parasites were removed by washing the plates with PBS and cells were cultured. Determination of the number of intracellular parasites was done by first fixing the cells in methanol and then staining with the nuclear stain DAPI (4',6'-diamidino-2-phenylindole; 1 μ g/ml) along with RNase A (10 μ g/ml) for 15 min (Gupta et al., 2016). Images were analysed using Olympus Fluoview software (version 3.1a, Tokyo, Japan).

Cytokine and chemokine analysis by ELISA

ELISA was performed using a sandwich ELISA kit (Quantikine M; R&D Systems, Minneapolis, MN, USA) for the various cytokines from PM ϕ and splenocytes. The detection limits of these assays were >5.1 , >2.5 , >2 , >4 , >4.6 and >2 pg/ml for the cytokines TNF- α , IL12p70, IFN- γ , IL-10, TGF- β and IL-4, respectively and >4.7 and >7.8 pg/ml for the chemokines CCL3 and CCL5, respectively.

Fluorescence microscopy

RAW 264.7 macrophages (2×10^5) were plated onto 18 mm² coverslips in DMEM medium and infected with *L. donovani* promastigotes (cell/parasite ratio 1:10). The cells were then washed twice in PBS and fixed with methanol at room temperature. The cells were permeabilised with 0.1% Triton X-100 and incubated with blocking solution (1% BSA) for 1 h at room temperature followed by overnight incubation with primary antibody against A₂B_R and

A₂B_R at 4°C. After washing, coverslips were incubated with both FITC- and Texas Red-conjugated secondary antibody for 1 h at room temperature. Cells were stained with DAPI (4',6'-diamidino-2-phenylindole, 1 μ g/ml) plus 10 μ g/ml RNase A in PBS to label the nucleus, mounted on slides and visualised under Olympus IX81 microscope equipped with a FV1000 confocal system using a 100 \times /60 \times oil immersion Plan Apo (N.A. 1.45) objectives. The images thus captured were analysed by Olympus Fluoview (version 3.1a; Tokyo, Japan) and mounted using Adobe Photoshop software.

In vivo infection

Animal maintenance and experiments were performed in accordance with the guidelines provided by the Committee for the Purpose of Control and Supervision of Experiments on Animals (New Delhi, India). The protocol was approved by the Institutional Animal Ethics Committee (IAEC). For *in vivo* infection, 6- to 8-week-old female BALB/c mice (~20 g) were injected via the tail vein with 1×10^7 stationary phase *L. donovani* promastigotes. Infection was assessed by removing the liver and spleen from infected mice up to 4 weeks after infection. Parasite burden was determined from Giemsa-stained impression smears (Srivastav et al., 2012). Liver and spleen parasite burdens, expressed as Leishman–Donovan units (LDU), were calculated as the number of amastigotes/1000 nucleated cells \times organ mass (in grams) (Murray et al., 1993). Splenocytes from BALB/c mice were isolated and cultured as described earlier (Arsenescu et al., 2005). Splenocytes were stimulated with soluble leishmanial antigen (20 mg/ml) and with adenosine (100 μ M) where applicable for 48 h, followed by measurement of pro-inflammatory cytokines by ELISA (Das et al., 2001).

Histopathology

Livers isolated from BALB/c mice were fixed in 10% formalin (Merck) and embedded in paraffin wax. Tissue sections (5 mm) were made with a microtome (Leica Biosystems) and stained with H&E to study the microarchitecture using light microscope (Nikon Microscope, Singapore). To quantify density of granuloma formation in the liver tissue, the total number of granulomas (immature and mature) were counted in 50 consecutive microscopic fields of sections derived from mice of the test groups.

Densitometric analysis

Densitometric analysis for all experiments was carried out using ImageJ software. Band intensities were quantitated densitometrically and the values obtained were normalised to endogenous control and expressed in arbitrary densitometric units. The ratios of OD of particular bands to endogenous control are represented as bar graphs adjacent to figures.

Statistical analysis

Most of the experiments were performed at least three times. All the data are shown as mean \pm s.d. of three independent experiments unless otherwise stated. *n* values are given in the legends of figure. Student's *t*-test was used to assess the statistical significance of intergroup comparisons using GraphPad Prism 7.0 and $P < 0.05$ was considered to be significant.

Acknowledgements

We thank Dr Sib Sankar Roy and Dr Sanghamitra Sengupta for the opportunity to carry out certain experiments in the Indian Institute of Chemical Biology (Kolkata, India) and University of Calcutta (Kolkata, India). We acknowledge DBT-CU-IPLS Core Facility for the confocal microscope facility and Dr Amrita Saha for helping with microscopy. We sincerely express our gratitude towards Subrata Modak, Avishek Nath, Dr Ranajoy Ghosh, and Dr Aniket Halder, Medical Technologists Laboratory, School of Digestive and Liver Diseases, the Institute of Post-Graduate Medical Education and Research (Kolkata, India) for histopathological facilities.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: M.B., P.G., A.U.; Methodology: M.B.; Software: M.B.; Validation: M.B., A.U.; Formal analysis: M.B., A.U.; Investigation: M.B., A.D., K.J.; Resources: A.U.; Data curation: M.B., A.U.; Writing - original draft: M.B., A.U.; Writing - review & editing: A.U.; Visualization: M.B.; Supervision: A.U.; Project administration: A.U.; Funding acquisition: A.U.

Funding

This work was supported by a Department of Biotechnology, Ministry of Science and Technology award (BT/HRD/NBA/38/03/2018), Indo Israel Grant, University Grants Commission [6-10/2016(IC)], Department of Science and Technology (SB/SO/BB-0055/2013), Department of Biotechnology (221/BT(Estt)/RD-40/2014), and University with Potential for Excellence II (UGC/148/UPE/ST1). M.B. received a fellowship from the University Grants Commission (New Delhi).

Supplementary information

Supplementary information available online at <http://jcs.biologists.org/lookup/doi/10.1242/jcs.239939.supplemental>

Peer review history

The peer review history is available online at <https://jcs.biologists.org/lookup/doi/10.1242/jcs.239939.reviewer-comments.pdf>

References

- Abu Kwak, Y. and Bumann, D. (2013). Microbial quest for food in vivo: 'nutritional virulence' as an emerging paradigm. *Cell. Microbiol.* **15**, 882-890. doi:10.1111/cmi.12138
- Alam, M. S., Kurtz, C. C., Wilson, J. M., Burnette, B. R., Wiznerowicz, E. B., Ross, W. G., Rieger, J. M., Figler, R. A., Linden, J., Crowe, S. E. et al. (2009). A2A adenosine receptor (AR) activation inhibits pro-inflammatory cytokine production by human CD4+ helper T cells and regulates Helicobacter-induced gastritis and bacterial persistence. *Mucosal Immunol.* **2**, 232-242. doi:10.1038/mi.2009.4
- Alam, M. S., Kuo, J. L., Ernst, P. B., Derr-Castillo, V., Pereira, M., Gaines, D., Costales, M., Bigley, E. and Williams, K. (2014). Ecto-5'-nucleotidase (CD73) regulates host inflammatory responses and exacerbates murine salmonellosis. *Sci. Rep.* **4**, 4486. doi:10.1038/srep04486
- Alvarez, C. L., Schachter, J., de Sá Pinheiro, A. A., Silva Lde, S., Verstraeten, S. V., Persechini, P. M. and Schwarzbaum, P. J. (2014). Regulation of extracellular ATP in human erythrocytes infected with Plasmodium falciparum. *PLoS ONE* **9**, e96216. doi:10.1371/journal.pone.0096216
- Antonoli, L., Pacher, P., Vizi, E. S. and Haskó, G. (2013). CD39 and CD73 in immunity and inflammation. *Trends Mol. Med.* **19**, 355-367. doi:10.1016/j.molmed.2013.03.005
- Arsenescu, R., Blum, A. M., Metwali, A., Elliott, D. E. and Weinstock, J. V. (2005). IL-12 induction of mRNA encoding substance P in murine macrophages from the spleen and sites of inflammation. *J. Immunol.* **174**, 3906-3911. doi:10.4049/jimmunol.174.7.3906
- Berrêdo-Pinho, M., Peres-Sampaio, C. E., Crispim, P. P. M., Belmont-Firpo, R., Lemos, A. P., Martiny, A., Vannier-Santos, M. A. and Meyer-Fernandes, J. R. (2001). A Mg-dependent ecto-ATPase in Leishmania amazonensis and its possible role in adenosine acquisition and virulence. *Arch. Biochem. Biophys.* **391**, 16-24. doi:10.1006/abbi.2001.2384
- Bouma, M. G., van den Wildenberg, F. A. and Buurman, W. A. (1996). Adenosine inhibits cytokine release and expression of adhesion molecules by activated human endothelial cells. *Am. J. Physiol.* **270**, C522-C529. doi:10.1152/ajpcell.1996.270.2.C522
- Bravo-Santano, N., Ellis, J. K., Mateos, L. M., Calle, Y., Keun, H. C., Behrends, V. and Letek, M. (2018). Intracellular staphylococcus aureus modulates host central carbon metabolism to activate autophagy. *mSphere* **3**, e00374-18. doi:10.1128/mSphere.00374-18
- Cassel, S. L. and Sutterwala, F. S. (2010). Sterile inflammatory responses mediated by the NLRP3 inflammasome. *Eur. J. Immunol.* **40**, 607-611. doi:10.1002/eji.200940207
- Cauwels, A., Rogge, E., Vandendriessche, B., Shiva, S. and Brouckaert, P. (2014). Extracellular ATP drives systemic inflammation, tissue damage and mortality. *Cell Death Dis.* **5**, e1102. doi:10.1038/cddis.2014.70
- Chaves, S. P., Torres-Santos, E. C., Marques, C., Figliuolo, V. R., Persechini, P. M., Coutinho-Silva, R. and Rossi-Bergmann, B. (2009). Modulation of P2X₇ purinergic receptor in macrophages by Leishmania amazonensis and its role in parasite elimination. *Microbes Infect.* **11**, 842-849. doi:10.1016/j.micinf.2009.05.001
- Chekeni, F. B., Elliott, M. R., Sandilos, J. K., Walk, S. F., Kinchen, J. M., Lazarowski, E. R., Armstrong, A. J., Penuela, S., Laird, D. W., Salvesen, G. S. et al. (2010). Pannexin 1 channels mediate 'find-me' signal release and membrane permeability during apoptosis. *Nature* **467**, 863-867. doi:10.1038/nature09413
- Chen, Y., Corriden, R., Inoue, Y., Yip, L., Hashiguchi, N., Zinkernagel, A., Nizet, V., Insel, P. A. and Junger, W. G. (2006). ATP release guides neutrophil chemotaxis via P2Y₂ and A₃ receptors. *Science* **314**, 1792-1795. doi:10.1126/science.1132559
- Chen, W., Zhu, S., Wang, Y., Li, J., Qiang, X., Zhao, X., Yang, H., D'Angelo, J., Becker, L., Wang, P. et al. (2019). Enhanced macrophage pannexin 1 expression and hemichannel activation exacerbates lethal experimental sepsis. *Sci. Rep.* **9**, 160. doi:10.1038/s41598-018-37232-z
- Corriden, R. and Insel, P. A. (2010). Basal release of ATP: an autocrine-paracrine mechanism for cell regulation. *Sci. Signal.* **3**, re1. doi:10.1126/scisignal.3104re1
- Costales, M. G., Alam, M. S., Cavanaugh, C. and Williams, K. M. (2018). Extracellular adenosine produced by ecto-5'-nucleotidase (CD73) regulates macrophage pro-inflammatory responses, nitric oxide production, and favors Salmonella persistence. *Nitric Oxide* **72**, 7-15. doi:10.1016/j.niox.2017.11.001
- Coutinho-Silva, R., Stahl, L., Raymond, M.-N., Jungas, T., Verbeke, P., Burnstock, G., Darville, T. and Ojcius, D. M. (2003). Inhibition of chlamydia infectious activity due to P2X₇R-dependent phospholipase D activation. *Immunology* **19**, 403-412. doi:10.1016/S1074-7613(03)00235-8
- Cronstein, B. N. (1994). Adenosine, an endogenous anti-inflammatory agent. *J. Appl. Physiol.* **76**, 5-13. doi:10.1152/jappl.1994.76.1.5
- Cunningham, A. C. (2002). Parasitic adaptive mechanisms in infection by leishmania. *Exp. Mol. Pathol.* **72**, 132-141. doi:10.1006/exmp.2002.2418
- Czyz, D. M., Willett, J. W. and Crosson, S. (2017). Brucella abortus induces a Warburg shift in host metabolism that is linked to enhanced intracellular survival of the pathogen. *J. Bacteriol.* **199**, e00227-17. doi:10.1128/JB.00227-17
- Das, L., Datta, N., Bandyopadhyay, S. and Das, P. K. (2001). Successful therapy of lethal murine visceral leishmaniasis with cystatin involves up-regulation of nitric oxide and a favorable T cell response. *J. Immunol.* **166**, 4020-4028. doi:10.4049/jimmunol.166.6.4020
- de Almeida Marques-da-Silva, E., de Oliveira, J. C., Figueiredo, A. B., de Souza Lima Junior, D., Carneiro, C. M., Rangel Fietto, J. L. and Crocco Afonso, L. C. (2008). Extracellular nucleotide metabolism in Leishmania: influence of adenosine in the establishment of infection. *Microbes Infect.* **10**, 850-857. doi:10.1016/j.micinf.2008.04.016
- de Menezes, J. P., Saraiva, E. M. and da Rocha-Azevedo, B. (2016). The site of the bite: Leishmania interaction with macrophages, neutrophils and the extracellular matrix in the dermis. *Parasit. Vectors* **9**, 264. doi:10.1186/s13071-016-1540-3
- de Souza, M. C., de Assis, E. A., Gomes, R. S., Marques da Silva Ede, A., Melo, M. N., Fietto, J. L. and Afonso, L. C. (2010). The influence of ecto-nucleotidases on Leishmania amazonensis infection and immune response in C57B/6 mice. *Acta Trop.* **115**, 262-269. doi:10.1016/j.actatropica.2010.04.007
- Dosch, M., Gerber, J., Jebbawi, F. and Beldi, G. (2018). Mechanisms of ATP release by inflammatory cells. *Int. J. Mol. Sci.* **19**, E1222. doi:10.3390/ijms19041222
- Figueiredo, A. B., Serafim, T. D., Marques-da-Silva, E. A., Meyer-Fernandes, J. R. and Afonso, L. C. (2012). Leishmania amazonensis impairs DC function by inhibiting CD40 expression via A2B adenosine receptor activation. *Eur. J. Immunol.* **42**, 1203-1215. doi:10.1002/eji.201141926
- Figueiredo, A. B., Souza-Testasica, M. C., Mineo, T. W. P. and Afonso, L. C. C. (2017). Leishmania amazonensis-induced cAMP triggered by adenosine A2B receptor is important to inhibit dendritic cell activation and evade immune response in infected mice. *Front. Immunol.* **8**, 849. doi:10.3389/fimmu.2017.00849
- Fontaine, K. A., Sanchez, E. L., Camarda, R. and Lagunoff, M. (2015). Dengue virus induces and requires glycolysis for optimal replication. *J. Virol.* **89**, 2358-2366. doi:10.1128/JVI.02309-14
- Fredholm, B. B., AP, I. J., Jacobson, K. A., Klotz, K. N. and Linden, J. (2001). International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol. Rev.* **53**, 527-552.
- Ghosh, K., Sharma, G., Saha, A., Kar, S., Das, P. K. and Ukil, A. (2013). Successful therapy of visceral leishmaniasis with curdlan involves T-helper 17 cytokines. *J. Infect. Dis.* **207**, 1016-1025. doi:10.1093/infdis/jis771
- Gross, A., Bouaboula, M., Casellas, P., Liautard, J.-P. and Dormand, J. (2003). Subversion and utilization of the host cell cyclic adenosine 5'-monophosphate/protein kinase A pathway by Brucella during macrophage infection. *J. Immunol.* **170**, 5607-5614. doi:10.4049/jimmunol.170.11.5607
- Guinamard, R., Simard, C. and Del Negro, C. (2013). Flufenamic acid as an ion channel modulator. *Pharmacol. Ther.* **138**, 272-284. doi:10.1016/j.pharmthera.2013.01.012
- Gupta, P., Srivastav, S., Saha, S., Das, P. K. and Ukil, A. (2016). Leishmania donovani inhibits macrophage apoptosis and pro-inflammatory response through AKT-mediated regulation of β-catenin and FOXO-1. *Cell Death Differ.* **23**, 1815-1826. doi:10.1038/cdd.2016.101
- Haschemi, A., Wagner, O., Marculescu, R., Wegiel, B., Robson, S. C., Gagliani, N., Gallo, D., Chen, J.-F., Bach, F. H. and Otterbein, L. E. (2007). Cross-regulation of carbon monoxide and the adenosine A_{2a} receptor in macrophages. *J. Immunol.* **178**, 5921-5929. doi:10.4049/jimmunol.178.9.5921
- Haskó, G. and Cronstein, B. (2013). Regulation of inflammation by adenosine. *Front. Immunol.* **4**, 85. doi:10.3389/fimmu.2013.00085
- Haskó, G. and Cronstein, B. N. (2004). Adenosine: an endogenous regulator of innate immunity. *Trends Immunol.* **25**, 33-39. doi:10.1016/j.it.2003.11.003
- Hoskin, D. W., Mader, J. S., Furlong, S. J., Conrad, D. M. and Blay, J. (2008). Inhibition of T cell and natural killer cell function by adenosine and its contribution to immune evasion by tumor cells (Review). *Int. J. Oncol.* **32**, 527-535. doi:10.3892/ijo.32.3.527
- Housley, G. D., Bringmann, A. and Reichenbach, A. (2009). Purinergic signaling in special senses. *Trends Neurosci.* **32**, 128-141. doi:10.1016/j.tins.2009.01.001
- Huang, S., Apasov, S., Koshiba, M. and Sitkovsky, M. (1997). Role of A_{2a} extracellular adenosine receptor-mediated signaling in adenosine-mediated

- inhibition of T-cell activation and expansion. *Blood* **90**, 1600-1610. doi:10.1182/blood.V90.4.1600
- Junger, W. G.** (2011). Immune cell regulation by autocrine purinergic signalling. *Nat. Rev. Immunol.* **11**, 201-212. doi:10.1038/nri2938
- Kar, S., Ukil, A., Sharma, G. and Das, P. K.** (2010). MAPK-directed phosphatases preferentially regulate pro- and anti-inflammatory cytokines in experimental visceral leishmaniasis: involvement of distinct protein kinase C isoforms. *J. Leukoc. Biol.* **88**, 9-20. doi:10.1189/jlb.0909644
- Kelly, B. and O'Neill, L. A. J.** (2015). Metabolic reprogramming in macrophages and dendritic cells in innate immunity. *Cell Res.* **25**, 771-784. doi:10.1038/cr.2015.68
- Khan, A. F., Thorne, P. R., Munoz, D. J., Wang, C. J. H., Housley, G. D. and Vljakovic, S. M.** (2007). Nucleoside transporter expression and adenosine uptake in the rat cochlea. *Neuroreport* **18**, 235-239. doi:10.1097/WNR.0b013e3280115918
- Kojima, S., Ohshima, Y., Nakatsukasa, H. and Tsukimoto, M.** (2017). Role of ATP as a key signaling molecule mediating radiation-induced biological effects. *Dose Response* **15**, 1559325817690638. doi:10.1177/1559325817690638
- Kreckler, L. M., Wan, T. C., Ge, Z.-D. and Auchampach, J. A.** (2006). Adenosine inhibits tumor necrosis factor- α release from mouse peritoneal macrophages via A2A and A2B but not the A3 adenosine receptor. *J. Pharmacol. Exp. Ther.* **317**, 172-180. doi:10.1124/jpet.105.096016
- Kronlage, M., Song, J., Sorokin, L., Isfort, K., Schwerdtle, T., Leipziger, J., Robaye, B., Conley, P. B., Kim, H. C., Sargin, S. et al.** (2010). Autocrine purinergic receptor signaling is essential for macrophage chemotaxis. *Sci. Signal.* **3**, ra55. doi:10.1126/scisignal.2000588
- Lammas, D. A., Stober, C., Harvey, C. J., Kendrick, N., Panchalingam, S. and Kumararatne, D. S.** (1997). ATP-induced killing of mycobacteria by human macrophages is mediated by purinergic P2Z(P2X7) receptors. *Immunity* **7**, 433-444. doi:10.1016/S1074-7613(00)80364-7
- Langston, H. P., Ke, Y., Gewirtz, A. T., Dombrowski, K. E. and Kapp, J. A.** (2003). Secretion of IL-2 and IFN- γ , but not IL-4, by antigen-specific T cells requires extracellular ATP. *J. Immunol.* **170**, 2962-2970. doi:10.4049/jimmunol.170.6.2962
- Lanzetta, P. A., Alvarez, L. J., Reinach, P. S. and Candia, O. A.** (1979). An improved assay for nanomole amounts of inorganic phosphate. *Anal. Biochem.* **100**, 95-97. doi:10.1016/0003-2697(79)90115-5
- Leite, P. M., Gomes, R. S., Figueiredo, A. B., Serafim, T. D., Tafuri, W. L., de Souza, C. C., Moura, S. A., Fietto, J. L., Melo, M. N., Ribeiro-Dias, F. et al.** (2012). Ecto-nucleotidase activities of promastigotes from *Leishmania* (Vianna) braziliensis relates to parasite infectivity and disease clinical outcome. *PLoS Negl. Trop. Dis.* **6**, e1850. doi:10.1371/journal.pntd.0001850
- Li, J. V., Wang, Y., Saric, J., Nicholson, J. K., Dirnhofer, S., Singer, B. H., Tanner, M., Wittlin, S., Holmes, E. and Utzinger, J.** (2008). Global metabolic responses of NMRI mice to an experimental *Plasmodium berghei* infection. *J. Proteome Res.* **7**, 3948-3956. doi:10.1021/pr800209d
- Lima, M. H. F., Sacramento, L. A., Quirino, G. F. S., Ferreira, M. D., Benevides, L., Santana, A. K. M., Cunha, F. Q., Almeida, R. P., Silva, J. S. and Carregaro, V.** (2017). *Leishmania infantum* parasites subvert the host inflammatory response through the adenosine A2A receptor to promote the establishment of infection. *Front. Immunol.* **8**, 815. doi:10.3389/fimmu.2017.00815
- Liu, D. and Uzonna, J. E.** (2012). The early interaction of *Leishmania* with macrophages and dendritic cells and its influence on the host immune response. *Front. Cell Infect. Microbiol.* **2**, 83. doi:10.3389/fcimb.2012.00083
- Mahamed, D. A., Toussaint, L. E. and Bynoe, M. S.** (2015). CD73-generated adenosine is critical for immune regulation during *Toxoplasma gondii* infection. *Infect. Immun.* **83**, 721-729. doi:10.1128/IAI.02536-14
- Maioli, T. U., Takane, E., Arantes, R. M. E., Fietto, J. L. and Afonso, L. C.** (2004). Immune response induced by New World *Leishmania* species in C57BL/6 mice. *Parasitol. Res.* **94**, 207-212. doi:10.1007/s00436-004-1193-6
- Meyer-Fernandes, J. R., Dutra, P. M. L., Rodrigues, C. O., Saad-Nehme, J. and Lopes, A. H. C. S.** (1997). Mg-dependent ecto-ATPase activity in *Leishmania tropica*. *Arch. Biochem. Biophys.* **341**, 40-46. doi:10.1006/abbi.1997.9933
- Moreira, D., Rodrigues, V., Abengoza, M., Rivas, L., Rial, E., Laforge, M., Li, X., Foretz, M., Viollet, B., Estaquier, J. et al.** (2015). *Leishmania infantum* modulates host macrophage mitochondrial metabolism by hijacking the SIRT1-AMPK axis. *PLoS Pathog.* **11**, e1004684. doi:10.1371/journal.ppat.1004684
- Murray, H. W., Miralles, G. D., Stoeckle, M. Y. and McDermott, D. F.** (1993). Role and effect of IL-2 in experimental visceral leishmaniasis. *J. Immunol.* **151**, 929-938.
- Nelson, M. M., Jones, A. R., Carmen, J. C., Sinai, A. P., Burchmore, R. and Wastling, J. M.** (2008). Modulation of the host cell proteome by the intracellular apicomplexan parasite *Toxoplasma gondii*. *Infect. Immun.* **76**, 828-844. doi:10.1128/IAI.01115-07
- North, R. A. and Verkhatsky, A.** (2006). Purinergic transmission in the central nervous system. *Pflugers Arch.* **452**, 479-485. doi:10.1007/s00424-006-0060-y
- O'Neill, L. A. J. and Pearce, E. J.** (2016). Immunometabolism governs dendritic cell and macrophage function. *J. Exp. Med.* **213**, 15-23. doi:10.1084/jem.20151570
- Olah, M. E. and Stiles, G. L.** (1995). Adenosine receptor subtypes: characterization and therapeutic regulation. *Annu. Rev. Pharmacol. Toxicol.* **35**, 581-606. doi:10.1146/annurev.pa.35.040195.003053
- Pelegrin, P. and Surprenant, A.** (2006). Pannexin-1 mediates large pore formation and interleukin-1 β release by the ATP-gated P2X7 receptor. *EMBO J.* **25**, 5071-5082. doi:10.1038/sj.emboj.7601378
- Peres, N. T. A., Cunha, L. C. S., Barbosa, M. L. A., Santos, M. B., de Oliveira, F. A., de Jesus, A. M. R. and de Almeida, R. P.** (2017). Infection of human macrophages by *leishmania infantum* is influenced by ecto-nucleotidases. *Front. Immunol.* **8**, 1954. doi:10.3389/fimmu.2017.01954
- Piccini, A., Carta, S., Tassi, S., Lasiglie, D., Fossati, G. and Rubartelli, A.** (2008). ATP is released by monocytes stimulated with pathogen-sensing receptor ligands and induces IL-1 β and IL-18 secretion in an autocrine way. *Proc. Natl. Acad. Sci. USA* **105**, 8067-8072. doi:10.1073/pnas.0709684105
- Rabhi, I., Rabhi, S., Ben-Othman, R., Rasche, A., Daskalaki, A., Trentin, B., Piquemal, D., Regnault, B., Descoteaux, A., Guieani-Tabbane, L. et al.** (2012). Transcriptomic signature of *Leishmania* infected mice macrophages: a metabolic point of view. *PLoS Negl. Trop. Dis.* **6**, e1763. doi:10.1371/journal.pntd.0001763
- Rai, A. K., Thakur, C. P., Velpandian, T., Sharma, S. K., Ghosh, B. and Mitra, D. K.** (2011). High concentration of adenosine in human visceral leishmaniasis despite increased ADA and decreased CD73. *Parasite Immunol.* **33**, 632-636. doi:10.1111/j.1365-3024.2011.01315.x
- Rai, A. K., Thakur, C. P., Kumar, P. and Mitra, D. K.** (2012). Impaired expression of CD26 compromises T-cell recruitment in human visceral leishmaniasis. *Eur. J. Immunol.* **42**, 2782-2791. doi:10.1002/eji.201141912
- Ribeiro, H. A. L., Maioli, T. U., de Freitas, L. M., Trieri, P. and Castiglione, F.** (2017). Modeling immune response to *leishmania* species indicates adenosine as an important inhibitor of th-cell activation. *Front. Cell Infect. Microbiol.* **7**, 309. doi:10.3389/fcimb.2017.00309
- Saha, A., Basu, M. and Ukil, A.** (2018). Recent advances in understanding *Leishmania donovani* infection: the importance of diverse host regulatory pathways. *IUBMB Life* **70**, 593-601. doi:10.1002/iub.1759
- Schaible, U. E. and Kaufmann, S. H. E.** (2007). Malnutrition and infection: complex mechanisms and global impacts. *PLoS Med.* **4**, e115. doi:10.1371/journal.pmed.0040115
- Schnurr, M., Then, F., Galambos, P., Scholz, C., Siegmund, B., Endres, S. and Eigler, A.** (2000). Extracellular ATP and TNF- α synergize in the activation and maturation of human dendritic cells. *J. Immunol.* **165**, 4704-4709. doi:10.4049/jimmunol.165.8.4704
- Schwiebert, L. M., Rice, W. C., Kudlow, B. A., Taylor, A. L. and Schwiebert, E. M.** (2002). Extracellular ATP signaling and P2X nucleotide receptors in monolayers of primary human vascular endothelial cells. *Am. J. Physiol. Cell Physiol.* **282**, C289-C301. doi:10.1152/ajpcell.01387.2000
- Shah, D., Romero, F., Stafstrom, W., Duong, M. and Summer, R.** (2014). Extracellular ATP mediates the late phase of neutrophil recruitment to the lung in murine models of acute lung injury. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **306**, L152-L161. doi:10.1152/ajplung.00229.2013
- Shi, L., Salamon, H., Eugenin, E. A., Pine, R., Cooper, A. and Gennaro, M. L.** (2015). Infection with *Mycobacterium tuberculosis* induces the Warburg effect in mouse lungs. *Sci. Rep.* **5**, 18176. doi:10.1038/srep18176
- Srivastav, S., Kar, S., Chande, A. G., Mukhopadhyaya, R. and Das, P. K.** (2012). *Leishmania donovani* exploits host deubiquitinating enzyme A20, a negative regulator of TLR signaling, to subvert host immune response. *J. Immunol.* **189**, 924-934. doi:10.4049/jimmunol.1102845
- Trautmann, A.** (2009). Extracellular ATP in the immune system: more than just a "danger signal". *Sci. Signal.* **2**, pe6. doi:10.1126/scisignal.256pe6
- Vasconcelos Rde, S., Mariotini-Moura, C., Gomes, R. S., Serafim, T. D., Firmino Rde, C., Silva, E. B. M., Castro, F. F., Oliveira, C. M., Borges-Pereira, L., de Souza, A. C. et al.** (2014). *Leishmania infantum* ecto-nucleoside triphosphate diphosphohydrolase-2 is an apyrase involved in macrophage infection and expressed in infected dogs. *PLoS Negl. Trop. Dis.* **8**, e3309. doi:10.1371/journal.pntd.0003309
- Vijayamahantesh, P. A., Amit, A., Kumar, S., Dikhit, M. R., Jha, P. K., Singh, A. K., Sinha, K. K., Pandey, K., Das, V. N. R., Das, P. et al.** (2016). Up regulation of A2B adenosine receptor on monocytes are crucially required for immune pathogenicity in Indian patients exposed to *Leishmania donovani*. *Cytokine* **79**, 38-44. doi:10.1016/j.cyto.2015.12.016
- Xu, S., Zhu, W., Shao, M., Zhang, F., Guo, J., Xu, H., Jiang, J., Ma, X., Xia, X., Zhi, X. et al.** (2018a). Ecto-5'-nucleotidase (CD73) attenuates inflammation after spinal cord injury by promoting macrophages/microglia M2 polarization in mice. *J. Neuroinflammation* **15**, 155. doi:10.1186/s12974-018-1183-8
- Xu, X., Zhu, Q., Niu, F., Zhang, R., Wang, Y., Wang, W., Sun, D., Wang, X. and Wang, A.** (2018b). A2BAR activation attenuates acute lung injury by inhibiting alveolar epithelial cell apoptosis both in vivo and in vitro. *Am. J. Physiol. Cell Physiol.* **315**, C558-C570. doi:10.1152/ajpcell.00294.2017
- Zhang, C., He, H., Wang, L., Zhang, N., Huang, H., Xiong, Q., Yan, Y., Wu, N., Ren, H., Han, H. et al.** (2017). Virus-triggered ATP release limits viral replication through facilitating IFN- β production in a P2X7-dependent manner. *J. Immunol.* **199**, 1372-1381. doi:10.4049/jimmunol.1700187
- Zhao, B., Gu, L., Liu, K., Zhang, M. and Liu, H.** (2017). Maxi-anion channels play a key role in glutamate-induced ATP release from mouse astrocytes in primary culture. *Neuroreport* **28**, 380-385. doi:10.1097/WNR.0000000000000759