

Corticosterone selectively decreases humoral immunity in female eiders during incubation

Sophie Bourgeon* and Thierry Raclot

Institut Pluridisciplinaire Hubert Curien (IPHC), Département Ecologie, Physiologie et Ethologie (DEPE), UMR 7178 CNRS-ULP, 23 rue Becquerel, F-67087 Strasbourg Cedex 2, France

*Author for correspondence (e-mail: sophie.bourgeon@c-strasbourg.fr)

Accepted 23 October 2006

Summary

Immunity is hypothesized to share limited resources with other physiological functions and this may partly account for the fitness costs of reproduction. Previous studies have shown that the acquired immunity of female common eider ducks (*Somateria mollissima*) is suppressed during their incubation, during which they entirely fast. Corticosterone was proposed to be an underlying physiological mechanism for such immunosuppression. Therefore, the current study aimed to assess the effects of exogenous corticosterone on acquired immunity in captive eiders. To this end, females were implanted with corticosterone pellets at different stages of their incubation fast. We measured total immunoglobulin levels, T-cell-mediated immune response, body mass and corticosterone levels in these females and compared them with those of control females prior to and after manipulation (i.e. corticosterone pellet implantation). To mimic corticosterone effects on body mass, we experimentally extended fasting duration in a group of females termed 'late fasters'. Implanted females had corticosterone levels 6 times higher and lost 35% more mass than control

females. Corticosterone levels in 'late fasters' were similar to those in control females but body mass was 8% lower in the former. The decrease in the immunoglobulin levels of corticosterone implanted females was twice as high as in control females, while the T-cell-mediated immune response was not significantly affected by the treatment. We found a decrease in the T-cell-mediated immune response only in 'late fasters' (by 60%), while the immunoglobulin level was not lower in this group than in corticosterone implanted or control females. Our study shows that in incubating eiders, exogenous corticosterone only decreased humoral immunity. We suggest that the immunosuppressive effect of corticosterone could be mediated through its effects on body reserves. Further experiments are required to determine the relationship between body condition and immune system in fasting birds.

Key words: bird, body reserve, fasting, glucocorticoid, immunosuppression.

Introduction

Life-history theory assumes that a major trade-off occurs between reproduction and survival, so that the costs associated with a given reproductive effort might have a deleterious impact on adult survival and future reproduction (Williams, 1966; Stearns, 1992). This concept is based on physiological trade-offs between resource-demanding activities within an individual. Recently, the costs of immune defenses have been emphasized (Gustafsson et al., 1994; Sheldon and Verhulst, 1996; Råberg et al., 1998). Namely, parental effort may induce immunosuppression in birds (Moreno et al., 1999). Acquired immunity can be classified into humoral immunity (mediated by B-lymphocytes) and cell-mediated immunity (mediated by T-lymphocytes) (Roitt et al., 1998). Deerenberg et al. (1997) showed in captive zebra finches *Taeniopygia guttata* that their humoral immunity was progressively reduced when the

reproductive effort is increased. Similarly, when raising experimentally enlarged broods, female pied flycatchers *Ficedula hypoleuca* exhibited reduced T-cell-mediated responses (Moreno et al., 1999). Previous studies have shown that the acquired immunity of female common eiders *Somateria mollissima* is suppressed during the incubation fast (Hanssen et al., 2004; Bourgeon et al., 2006), while its experimental activation has strong negative effects on the fitness of female eiders (Hanssen et al., 2004). Whatever the ultimate factors explaining such an immunosuppression in breeding eider ducks, the underlying physiological mechanisms still remain unknown. Among potential proximate factors underlying immunosuppression during reproduction, a link between immunocompetence and hormonal changes has been proposed (see Deerenberg et al., 1997).

Glucocorticoids are an essential component of the

endogenous immunoregulatory network, while also being associated with stress. Hence, these hormones establish a close endocrine link between immunocompetence and stress (Apanius, 1998). Råberg et al. (Råberg et al., 1998) hypothesized that corticosterone, secreted during stressful activities, reduces the acquired immune function. However, while the T-cell-mediated immune response was suppressed by experimentally elevated corticosterone levels in non-breeding New Jersey house sparrows *Passer domesticus* (Martin II et al., 2005), it did not significantly covary with natural corticosterone concentrations in breeding barn swallows *Hirundo rustica* (Saino et al., 2002). Similarly in female common eiders both components of the acquired immunity decreased independently of plasma corticosterone level, which itself did not vary significantly over the incubation period (Bourgeon et al., 2006). However, in several bird species, organisms are metabolically prepared for a long-term fast (Le Maho et al., 1981; Cherel et al., 1988; Lindgård et al., 1992). Indeed, fasting is first characterized by exhaustion of glycogen reserves (phase I) before a long period of protein sparing and preferential mobilization of fat stores (phase II), which is followed by a period of increased net protein catabolism (phase III). While phases I/II are characterized by the maintenance of low plasma levels of corticosterone, phase III of fasting is associated with an increase in plasma corticosterone levels. Such an elevation could be responsible for the increase in protein catabolism (proteolysis), because corticosterone is known to mobilize peripheral calorie stores for glucose production and energy utilization (Robin et al., 1998). Depending on conditions during the breeding season, incubating female eiders can enter phase III of fasting (see Le Maho, 1981; Le Maho et al., 1983).

Our main objective in this study was to examine the effects of increased plasma corticosterone levels on both components of the acquired immunity in wild female common eiders at different stages of their incubation fast. To this end, female eiders, nesting in the high Arctic, were implanted with corticosterone pellets both at the beginning and at the end of incubation. Subsequently, female total immunoglobulin levels, T-cell-mediated immune response, body mass and plasma corticosterone levels were measured and compared with those in control birds prior to and after manipulation. To mimic corticosterone effects on body mass, fasting duration in a group of females (termed 'late fasters') was experimentally extended. Body mass loss in these females was therefore increased, while corticosterone levels remained unaltered (birds remained in phase II and did not reach phase III). This allowed us to discriminate the effects of increased body mass loss on the acquired immunity from those of elevated corticosterone levels. We predicted that an increased plasma corticosterone level, which increases proteolysis during the female fast, should have immunosuppressive effects. Implanted females should therefore show T-cell-mediated immune responses and/or an immunoglobulin level lower than in control females but similar to that of 'late fasters'.

Materials and methods

The study was conducted in a colony of common eider ducks *Somateria mollissima* L. on Prins Heinrich Island, Kongsfjorden, Svalbard Archipelago (78°55'N, 20°07'E) between June and July 2005. The breeding colony in the study site contained about 105 nests. Females in the study area laid between one and five eggs, but a clutch size of 3–4 eggs was most common (20% and 61%, respectively, $N=105$). Eider ducklings are precocial and are cared for by the female only. Incubation lasts between 24 and 26 days (Korschgen, 1977). All birds started their incubation between June 07 and June 15, the main laying period for the colony. Ducks that laid their eggs after this period were not considered in this study. Ambient temperatures in June and July ranged from 2 to 10°C.

Sampling protocol

Nests were checked at least every second day throughout the study period. This was done to determine initial clutch size but also to investigate the rate of egg predation and nest desertion. A clutch of eggs was considered complete when no additional egg was laid during a 2-day period (Yoccoz et al., 2002). Females that suffered partial egg predation were excluded from this study. Female eiders were caught on their nests using a bamboo pole with a nylon snare. Blood was collected from the brachial vein within 3 min of capture, stored in tubes containing EDTA (an anticoagulant agent) and kept on ice until being centrifuged in the laboratory (at 4472 g), for 5 min, at 4°C. Plasma samples were stored at -20°C and subsequently used to measure immunoglobulin and corticosterone levels. After blood sampling, body size was recorded (wing and tarsus lengths) and birds were weighed using a portable electronic balance (to ± 2 g).

Experimental groups and corticosterone implantation

A total of 36 females with mean clutch sizes of 3–4 eggs (3.69 ± 0.08 eggs, $N=36$) (mean \pm s.e.m.) was used in this study. Freely incubating females were classified into two experimental groups: corticosterone ($N=18$) and control females ($N=18$). Previous results have shown that both components of the acquired immune system are decreased during incubation in eider ducks (Bourgeon et al., 2006), so females from both groups were caught during the first part of incubation (10.64 ± 0.62 days, $N=22$) (mean \pm s.e.m.) (11 females from each group) and near the end of incubation (20.07 ± 0.24 days, $N=14$) (7 females from each group). After capture, a blood sample was taken, and body size and mass were recorded. Half of the females were then implanted with corticosterone pellets (see below), while the others underwent the same procedure without actual implantation. Birds were held in cages for the following 5 days at ambient temperatures, with snow given as fresh water *ad libitum*. Four days after the treatment (18.31 ± 0.87 days into their incubation, $N=36$) (mean \pm s.e.m.), another blood sample was taken, a phytohemagglutinin (PHA) skin test conducted (see below), and body mass determined. Birds were released 24 h later, after the PHA skin test had been read. Additionally, in a group

termed 'late fasters', which consisted of seven captive females with mean clutch sizes from 2–5 eggs (3.43 ± 0.53 eggs, $N=7$) (mean \pm s.e.m.), fasting duration was experimentally extended. Birds remained in phase II of fasting during experimentation and never entered into phase III. This protocol allowed discrimination of the effects of increased body mass loss on the acquired immunity from those of elevated corticosterone levels. To this end, these females had been incubating eggs for at least 23 days (24.86 ± 0.40 days, $N=7$) (mean \pm s.e.m.), i.e. eggs were close to hatching. We took a blood sample and recorded body size and mass. Females were then held in captivity for 5 days (29.86 ± 0.40 days into their incubation, $N=7$) (mean \pm s.e.m.), at which point a further blood sample was taken, a PHA skin test performed (see below), and body mass recorded. Females were released 24 h after this, when the PHA skin test had been read.

Corticosterone pellets (100 mg, 21 day release, G-111) were obtained from Innovative Research of America (Sarasota, FL, USA). The implants slowly release the hormone, which enters the bloodstream. In preliminary trials we found that this dose was sufficient to induce marked increases in the level of plasma corticosterone and accelerate body mass loss as early as 2 days after the start of treatment (S.B. and T.R., unpublished data). For the implantation of the corticosterone pellets we followed the recommendations of the manufacturer. Briefly, a small patch of skin at the back side of the bird's neck was shaved and disinfected using alcohol and betadine (iodine solution). A small incision equal to the size of the pellet was made and the implant was inserted underneath the skin. The skin was closed with a single stitch, using surgical thread. The wound was cleaned with betadine and sprayed with an aluminium powder. The surgical procedure required less than 10 min. Control animals underwent the identical procedure, without actual insertion of a pellet.

T-cell-mediated immune response: PHA skin test

For this test we challenged one wing-web with the mitogenic PHA, while the other wing-web (control) was injected with phosphate buffered saline (PBS). Briefly, 100 μ l of 5 mg ml⁻¹ PHA (Sigma L 8754) in PBS were injected intradermally into the right wing-web, while the left wing-web was injected with an equal volume of PBS. This procedure was shown to induce little physiological stress in birds (Merino et al., 1999). Injection sites on the wing-web were measured using a micrometer calliper (three readings) just before and 24 h after injection with PHA or PBS. The T-cell-mediated immune response was taken as the difference between the two wing-web swellings.

Immunoglobulin levels: ELISA test

A sensitive ELISA method was used to determine the amount of serum immunoglobulins in eider duck blood. This method using commercial anti-chicken antibodies has so far been validated in six wild avian species (Martinez et al., 2003). Despite the fact that Anseriforms have an additional immunoglobulin isotype (IgY), which is not found in other

birds (Parham, 1995), we assumed linear cross-reactivity. Accordingly, the values obtained were used as relative immunoglobulin levels.

To determine the linear range of the sigmoid curve, ELISA plates were coated with serial dilutions of serum (100 μ l) in carbonate–bicarbonate buffer (0.1 mol l⁻¹, pH 9.6) and incubated overnight at 4°C. We selected the data obtained from trials using the serum dilution nearest to the centre of its linear range. ELISA plates were then coated with 100 μ l of diluted serum samples from female eiders (two samples per female diluted to 1/32 000 in carbonate–bicarbonate buffer) and incubated for 1 h at 37°C. After a second incubation overnight at 4°C, the plates were washed with a solution (200 μ l) of phosphate buffer saline and Tween (PBS-Tween), and a diluent (100 μ l), containing 5% powdered milk in PBS was added. Following incubation for 1 h at 37°C, the plates were washed with PBS-Tween buffer. 100 μ l of anti-chicken conjugate (Sigma A 9046) was added at 1:250 (v/v) and the plates were incubated for 2 h at 37°C. After three washes, the plates were filled with 100 μ l of a solution consisting of 2,2'-azino-bis-(3-ethylbenzthiazoline-6-sulphonic acid (ABTS) and concentrated hydrogen peroxide diluted to 1:1000. Following incubation for 1 h at 37°C, absorbance was measured at 405 nm using a plate spectrophotometer (Awareness Technology, Inc., Palm City, FL, USA).

Assessment of the corticosterone levels

Corticosterone concentrations were determined by radioimmunoassay (RIA) in our laboratory using an ¹²⁵I RIA double antibody kit from ICN Biomedicals (Costa Mesa, CA, USA). The corticosterone RIA had an intra-assay variability of 7.1% ($N=10$ duplicates) and an inter-assay variability of 6.5% ($N=15$ duplicates).

Statistical analysis

Statistical analysis was conducted with SPSS 12.0.1 (SPSS Inc., Chicago, IL, USA). Values are means \pm standard error (s.e.m.). Corticosterone levels were not normally distributed (Kolmogorov–Smirnov test, $P<0.05$). Hence, data were log transformed to meet parametric assumptions, before parametric tests were used. Repeated measures (RM) two-way ANOVA was used to test for the effects of the treatment and incubation stage on corticosterone levels, body mass and immunoglobulin level. Two-way ANOVA was used to test for the effects of the treatment and incubation stage on the T-cell-mediated immune response. Linear regression analysis was used to assess the relationships between all parameters measured.

Results

Tables 1 and 2 provide biological, hormonal and immune profiles for the female eider ducks (before and after implantation) sampled during the first part and near the end of their incubation period, respectively. Initial clutch size, tarsus and wing lengths, days of incubation, body mass, immunoglobulin level and corticosterone levels were not

Table 1. Profiles for both experimental groups of captive female eiders (corticosterone and control group), sampled during their early phase of incubation

	Female group	
	1: Corticosterone females	2: Control females
Before implantation		
Initial clutch size (eggs)	3.82±0.12 ^a	3.54±0.16 ^a
Tarsus length (cm)	6.15±0.05 ^a	6.21±0.04 ^a
Wing length (cm)	29.14±0.15 ^a	29.45±0.17 ^a
Time of sampling (days)	10.27±0.76 ^a	11.00±1.00 ^a
Body mass at sampling (g)	1687±39 ^a	1753±34 ^a
Immunoglobulin level (absorbance units)	1.04±0.06 ^a	0.94±0.07 ^a
[Corticosterone] (ng ml ⁻¹)	14.34±2.17 ^a	14.11±1.63 ^a
After implantation		
Time of sampling (days)	14.27±0.76 ^a	15.00±1.00 ^a
Body mass at sampling (g)	1430±33 ^a	1575±32 ^b
Total body mass loss (g)	257.09±13.12 ^a	178.55±8.42 ^b
Immunoglobulin level (absorbance units)	0.57±0.05 ^a	0.70±0.05 ^a
T-cell-mediated immune response (mm)	0.72±0.14 ^a	0.97±0.18 ^a
[Corticosterone] (ng ml ⁻¹)	109.90±8.16 ^a	14.70±1.85 ^b

Time of sampling g' indicates incubation stage.
 Values are means ± s.e.m., N=11 for each group.
 Different superscript letters indicate a significant difference between groups (T-tests).

significantly different between the experimental and control group before manipulation (i.e. corticosterone pellet implantation) at both incubation stages.

Effects of corticosterone implants on body mass

As expected, implants induced a significant increase in

corticosterone levels (RM two-way ANOVA: effects of repetition: $F_{1,32}=65.95$, $P<0.0001$; effects of treatment: $F_{1,32}=54.16$, $P<0.0001$; effects of incubation stage: $F_{1,32}=0.06$, $P=0.81$; interaction: $F_{1,32}=0.73$, $P=0.40$). Four days after implantation, corticosterone levels in implanted birds were 6 times higher than in control females, independent of incubation

Table 2. Profiles for the three experimental groups of captive female eiders (corticosterone, control and 'late fasters'), sampled near the end of their incubation

	Female group		
	1: Corticosterone	2: Control	3: Late fasting
Before implantation			
Initial clutch size (eggs)	3.71±0.18 ^a	3.71±0.18 ^a	3.43±0.53 ^a
Tarsus length (cm)	6.16±0.05 ^a	6.07±0.04 ^a	6.01±0.06 ^a
Wing length (cm)	29.19±0.24 ^a	29.14±0.20 ^a	29.29±0.18 ^a
Time of sampling (days)	20.29±0.29 ^a	19.86±0.40 ^a	24.86±0.40 ^b
Body mass at sampling (g)	1549±27 ^a	1571±37 ^a	1441±36 ^b
Immunoglobulin level (absorbance units)	0.70±0.06 ^a	0.70±0.04 ^a	0.82±0.07 ^a
[Corticosterone] (ng ml ⁻¹)	18.36±3.09 ^a	18.35±4.45 ^a	13.26±1.99 ^a
After implantation			
Time of sampling (days)	24.29±0.29 ^a	23.86±0.40 ^a	29.86±0.40 ^b
Body mass at sampling (g)	1315±23 ^{a,b}	1394±36 ^a	1285±27 ^b
Total body mass loss (g)	234.00±15.93 ^a	177.14±9.54 ^b	155.43±10.45 ^b
Immunoglobulin level (absorbance units)	0.47±0.03 ^a	0.52±0.04 ^a	0.57±0.05 ^a
T-cell-mediated immune response (mm)	1.00±0.13 ^a	1.27±0.18 ^a	0.47±0.08 ^b
[Corticosterone] (ng ml ⁻¹)	111.10±18.06 ^a	22.33±5.29 ^b	16.72±2.21 ^b

'Time of sampling' indicates incubation stage.
 Values are means ± s.e.m., N=7 for each group.
 Different superscript letters indicate a significant difference between groups (one-way ANOVA).

Table 3. Results for linear regressions between immune parameters, body mass and corticosterone level in captive female eiders, sampled during the first part of incubation

	Immunoglobulin level (absorbance units)	T-cell-mediated immune response (mm)	[Corticosterone] (ng ml ⁻¹)
Before implantation			
Body mass (g)	$F_{1,21}=1.19, P=0.29 (+)$	–	$F_{1,21}=0.07, P=0.80 (+)$
Immunoglobulin level (absorbance units)	–	–	$F_{1,21}=3.99, P=0.06 (+)$
[Corticosterone] (ng ml ⁻¹)	–	–	–
After implantation			
Body mass (g)	$F_{1,21}=2.37, P=0.14 (+)$	$F_{1,21}=5.86, P=0.02 (+)$	$F_{1,21}=6.80, P=0.02 (-)$
Immunoglobulin level (absorbance units)	–	$F_{1,21}=3.58, P=0.07 (+)$	$F_{1,21}=2.68, P=0.12 (-)$
T-cell-mediated immune response (mm)	–	–	$F_{1,21}=1.13, P=0.30 (-)$
[Corticosterone] (ng ml ⁻¹)	–	–	–

(+), (–) indicate positive or negative relationships, respectively.

stage. Moreover, corticosterone levels in ‘late fasters’ were far below the levels of implanted females. In fact, they were not significantly different from the levels in control females sampled at the end of their incubation period (*T*-test: $t=-0.03$; $N=25$; $P=0.97$). Body mass in corticosterone implanted females was significantly decreased by 18%, while it only decreased by 12% in control females (RM two-way ANOVA: effects of repetition: $F_{1,32}=1191.35, P<0.0001$; effects of treatment: $F_{1,32}=30.46, P<0.0001$; effects of incubation stage: $F_{1,32}=0.10, P=0.33$; interaction: $F_{1,32}=0.78, P=0.38$). Four days after implantation, implanted females were significantly lighter than control females (by 8%), independent of the incubation stage. In fact, body mass loss per day in implanted females was 35% higher than in control females, regardless of the incubation stage (two-way ANOVA: effects of treatment: $F_{1,32}=25.89, P<0.0001$; effects of incubation stage: $F_{1,32}=1.29, P=0.26$; interaction: $F_{1,32}=0.01, P=0.92$). Corticosterone levels in birds during early incubation were negatively related to body mass only after the implantation (Table 3), when high corticosterone levels were associated with low body mass values. However, there was no significant relationship between

these parameters in birds near the end of their incubation period, neither before nor after implantation (Table 4). Finally, body mass of ‘late fasters’ was not significantly different from that of implanted females but was significantly lower than in control females sampled at the end of their incubation (8%; Table 2) (RM one-way ANOVA: effects of repetition: $F_{1,18}=707.63, P<0.0001$; effects of treatment: $F_{2,18}=10.89, P=0.001$).

Effects of corticosterone on the T-cell-mediated immune response and immunoglobulin level

Corticosterone implants had no significant effect on the T-cell-mediated immune response (two-way ANOVA: effects of treatment: $F_{1,32}=2.49, P=0.12$; effects of incubation stage: $F_{1,32}=2.94, P=0.10$; interaction: $F_{1,32}=0.003, P=0.95$; Fig. 1). Responses in implanted females were similar to that of control females, independent of the incubation stage. However, the immune response in ‘late fasters’ was significantly reduced when compared with implanted and control females at the end of their incubation (53% and 63%, respectively) (one-way ANOVA: effects of treatment: $F_{2,18}=9.39, P=0.002$; Fig. 1).

Table 4. Results for linear regressions between immune parameters, body mass and corticosterone level in captive female eiders, sampled near the end of incubation

	Immunoglobulin level (absorbance units)	T-cell-mediated immune response (mm)	[Corticosterone] (ng ml ⁻¹)
Before implantation			
Body mass (g)	$F_{1,20}=3.00, P=0.10 (-)$	–	$F_{1,20}=0.26, P=0.61 (-)$
Immunoglobulin level (absorbance units)	–	–	$F_{1,20}=7.70, P=0.01 (-)$
[Corticosterone] (ng ml ⁻¹)	–	–	–
After implantation			
Body mass (g)	$F_{1,20}=4.16, P=0.05 (-)$	$F_{1,20}=0.17, P=0.68 (+)$	$F_{1,20}=0.43, P=0.52 (-)$
Immunoglobulin level (absorbance units)	–	$F_{1,20}=1.16, P=0.29 (-)$	$F_{1,20}=2.76, P=0.11 (-)$
T-cell-mediated immune response (mm)	–	–	$F_{1,20}=1.04, P=0.32 (+)$
[Corticosterone] (ng ml ⁻¹)	–	–	–

(–), (–) indicate positive or negative relationships, respectively.




Fig. 1. Effects of treatment on wing-web swelling in female eiders: corticosterone implanted (hatched bars), sham implanted (open bars), and 'late fasting' females (cross hatched bars). Values are means \pm s.e.m. a,b indicate a significant difference between groups (LSD *post-hoc* tests).

There was no relationship between the T-cell-mediated immune response and plasma corticosterone levels at any stage (Tables 3 and 4). However, there was a positive significant relationship between the T-cell-mediated immune response and body mass but only during early incubation (Tables 3 and 4), so that the immune response was stronger in heavier females.

After 4 days, the immunoglobulin level in implanted females was significantly decreased by 45% and 33%, when sampled during early and late incubation, respectively. The corresponding decline in immunoglobulin level of control females was only 25% and 25% (RM two-way ANOVA: effects of repetition: $F_{1,32}=105.10$, $P<0.0001$; effects of treatment: $F_{1,32}=7.34$, $P=0.01$; effects of incubation stage: $F_{1,32}=7.14$, $P=0.01$; interaction: $F_{1,32}=2.77$, $P=0.11$; Fig. 2), indicating that corticosterone pellet implantation had a negative effect on the female immunoglobulin level. This effect was stronger when implantation occurred during early incubation rather than during late incubation. The immunoglobulin level in 'late fasters' was not significantly different from that of implanted or control females sampled near the end of their incubation (RM one-way ANOVA: effects of repetition: $F_{1,18}=78.15$, $P<0.0001$; effects of treatment: $F_{2,18}=0.74$, $P=0.49$; Fig. 2). We found no relationship between the immunoglobulin and plasma corticosterone levels, either before or after the treatment during early incubation. The same held true for the period after implantation, when birds were near the end of incubation. By contrast, both parameters were negatively related before implantation of ducks that were near the end of the incubation period. High corticosterone levels were then associated with a low immunoglobulin level (Table 4; Fig. 3). Furthermore, there was no relationship between the immunoglobulin level and body mass, neither before nor after the treatment at any stage of incubation (Tables 3 and 4). Finally, we did not find a relationship between




Fig. 2. Effects of treatment on immunoglobulin level before (hatched bars) and after (open bars) manipulation in female eiders. Values are means \pm s.e.m. a-d indicate a significant difference between groups (LSD *post-hoc* tests).

both components of the acquired immunity after hormone implantation at any stage of the eider duck incubation period (Tables 3 and 4). Hence, the immunoglobulin level was independent of the T-cell-mediated immune response.

Discussion

It has previously been shown that the acquired immunity is significantly decreased during the incubation fast of female common eiders (Hanssen et al., 2004; Bourgeon et al., 2006). Among other potential scenarios, reduced immunocompetence during reproduction has been attributed to hormonal regulation, in particular through the action of glucocorticoids (Deerenberg




Fig. 3. Relationship between corticosterone level and immunoglobulin level before implantation in female eiders sampled near the end of incubation.

et al., 1997; Råberg et al., 1998). Hence, the main objective of the current study was to investigate the potential physiological mechanisms underlying such immunosuppression. We therefore assessed the effects of exogenous corticosterone on both components of the acquired immune system in female eiders during different stages of their incubation fast. In addition, a group of females whose fasting duration was experimentally extended was used to discriminate the effects of increased body mass loss on the acquired immunity from those of elevated corticosterone levels.

Experimentally increased plasma corticosterone levels only affected one of the two components of the acquired immunity. The immunoglobulin level in implanted females was significantly decreased when compared with that of control females. This response was strongest when birds were implanted at the beginning of their incubation fast. However, the T-cell-mediated immune response was not significantly affected by the treatment. Paradoxically, there was no significant relationship between plasma corticosterone and immunoglobulin levels after the implantation at any incubation stage. By contrast, before the treatment in ducks that were near the end of their incubation, high corticosterone levels seemed to be associated with a low immunoglobulin level.

Whatever the effects of exogenous corticosterone, and similar to an earlier investigation (Bourgeon et al., 2006), we did not find a significant relationship between both components of the acquired immunity. This lends support to the view that variations in one component of the acquired immunity are not necessarily a reliable indicator of changes in the other (Norris and Evans, 2000). In fact, in the present study, the immunoglobulin level seemed to be more sensitive to corticosterone treatment than the T-cell-mediated immune response. In control females, the immunoglobulin level significantly decreased throughout incubation, while the T-cell-mediated immune response did not vary significantly. This would seem to contrast with the finding of a previous investigation (Bourgeon et al., 2006) suggesting that both components significantly decrease throughout the incubation fast of eider ducks. This apparent discrepancy could lie in the fact that smaller sample sizes have been used, which is reinforced by high variances observed in this immune response. Moreover, we cannot exclude the possibility that effects of corticosterone on the T-cell-mediated immune response might require more time (see Dhabhar and McEwen, 1997) and/or higher corticosterone concentrations. In the current study, a significant decrease in the T-cell-mediated immune response was only observed in 'late fasters', while their immunoglobulin level was not lower than that of corticosterone implanted or control females. Implanted females significantly lost more weight than control females, which is consistent with the observation (Cherel et al., 1988) that high levels of corticosterone increase proteolysis in fasting birds. In a preliminary study on captive female eiders, high doses of exogenous corticosterone, administered for a few days, induced a rise in plasma levels of uric acid, indicating protein breakdown (Crisuolo et al., 2005).

Corticosterone levels of 'late fasters' in the current study were similar to that of control females but body mass was 8% lower in the former. Hence, the T-cell-mediated immune response appears to be more sensitive to body mass loss than to elevated levels of corticosterone, supporting the view of an indirect effect of corticosterone on this immune parameter. Accordingly, in the present study we found a positive relationship between body mass and T-cell-mediated immune response, where the response was stronger in females with a greater body mass. By contrast, the immunoglobulin level appears to be more sensitive to high corticosterone levels than to body mass loss. Nevertheless, corticosterone levels and body mass were negatively related in the current study, so that high levels of corticosterone were associated with low body mass. Our results agree with previous findings from breeding black-legged kittiwakes *Rissa tridactyla*, where high corticosterone levels were associated with a marked decline in body condition (Kitaysky et al., 1999). In this context, it would be interesting for future studies (1) to experimentally extend the fasting duration of eiders until they reach phase III of fasting or (2) to find free-ranging birds that spontaneously shift from lipid to protein utilization, so that the effects of both elevated corticosterone levels and decreased body mass on the acquired immunity can be examined.

The present study showed that an increasing body mass loss, caused either by corticosterone administration, or by an experimental extension of fasting duration, negatively affected the birds' immunoglobulin level and their T-cell-mediated immune response, respectively. This result lends support to the resource-limitation hypothesis, which predicts that investment in costly behaviours, such as reproduction, reduces the amount of resources available to other systems, such as the immune system (Sheldon and Verhulst, 1996; Råberg et al., 1998). However, evidence for an energetically costly immune response is still equivocal (Råberg et al., 1998; Eraud et al., 2005; Verhulst et al., 2005). Whatever the ultimate factors explaining immunosuppression during reproduction, corticosterone was proposed to regulate immunosuppression in incubating birds (Deerenberg et al., 1997; Råberg et al., 1998; Saino et al., 2003). However, during fasting corticosterone should be maintained at low levels to avoid metabolic disorders, such as protein catabolism (Cherel et al., 1988). In the present study we did not find marked variations in corticosterone levels during the incubation period of control and 'late fasting' females. However, 'late fasters' had a lower T-cell-mediated immune responses than corticosterone implanted or control females. This would support the view that fasting duration and/or body composition (see below) might be relevant parameters for immunosuppression.

Other aspects of immunosuppression, namely humoral immunity, could be mediated by factors related to fuel utilization or body mass loss. Such a relationship between energy storage/mobilization and immunocompetence might plausibly be mediated through nutritional and/or endocrine factors (Apanius, 1998). Exogenous administration of corticosterone is likely to increase proteolysis. Consequently,

lean body mass will be decreased, while the energy stored as lipids within the body will be spared. Hence, for the same final body mass, adiposity of 'late fasters' should be lower than for corticosterone implanted females, as reported to be the case in dark-eyed juncos *Junco hyemalis* treated with corticosterone (Gray et al., 1990). Currently, adipose tissue is perceived as an active participant in the regulation of essential and prominent body processes such as immune homeostasis (Matarese and La Cava, 2004). This raises the question of how body reserves might control the immune system (Demas and Sakaria, 2005). Some adipose humoral signals, such as leptin, are generated in proportion to fat stores and act on feedback control systems to influence numerous biological processes (Löhmus and Sundström, 2004; Matarese et al., 2005). In fact, leptin is secreted primarily by adipose tissue and has been shown to enhance a variety of immunological parameters in mammals (Lord et al., 1998; Faggioni et al., 2001; Demas et al., 2003) and birds (Löhmus et al., 2004). Consequently, leptin levels might be lower in 'late fasters' when compared with corticosterone implanted eiders. To gain further insight into the role that leptin plays for the immune system of fasting birds, plasma measurements of leptin and manipulation of its circulating concentrations would be useful.

In conclusion, exogenous corticosterone decreased only one component of the acquired immune system in incubating female eider ducks. While the treatment significantly decreased their immunoglobulin level, their T-cell-mediated immune response was not affected. Implanted females lost significantly more weight than control birds. Females, whose fasting duration was experimentally extended, increasing body mass loss, displayed lower T-cell-mediated immune responses than implanted females, while their corticosterone levels remained at baseline values. Consequently, the immunosuppressive effect of corticosterone appears to be mediated by its effect on body reserves, which have been shown to play an important role in the regulation of the immune system. For example, leptin, which conveys information on energy availability, could be involved in the observed immunosuppression. Further experiments are required to determine the relationship between body condition and immune function in incubating female eiders. Our results raise the question of the physiological mechanisms that can explain the effects of corticosterone on the immune response. Furthermore, could it be that depending on its concentration, this hormone is able to trigger different responses, as has already been reported in the context of foraging behaviour (Wingfield et al., 1998).

Financial support for this study was provided by the Institut Polaire Français Paul Emile Victor (IPEV) and through a fellowship from the French MENRT to S. Bourgeon. The study was approved by the Norwegian Animal Research Authority and the Ethic Committee of the IPEV. Permission to work on common eiders was also granted by the Governor of Svalbard.

References

- Apanius, V. (1998). Stress and immune defense. *Adv. Stud. Behav.* **27**, 133-153.
- Bourgeon, S., Criscuolo, F., Le Maho, Y. and Raclot, T. (2006). Phytohemagglutinin response and immunoglobulin level decrease during incubation fasting in female Common Eiders. *Physiol. Biochem. Zool.* **79** (4), 793-800.
- Cherel, Y., Robin, J.-P., Walch, O., Karmann, H., Netchitailo, P. and Le Maho, Y. (1988). Fasting in king penguins I. Hormonal and metabolic changes during breeding. *Am. J. Physiol.* **254**, R170-R177.
- Criscuolo, F., Chatsel, O., Berile, F., Gabrielsen, G. W., Le Maho, Y. and Raclot, T. (2005). Corticosterone alone does not trigger a short term behavioural shift in incubating female common eiders *Somateria mollissima*, but does modify long term reproductive success. *J. Avian Biol.* **36**, 306-312.
- Deerenberg, C., Arpanius, V., Daan, S. and Bos, N. (1997). Reproductive effort decreases antibody responsiveness. *Proc. R. Soc. Lond. B* **264**, 1021-1029.
- Demas, G. E. and Sakaria, S. (2005). Leptin regulates energetic tradeoffs between body fat and humoral immunity. *Proc. R. Soc. Lond. B* **272**, 1845-1850.
- Demas, G. E., Drazen, D. L. and Nelson, R. J. (2003). Reductions in total body fat decrease humoral immunity. *Proc. R. Soc. Lond. B* **270**, 905-911.
- Dhabhar, F. S. and McEwen, B. S. (1997). Acute stress enhances while chronic stress suppresses cell-mediated immunity in vivo: a potential role for leukocyte trafficking. *Brain Behav. Immun.* **11**, 286-306.
- Eraud, C., Duriez, O., Chastel, O. and Faivre, B. (2005). The energetic cost of humoral immunity in the collared dove, *Streptopelia decaocto*: is the magnitude sufficient to force energy-based trade-offs? *Funct. Ecol.* **19**, 110-118.
- Faggioni, R., Feingold, K. R. and Grunfeld, C. (2001). Leptin regulation of the immune response and the immuno-deficiency of malnutrition. *FASEB J.* **15**, 2565-2571.
- Gray, J. M., Yarian, Y. and Ramenofsky, M. (1990). Corticosterone, foraging behavior, and metabolism in dark-eyed juncos, *Junco hyemalis*. *Gen. Comp. Endocrinol.* **79**, 375-384.
- Gustafsson, L., Nordling, D., Andersson, M. S., Sheldon, B. C. and Qvarnström, A. (1994). Infectious diseases, reproductive effort and the cost of reproduction in birds. *Philos. Trans. R. Soc. B* **346**, 323-331.
- Hanssen, S. A., Hasselquist, D., Folstad, I. and Erikstad, K. J. (2004). Costs of immunity: immune responsiveness reduces survival in a vertebrate. *Proc. R. Soc. Lond. B* **271**, 925-930.
- Hollmén, T., Franson, J. C., Hario, M., Sankari, S., Kilpi, M. and Lindström, K. (2001). Use of serum biochemistry to evaluate nutritional status and health of incubating Common Eiders (*Somateria mollissima*) in Finland. *Physiol. Biochem. Zool.* **74**, 333-342.
- Kitaysky, A. S., Wingfield, J. C. and Piatt, J. F. (1999). Dynamics of food availability, body condition and physiological response in breeding black-legged kittiwakes. *Funct. Ecol.* **13**, 577-585.
- Korschgen, C. E. (1977). Breeding stress of female eiders in Maine. *J. Wildlife Manage.* **41**, 360-373.
- Le Maho, Y. (1983). Metabolic adaptations to long-term fasting in Antarctic penguins and domestic geese. *J. Therm. Biol.* **8**, 91-96.
- Le Maho, Y., Vu Van Kha, H., Koubi, H., Dewasmes, G., Girard, J., Ferre, P. and Cagnard, M. (1981). Body composition, energy expenditure, and plasma metabolites in long-term fasting geese. *Am. J. Physiol.* **241**, E342-E354.
- Lindgård, K., Stokkan, K. A., Le Maho, Y. and Groscolas, R. (1992). Protein utilization during starvation in fat and lean Svalbard ptarmigan (*Lagopus mutus hyperboreus*). *J. Comp. Physiol. B* **162**, 607-613.
- Löhmus, M. and Sundström, L. F. (2004). Leptin and social environment influence the risk-taking and feeding behaviour of Asian blue quail. *Anim. Behav.* **68**, 607-612.
- Löhmus, M., Olin, M., Sundström, L. F., Troedsson, M. H. T., Molitor, T. W. and El Halawani, M. (2004). Leptin increases T-cell immune response in birds. *Gen. Comp. Endocrinol.* **139**, 245-250.
- Lord, G. M., Matarese, G., Howard, J. K., Baker, R. J., Bloom, S. R. and Lechler, R. I. (1998). Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* **394**, 897-901.
- Martin, II, L. B., Gilliam, J., Han, P., Lee, K. and Wilewski, M. (2005). Corticosterone suppresses cutaneous immune function in temperate but not tropical House Sparrows, *Passer domesticus*. *Gen. Comp. Endocrinol.* **140**, 126-135.
- Martinez, J., Tomás, G., Merino, S., Arriero, E. and Moreno, J. (2003).

- Detection of serum immunoglobulins in wild birds by direct ELISA: a methodological study to validate the technique in different species using antichickens antibodies. *Funct. Ecol.* **17**, 700-706.
- Matarese, G. and La Cava, A.** (2004). The intricate interface between immune system and metabolism. *Trends Immunol.* **25**, 193-200.
- Matarese, G., Moschos, S. and Mantzoros, C. S.** (2005). Leptin in immunology. *J. Immunol.* **173**, 3137-3142.
- Merino, S., Martinez, J., Møller, A. P., Sanabria, L., De Lope, F., Perez, J. and Rodriguez-Caabeiro, F.** (1999). Phytohaemagglutinin injection assay and physiological stress in nestling house martins. *Anim. Behav.* **58**, 219-222.
- Moreno, J., Merino, S., Potti, J., de Leon, A. and Rodriguez, R.** (1999). Maternal energy expenditure does not change with flight costs or food availability in the pied flycatcher (*Ficedula hypoleuca*): costs and benefits for nestlings. *Behav. Ecol. Sociobiol.* **46**, 244-251.
- Norris, K. and Evans, M. R.** (2000). Ecological immunology: life history trade-offs and immune defense in birds. *Behav. Ecol.* **11**, 19-26.
- Parham, P.** (1995). The duck's dilemma. *Nature* **374**, 16-17.
- Råberg, L., Grahn, M., Hasselquist, D. and Svensson, E.** (1998). On the adaptive significance of stress-induced immunosuppression. *Proc. Roy. Soc. Lond. B* **265**, 1637-1641.
- Robin, J.-P., Boucontet, L., Chillet, P. and Groscolas, R.** (1998). Behavioral changes in fasting emperor penguins: evidence for a 'refeeding signal' linked to a metabolic shift. *Am. J. Physiol.* **274**, R746-R753.
- Roitt, I. M., Brostoff, J. and Male, D. K.** (1998). *Immunology*. London: Mosby.
- Saino, N., Incagli, M., Martinelli, R. and Møller, A. P.** (2002). Immune response of male barn swallows in relation to parental effort, corticosterone plasma levels, and sexual ornamentation. *Behav. Ecol.* **13**, 169-174.
- Saino, N., Suffritti, C., Martinelli, R., Rubolini, D. and Møller, A. P.** (2003). Immune response covaries with corticosterone plasma levels under experimentally stressful conditions in nestling barn swallows (*Hirundo rustica*). *Behav. Ecol.* **14**, 318-325.
- Sheldon, B. C. and Verhulst, S.** (1996). Ecological immunology: costly parasite defenses and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* **11**, 317-321.
- Stearns, S. C.** (1992). *The Evolution of Life Histories*. Oxford: Oxford University Press.
- Verhulst, S., Riedstra, B. and Wiersma, P.** (2005). Brood size and immunity costs in zebra finches *Taeniopygia guttata*. *J. Avian Biol.* **36**, 22-30.
- Williams, G. C.** (1966). Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am. Nat.* **100**, 687-690.
- Wingfield, J. C., Maney, D. L., Breuner, C. W., Jacobs, J. D., Lynn, S., Ramenofsky, M., and Richardson, R. D.** (1998). Ecological bases of hormone-behavior interactions: the 'emergency life history stage'. *Am. Zool.* **38**, 191-206.
- Yoccoz, N. G., Erikstad, K. E., Bustnes, J. O., Hanssen, S. A. and Tveraa, T.** (2002). Costs of reproduction in common eiders (*Somateria mollissima*): an assessment of relationships between reproduction based on observational and experimental studies. *J. Appl. Stat.* **29**, 57-64.