

RESEARCH ARTICLE

Visual system development of the spotted unicornfish, *Naso brevirostris* (Acanthuridae)

Valerio Tettamanti^{1,2}, Fanny de Busserolles¹, David Lecchini^{3,4}, N. Justin Marshall¹ and Fabio Cortesi^{1,*}

ABSTRACT

Ontogenetic changes of the visual system are often correlated with shifts in habitat and feeding behaviour of animals. Coral reef fishes begin their lives in the pelagic zone and then migrate to the reef. This habitat transition frequently involves a change in diet and light environment as well as major morphological modifications. The spotted unicornfish, *Naso brevirostris*, is known to shift diet from zooplankton to algae and back to mainly zooplankton when transitioning from larval to juvenile and then to adult stages. Concurrently, *N. brevirostris* also moves from an open pelagic to a coral-associated habitat before migrating up in the water column when reaching adulthood. Using retinal mapping techniques, we discovered that the distribution and density of ganglion and photoreceptor cells in *N. brevirostris* changes primarily during the transition from the larval to the juvenile stage, with only minor modifications thereafter. Similarly, visual gene (opsin) expression based on RNA sequencing, although qualitatively similar between stages (all fish mainly expressed the same three cone opsins; *SWS2B*, *RH2B*, *RH2A*), also showed the biggest quantitative difference when transitioning from larvae to juveniles. The juvenile stage in particular seems mismatched with its reef-associated ecology, which may be due to this stage only lasting a fraction of the lifespan of these fish. Hence, the visual ontogeny found in *N. brevirostris* is very different from the progressive changes found in other reef fishes, calling for a thorough analysis of visual system development of the reef fish community.

KEY WORDS: Visual ecology, Teleosts, Fish, Ontogeny, Opsin, Retina, Visual acuity

INTRODUCTION

Many animals use vision to perform important behavioural tasks such as feeding, mating, avoiding predators and finding a suitable home (Cronin et al., 2014). At the core of the vertebrate visual system is the retina, an extrusion of the brain which is subdivided into various functional layers, two of which are at the centre of this study, the photoreceptor layer and the ganglion cell layer.

The photoreceptor layer is the first stage of visual processing and is composed of morphologically diverse cone and rod photoreceptor cells which absorb light, transform it into an electrical signal, and send the information downstream to various neural cells via the phototransduction cascade. Cones mediate vision in bright-light

conditions and colour vision, while rods mediate vision in dim-light conditions (Walls, 1942). Cones can further be classified into different types depending on their morphology and/or the type of photopigment (an opsin protein covalently bound to a light-absorbing chromophore) they possess (Hunt et al., 2014). Morphologically, cones can be classified as single, double, triple or quadruple, although only the first two configurations are common and are often arranged in regular and specific patterns or mosaics (Bowmaker and Kunz, 1987; Peichl et al., 2004). Molecularly, cones are also classified into four types based on the opsin genes they express, which encode different protein classes that are sensitive to different parts of the visible light spectrum: a short-wavelength protein class 1 opsin (SWS1) maximally sensitive to UV–violet wavelengths (355–450 nm peak spectral sensitivity, λ_{\max}), a second short-wavelength class opsin (SWS2) maximally sensitive to the violet–blue part of the spectrum (410–490 nm λ_{\max}), a middle-wavelength class 2 opsin (RH2) maximally sensitive to blue–green wavelengths (435–535 nm λ_{\max}) and a long-wavelength class opsin (LWS) maximally sensitive to the green–red part of the light spectrum (490–625 nm λ_{\max}) (Bowmaker, 2008; de Busserolles et al., 2017; Torres-Dowdall et al., 2017). In percomorph fishes, SWS proteins are found in single cones, whereas RH2 and LWS opsins occur in double cones (Hunt et al., 2014). Most vertebrates possess a single type of rod photoreceptor expressing the rod opsin protein (RH1; 460–530 nm λ_{\max} ; Bowmaker, 2008; but see Musilova et al., 2019, for deep-sea fishes with multiple RH1s that extend their sensitivities to ~440 nm λ_{\max}).

The ganglion cell layer is the last stage of visual processing in the retina and is composed of ganglion cells that possess axons that reach to the inner surface of the retina and converge into the optic nerve to send the information into the central nervous system (Walls, 1942). Therefore, the arrangement and spacing between ganglion cells is one of the determining factors of visual acuity (or resolution) (Fernald, 1988).

In order to perform at its optimum, the visual system of a particular species is adapted to the type of habitat they live in and to the prevailing surrounding light conditions (Lythgoe, 1979). In general, vertebrates range from cone monochromats with a single spectral class of cone photoreceptor (e.g. sharks and many rays), over dichromats and trichromats (e.g. most mammals and many marine fishes), to tetrachromats (e.g. most birds and many freshwater fishes; Bowmaker, 2008). Cone photoreceptors and their respective opsin repertoires are particularly diverse in teleost fishes (e.g. Lin et al., 2017; Musilova et al., 2019). This is thought to primarily be due to the different light environments that fish inhabit (Cronin et al., 2014; Lythgoe, 1979), but in some instances may also be driven by sexual selection (Endler, 1990) and/or the feeding habits of species. For example, UV photoreception increases feeding efficiency in some fishes eating UV-absorbing or scattering zooplankton (Novales-Flamarique, 2016; Loew et al., 1993; Browman et al., 1994), while herbivorous fishes may profit from visual systems tuned to longer wavelengths as a result of the

¹Queensland Brain Institute, The University of Queensland, 4072 Brisbane, QLD, Australia. ²Swiss Federal Institute of Technology Zurich, 8092 Zurich, Switzerland. ³PSL Research University: EPHE-UPVD-CNRS, USR3278 CRIOBE, BP 1013, 98729 Papetoai, Moorea, French Polynesia. ⁴Laboratoire d'Excellence 'CORAIL', 75014 Paris, France.

*Author for correspondence (fabio.cortesi@uqconnect.edu.au)

© V.T., 0000-0002-9078-734X; F.d.B., 0000-0002-4602-9840; D.L., 0000-0002-6347-1112; N.J.M., 0000-0001-9006-6713; F.C., 0000-0002-7518-6159

List of symbols and abbreviations

| | |
|---------------------------|---|
| bp | base pairs |
| CE | Shaeffer's coefficient of error |
| DC | double cone |
| LWS | long-wavelength sensitive |
| LWS | long-wavelength-sensitive opsin gene |
| MSP | microspectrophotometry |
| PDG | peak density of ganglion cells |
| <i>RH1</i> | rhodopsin 1 opsin gene |
| <i>RH2A</i> , <i>RH2B</i> | rhodopsin-like 2 opsin genes |
| SC | single cone |
| SL | standard length |
| SNP | single nucleotide polymorphism |
| SRP | spatial resolving power |
| SWS | short-wavelength sensitive |
| <i>SWS1</i> | short-wavelength-sensitive 1 opsin gene |
| <i>SWS2</i> | short-wavelength-sensitive 2 opsin gene |
| λ_{\max} | peak spectral sensitivity |

red-reflecting properties of chlorophyll (Cortesi et al., 2018 preprint; Marshall et al., 2003; Stieb et al., 2017).

Moreover, the density of photoreceptors and ganglion cells can also vary not only between species but also within an individual's retina over its lifespan (Shand et al., 1999, 2000). The study of their distribution using the wholemount technique (Coimbra et al., 2006; Stone and Johnston, 1981; Ullmann et al., 2012) provides useful information on the visual ecology of a species, which usually reflects its habitat and behavioural ecology (Bozzano and Collin, 2000; Collin and Pettigrew, 1988a,b, 1989; Hughes, 1977). Two main specializations can be found in vertebrates: area and streaks (Collin and Pettigrew, 1988a,b). Both specializations have higher densities of cells compared with the rest of the retina, resulting in regions of acute vision in the corresponding field of view. An area is a concentric increase in cell density in a particular region of the retina; in some vertebrates, it is termed a fovea because of other structural adaptations (Walls, 1942). In teleost fishes, areas are often located temporally (i.e. area temporalis) and found in species that live in enclosed environments such as caves or coral structures, and/or coral overhangs (Collin and Pettigrew, 1989; Collin and Shand, 2003). The temporal area receives the visual information from the frontal field of view, corresponding to the natural swimming direction of fish. Nevertheless, multiple area centralis can also be found in a single retina (Collin and Pettigrew, 1989). For example, triggerfishes (Balistidae) possess an area in both the nasal and temporal part of the retina, which correlates with two main visual tasks: feeding (temporal) and predator avoidance (nasal) (Collin and Pettigrew, 1988b; Ito and Murakami, 1984). A horizontal streak is defined by an increase in cell density along the meridian. Most horizontal streaks are found in the central meridian, but sometimes they can also be located more ventrally or dorsally (Collin and Shand, 2003). The streak maintains a high spatial resolving power (SRP) throughout the horizontal section of the retina and is thought to be used to scan the horizon. It leads to an elongated sampling of the visual environment without continuous eye movements (Collin and Shand, 2003). Teleost fishes possessing a horizontal streak are commonly found in open water environments such as sandy bottoms or pelagic open ocean environments (Collin and Pettigrew, 1988b).

Variability in retinal structure and opsin gene repertoire does not only exist between species; both visual features may also change throughout the life of an individual. This is especially true for species that undergo substantial habitat changes during ontogeny

such as coral reef fishes. The life of most coral reef fishes starts in the shallower zone of the open ocean as larvae (Helfman et al., 2009; Job and Bellwood, 2000), where resources may be high and the risk of predation is low (Fortier and Harris, 1989). At this stage, pelagic fish larvae feed typically on zooplankton (Boehlert, 1996) and rely on vision primarily for fundamental tasks such as predator avoidance and feeding (Leis and Carson-Ewart, 1999). After their oceanic phase, reef fish larvae typically find a suitable coral reef patch to settle on and again vision is one of the main senses used (Lecchini et al., 2005a,b). During this settlement phase, reef fish larvae undergo metamorphosis and reach the juvenile stage in which they usually already possess all basic morphological features of the adult form (Holzer et al., 2017). Following settlement on the reef, juvenile fishes are challenged with visual cues that are much more complex than in the open ocean, varying in both chromaticity and luminance. Hence, at this stage (or slightly before; Cortesi et al., 2016) the visual system of coral reef fishes is expected to undergo changes in both morphology and physiology (Helfman et al., 2009).

To date, changes in arrangement (i.e. mosaic) and distribution of the photoreceptor cells throughout ontogeny have been documented only in a few coral reef fishes (Shand, 1997). These changes are thought to enhance survival by increasing feeding success and facilitate predator avoidance in reef stages (juveniles and adults; Shand, 1997). Along with changes in morphology, ontogenetic changes in opsin gene expression have also been reported for a handful of species (Cortesi et al., 2015b, 2016). For example, in the dottyback *Pseudochromis fuscus*, the number and type of opsin genes that are expressed differs between larval, juvenile and adult stages. Along with the change in opsin gene expression, the visual system may also transform to a more complex colour-processing capability, such as dichromacy to trichromacy or even up to tetrachromacy. This increase in chromaticity ultimately requires behavioural testing to confirm and is likely to reflect major habitat transitions throughout development equipping fishes such as dottybacks with a more complex visual system as they grow and mature (Cortesi et al., 2015b, 2016). In comparison, while some freshwater cichlid species show a similarly dynamic change in opsin gene expression through ontogeny, other species do not change gene expression much (neoteny) or directly develop from the larval to the adult gene-expression pattern (Carleton et al., 2008; Härer et al., 2017). We currently do not know whether a progressive developmental change of the visual system, as found in the dottyback (Cortesi et al., 2016), for example, is a common feature shared among coral reef fishes, or whether some species also show different developmental modes, similar to what is found in cichlid fishes.

In this study, we investigated ontogenetic changes in retinal topography and opsin gene expression in three life stages (larval, juvenile, adult) of the spotted unicornfish, *Naso brevirostris* (Cuvier 1829), from the surgeonfish family (Acanthuridae) (Fig. 1). *Naso brevirostris* is known to shift both diet and habitat during ontogeny (Choat et al., 2002; Randall et al., 1997). Pelagic larvae feed on zooplankton before settling on the reef, where they mainly feed on algae as juveniles. As adults, *N. brevirostris* migrate to the reef slope, returning to a mostly zooplanktivorous diet (Choat et al., 2002, 2004; Randall et al., 1997). We therefore hypothesized that the visual system of *N. brevirostris* would show a 'classic' developmental mode, linked to either changes in habitat or diet, or both, and with a progression from larval, to juvenile and finally adult traits. Moreover, *N. brevirostris* develops an elongated rostral snout during maturation, and this prominent morphological change may also affect its visual requirements as it might obstruct the visual field of the fish.




Fig. 1. *Naso brevirostris* developmental stages. The spotted unicornfish, *N. brevirostris*, shows pronounced ontogenetic changes in habitat, diet and morphology. (A) A 'transparent' zooplanktivorous larva at the settlement stage (i.e. when returning from the pelagic zone to the reef). (B) An algivorous juvenile that lives in close proximity to the reef. (C) A zooplanktivorous adult that lives in the water column above the reef. Note the growth of the prominent snout throughout development.

MATERIALS AND METHODS

Study species and collection

Individuals of *N. brevirostris* were collected on the Northern Great Barrier Reef, Australia, under the Great Barrier Reef Marine Park Association (GBRMPA) permits G17/38160.1 and G16/38497.1, Queensland Fisheries permit no. 180731, or in French Polynesia. Adults [$n=5$; standard length (SL): range 27.7–33 cm, mean \pm s.d. 30.4 \pm 1.9 cm] were collected with a spear gun from No Name Reef (14°65'S, 145°65'E) on the outer Great Barrier Reef, Australia, in February 2018. Juveniles ($n=8$; SL, 6.5–19.5 cm, 14.5 \pm 3.7 cm) were collected using barrier nets, spear guns or clove oil and hand nets from reefs surrounding Lizard Island (14°40'S, 145°27'E) on the Great Barrier Reef between February 2016 and February 2018, and one juvenile fish was acquired through the aquarium supplier Cairns Marine (<http://www.cairnsmarine.com/>). Larval fish ($n=5$; SL: 3.1–3.3 cm, 3.1 \pm 0.1 cm) were captured using a crest net on Tetiaroa Island, French Polynesia (16°99'S, 149°58'W) in March 2018 (Besson et al., 2017; Lecchini et al., 2004). All animals were anaesthetized using an overdose of clove oil (10% clove oil, 40% ethanol, 50% seawater) and killed by decapitation under an animal ethics protocol of The Queensland Brain Institute (QBI/236/13/ARC/US AIRFORCE and QBI/304/16).

Each individual was photographed with a ruler in the frame, to enable us to extract the SL later on using Fiji v.1.0 (Schindelin et al., 2012). The eyes were enucleated and the cornea and lens removed using micro-dissection scissors. A small dorsal cut was made to keep track of the eye's orientation. The samples collected for retinal mapping were fixed in 4% paraformaldehyde in 0.1 mol l⁻¹ phosphate-buffered saline (PBS; pH 7.4) and stored at 4°C, and the eyes used for RNA sequencing were kept in RNAlater (Sigma) and stored at -20°C. For each eye, the lens diameter was measured after dissection and fixation.

Preparation of retinal wholemounts

Retinal wholemounts were prepared according to standard protocols (Coimbra et al., 2006, 2012; Stone and Johnston, 1981). Briefly, each eye cup was cut radially multiple times, to enable us to flatten it on a microscopy glass slide without damaging the tissue. The retina was oriented using the falciform process that extends ventrally. The sclera and choroid were gently removed and the retinas were bleached overnight in the dark at ambient temperature in a 3% hydrogen peroxide solution (in PBS). Large-sized adult retinas, which have a more developed retinal pigment epithelium, were bleached in the same solution but with a few drops of potassium hydroxide (Ullmann et al., 2012). While potassium hydroxide accelerates the bleaching process by

increasing the pH of the solution, this type of bleaching is quite aggressive for the tissue. Therefore, these retinas were only bleached for 2–3 h in the dark.

For ganglion cell analyses, retinas were mounted with the ganglion cell layer facing up on a gelatinized slide and left to dry overnight at room temperature in formalin vapours (Coimbra et al., 2006, 2012). Wholemounts were then stained in 0.1% Cresyl Violet (Nissl staining) following the protocol of Coimbra et al. (2006) and then mounted with Entellan New (Merck). Shrinkage of the retina using this technique is usually deemed negligible and, if present, restricted to the borders of the retina (Coimbra et al., 2006). In this study, however, the retinas did not show equal shrinkage because of major differences in retina size between developmental stages. As such, the smaller retinas (larval stage) were more affected by shrinkage, because of their smaller surface (i.e. higher proportion of retinal borders), than those from the other stages (adult and juvenile stages). Shrinkage in these retinas was easily identified under the microscope and was taken into consideration in the data interpretation.

For photoreceptor analyses, retinas were wholemounted in 80% glycerol in PBS, on non-gelatinized slides with the inner (vitreal) surface facing downwards. Contrary to ganglion cell mounting, photoreceptor mounting shows negligible shrinkage as it takes place in an aqueous medium (Peichl et al., 2004).

Stereological analyses and construction of topographic maps

The topographic distribution and the total number of ganglion cells, single cones, double cones and total cones in the three life stages of *N. brevirostris* were assessed using the optical fractionator technique (West et al., 1991), modified for wholemount retina use by Coimbra et al. (2009, 2012). A computer running Stereo Investigator software (v2017.01.1, 64-bit, Microbrightfield) coupled to a compound microscope (Zeiss Imager.Z2) equipped with a motorized stage (MAC 6000 System, Microbrightfield) and a digital colour camera (Microbrightfield) was used for the analysis. The contour of each retina wholemount was digitized using a $\times 5$ objective (numerical aperture 0.16) and cells were counted randomly and systematically, using a $\times 63$ oil immersion objective (numerical aperture 1.40) and the parameters summarized in Table S1. The total number of cells for each sample was then estimated by multiplying the sum of the neurons (ganglion cells or photoreceptors) counted by the area of the sampling fraction (Coimbra et al., 2009; West et al., 1991).

The counting frame and grid size were chosen carefully in order to achieve an acceptable Shaeffer's coefficient of error (CE), while maintaining the highest level of sampling. The CE measures the

accuracy of the estimation of the total cell number and is deemed acceptable below 0.1 (Glaser and Wilson, 1998; Slomianka and West, 2005). The counting frame was adjusted between life stages to reach an average count of around 40 and 80 cells per sampling site for ganglion cells and photoreceptors, respectively, but was kept identical for individuals of the same life stage (Table S1). As fish of similar life stages can have a wide variation of SL, the grid size was adjusted for all individuals to allow sampling of around 200 sites ($\pm 10\%$) (de Busserolles et al., 2014a,b).

Three cell types can be found in the ganglion cell layer: ganglion cells, displaced amacrine cells and glial cells. These can usually be distinguished based on cytological criteria (Collin, 1988; Collin and Pettigrew, 1988c; Hughes, 1975), with ganglion cells having an irregular shape, an extensive nucleus and a larger size compared with the smaller, rounder amacrine cells, which have a darker stained appearance, and glial cells, which have an elongated shape relative to the other two cell types (Fig. 2A). However, as amacrine cells were often difficult to distinguish from ganglion cells in *N. brevirostris*, especially in high-density areas, amacrine cells were included in all counts and only glial cells were excluded. The inclusion of amacrine cells in the analysis should not interfere with the overall topography, as the distribution of amacrine cells has been shown to match the ganglion cell distribution in other animals (Bailes et al., 2006; Coimbra et al., 2006; Collin, 2008; Collin and Pettigrew, 1988c; Wong and Hughes, 1987), and the density of displaced amacrine cells in *N. brevirostris* was relatively low. However, the inclusion of amacrine cells in the ganglion cell counts will contribute to a slight overestimation of ganglion cell density and ultimately to a slight overestimation of SRP. For ganglion cell

analysis, sub-sampling was performed in the regions of highest cell density to allow a more accurate estimation of the peak ganglion cell density. The same counting frame parameters as for the whole retina were used for sub-sampling, but the grid size was reduced by half.

Photoreceptor cells, in contrast, could be distinguished unambiguously as single and double cones (Fig. 2B). The two cone types were counted separately and simultaneously using two different markers to acquire data for single cones alone, double cones alone and total cones (single and double cones).

Topographic maps were created using the statistical program R v.3.4.1 (R Foundation for Statistical Computing, 2012) with the results exported from Stereo Investigator and the R script provided by Garza-Gisholt et al. (2014). As for previous retinal topography studies on teleost fishes (Dalton et al., 2016; de Busserolles et al., 2014a,b), the Gaussian Kernel Smoother from the Spatstat package (Baddeley and Turner, 2005) was chosen and the sigma value was adjusted to the distance between points (i.e. grid size) for each map (Fig. 3).

SRP estimation

The upper limit of the SRP in cycles per degree was estimated for each individual using the ganglion cell peak density as described by Collin and Pettigrew, 1989. The following formula was used:

$$\alpha = \arctan(1/f), \quad (1)$$

where α is the angle subtending 1 mm on the retina and calculated assuming that f , the focal length of the fish, is 2.55 times the lens radius, the standard for teleost fishes according to the Matthiessen ratio (Matthiessen, 1882). Then, knowing α , the peak density of ganglion cells (PDG in cells mm^{-1}) and the fact that two ganglion cells are needed to distinguish a visual element from its neighbouring element, the SRP in cycles per degree can be calculated as follows:

$$\text{SRP} = (\text{PDG}/\alpha)/2. \quad (2)$$

Transcriptome sequencing, quality filtering and *de novo* assembly

The retinas from different life stages of *N. brevirostris* (adult, $n=3$; juvenile, $n=6$; larvae, $n=3$) were dissected out of the eye cup, total RNA was extracted, and their retinal transcriptomes were sequenced according to Musilova et al. (2019). Briefly, total RNA of larval and smaller juvenile retinas was extracted using the RNeasy Mini Kit (Qiagen), and for larger juvenile and adult retinas the RNeasy Midi Kit (Qiagen), according to the manufacturer's instructions, which included a DNase treatment. Total RNA concentration and quality were determined using a Eukaryotic Total RNA NanoChip on an Agilent 2100 BioAnalyzer (Agilent Technologies). Juvenile transcriptomes were sequenced in-house at the Queensland Brain Institute's sequencing facility. For these samples, strand-specific libraries were barcoded and pooled at equimolar ratios and sequenced at PE125 on a HiSeq 2000 using Illumina's SBS chemistry version 4. Library preparation (strand-specific, 300 bp insert) and transcriptome sequencing (RNAseq HiSeq PE150) for larval and adult individuals was outsourced to Novogene (<https://en.novogene.com/>).

Retinal transcriptomes were filtered, and *de novo* assembled following the protocol described in de Busserolles et al. (2017). Briefly, raw-reads of transcriptomes were uploaded to the Genomics Virtual Laboratory (GVL 4.0.0) (Afgan et al., 2015) on the Galaxy Australia server (<https://usegalaxy.org.au/>), filtered by quality using Trimmomatic (Galaxy version 0.36.4) (Bolger et al., 2014) and then




Fig. 2. Light micrographs of various retinal layers as found in an adult *N. brevirostris*. (A) Micrographs of the Nissl-stained ganglion cell layer taken in a low-density area (nasal part) and a peak-density area (central part) of the retina. Ganglion cells (gc) could clearly be distinguished from glial cells (g) by their round shape and difference in size. Distinguishing amacrine cells (ac) from ganglion cells, however, was more difficult. (B) Micrographs of the photoreceptor layer taken in a low-density area (nasal part) and a peak-density area (temporal part). Photoreceptors formed a square mosaic with a central single cone surrounded by four double cones. Scale bars: 50 μm .




Fig. 3. Topographic heat maps of ganglion and photoreceptor cell distribution. (A) Topographic distribution of retinal ganglion cells revealed a pronounced horizontal streak with a peak cell density in the central area in adult and juvenile individuals. In larvae, the streak was less pronounced and peak cell densities were found in the nasal and temporal parts instead (see Fig. S1 for maps of additional individuals). (B) Topographic distribution of total photoreceptors (double and single cones) revealed an increase in specialization with life stage from a weak horizontal streak with two large areas of high cell density in larvae to the formation of a pronounced horizontal streak and a weak dorsal vertical streak in juveniles. A more pronounced dorsal vertical streak was found in adults [see Fig. S2 and Dryad (<https://doi.org/10.5061/dryad.k6djh9w37>) for single and double cone maps as well as maps of additional individuals]. Black lines represent isodensity contours, and values are expressed in densities of $\times 10^3$ cells mm^{-2} . V, ventral; T, temporal. Scale bars: 1 mm.

de novo assembled using Trinity (Galaxy version 2.4.0.0) (Haas et al., 2013).

Opsin gene mining and phylogenetic reconstruction

Following the protocol in de Busserolles et al. (2017), the *N. brevirostris* transcriptomes were mined for their visual opsin genes. Briefly, using the opsin coding sequences from the dusky dotyback, *Pseudochromis fuscus* (Cortesi et al., 2016), we searched for the *N. brevirostris* opsin genes by mapping the *de novo* assembled transcripts to the *P. fuscus* reference genes using Geneious v.11.1.3 (www.geneious.com). *Pseudochromis fuscus* was chosen because it is relatively closely related to *N. brevirostris*, and because it possesses orthologues from all of the ancestral vertebrate opsin genes (Cortesi et al., 2016).

Assemblies based on short-read libraries tend to overlook genes expressed at low levels and similar gene copies, and/or short-reads may be misassembled (chimeric sequences); for that reason, a second approach was used to confirm the visual opsin genes of *N. brevirostris*. A manual extraction of the gene copies was performed by mapping raw-reads against the *P. fuscus* references and then moving from single nucleotide polymorphism (SNP) to SNP along the gene, taking advantage of paired-end information to bridge the gaps between SNPs. The extracted reads were then *de novo* assembled and their consensus was used as a template against which unassembled reads were re-mapped to elongate the region of interest; this approach eventually led to a reconstruction of

the whole coding region (for details on this approach, see de Busserolles et al., 2017; Musilova et al., 2019).

Opsin gene identity was then confirmed using BLAST (<http://blast.ncbi.nlm.nih.gov/>) and by phylogenetic reconstruction to a reference dataset obtained from GenBank (www.ncbi.nlm.nih.gov/genbank/) and Ensembl (www.ensembl.org/) (as per de Busserolles et al., 2017; Fig. 4; Fig. S4). The opsin gene phylogeny was obtained by first aligning all opsin genes, i.e. the reference dataset and *N. brevirostris* genes, using the L-INS-I settings as part of the Geneious MAFFT plugin v.1.3.7 (Kato and Standley, 2013). jModeltest v.2.1.10 (Darriba et al., 2012) was subsequently used to select the most appropriate model of sequence evolution based on the Akaike information criterion. MrBayes v.3.2.6 (Ronquist et al., 2012) as part of the CIPRES platform (Miller et al., 2010) was then used to infer the phylogenetic relationship between opsin genes using the following parameter settings: GTR+I+G model; two independent MCMC searches with four chains each; 10 million generations per run; 1000 generations sample frequency; and 25% burn-in.

Opsin gene mining and phylogenetic reconstruction revealed, amongst a number of other visual opsin genes, two *N. brevirostris* *RH2* paralogues, of which one clustered within the *RH2A* clade of other percomorph fishes. However, the phylogenetic placement of the second paralogue could not be fully resolved using this approach alone (Fig. 4). Therefore, in order to resolve a more detailed relationship between the two *N. brevirostris* *RH2* paralogues, we took advantage of the phylogenetic signal within the single exons of




Fig. 4. Vertebrate visual opsin gene phylogeny. The different *N. brevirostris* opsin genes which were mined from the retinal transcriptomes are highlighted in bold and belong to four of the five major visual opsin classes. Black circles indicate Bayesian posterior probabilities >0.8. Note that the *N. brevirostris* RH2B gene is placed in between the percomorph RH2A and RH2B clades (also see Fig. 5). RH1, rhodopsin 1 (rod opsin); RH2, rhodopsin-like 2; SWS2, short-wavelength sensitive 2; LWS, long-wavelength sensitive; va, vertebrate ancient opsin (outgroup). Scale bar: 0.2 substitutions per site. Detailed phylogeny and GenBank accession numbers are shown in Fig. S4.

the displaced paralogue (as per Cortesi et al., 2015b). The five *N. brevirostris* RH2-2(B) exons were obtained by annotating the coding regions of the gene with a *P. fuscus* RH2 orthologue. The single exons were separated from one another and inserted as ‘single genes’ in the alignment in a reduced (RH2 genes only) reference dataset, along with the *N. brevirostris* RH2A gene. The RH2-specific phylogeny was then reconstructed using MrBayes on the CIPRES platform using the same parameters as before (Fig. 5A).

To support our phylogenetic analyses, the amino acid sequences of the RH2 paralogues were aligned to the amino acid sequence of bovine rhodopsin (GenBank accession no. NM001014890) using MAFFT in Geneious with parameters: L-INS-I, BLOSUM62, gap open penalty 1.53, offset value 0.123. Referring to bovine rhodopsin as the reference, this allowed us to compare site 122 between RH2s, which has been found to be key to distinguish between longer-shifted (RH2A in percomorphs) and shorter-shifted (RH2B in percomorphs) RH2 copies (Chinen et al., 2005; Fig. 5C).

Opsin gene expression

Quantitative opsin gene expression was assessed by mapping the filtered reads to the assembled coding regions of the *N. brevirostris* opsin genes as per de Busserolles et al. (2017). This methodology was used for each individual of the three life stages. Proportional

opsin expression for single cones ($p_{i,SC}$) and double cones ($p_{i,DC}$) for each gene (i) was then calculated by first normalizing the number of reads of each gene (R_i) to the length of each gene-specific coding region/sequence:

$$R_{i,normalized} = (R_i/bp_i), \quad (3)$$

where bp_i is the number of base pairs in the coding sequence of gene i , which was used to normalize the data between the opsins. The proportion of opsin expressed, out of the total normalized expression for single (T_{SC}) and double cones (T_{DC}), was then calculated separately. The following formulas were used, depending on the type of cone that gene i was expressed in:

$$p_{i,SC} = R_{i,normalized}/T_{SC} \quad (4)$$

or

$$p_{i,DC} = R_{i,normalized}/T_{DC}. \quad (5)$$

We also calculated the proportional expression of the rod opsin compared with total normalized opsin expression (T_{opsin}):

$$p_{i,rod} = R_{i,normalized}/T_{opsin}. \quad (6)$$

RESULTS

Topographic distribution of ganglion cells and SRP

Topographic maps of ganglion cells (including amacrine cells) for the three life stages of *N. brevirostris* were constructed from Nissl-stained retinal wholemounts. Little variation in topographic distribution of ganglion cells was observed within the same ontogenetic stage. Therefore, the topographic map of only one individual per life stage is presented here (Fig. 3A), and the results for the remaining individuals are shown in Fig. S1.

Differences in retinal topography were mainly found between the larval stage and the two later stages (Fig. 3A). In general, the larval retina showed less specialization compared with juvenile and adult retinas. In the larval retina, the onset of a horizontal streak was observed with the highest cell density found in the central meridian of the retina (1.5 times increase compared with areas with the lowest cell densities). Within this weak streak, three areas of high cell density were found: in the nasal, central and temporal parts of the retina. However, these areas of high cell density should be taken with caution because of the limitations of the Nissl-staining protocol for very small retinas. Larval retinas were challenging to prepare and analyse because of their small size and thus the higher amount of shrinkage present after staining. After several attempts with different larvae, only one larval retina was deemed acceptable for analysis. Even for this individual, the areas of high cell density in the nasal and temporal part of the retina are questionable, as they are very close to the retinal borders and therefore could be the result of shrinkage. A prominent horizontal streak along with a centralized area centralis (the area centralis had a 2.5–3 times increase in cell density compared with areas with the lowest cell densities) was present in the juvenile and adult individuals. Similar to the larvae, the streak in juveniles and adults was located on the central meridian of the retina, extending to the nasal and temporal margins. Although slightly different patterns were found for each life stage, they all showed a higher ganglion cell density in the central area close to the optic nerve, accompanied by a horizontal streak (Fig. 3A).

The total number of ganglion cells increased with the size of fish and ranged from 208,975 cells for the larval individual, to ~1,600,000 cells for juveniles and ~2,100,000 cells for adults (Table 1). Conversely, the mean cell density decreased with the size




Fig. 5. Single-exon green opsin (RH2) phylogeny. (A) Using the single exons (green) of the *N. brevisrostris* RH2B gene revealed that exons 3–5 cluster within or close to the percomorph RH2B clade, whereas exons 1 and 2 cluster close to or within the RH2A clade (*N. brevisrostris* RH2A in orange). Note that *Oryzias latipes* genes have a different nomenclature from the other fish opsin genes. Nodes denote Bayesian posterior probabilities. (B) Illustration of the relationship between the two RH2 paralogues of *N. brevisrostris* based on the single-exon RH2B phylogeny. The suggested gene conversion from RH2A into exons 1–3 of RH2B makes it near impossible to resolve its phylogenetic position if considering the whole coding region of the gene (see also Fig. 4; Fig. S4). The asterisk marks the approximate position of the amino acid residue 122 in RH2B. (C) RH2A and RH2B amino acid sequence alignments for residues 113–131 (according to bovine rhodopsin). Note that site 122 in RH2B, despite lying with the gene conversion region, shows the RH2B-specific, shorter-shifted 122Q (Chinen et al., 2005).

of the fish, ranging from 19,439 cells mm⁻² in the larval individual, to ~8,500 cells mm⁻² in juveniles and ~5000 cells mm⁻² in adults. Peak cell density also decreased through development, from 30,400 cells mm⁻² in the larval individual, to ~23,000 cells mm⁻² in juveniles and ~20,500 cells mm⁻² in adults.

Based on the peak ganglion cell density, the SRP of *N. brevisrostris* ranged from 2.98 cycles deg⁻¹ in the larval individual, to ~8.0 cycles deg⁻¹ in juveniles and a maximum of 11.0 cycles deg⁻¹ in adults (Table 1). Overall, SRP or visual acuity in *N. brevisrostris* increased with the size of the fish with very little variation found within ontogenetic stages (Fig. S2).

Topographic distribution of cone photoreceptors

The density and topographic distribution of cone photoreceptors (double and single cones) was assessed in the three life stages of

N. brevisrostris. Double and single cones were arranged in a square mosaic, with a single cone at the centre of four double cones (Fig. 2B). This pattern was consistent throughout the entire retina, thus providing a ratio of double cones to single cones of 2:1. As a consequence of this regular arrangement, the topographic distribution of single cones, double cones and total cones was identical. Moreover, similar to the ganglion cell topography, little variation in topographic distribution of cone photoreceptors was observed within the same ontogenetic stage. Therefore, the total cone topographic map of only one individual per life stage is presented here (Fig. 3B). The remaining maps (i.e. for single and double cones separately, and maps of all individuals) are provided in Fig. S3 and in Dryad (<https://doi.org/10.5061/dryad.k6djh9w37>).

The topographic distribution of cone photoreceptors varied between stages, with a continuous increase in specialization from

Table 1. Summary of ganglion cell quantitative data obtained using the optical fractionator method on the wholemount retinas of three developmental stages of *Naso brevisrostris*

| Stage | Individual | SL (cm) | Peak cell density (cells mm ⁻²) | Mean cell density (cells mm ⁻²) | Total cells | Lens \varnothing (mm) | SRP (cycles deg ⁻¹) |
|----------|------------|---------|---|---|-------------|-------------------------|---------------------------------|
| Adult | ID1 | 30.6 | 20,617 | 5340 | 2,034,000 | 6.6 | 10.6 |
| | ID2 | 30.8 | 20,370 | 4772 | 2,100,825 | 6.9 | 11.0 |
| Juvenile | ID1 | 13.8 | 23,750 | 8130 | 1,617,968 | 4.7 | 8.1 |
| | ID2 | 15.2 | 24,531 | 8688 | 1,737,656 | 4.9 | 8.6 |
| | ID3 | 15.7 | 21,875 | 8584 | 1,450,625 | 4.5 | 7.5 |
| Larvae | ID1 | 3.18 | 30,400 | 19,439 | 208,975 | 1.4 | 2.98 |

SL, standard length; SRP, spatial resolving power; \varnothing , diameter.

Table 2. Summary of photoreceptor quantitative data obtained using the optical fractionator method on the wholemount retinas of three developmental stages of *N. brevis*

| Stage | Individual | SL (cm) | Total DC | Peak DC density (cells mm ⁻²) | Total SC | Peak SC density (cells mm ⁻²) | Total cones | Peak total cone density (cells mm ⁻²) |
|----------|------------|---------|-----------|---|-----------|---|-------------|---|
| Adult | ID3 | 29.7 | 3,738,296 | 23,703 | 1,926,513 | 12,098 | 5,664,809 | 35,801 |
| | ID4 | 27.7 | 3,820,839 | 21,208 | 1,972,975 | 12,345 | 5,793,814 | 33,553 |
| Juvenile | ID1 | 13.8 | 2,466,222 | 34,218 | 1,283,524 | 18,125 | 3,749,746 | 52,343 |
| | ID4 | 19.5 | 3,212,099 | 32,968 | 1,667,475 | 17,187 | 4,879,574 | 50,155 |
| Larvae | ID2 | 3.08 | 452,010 | 45,432 | 231,371 | 23,703 | 683,381 | 69,135 |
| | ID3 | 3.30 | 413,476 | 45,432 | 211,105 | 23,704 | 624,581 | 69,136 |

SL, standard length; DC, double cone; SC, single cone.

the larval to the adult stage (Fig. 3B). Larvae had a weak horizontal streak in the central meridian with two large areas of high cell density in the nasal part and the temporal part of the retina. One of the two analysed larval individuals also showed a dorsal increase in cell density (Fig. S3f). However, this apparent increase in cell density was probably caused by an artefact from not properly flattening the dorsal part of the retina during mounting and should therefore be disregarded. Compared with the larvae, juveniles had a more pronounced horizontal streak in the central meridian with a peak density of cells in the temporal part of the retina. Moreover, a weak vertical streak could be seen in the temporal part of the retina, extending from the dorso-temporal area to the ventro-temporal area. In adults, the horizontal streak in the central meridian was still present but did not extend as far into the nasal part as in the juveniles. Moreover, the vertical streak was more prominent compared with that found in juveniles, resulting in a very large area of high cell density in the temporal region (Fig. 3B). The continuous nature of the transition between juvenile and adult specializations is highlighted by the topography of individuals of intermediate sizes (Fig. S3). For example, the horizontal streak was less pronounced in the nasal part of a larger (SL 19.5 cm; Fig. S3c) compared with a smaller juvenile (SL 13.8 cm; Fig. S3d). Conversely, the vertical streak in a smaller adult (SL 27.7 cm; Fig. S3b) was still developing compared with that found in a larger adult (SL 29.7 cm; Fig. S3a).

Similar to the ganglion cells (Table 1), the total number of photoreceptors increased with the size of the fish, ranging from ~650,000 cells in larvae, to ~4,300,000 cells in juveniles and ~5,700,000 cells in adults (Table 2). A large difference in the total number of photoreceptors was found between the two juvenile individuals. This difference is probably due to the size difference between these individuals. Photoreceptor peak cell density decreased with the size of the fish, ranging from ~69,000 cells mm⁻² in larvae, to ~51,000 cells mm⁻² in juveniles and ~34,000 cells mm⁻² in adults (Table 2).

The total number of cone photoreceptors was greater than the total number of ganglion cells, indicating a high summation ratio between the two cell types. For one individual (juvenile ID1), the distribution of both ganglion cells and photoreceptors was analysed, which allowed us to estimate the summation ratio between photoreceptors and ganglion cells in low- and high-density areas. For this individual, the summation ratio was found to be as low as 2.3 in the central part and as high as 5.4 in the ventro-temporal part of the retina.

Visual opsin genes and their expression in *N. brevis*

Naso brevis were found to mainly express four opsin genes in their retinas. Independent of ontogeny, these were the blue-violet opsin *SWS2B*, the green opsins *RH2B* and *RH2A*, and the rod opsin *RH1*. The red opsin *LWS* was also found to be expressed, albeit at very low levels in all stages (0.1–6.5% of total double cone opsin




Fig. 6. Proportional expression of *N. brevis* opsin genes. (A) Independent of ontogenetic stage, *N. brevis* expressed the single cone opsin gene *SWS2B* (100% of single cone expression; see Table S2 for details) and three double cone opsin genes: *RH2B*, *RH2A* and *LWS*. The proportional expression of double cone opsins revealed a change in expression of the two *RH2* genes between the larval and the juvenile stage as well as a steady increase in the expression of *LWS* with development. (B) The proportional expression of cone (*SWS2B*, *RH2B*, *RH2A*, *LWS*) versus rod opsin (*RH1*) remained similar throughout ontogeny. The box boundaries indicate the second and third quartiles, the central line is the median and the whiskers indicate the first and fourth quartiles of the data.

expression; Fig. 6A; Table S2). The phylogenetic reconstruction based on the full coding regions of the genes confirmed the positioning of all genes within their respective opsin class (Fig. 4). However, for *RH2B* in particular, the resolution between *RH2*-specific clades was poor (Fig. 4; Fig. S4). This was resolved using the exon-based approach, which showed the placement of some of the *N. brevirostris RH2B* exons within a greater percomorph *RH2B* clade (Fig. 5A). We also found evidence for substantial gene conversion affecting this gene, with the placement of exons 1 and 2 close to, or within, the *RH2A* clade (Fig. 5B). Amino acid comparison between the two *RH2* paralogues corroborated our phylogenetic placement of genes and revealed the conserved sites E122 and Q122 for the longer- (*RH2A*) and shorter-shifted (*RH2B*) copies, respectively (Fig. 5C).

Quantitative opsin gene expression revealed that *SWS2B* was the only single cone gene and thus was expressed at 100% in all developmental stages (Table S2). Of the double cone opsins, there was a change in expression for the *RH2* genes with ontogeny. During the larval stage ($n=3$), *RH2B* (mean \pm s.e.m., 36.2 \pm 4.8%) was less highly expressed compared with *RH2A* (63.6 \pm 4.8%). The opposite pattern was found in the juvenile ($n=6$) and adult stages ($n=3$), where *RH2B* was the most highly expressed of all double cone opsins genes (juvenile: 56 \pm 1.3%, adult: 56.1 \pm 1.9%). *RH2A* in the juvenile (41.2 \pm 1.4%) and adult (38.1 \pm 1.6%) stages was less highly expressed. Despite *LWS* being expressed at low levels in all stages, there was a noticeable increase in expression with development (larva: 0.2 \pm 0.0%, juvenile: 2.8 \pm 0.7%, adult 5.8 \pm 0.4%; Fig. 6A). Rod opsin (*RH1*) expression was substantially higher compared with cone opsin expression in all stages (82–86% for all stages; Fig. 6B).

DISCUSSION

The visual systems of fishes often change through development when transitioning from one habitat to another. These changes are usually associated with a shift in light environment, e.g. when moving from the open ocean to a coral reef, but possibly also with changes in diet and predation pressure (Sale, 2013). Our objective was to assess the visual system development in the spotted unicornfish, *N. brevirostris*. *Naso brevirostris* experiences multiple changes in habitat, diet and morphology throughout ontogeny (from larval to adult stages; Fig. 1) making it a prime candidate to study visual system changes on the reef.

Ganglion cell topography

Retinal topography is an effective method to identify visual specializations and recognize the area of the visual field of greatest importance to a species as it visually samples its environment (Collin, 2008; Collin and Pettigrew, 1988a; Hughes, 1977). In marine fishes, visual specializations have been found to correlate with the structure and symmetry of the environment they live in and/or with their feeding behaviour (Caves et al., 2017; Collin and Pettigrew, 1988a,b; Ito and Murakami, 1984; Shand, 1997). In this study, we show that the *N. brevirostris* eye possesses a horizontal streak in all life stages (Fig. 3A). This type of specialization has previously been found in species living in open environments where an uninterrupted view of the horizon, defined by the sand–water or air–water interface, is present (Collin and Pettigrew, 1988b). As *N. brevirostris* spends much of its life (larval and adult stages) searching for prey in the water column, having a pronounced horizontal streak is likely to increase feeding and predator surveillance capabilities by allowing it to scan the horizon without using excessive eye movements (Collin and Shand, 2003). Moreover, at the larval stage, this type of specialization may also enable fish to scan the environment when searching for a reef habitat to settle on. In contrast, a horizontal streak does not seem

to match the visual needs at the juvenile stage during which *N. brevirostris* lives in close association with the reef, i.e. in a more enclosed 3D environment. At this life stage, we would have expected to find one (or multiple) area centralis and no horizontal streak – a common feature in fish that live close to or within the reef matrix (Collin and Pettigrew, 1988a,c). Compared with the lifespan of these fishes (up to 20 years), the juvenile stage is relatively short (~3 years; Choat and Axe, 1996), which may explain the maintenance of the horizontal streak throughout development.

The peak ganglion cell density in the juvenile and adult stages is in the central part of the retina (i.e. at the centre of the streak; Fig. 3A). This is quite unusual, as the area of high cell density in coral reef fishes is normally found in the temporal zone, which receives information from the nasal visual field, and thus is usually correlated with feeding and predator avoidance in front of the fish (Collin and Pettigrew, 1988a, 1989; Fritsches et al., 2003; Fritsch et al., 2017; Shand et al., 2000). The type of specialization found in *N. brevirostris* could potentially be associated with its unusual visual behaviour, as these fish examine objects side-on (V.T., unpublished observations). A possible explanation for this peculiar behaviour is that because of its protruded snout, which grows through development, the frontal image might be partially blocked and stereoscopic vision may be impaired or even impossible (Purcell and Bellwood, 1993). Although the visual field of *N. brevirostris* was not investigated in their study, Brandl and Bellwood (2013) suggest that the protruding snout found in many *Naso* species indeed prevents an overlap of their horizontal field of view. Similar to the monocular vision found in hammerhead sharks (Lisney and Collin, 2008; McComb et al., 2009), increasing visual acuity in the central part of the retina would thus maximize a sideward oriented visual field. Together with a visual streak, a high cell density in the central part of the retina is likely to enable *N. brevirostris* to accurately navigate both within the complexity of the reef and in open water.

Photoreceptor topography

The photoreceptor topography showed a similar trend to the ganglion cell topography, with bigger changes occurring between the larval and juvenile stage and only minor adjustments thereafter (Fig. 3B). In addition to a weak horizontal streak, larval fish had two peaks of high cell density in the nasal and temporal zones, which comply with two of their main ecological needs: feeding (temporal; looking forward) and predator avoidance (nasal; looking backwards) (Boehlert, 1996; Collin and Pettigrew, 1988a; Fortier and Harris, 1989). These high-density regions were not matched by the ganglion cell topography and, as such, are likely to provide areas of higher sensitivity (i.e. areas of high photoreceptor to ganglion cell ratio; Walls, 1942). Moreover, although larval fishes rely mainly on olfactory cues to zoom in on a suitable habitat for settlement (Lecchini et al., 2005b), the area of high cell density in the temporal zone in particular might also assist when searching for such habitats over longer distances (Mouritsen et al., 2013).

In addition to the horizontal streak becoming more prominent in juvenile and adult stages, *N. brevirostris* showed a temporal vertical specialization appearing at the juvenile stage and being more pronounced in adults (Fig. 3B). Such a double streak specialization, with a vertical and a horizontal component, is the first to be found in coral reef fishes. *Naso brevirostris* adults live on the coral reef slope/wall, and move up and down the wall (from 2 to 122 m) while foraging and searching for mates (Mundy, 2005). As such, in line with the terrain hypothesis (Hughes, 1977), the evolution of this vertical specialization is likely to be a result of the vertical component in their visual environment.

A difference in the topography of ganglion cells and photoreceptors means that the summation ratio between the cell types, i.e. the sensitivity and spatial resolution of the retina, also differs depending on the visual field in question. For example, high photoreceptor densities and comparable low ganglion cell densities in the ventro-temporal and dorso-temporal parts of the vertical streak confer higher sensitivity to these two areas (Walls, 1942). Theoretically, this enables juvenile and adult *N. brevirostris* to detect even small differences in luminance, which may help to reveal well-camouflaged predators against the reef wall. A high density of both photoreceptor and ganglion cells in the centre of the retina, in contrast, confers a low summation ratio which leads to an increase in visual acuity (Walls, 1942). This area of high acuity may help fish to identify conspecifics and also to distinguish between food items (Cronin et al., 2014).

To summarize, the retinal topography in *N. brevirostris* may be adapted to the habitat in both the larval and the adult stage. Juveniles live in a more enclosed, 3D coral reef environment compared with the other two life stages. Therefore, we would have expected the juvenile visual system to reflect its habitat by having a less-developed streak and one or more area centralis instead. The unexpected topography found in the juvenile stage may be explained by this stage only lasting a fraction of the lifespan of *N. brevirostris* (Choat and Axe, 1996). The relatively short period of time spent in a habitat rich in shelter and food enables the fish to grow big enough to avoid most predators (Barnes and Hughes, 1999; Lasiak, 1986). During this time, juvenile *N. brevirostris* mostly feed on benthic algae, which do not require a highly specialized visual system in terms of retinal topography (Caves et al., 2017; Collin and Pettigrew, 1988a,b; Randall et al., 1997). As such, instead of changing the visual system multiple times, it is probably more energy efficient to maintain (or slightly adjust) a visual system that is optimally adapted for both the larval and adult stages.

Visual acuity

The visual acuity of *N. brevirostris* was found to increase through development (Fig. S2; Table 1). This seems to be a common feature in coral reef fishes, as a higher acuity often correlates with an increase in eye size during growth. The benefit of having a higher visual acuity is that, as fish grow and expand their home ranges, it increases the distance at which visual objects such as predators, conspecifics and food can be detected (Caves et al., 2017; Shand, 1997). Accordingly, as in other coral reef fish larvae (Shand, 1997), the acuity of *N. brevirostris* larvae was relatively poor (2.98 cycles deg⁻¹). The overabundance of zooplankton in their habitat means that larval coral reef fishes can wander instead of using a lock-and-pursuit feeding behaviour, i.e. they do not need to spot their food from a distance, but rather bump into it while floating in the plankton (Evans and Fernald, 1990; Fortier and Harris, 1989). Once settled on the reef, the visual acuity of *N. brevirostris* starts to increase in line with their growth (Fig. S2). Adult *N. brevirostris* were found to have a similar visual acuity (~11 cycles deg⁻¹) to that of other reef fishes of comparable size, such as the clown triggerfish, *Balistoides conspicillus*, a species that also inhabits the reef slope and shows a pronounced horizontal streak (Collin and Pettigrew, 1989).

Opsin gene evolution and expression

Phylogenetic reconstruction showed that the *N. brevirostris* visual opsins belong to the opsin gene clades usually found within percomorph fishes (Fig. 4). However, within the *RH2* genes, an exon-based phylogeny revealed that the *N. brevirostris RH2B* gene is likely to have undergone substantial gene conversion (Fig. 5A). As such, it seems that parts of its first and second exon have been

acquired from the *RH2A* paralogue, explaining its phylogenetic uncertainty when using whole coding region-based reconstructions (Fig. 4; Fig. S4). This is not that surprising, as *RH2* duplicates in teleosts are commonly found in tandem (e.g. Musilova et al., 2019) and, as is the case for other teleost opsin genes (Cortesi et al., 2015b; Hofmann and Carleton, 2009), frequently experience gene conversion (Cortesi et al., 2015b; Escobar-Camacho et al., 2016; Hofmann et al., 2012). This phenomenon is thought to be one of the main mechanisms for concerted evolution in small gene families, which often originate from tandem duplications (Li, 1997; Ohta, 1983) and could help to preserve gene function by repairing null mutations (Innan, 2009) or by resurrecting previously pseudogenized gene copies (Cortesi et al., 2015b). As the *RH2* opsin genes are highly expressed in *N. brevirostris*, they seem rather important for their visual ecology, and it is therefore likely that gene conversion played a major evolutionary role in maintaining their function.

Interestingly, the key spectral tuning site 122, despite being within the area of gene conversion, showed the *RH2B*-specific 122Q (Fig. 5B,C). As the remainder of the exon is almost identical to *RH2A*, it is likely that this site was re-gained post-gene conversion, suggesting that strong selection is maintaining two spectrally diverse *RH2* copies in this species. As such, based on opsin gene expression, *N. brevirostris* could be behaviourally trichromatic (i.e. has three spectral sensitivities) for all three developmental stages, with the violet opsin *SWS2B* and the blue-green opsin *RH2B* and *RH2A* genes being expressed in sufficient quantity to enable this level of chromatic analysis. Supporting these findings, microspectrophotometry (MSP) in adults of two closely related *Naso* species (*Naso literatus* and *Naso unicornis*; Sorenson et al., 2013) found three cone photoreceptors with spectral sensitivities of ~420 nm λ_{\max} for single cones, and ~490 nm λ_{\max} and ~515 nm λ_{\max} for the accessory and principal members of double cones, respectively (Losey et al., 2003). A short-shifted visual system with high sensitivity in the violet to green range might benefit feeding on zooplankton and gelatinous prey during the larval and adult stages of *N. brevirostris* (Marshall et al., 2003). However, it seems at odds with the mainly algivorous diet of the juvenile stage, where a red-shifted visual system would be of advantage (Cortesi et al., 2018 preprint; Stieb et al., 2017).

Rather than the anticipated long-wavelength shift in juveniles, the most notable change in opsin gene expression was found in the *RH2* copies between the larval and later *N. brevirostris* stages (Fig. 6A). A change in expression levels of *RH2B* and *RH2A* has also been found in coral reef damselfishes (Pomacentridae) (Stieb et al., 2016). On shallow, clear coral reefs, a broad spectrum of light is available (Marshall et al., 2003). However, with increasing depth, the long and short ends of the spectrum are cut off as a result of absorption and scattering through interfering particles, resulting in a blue mid-wavelength saturated light environment (Smith and Baker, 1981). Consequently, in an attempt to maximize photon catch, some damselfish species have been found to increase the expression of the blue-sensitive *RH2B* gene and simultaneously decrease the expression of the green-sensitive *RH2A* gene with increasing depth (Stieb et al., 2016). In *N. brevirostris*, the shift in expression of *RH2* genes occurs between the larval and juvenile stages where depth differences do not seem that relevant. In lieu of depth, individuals migrate from a pelagic blue-shifted open water environment to the more green-shifted light environment of the coral reef (Marshall et al., 2003). This could in theory explain the high *RH2A* expression in larval fish at the settlement stage; however, it does not explain the increase in *RH2B* expression post-settlement (Fig. 6A). An increasing number of fishes are being found to change their opsin

gene expression to tune photoreceptors to the prevailing photic environment (e.g. Fuller et al., 2004; Härer et al., 2017; Hofmann et al., 2010; Luehrmann et al., 2018; Nandamuri et al., 2017; Shand et al., 2008; Spady et al., 2016; Stieb et al., 2016). At the opposite end of the spectrum, opsin gene expression might be pre-programmed either by phylogeny or on a species by species basis, as exemplified by only some damselfishes changing expression with depth (Stieb et al., 2016). It is therefore possible that opsin gene expression in *N. brevirostris* is under phylogenetic control and that changes in photic environment contribute very little to opsin gene expression in this case. Also, it is important to note that we currently do not know whether and how quantitative differences in opsin gene expression translate to the behaviour of *N. brevirostris*. Hence, until these results become available, it remains speculative as to what impact the changes observed here might have on the visual ecology of these fishes.

Naso brevirostris was not found to express the UV-sensitive *SWS1* gene at any developmental stage. *SWS1* expression is often found in larval fish and more generally in fishes feeding on zooplankton, with UV-vision thought to increase the detectability of this food source (Browman et al., 1994; Sabbah et al., 2010). As *N. brevirostris* feeds on zooplankton at both larval and adult stages (Choat et al., 2002, 2004), the lack of *SWS1* expression seems striking. However, it does support ocular media measurements, which revealed UV-blocking lenses in both larval and adult *N. brevirostris* (Siebeck and Marshall, 2007). UV-blocking lenses seem common in many bigger coral reef fishes, which is thought to enhance sighting distance by reducing chromatic aberration and scatter, as well as protecting the eye from the damage caused by these high-intensity wavelengths (Siebeck and Marshall, 2001). Instead, the expression of the violet-sensitive *SWS2B* gene, as its spectral absorption curve reaches into the near-UV (based on MSP in closely related species; Losey et al., 2003), may be sufficient to increase the discrimination of zooplankton from the water background while foraging.

We furthermore found low expression of the *LWS* gene (<6%) at all developmental stages. This suggests that *LWS* expression is either restricted to certain areas of the retina or interspersed at low frequency across the retina, or some photoreceptors might co-express *LWS* with an *RH2* gene (e.g. Cortesi et al., 2016; Dalton et al., 2014; Torres-Dowdall et al., 2017). MSP in related *Naso* species did not show any long-wavelength-sensitive photoreceptors, nor did it show any evidence for opsin co-expression (i.e. red-shifted unusually broad absorbance peaks; Losey et al., 2003). As this technique only samples a subset of the photoreceptors across the retina, it might be that the photoreceptors containing this pigment were missed because of their low number or that *LWS* was simply not expressed in these fishes.

It is possible that the *LWS* expression found here is just a by-product of the way opsin gene expression is controlled and that it does not serve any ecological function. Nevertheless, *LWS* expression did increase steadily with development (Fig. 6A). Hence, an alternative explanation might be that *LWS* is co-expressed with an *RH2* gene, which has been shown to increase achromatic discrimination in cichlids (Dalton et al., 2014). Moreover, *LWS* expression has recently been shown to be correlated with algal feeding in damselfishes (Stieb et al., 2017), and blennies (Cortesi et al., 2018 preprint). As *N. brevirostris* juveniles, and to a certain extent also adults (Choat et al., 2002), feed on algal turf, an increase in *LWS* expression in later stages may improve feeding efficiency through the increased contrast of algae against the reef background (Cortesi et al., 2015a; Marshall et al., 2003; Stieb et al., 2017).

In situ hybridization studies (e.g. Dalton et al., 2014, 2016; Stieb et al., 2019; Torres-Dowdall et al., 2017) coupled with behavioural colour vision experiments (e.g. Cheney et al., 2019) will be needed in the future to assess the distribution and function of the various opsin genes and ultimately the colour vision system of *N. brevirostris*.

Conclusion

Using a multidisciplinary approach, we analysed the ontogeny of the visual system of *N. brevirostris*. Changes in retinal topography and opsin gene expression were mainly found between the larval and juvenile stages, with few minor modifications thereafter. The juvenile stage in particular did not seem to match its ecology very well, which may be explained by the short duration of this stage compared with the lifespan of these fishes. This is different from the steady development that was found in other reef fishes (Cortesi et al., 2016; Shand et al., 2008; Job and Julia, 2001) and highlights the need for a comprehensive analysis of visual ontogeny across the reef fish community.

Acknowledgements

We would like to thank Sara M. Stieb, Vivien Rothenberger, Simon Dunn and the staff of Tetiaroa Society (Moana LeRohellec) and of CRIOBE (Camille Gache) for assistance with specimen collection. We furthermore thank the staff at the Lizard Island Research Station for logistical support, and Janette Edson from the Queensland Brain Institute (QBI) sequencing facility for library preparation and RNA sequencing. We also acknowledge QBI Advanced Microscopy Facility for the use of the Stereo Investigator (software v.2017.01.1), generously supported by the Australian Government through the ARC LIEF grant LE100100074.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: F.d.B., N.J.M., F.C.; Methodology: V.T., F.d.B., F.C.; Software: V.T.; Validation: F.d.B., F.C.; Formal analysis: V.T., F.C.; Investigation: V.T., F.C.; Resources: F.d.B., D.L., N.J.M., F.C.; Data curation: F.C.; Writing - original draft: V.T., F.d.B., F.C.; Writing - review & editing: V.T., F.d.B., D.L., N.J.M., F.C.; Visualization: V.T., F.d.B.; Supervision: F.d.B., N.J.M., F.C.; Project administration: F.C.; Funding acquisition: D.L., N.J.M., F.C.

Funding

This study was funded by the Sea World Research and Rescue Foundation, an Australian Research Council Discovery Project Grant (ARC DP180102363), and by the Tetiaroa Society, the Leonardo Di Caprio Foundation and Mission Blue for the CRIOBE study at Tetiaroa. F.d.B. was supported by an Australian Research Council DECRA Fellowship (DE180100949), N.J.M. by an Australian Research Council Laureate Fellowship, and F.C. by a University of Queensland Development Fellowship.

Data availability

Raw-read transcriptomes (PRJNA590573, SAMN13340708–13340719) and single gene sequences (MN720499–MN720503) are available through GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>). Gene alignments, phylogenies, transcriptome assemblies and retinal topography maps of single and double cones for each individual are available from the Dryad digital repository (Tettamanti et al., 2019): [dryad.k6dj9w37](https://doi.org/10.2202/dryad.k6dj9w37)

Supplementary information

Supplementary information available online at <http://jeb.biologists.org/lookup/doi/10.1242/jeb.209916.supplemental>

References

- Afgan, E., Sloggett, C., Goonasekera, N., Makunin, I., Benson, D., Crowe, M., Gladman, S., Kowsar, Y., Pheasant, M., Horst, R. et al. (2015). Genomics virtual laboratory: a practical bioinformatics workbench for the cloud. *PLoS ONE* **10**, e0140829. doi:10.1371/journal.pone.0140829
- Baddeley, A. and Turner, R. (2005). Spatstat: an R package for analyzing spatial point patterns. *J. Stat. Softw.* **12**, 1–42. doi:10.18637/jss.v012.i06

- Bailes, H. J., Trezise, A. E. and Collin, S. P. (2006). The number, morphology, and distribution of retinal ganglion cells and optic axons in the Australian lungfish *Neoceratodus forsteri* (Krefft 1870). *Vis. Neurosci.* **23**, 257-273. doi:10.1017/S0952523806232103
- Barnes, R. S. K. and Hughes, R. N. (1999). *An Introduction to Marine Ecology*. Oxford, UK: Blackwell Science Ltd, a Blackwell Publishing company.
- Besson, M., Gache, C., Brooker, R. M., Moussa, R. M., Waqalevu, V. P., LeRohellec, M., Jaouen, V., Peyrusse, K., Berthe, C., Bertucci, F. et al. (2017). Consistency in the supply of larval fishes among coral reefs in French Polynesia. *PLoS ONE* **12**, e0178795. doi:10.1371/journal.pone.0178795
- Boehlert, G. W. (1996). Larval dispersal and survival in tropical reef fishes. In *Reef Fisheries* (ed. N. V. C. Polunin and C. M. Roberts), pp. 61-84. Dordrecht, NL: Springer Netherlands.
- Bolger, A. M., Lohse, M. and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120. doi:10.1093/bioinformatics/btu170
- Bowmaker, J. K. (2008). Evolution of vertebrate visual pigments. *Vision Res.* **48**, 2022-2041. doi:10.1016/j.visres.2008.03.025
- Bowmaker, J. K. and Kunz, Y. W. (1987). Ultraviolet receptors, tetrachromatic colour vision and retinal mosaics in the brown trout (*Salmo trutta*): age-dependent changes. *Vision Res.* **27**, 2101-2108. doi:10.1016/0042-6989(87)90124-6
- Bozzano, A. and Collin, S. P. (2000). Retinal ganglion cell topography in elasmobranchs. *Brain Behav. Evol.* **55**, 191-208. doi:10.1159/000006652
- Brandl, S. J. and Bellwood, D. R. (2013). Morphology, sociality, and ecology: can morphology predict pairing behavior in coral reef fishes? *Coral Reefs*. **32**, 835-846. doi:10.1007/s00338-013-1042-0
- Browman, H. I., Novales-Flamarique, I. and Hawryshyn, C. (1994). Ultraviolet photoreception contributes to prey search behaviour in two species of zooplanktivorous fishes. *J. Exp. Biol.* **186**, 187-198.
- Carleton, K. L., Spady, T. C., Streelman, J. T., Kidd, M. R., McFarland, W. N. and Loew, E. R. (2008). Visual sensitivities tuned by heterochronic shifts in opsin gene expression. *BMC Biol.* **6**, 22. doi:10.1186/1741-7007-6-22
- Caves, E. M., Sutton, T. T. and Johnsen, S. (2017). Visual acuity in ray-finned fishes correlates with eye size and habitat. *J. Exp. Biol.* **220**, 1586-1596. doi:10.1242/jeb.151183
- Cheney, K. L., Green, N. F., Vibert, A. P., Vorobyev, M., Marshall, N. J., Osorio, D. C. and Endler, J. A. (2019). An Ishihara-style test of animal colour vision. *J. Exp. Biol.* **222**, jeb189787. doi:10.1242/jeb.189787
- Chinen, A., Matsumoto, Y. and Kawamura, S. (2005). Reconstitution of ancestral green visual pigments of zebrafish and molecular mechanism of their spectral differentiation. *Molecular biology and evolution*. *Mol. Biol. Evol.* **22**, 1001-1010. doi:10.1093/molbev/msi086
- Choat, J. H. and Axe, L. M. (1996). Growth and longevity in acanthurid fishes; an analysis of otolith increments. *Mar. Ecol. Prog. Ser.* **134**, 15-26. doi:10.3354/meps134015
- Choat, J. H., Clements, K. D. and Robbins, W. D. (2002). The trophic status of herbivorous fishes on coral reefs. I. Dietary analyses. *Mar. Biol.* **140**, 613-623. doi:10.1007/s00227-001-0715-3
- Choat, J. H., Robbins, W. D. and Clements, K. D. (2004). The trophic status of herbivorous fishes on coral reefs. II. Food processing modes and trophodynamics. *Mar. Biol.* **145**, 445-454. doi:10.1007/s00227-004-1341-7
- Coimbra, J. P., Marceliano, M. L., da Silveira Andrade-Da-Costa, B. L. and Yamada, E. S. (2006). The retina of tyrant flycatchers: topographic organization of neuronal density and size in the ganglion cell layer of the great kiskadee *Pitangus sulphuratus* and the rusty margined flycatcher *Myiozetetes cayanensis* (Aves: Tyrannidae). *Brain Behav. Evol.* **68**, 15-25. doi:10.1159/000092310
- Coimbra, J. P., Trévia, N., Videira Marceliano, M. L., da Silveira Andrade-Da-Costa, B. L., Picanço-Diniz, C. W. and Yamada, E. S. (2009). Number and distribution of neurons in the retinal ganglion cell layer in relation to foraging behaviors of tyrant flycatchers. *J. Comp. Neurol.* **514**, 66-73. doi:10.1002/cne.21992
- Coimbra, J. P., Nolan, P. M., Collin, S. P. and Hart, N. S. (2012). Retinal ganglion cell topography and spatial resolving power in penguins. *Brain Behav. Evol.* **80**, 254-268. doi:10.1159/000341901
- Collin, S. P. (1988). The retina of the shovel-nosed ray, *Rhinobatos batillum* (Rhinobatidae): morphology and quantitative analysis of the ganglion, amacrine and bipolar cell populations. *Exp. Biol.* **47**, 195-207.
- Collin, S. P. (2008). A web-based archive for topographic maps of retinal cell distribution in vertebrates. *Clin. Exp. Optom.* **91**, 85-95. doi:10.1111/j.1444-0938.2007.00228.x
- Collin, S. P. and Pettigrew, J. D. (1988a). Retinal topography in reef teleosts. I. Some species with well-developed areae but poorly-developed streaks. *Brain Behav. Evol.* **31**, 269-282. doi:10.1159/000116594
- Collin, S. P. and Pettigrew, J. D. (1988b). Retinal topography in reef teleosts. II. Some species with prominent horizontal streaks and high-density areae. *Brain Behav. Evol.* **31**, 283-295. doi:10.1159/000116595
- Collin, S. P. and Pettigrew, J. D. (1988c). Retinal ganglion cell topography in teleosts: a comparison between nissl-stained material and retrograde labelling from the optic nerve. *J. Comp. Neurol.* **276**, 412-422. doi:10.1002/cne.902760306
- Collin, S. P. and Pettigrew, J. D. (1989). Quantitative comparison of the limits on visual spatial resolution set by the ganglion cell layer in twelve species of reef teleosts. *Brain Behav. Evol.* **34**, 184-192. doi:10.1159/000116504
- Collin, S. P. and Shand, J. (2003). Retinal sampling and the visual field in fishes. In *Sensory Processing in Aquatic Environments* (ed. S. P. Collin and N. J. Marshall), pp. 139-169. New York, NY: Springer New York.
- Cortesi, F., Feeney, W. E., Ferrari, M. C., Waldie, P. A., Phillips, G. A., McClure, E. C., Sköld, H. N., Salzburger, W., Marshall, N. J. and Cheney, K. L. (2015a). Phenotypic plasticity confers multiple fitness benefits to a mimic. *Curr. Biol.* **25**, 949-954. doi:10.1016/j.cub.2015.02.013
- Cortesi, F., Musilová, Z., Stieb, S. M., Hart, N. S., Siebeck, U. E., Malmström, M., Tørresen, O. K., Jentoft, S., Cheney, K. L., Marshall, N. J. et al. (2015b). Ancestral duplications and highly dynamic opsin gene evolution in perciform fishes. *Proc. Natl. Acad. Sci. USA* **112**, 1493-1498. doi:10.1073/pnas.1417803112
- Cortesi, F., Musilová, Z., Stieb, S. M., Hart, N. S., Siebeck, U. E., Cheney, K. L., Salzburger, W. and Marshall, N. J. (2016). From crypsis to mimicry: changes in colour and the configuration of the visual system during ontogenetic habitat transitions in a coral reef fish. *J. Exp. Biol.* **219**, 2545-2558. doi:10.1242/jeb.139501
- Cortesi, F., Cheney, K. L., Cooke, G. M. and Ord, T. (2018). Opsin gene evolution in amphibious and terrestrial combtooth blennies (Blenniidae). *bioRxiv* doi:10.1101/503516
- Cronin, T. W., Johnsen, S., Marshall, N. J. and Warrant, E. J. (2014). *Visual Ecology*. Princeton, NJ: Princeton University Press.
- Dalton, B. E., Loew, E. R., Cronin, T. W. and Carleton, K. L. (2014). Spectral tuning by opsin coexpression in retinal regions that view different parts of the visual field. *Proc. R. Soc. B Biol. Sci.* **281**, 20141980. doi:10.1098/rspb.2014.1980
- Dalton, B. E., de Busserolles, F., Marshall, N. J. and Carleton, K. L. (2016). Retinal specialization through spatially varying cell densities and opsin coexpression in cichlid fish. *J. Exp. Biol.* **220**, 266-277. doi:10.1242/jeb.149211
- Darriba, D., Taboada, G. L., Doallo, R. and Posada, D. (2012). jModelTest 2: more models, new heuristics and parallel computing. *Nat. Methods* **9**, 772. doi:10.1038/nmeth.2109
- de Busserolles, F., Marshall, N. J. and Collin, S. P. (2014a). Retinal ganglion cell distribution and spatial resolving power in deep-sea lanternfishes (Myctophidae). *Brain Behav. Evol.* **84**, 262-276. doi:10.1159/000365960
- de Busserolles, F., Fitzpatrick, J. L., Marshall, N. J. and Collin, S. P. (2014b). The influence of photoreceptor size and distribution on optical sensitivity in the eyes of lanternfishes (Myctophidae). *PLoS ONE* **9**, e99957. doi:10.1371/journal.pone.0099957
- de Busserolles, F., Cortesi, F., Helvik, J. V., Davies, W. I. L., Templin, R. M., Sullivan, R. K. P., Michell, C. T., Mountford, J. K., Collin, J. K., Irgoien, X. et al. (2017). Pushing the limits of photoreception in twilight conditions: the rod-like cone retina of the deep-sea pearlshades. *Sci. Adv.* **3**, eaao4709. doi:10.1126/sciadv.aao4709
- Endler, J. A. (1990). On the measurement and classification of colour in studies of animal colour patterns. *Biol. J. Linn. Soc.* **41**, 315-352. doi:10.1111/j.1095-8312.1990.tb00839.x
- Escobar-Camacho, D., Ramos, E., Martins, C. and Carleton, K. L. (2016). The opsin genes of amazonian cichlids. *Mol. Ecol.* **26**, 1343-1356. doi:10.1111/mec.13957
- Evans, B. I. and Fernald, R. D. (1990). Metamorphosis and fish vision. *J. Neurobiol.* **21**, 1037-1052. doi:10.1002/neu.480210709
- Fernald, R. D. (1988). Aquatic adaptations in fish eyes. In *Sensory Biology of Aquatic Animals* (ed. J. Atema, R. R. Fay, A. N. Popper and W. N. Tavolga), pp. 435-466. New York, NY: Springer New York.
- Fortier, L. and Harris, R. P. (1989). Optimal foraging and density-dependent competition in marine fish larvae. *Mar. Ecol. Prog. Ser.* **51**, 19-33. doi:10.3354/meps051019
- Fritsch, R., Collin, S. P. and Michiels, N. K. (2017). Anatomical analysis of the retinal specializations to a crypto-benthic, micro-predatory lifestyle in the Mediterranean triplefin blenny *Tripterygion delaisi*. *Front. Neuroanat.* **11**, 122. doi:10.3389/fnana.2017.00122
- Fritsches, K. A., Marshall, N. J. and Warrant, E. (2003). Retinal specializations in the blue marlin: eyes designed for sensitivity to low light levels. *Mar. Freshw. Res.* **54**, 333-341. doi:10.1071/MF02126
- Fuller, R. C., Carleton, K. L., Fadool, J. M., Spady, T. C. and Travis, J. (2004). Population variation in opsin expression in the bluefin killifish, *Lucania goodei*: a real-time PCR study. *J. Comp. Physiol. A* **190**, 147-154. doi:10.1007/s00359-003-0478-z
- Garza-Gisholt, E., Hemmi, J. M., Hart, N. S. and Collin, S. P. (2014). A comparison of spatial analysis methods for the construction of topographic maps of retinal cell density. *PLoS ONE* **9**, e93485. doi:10.1371/journal.pone.0093485
- Glaser, E. M. and Wilson, P. D. (1998). The coefficient of error of optical fractionator population size estimates: a computer simulation comparing three estimators. *J. Microsc.* **192**, 163-171. doi:10.1046/j.1365-2818.1998.00417.x
- Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B., Eccles, D., Li, B. and Lieber, M. (2013). De novo transcript

- sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494. doi:10.1038/nprot.2013.084
- Härer, A., Torres-Dowdall, J. and Meyer, A. (2017). Rapid adaptation to a novel light environment: the importance of ontogeny and phenotypic plasticity in shaping the visual system of Nicaraguan Midas cichlid fish (*Amphilophus citrinellus* spp.). *Mol. Ecol.* **26**, 5582–5593. doi:10.1111/mec.14289
- Helfman, G., Collette, B. B., Facey, D. E. and Bowen, B. W. (2009). *The Diversity of Fishes: Biology, Evolution, and Ecology*. West Sussex, UK: Wiley-Blackwell.
- Hofmann, C. M. and Carleton, K. L. (2009). Gene duplication and differential gene expression play an important role in the diversification of visual pigments in fish. *Integr. Comp. Biol.* **49**, 630–643. doi:10.1093/icb/icp079
- Hofmann, C. M., O'Quin, K. E., Smith, A. R. and Carleton, K. L. (2010). Plasticity of opsin gene expression in cichlids from Lake Malawi. *Mol. Ecol.* **19**, 2064–2074. doi:10.1111/j.1365-294X.2010.04621.x
- Hofmann, C. M., Marshall, N. J., Abdilleh, K., Patel, Z., Siebeck, U. E. and Carleton, K. L. (2012). Opsin evolution in damselfish: convergence, reversal, and parallel evolution across tuning sites. *J. Mol. Evol.* **75**, 79–91. doi:10.1007/s00239-012-9525-0
- Holzer, G., Besson, M., Lambert, A., Francois, L., Barth, P., Gillet, B., Hughes, S., Piganeau, G., Leulier, F., Viriot, L. et al. (2017). Fish larval recruitment to reefs is a thyroid hormone-mediated metamorphosis sensitive to the pesticide chlorpyrifos. *eLife* **6**, e27595. doi:10.7554/eLife.27595
- Hughes, A. (1975). A quantitative analysis of the cat retinal ganglion cell topography. *J. Comp. Neurol.* **163**, 107–128. doi:10.1002/cne.901630107
- Hughes, A. (1977). The topography of vision in mammals of contrasting life style: comparative optics and retinal organisation. In *The Visual System in Vertebrates* (ed. F. Crescitelli), pp. 613–756. Berlin, DE: Springer Berlin Heidelberg.
- Hunt, D. M., Hankins, M. W., Collin, S. P. and Marshall, N. J. (2014). *Evolution of Visual and Non-Visual Pigments*. New York, NY: Springer New York.
- Innan, H. (2009). Population genetic models of duplicated genes. *Genetica* **137**, 19. doi:10.1007/s10709-009-9355-1
- Ito, H. and Murakami, T. (1984). Retinal ganglion cells in two teleost species, *Sebastiscus marmoratus* and *Navodon modestus*. *J. Comp. Neurol.* **229**, 80–96. doi:10.1002/cne.902290107
- Job, S. D. and Bellwood, D. R. (2000). Light sensitivity in larval fishes: implications for vertical zonation in the pelagic zone. *Limnol. Oceanogr.* **45**, 362–371. doi:10.4319/lo.2000.45.2.0362
- Job, S. D. and Julia, S. (2001). Spectral sensitivity of larval and juvenile coral reef fishes: implications for feeding in a variable light environment. *Mar. Ecol. Prog. Ser.* **214**, 267–277. doi:10.3354/meps214267
- Katoh, K. and Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780. doi:10.1093/molbev/mst010
- Lasiak, T. A. (1986). Juveniles, food and the surf zone habitat: implications for teleost nursery areas. *South African J. Zool.* **21**, 51–56. doi:10.1080/02541858.1986.11447956
- Lecchini, D., Dufour, V., Carleton, J., Strand, S. and Galzin, R. (2004). Estimating the patch size of larval fishes during colonization on coral reefs. *J. Fish Biol.* **65**, 1142–1146. doi:10.1111/j.0022-1112.2004.00493.x
- Lecchini, D., Planes, S. and Galzin, R. (2005a). Experimental assessment of sensory modalities of coral-reef fish larvae in the recognition of their settlement habitat. *Behav. Ecol. Sociobiol.* **58**, 18–26. doi:10.1007/s00265-004-0905-3
- Lecchini, D., Shima, J., Banaigs, B. and Galzin, R. (2005b). Larval sensory abilities and mechanisms of habitat selection of a coral reef fish during settlement. *Oecologia* **143**, 326–334. doi:10.1007/s00442-004-1805-y
- Leis, J. M. and Carson-Ewart, B. M. (1999). In situ swimming and settlement behaviour of larvae of an Indo-Pacific coral-reef fish, the coral trout *Plectropomus leopardus* (Pisces: Serranidae). *Mar. Biol.* **134**, 51–64. doi:10.1007/s002270050524
- Li, W. (1997). *Molecular Evolution*. Sunderland, USA: Sinauer Associates Incorporated.
- Lin, J.-J., Wang, F.-Y., Li, W.-H. and Wang, T.-Y. (2017). The rises and falls of opsin genes in 59 ray-finned fish genomes and their implications for environmental adaptation. *Sci. Rep.* **7**, 15568. doi:10.1038/s41598-017-15868-7
- Lisney, T. J. and Collin, S. P. (2008). Retinal ganglion cell distribution and spatial resolving power in elasmobranchs. *Brain Behav. Evol.* **72**, 59–77. doi:10.1159/000146082
- Loew, E. R., McFarland, W. N., Mills, E. L. and Hunter, D. (1993). A chromatic action spectrum for planktonic predation by juvenile yellow perch, *Perca flavescens*. *Can. J. Zool.* **71**, 384–386. doi:10.1139/z93-053
- Losey, G. S., McFarland, W. N., Loew, E. R., Zamzow, J. P., Nelson, P. A. and Marshall, N. J. (2003). Visual biology of Hawaiian coral reef fishes. I. Ocular transmission and visual pigments. *Copeia* **2003**, 433–454. doi:10.1643/01-053
- Luehrmann, M., Stieb, S. M., Carleton, K. L., Pietzker, A., Cheney, K. L. and Marshall, N. J. (2018). Short-term colour vision plasticity on the reef: changes in opsin expression under varying light conditions differ between ecologically distinct fish species. *J. Exp. Biol.* **221**, jeb175281. doi:10.1242/jeb.175281
- Lythgoe, J. N. (1979). *Ecology of Vision*. New York, US: Oxford University Press.
- Marshall, N. J., Jennings, K., McFarland, W. N., Loew, E. R. and Losey, G. S. (2003). Visual biology of Hawaiian coral reef fishes. III. Environmental light and an integrated approach to the ecology of reef fish vision. *Copeia* **2003**, 467–480. doi:10.1643/01-056
- Matthiessen, L. (1882). Ueber die Beziehungen, welche zwischen dem Brechungsindex des Kerncentrums der Krystalllinse und den Dimensionen des Auges bestehen. *Archiv für die gesamte Physiologie des Menschen und der Tiere* **27**, 510–523. doi:10.1007/BF01802978
- McComb, D. M., Tricas, T. C. and Kajjira, S. M. (2009). Enhanced visual fields in hammerhead sharks. *J. Exp. Biol.* **212**, 4010–4018. doi:10.1242/jeb.032615
- Miller, M. A., Pfeiffer, W. and Schwartz, T. (2010). *Creating the CIPRES Science Gateway for Inference of Large Phylogenetic Trees*, pp. 1–8. Gateway Computing Environments Workshop (GCE), IEEE.
- Mouritsen, H., Atema, J., Kingsford, M. J. and Gerlach, G. (2013). Sun compass orientation helps coral reef fish larvae return to their natal reef. *PLoS ONE* **8**, e66039. doi:10.1371/journal.pone.0066039
- Mundy, B. C. (2005). Checklist of the fishes of the Hawaiian Archipelago. *Bishop Mus. Bull. Zool.* **6**, 1–704.
- Musilova, Z., Cortesi, F., Matschiner, M., Davies, W. I. L., Patel, J. S., Stieb, S. M., de Busserolles, F., Malmström, M., Tørresen, O. K., Brown, C. J. et al. (2019). Vision using multiple distinct rod opsins in deep-sea fishes. *Science* **364**, 588–592. doi:10.1126/science.aav4632
- Nandamuri, S. P., Yourick, M. R. and Carleton, K. L. (2017). Adult plasticity in African cichlids: rapid changes in opsin expression in response to environmental light differences. *Mol. Ecol.* **26**, 6036–6052. doi:10.1111/mec.14357
- Novales-Flamarique, I. (2016). Diminished foraging performance of a mutant zebrafish with reduced population of ultraviolet cones. *Proc. R. Soc. B Biol. Sci.* **283**, 20160058. doi:10.1098/rspb.2016.0058
- Ohta, T. (1983). On the evolution of multigene families. *Theoretical Population Biology* **23**, 216–240.
- Peichl, L., Nemeč, P. and Burda, H. (2004). Unusual cone and rod properties in subterranean African mole-rats (Rodentia, Bathyergidae). *Eur. J. Neurosci.* **19**, 1545–1558. doi:10.1111/j.1460-9568.2004.03263.x
- Purcell, S. W. and Bellwood, D. R. (1993). A functional analysis of food procurement in two surgeonfish species, *Acanthurus nigrofasciatus* and *Ctenochaetus striatus* (Acanthuridae). *Environ. Biol. Fishes* **37**, 139–159. doi:10.1007/BF00000589
- Randall, J. E., Gerald, R. A. and Steene, R. C. (1997). *Fishes of the Great Barrier Reef and Coral Sea*. Honolulu, USA: University of Hawaii Press.
- Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M. A. and Huelsenbeck, J. P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* **61**, 539–542. doi:10.1093/sysbio/sys029
- Sabbah, S., Laria, R. L., Gray, S. M. and Hawryshyn, C. W. (2010). Functional diversity in the color vision of cichlid fishes. *BMC Biol.* **8**, 133. doi:10.1186/1741-7007-8-133
- Sale, P. F. (2013). *The Ecology of Fishes on Coral Reefs*. Elsevier.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B. et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682. doi:10.1038/nmeth.2019
- Shand, J. (1997). Ontogenetic changes in retinal structure and visual acuity: a comparative study of coral-reef teleosts with differing post-settlement lifestyles. *Environ. Biol. Fishes.* **49**, 307–322. doi:10.1023/A:1007353003066
- Shand, J., Archer, M. A. and Collin, S. P. (1999). Ontogenetic changes in the retinal photoreceptor mosaic in a fish, the black bream, *Acanthopagrus butcheri*. *J. Comp. Neurol.* **412**, 203–217. doi:10.1002/(SICI)1096-9861(19990920)412:2<203::AID-CNE2>3.0.CO;2-3
- Shand, J., Chin, S. M., Harman, A. M., Moore, S. and Collin, S. P. (2000). Variability in the location of the retinal ganglion cell area centralis is correlated with ontogenetic changes in feeding behavior in the black bream, *Acanthopagrus butcheri* (Sparidae, Teleostei). *Brain Behav. Evol.* **55**, 176–190. doi:10.1159/00006651
- Shand, J., Davies, W. L., Thomas, N., Balmer, L., Cowing, J. A., Pointer, M., Carvalho, L. S., Trezise, A. E. O., Collin, S. P. and Beazley, L. D. (2008). The influence of ontogeny and light environment on the expression of visual pigment opsins in the retina of the black bream, *Acanthopagrus butcheri*. *J. Exp. Biol.* **211**, 1495–1503. doi:10.1242/jeb.012047
- Siebeck, U. E. and Marshall, N. J. (2001). Ocular media transmission of coral reef fish — can coral reef fish see ultraviolet light? *Vision Res.* **41**, 133–149. doi:10.1016/S0042-6989(00)00240-6
- Siebeck, U. E. and Marshall, N. J. (2007). Potential ultraviolet vision in pre-settlement larvae and settled reef fish—A comparison across 23 families. *Vision Res.* **47**, 2337–2352. doi:10.1016/j.visres.2007.05.014
- Slomińska, L. and West, M. J. (2005). Estimators of the precision of stereological estimates: an example based on the CA1 pyramidal cell layer of rats. *Neuroscience* **136**, 757–767. doi:10.1016/j.neuroscience.2005.06.086
- Smith, R. C. and Baker, K. S. (1981). Optical properties of the clearest natural waters (200–800 nm). *Appl. Opt.* **20**, 177–184. doi:10.1364/AO.20.000177
- Sorenson, L., Santini, F., Carnevale, G. and Alfaro, M. E. (2013). A multi-locus timetree of surgeonfishes (Acanthuridae, Percomorpha), with revised family taxonomy. *Mol. Phylogenet. Evol.* **68**, 150–160. doi:10.1016/j.ympev.2013.03.014

- Spady, T. C., Parry, J. W. L., Robinson, P. R., Hunt, D. M., Bowmaker, J. K. and Carleton, K. L.** (2006). Evolution of the cichlid visual palette through ontogenetic subfunctionalization of the opsin gene arrays. *Mol. Biol. Evol.* **23**, 1538-1547. doi:10.1093/molbev/msl014
- Stieb, S. M., Carleton, K. L., Cortesi, F., Marshall, N. J. and Salzburger, W.** (2016). Depth-dependent plasticity in opsin gene expression varies between damselfish (Pomacentridae) species. *Mol. Ecol.* **25**, 3645-3661. doi:10.1111/mec.13712
- Stieb, S. M., Cortesi, F., Sueess, L., Carleton, K. L., Salzburger, W. and Marshall, N. J.** (2017). Why UV vision and red vision are important for damselfish (Pomacentridae): structural and expression variation in opsin genes. *Mol. Ecol.* **26**, 1323-1342. doi:10.1111/mec.13968
- Stieb, S. M., de Busserolles, F., Carleton, K. L., Cortesi, F., Chung, W. S., Dalton, B. E., Hammond, L. A. and Marshall, N. J.** (2019). A detailed investigation of the visual system and visual ecology of the Barrier Reef anemonefish, *Amphiprion akindynos*. *Sci. Rep.* **9**, 1-4.
- Stone, J. and Johnston, E.** (1981). The topography of primate retina: a study of the human, bushbaby, and new-and old-world monkeys. *J. Comp. Neurol.* **196**, 205-223. doi:10.1002/cne.901960204
- Tettamanti, V., de Busserolles, F., Lecchini, D., Marshall, N. J. and Cortesi, F.** (2019). Data from: Visual system development of the spotted unicornfish, *Naso brevirostris* (Acanthuridae), v3. Dryad Digital Repository. <https://doi.org/10.5061/dryad.k6djh9w37>
- Torres-Dowdall, J., Pierotti, M. E. R., Härer, A., Karagic, N., Woltering, J. M., Henning, F., Elmer, K. R. and Meyerm, A.** (2017). Rapid and parallel adaptive evolution of the visual system of Neotropical Midas cichlid fishes. *Mol. Biol. Evol.* **34**, 2469-2485. doi:10.1093/molbev/msx143
- Ullmann, J. F., Moore, B. A., Temple, S. E., Fernández-Juricic, E. and Collin, S. P.** (2012). The retinal wholemount technique: a window to understanding the brain and behaviour. *Brain Behav. Evol.* **79**, 26-44. doi:10.1159/000332802
- Walls, G. L.** (1942). *The Vertebrate Eye and its Adaptive Radiation*. Bloomfield Hills, Michigan, USA: The Cranbrook Press.
- West, M. J., Slomianka, L. H. J. G. and Gundersen, H. J. G.** (1991). Unbiased stereological estimation of the total number of neurons in the subdivisions of the rat hippocampus using the optical fractionator. *Anat. Rec.* **231**, 482-497. doi:10.1002/ar.1092310411
- Wong, R. O. L. and Hughes, A.** (1987). Developing neuronal populations of the cat retinal ganglion cell layer. *J. Comp. Neurol.* **262**, 473-495. doi:10.1002/cne.902620403