

RESEARCH ARTICLE

Interactions between developmental and adult acclimation have distinct consequences for heat tolerance and heat stress recovery

Quentin Willot^{1,*}, Ben Loos² and John S. Terblanche¹

ABSTRACT

Developmental and adult thermal acclimation can have distinct, even opposite, effects on adult heat resistance in ectotherms. Yet, their relative contribution to heat-hardiness of ectotherms remains unclear despite the broad ecological implications thereof. Furthermore, the deterministic relationship between heat knockdown and recovery from heat stress is poorly understood but significant for establishing causal links between climate variability and population dynamics. Here, using *Drosophila melanogaster* in a full-factorial experimental design, we assessed the heat tolerance of flies in static stress assays, and document how developmental and adult acclimation interact with a distinct pattern to promote survival to heat stress in adults. We show that warmer adult acclimation is the initial factor enhancing survival to constant stressful high temperatures in flies, but also that the interaction between adult and developmental acclimation becomes gradually more important to ensure survival as the stress persists. This provides an important framework revealing the dynamic interplay between these two forms of acclimation that ultimately enhance thermal tolerance as a function of stress duration. Furthermore, by investigating recovery rates post-stress, we also show that the process of heat-hardening and recovery post-heat knockdown are likely to be based on set of (at least partially) divergent mechanisms. This could bear ecological significance as a trade-off may exist between increasing thermal tolerance and maximizing recovery rates post-stress, constraining population responses when exposed to variable and stressful climatic conditions.

KEY WORDS: Insects, Thermal biology, Acclimation, Heat stress, Heat coma, Thermal tolerance

INTRODUCTION

In ectotherms, thermal acclimation has long been recognized to occur at different time scales. Transient increase in heat tolerance or heat resistance can be achieved through a heat-hardening response (i.e. a brief exposure to sublethal temperatures enhancing an individual's ability to cope with a subsequent heat stress; Levins, 1969; Dahlgard et al., 1998). In turn, developmental acclimation (i.e. temperature of embryonic development or rapidly developing life stages prior to sexual maturity) also contributes to the plasticity of individuals' thermal limits, through additive or interactive effects with thermal conditions or acclimation occurring later in life (Crill et al., 1996; Angilletta, 2009; Slotsbo et al., 2016; Beaman

et al., 2016; Kellermann et al., 2017). In consequence, disentangling the relative contribution of short-term acclimation (i.e. heat-hardening) and long-term developmental and adult acclimation responses to the phenotypic plasticity of thermal limits has proven a complex task in ectotherms. Because thermal tolerance correlates strongly with geographic distribution and population abundance or viability in insects (Sørensen et al., 2005a,b; Mitchell et al., 2011; Kellermann et al., 2012; Vorhees et al., 2013; Overgaard et al., 2014; Andersen et al., 2015; Bush et al., 2016; Liu et al., 2020), modeling the contribution of different forms of acclimation to thermal plasticity might prove pivotal in our ability to predict species' responses to climate change (Allen et al., 2016; Sinclair et al., 2016; González-Tokman et al., 2020; Bräschler et al., 2020).

In a variety of insects, and especially *Drosophila melanogaster*, the relationship between thermal acclimation (i.e. the exposure to new thermal conditions), induced phenotypic plasticity (i.e. the response) and the molecular processes responsive to heat stress has been a subject of intense research for several decades (Sørensen et al., 2005a, b; Overgaard et al., 2005; Malmendal et al., 2006; Colinet et al., 2013; MacMillan et al., 2016; Kristensen et al., 2016; Schou et al., 2017; Somero, 2020). These processes include, among many others, the evolutionary conserved heat-shock response that mitigate the effects of proteotoxic stresses (Richter et al., 2010). However, quantifying the nature and relative contributions that each form of acclimation brings to heat tolerance has proven challenging at least partly because the outcomes of these stress assays depend on choice of methodology (Mitchell and Hoffmann, 2010; Terblanche et al., 2011). Metrics of heat tolerance in *D. melanogaster* have historically been recorded using a few major approaches that include static (constant controlled temperatures) or ramping (increasing controlled temperatures) assays, and the specific assay conditions can greatly impact any test outcomes and interpretations thereof. Using acclimation temperatures ranging from 12 to 32°C as well as a variety of ramping rates, a number of studies highlighted shifts in CT_{max} (critical thermal maximum; often described as the temperature that results in loss of muscle coordination or onset of muscle spasms in a heating assay) of approximately 1°C owing to warmer adult acclimation, and up to 3°C for its developmental counterpart (Crill et al., 1996; Sejerikilde et al., 2003; Slotsbo et al., 2016; van Heerwaarden et al., 2016; Kellermann et al., 2017; Schou et al., 2017; Salachan et al., 2019). Consistently, all forms of acclimation also increase the time flies could tolerate milder but constant heat stress in static assays (time to knockdown; Levins, 1969; Castañeda et al., 2015; Salachan et al., 2019). Furthermore, a significant interactive effect between developmental and adult acclimation has been highlighted (Slotsbo et al., 2016; Kellermann et al., 2017), the magnitude and contribution of which to the heat-hardening process (i.e. acquired thermotolerance) remains rather poorly known.

Building on this background, this work primarily aims to further investigate the impact of the interactive effects between the two

¹Center for Invasion Biology, Department of Conservation Ecology & Entomology, Stellenbosch University, Stellenbosch 7602, South Africa. ²Department of Physiological Sciences, Stellenbosch University, Stellenbosch 7602, South Africa.

*Author for correspondence (qwillot@sun.ac.za)

 Q.W., 0000-0001-5955-9456; J.S.T., 0000-0001-9665-9405

major forms of thermal acclimation, and how they dynamically contribute to enhancing *D. melanogaster* heat limits. To do so, we explored the combinations of three developmental and three adult acclimation temperatures on time to heat knockdown (HKD) and recovery post-knockdown of flies in static assays. Interactive effects between developmental and adult acclimation in *D. melanogaster* have been demonstrated through measures of CT_{max} using controlled ramping protocols (Slotsbo et al., 2016; Kellermann et al., 2017). These studies also emphasized a greater impact of developmental acclimation in enhancing the upper thermal limits of flies. We therefore hypothesized that in our static assays, warmer developmental acclimation would be the dominant factor increasing flies' time to knockdown. Second, we predicted that the magnitude of this interaction would be dynamic and follow a temporal pattern, based on the assumption that the two forms of acclimation may leverage sets of partially different molecular-level processes with distinct time scales of dynamic responses (e.g. Telonis-Scott et al., 2014). Finally, we investigated how these two forms of thermal acclimation may potentially drive faster recoveries post heat knockdown, as this metric has seldom been reported before, but might prove important to survive transient stress exposure under more variable microclimatic conditions (Ma et al., 2021).

MATERIALS AND METHODS

Fly stocks, rearing, and acclimation

Wild *Drosophila* flies were bait-trapped in April 2019 around the Stellenbosch University campus, Western Cape, South Africa. Single female flies were isolated into 250 ml rearing flasks supplied with 50 ml of Bloomington cornmeal diet (Lewis, 1960) and left to lay eggs for 1 day. Between two and five of the first emerging adults were used for species determination following the key of Markow and Grady (2005). The remaining first-generation individuals from 16 confirmed *D. melanogaster* Meigen 1830 isofemale lines were pooled and mass-reared together for five generations prior to experiments. An overview of the rearing and acclimation procedures used in experiments is given in Fig. S1. Stocks were kept in separated incubators at either 15, 25 or 30°C in the form of three 250 ml flasks supplied with Bloomington cornmeal diet, at a density of ~100 adult flies per flask. Before the emergence of a new

generation (~7 days at 30°C, ~10 days at 25°C, and ~21 days at 15°C), older individuals were discarded. Less than 24 h after their emergence, ~100 newly hatched flies were isolated in a fresh flask to replenish stocks, while ~50 others were isolated for experimental assays. These experimental batches were further left to mature for 10 additional days at 15, 25 or 30°C prior to experiments, and their offspring were discarded. This effectively created nine unique thermal history conditions, in which batches of adult flies emerged within 24 h under developmental acclimation conditions (15, 25 or 30°C; adults therefore experienced developmental acclimation conditions for a maximum of 1 day or ≤10% of their adult life) and were subsequently exposed to either one of the three adult acclimation temperatures (15, 25 or 30°C) until they reached the age of 10 days old.

Heat knockdown dynamic

For each of our nine unique thermal history conditions, between seven and 12 replicates each containing 15–25 flies, with a 1:1 sex ratio, were isolated into glass vials with a moist cotton ball. The vials were then submerged in a digital water bath (GD120 series stirred water bath, Grant Instrument Ltd) and the temperature inside the vials was monitored using 0.075 mm diameter thermocouples [Type T Thermocouple (Copper/Constantan), OMEGA Engineering, CT, USA] connected to a digital thermometer (Fluke 52-II Dual Input Digital Thermometer, WA, USA). Flies were exposed to a constant 37°C, and the proportion of heat knocked-down flies in each batch was monitored every 15 min. HKD was considered positive for an individual fly when it adopted an immobile curled up position on its back, without any further response to external stimulation (vial shaking). Downstream analysis was performed by fitting a logistic regression model on each replicate of our nine thermal history conditions, through a best-fit model approach. This allowed us to extract five percentiles per thermal history condition, based on 7–12 replicated curves per thermal history, corresponding to the time needed on average to reach 10% (lethal time; LT10), 25% (LT25), 50% (LT50), 75% (LT75) and 90% (LT90) of heat knocked-down flies. Finally, we further tested for statistical significance of treatments on LT values, and quantified the relationship between developmental acclimation, adult acclimation and extracted LT values using an effect size ω^2 statistic (Olejnik and Algina, 2003).

Table 1. Analysis of the impact of development and adult acclimation on time for *Drosophila melanogaster* flies to reach a set proportion of knocked-down individuals

Lethal time (%)	Fixed effects	Effect size	Effect size 90% CI	P-value
LT10	Adult acclimation	0.66	0.56–0.73	0.032*
	Developmental acclimation	0.07	0.01–0.18	0.841
	Developmental×adult acclimation	6.95 ^{−04}	0.00–0.00	0.331
LT25	Adult acclimation	0.76	0.68–0.81	0.001**
	Developmental acclimation	0.12	0.04–0.26	0.551
	Developmental×adult acclimation	0.03	0.00–0.12	0.763
LT50	Adult acclimation	0.79	0.72–0.83	0.004**
	Developmental acclimation	0.17	0.06–0.29	0.485
	Developmental×adult acclimation	0.07	0.01–0.18	0.065
LT75	Adult acclimation	0.77	0.70–0.82	0.035*
	Developmental acclimation	0.17	0.06–0.29	0.117
	Developmental×adult acclimation	0.09	0.02–0.21	0.006**
LT90	Adult acclimation	0.73	0.65–0.79	0.132
	Developmental acclimation	0.15	0.02–0.21	0.036*
	Developmental×adult acclimation	0.10	0.02–0.21	0.001**

Overall, adult acclimation accounted for the majority of variation in lethal times (LTs) across thermal histories but became less significant later stage of the experiment (LT90). As opposed, developmental acclimation accounted for an initial minority of variation in LTs, but its significance increased with knockdown proportion. Finally, both the significance and size effect of the interaction between developmental and adult acclimation increased proportionally with stress duration. * $P < 0.05$; ** $P < 0.01$.

Recovery experiments

Single flies were isolated into glass vials containing a moist cotton ball. The vials were then submerged in a programmable circulating water bath and flies were continuously exposed to a constant 41°C and until HKD, as described above in the heat knockdown assay (i.e. each adult fly was exposed to 41°C for a specific amount of time until HKD, resulting in a different exposure time for each fly). Each heat knocked-down fly was immediately placed back at room temperature (21°C) and continuously monitored until recovery, defined as the time at which it could stand on its legs again, without stumbling over from external stimulation (vial shaking). Both times to HKD and recovery were carefully recorded for 30 flies coming from three different rearing bottle replicates (10 flies taken from each bottle), tested with a 1:1 sex ratio per thermal history condition (nine conditions; 270 individuals in total). Mortality was typically

low in these assays, amounting to two flies (99.23% survival rate), and thus not considered further in subsequent analyses. Kaplan–Meier knockdown curves were drawn from the raw observed data, and the impact of thermal histories on time to HKD and time to recovery was assessed using two complementary analyses. First, a log-rank analysis was performed to test for significant differences between knockdown curves. Second, we used a Cox proportional hazards model to test for significant effects of our treatments on HKD curves (developmental acclimation, adult acclimation and the interaction between the two).

Relationship between time to heat knockdown and time to recovery

Each fly used in HKD experiments at 41°C also had a linked recovery time. These times were plotted against each other and




Fig. 1. Best-fit model of heat knockdown (HKD) curves for *Drosophila melanogaster* flies exposed to a constant 37°C, as a function of their thermal histories. HKD curves are presented for flies under the nine conditions of our full-factorial analysis, sorted by developmental/adult acclimation temperatures: (A) 15/15°C; (B) 15/25°C; (C) 15/30°C; (D) 25/15°C; (E) 25/25°C; (F) 25/30°C; (G) 30/15°C; (H) 30/25°C; and (I) 30/30°C. Adult acclimation for a fixed developmental acclimation temperature thus increases from left to right. Developmental acclimation for a fixed adult acclimation thus increases from top to bottom ($N=172\pm32$ flies over 7–12 replicates per curve). The central red dots allow for quicker visual comparison of curve shifts as result of treatments. Increased adult acclimation temperatures (from left to right) consistently increased time for flies to enter knockdown and shifted the curves to the right regardless of developmental acclimation. Increased developmental acclimation temperatures (from top to bottom) had a more marginal and less consistent effect on curve shapes. Statistical analysis is presented in Table 1.

screened for significant ordinary least-square linear regressions both as a function of their thermal history and on the global dataset. In addition, because Kaplan–Meier curves do not integrate slope data, we refitted polynomial curves on time to HKD and recovery at 41°C and extracted slope values. These slopes were plotted against time to recovery slopes and also screened for significant linear regressions.

Statistical analysis

Log-rank tests of Kaplan–Meier curves were performed using GraphPad Prism 6.01. Cox proportional hazards and best-fit models as well as effect size statistics were performed using R v.3.6.3 (<https://www.r-project.org/>; <https://CRAN.R-project.org/package=survival> and <https://CRAN.R-project.org/package=effsize> libraries).

RESULTS

Adult and developmental acclimation contribute in unique ways to heat stress dynamics

Overall, warmer adult acclimation had a strong and consistent effect in delaying time to HKD in flies (Table 1) at 37°C, shifting HKD curves to the right (Fig. 1, from left to right). Adult acclimation accounted for most of the variation across all LTs, but its effect size and thus also statistical significance decreased at later stages of the experiment (Table 1). Warmer developmental acclimation (Fig. 1,

from top to bottom) had an overall weaker effect in delaying time to HKD. However, as opposed to adult acclimation, its statistical significance and effect size contribution increased as a function of time under heat stress, i.e. as knockdown progressed (Table 1). Finally, although we detected no initial effect of the interaction between developmental and adult acclimation in delaying time to HKD in flies, its statistical significance and effect size contribution increased proportionally with the duration of the stress (Table 1). To summarize, the two forms of acclimation did not contribute equally to variation in the shape of the HKD curve. Overall, adult acclimation was more significant in delaying the onset of HKD at the beginning of the stress assay, while developmental acclimation and the interactive effect of developmental and adult acclimation became gradually more significant at later stages.

Impact of developmental and adult acclimation on recovery dynamics

To further test how developmental and adult acclimations may contribute differently to heat tolerance and especially HKD recoveries, we performed static knockdown and recovery assays by exposing single *D. melanogaster* flies to a constant 41°C. Time to HKD was carefully monitored for each individual, and each knocked-down fly was then immediately removed from the experimental setup and placed back at room temperature (21°C)




Fig. 2. Heat knockdown curves of *D. melanogaster* flies as a function of their thermal histories. Flies were exposed to a constant 41°C until knockdown, and curves are provided with fixed developmental (A–C) or adult (D–F) acclimation temperatures ($N=30$ per condition). Time to HKD was measured for 30 individual flies per thermal history condition, and data was pooled to form a single HKD curve per thermal history. The central red dots allow for quicker visual comparison of curve shifts as result of treatments. Warmer adult acclimation consistently increased times to HKD (A–C). Warmer developmental acclimation also had a significant but less consistent effect on times to HKD (D–F). Different lowercase letters indicate a significant difference between treatments (log-rank test, $P<0.01$).

until recovery. Consistent with data from HKD dynamics at 37°C (Table 1, Fig. 1), a log-rank analysis of HKD curves at 41°C confirmed that warmer adult acclimation was the dominant effect delaying the onset of knockdown (Fig. 2A–C), with developmental acclimation having a milder impact (Fig. 2C–E). Concerning recoveries from HKD, we did not detect adult acclimation to have an impact (Fig. 3A–C), whereas colder developmental acclimation consistently decreased time to recovery (Fig. 3D–F). A second layer of analysis using a Cox proportional hazards model confirmed these observations. Both adult ($P<0.001$) and developmental ($P<0.01$) acclimation had a strong effect on flies' time to HKD at 41°C, with adult acclimation being the far greater effect of the two. Adult acclimation did not affect recovery times, whereas developmental acclimation had a strong significant effect on time to recovery ($P<0.001$).

Relationship between heat-hardiness and recovery rates

Each *D. melanogaster* used in static assays at 41°C had a single time to HKD linked to its time to recovery. These values were plotted against each other for each individual *D. melanogaster*, and significant regressions were screened both as a function of flies' thermal history (Fig. 4A) and on the global dataset (Fig. 4B). No significant positive or negative correlations were found when considering the thermal history of individuals, with the exception of a weak but significant

linear relationship in flies reared at 25°C and adults acclimated at 30°C (Table S1). A negligible but statistically significant linear correlation between time to HKD and recovery was also present at the level of the global dataset ($R^2=0.07$, $P<0.001$; Fig. 3B, Table S1). Finally, to test for correlations between time to HKD and time to recovery, we fitted polynomial curves on time to HKD (Fig. S3A) and time to recovery (Fig. S3B) data and extracted curve slopes as a function of thermal histories of individuals (Table S2). No significant correlation between rates to HKD slopes and rates to recovery slopes were found on the global dataset (Fig. S4A). In line with analysis from Kaplan–Meier curves (Fig. 3), recovery rates were comparatively higher for colder developmentally acclimated flies than for warmer ones (Fig. S4B).

DISCUSSION

Our results first show that warmer adult acclimation was the dominant factor delaying the onset of heat knockdown in static assays, for flies exposed to 37°C (Figs 1 and 2, Table 1). We measured the effect size of adult acclimation to be at least several fold greater than that of developmental acclimation at lower knockdown percentages (Table 1). This was somewhat unexpected, as it contrasts with both our predictions and results of previous studies (Slotsbo et al., 2016; Kellermann et al., 2017; Schou et al., 2017), which showed that developmental acclimation was found to increase the CT_{max} of flies to a greater extent than adult




Fig. 3. Recovery curves of *D. melanogaster* flies as a function of their thermal histories. Flies were exposed to 41°C until knockdown and immediately put back at 21°C for recovery, as a function of their thermal histories, with either fixed developmental (A–C) or adult (D–F) acclimation temperatures ($N=30$ per condition). Time to recovery was measured for 30 individual flies per thermal history conditions, and data was pooled to extract a single HKD curve per thermal history. The central red dots allow for quicker visual comparison of curve shifts as result of treatments. No effect of adult acclimation was found on recovery rates (A–C), whereas colder developmental temperatures significantly sped up recoveries (D–F). Different lowercase letters indicate a significant difference between treatments (log-rank test, $P<0.01$).




Fig. 4. Time to recovery as a function of time to HKD in individual *D. melanogaster* flies. (A) Linear regression for individuals pooled as a function of their thermal histories ($N=30$ per thermal history conditions). No significant correlation was observed, with exception of a weak linear relationship in flies reared at 25°C and adults acclimated at 30°C (Table S1). (B) Regression on the global dataset ($N=270$). A weak but significant linear correlation was present ($R^2=0.07$, $P<0.001$; Table S1), indicating that time to heat knockdown was overall a poor predictor of time to recovery.

acclimation did. Such discrepancies in results could first be the outcome of methodological artefacts. Indeed, owing to their longer duration, milder static assays perhaps can offer a better temporal resolution of a heat-hardening process than a ramping one (Salachan et al., 2019). However, reaching time to heat knockdown in milder conditions could incur other physiological costs or induce other processes (such as dehydration or starvation that could lead to a bias in heat-tolerance metrics; see Terblanche et al., 2011; Manenti et al., 2018; but also see Overgaard et al., 2012 and Mitchell et al., 2017 for evidence to the contrary). The inherent methodological trade-offs between static and ramping assays have long been argued to induce diverse molecular stress responses and therefore perhaps constitute distinct forms of genetic variation, with different dynamics, in exposed animals (Cooper et al., 2008; Mitchell and Hoffmann, 2010; Sgrò et al., 2010; Santos et al., 2011; Terblanche et al., 2011; Rezende et al., 2014). Thus, stress duration, intensity and potential ramping rates are critical parameters to interpret heat-tolerance metrics in a biologically relevant context (Kovacevic et al., 2019; Kingsolver and Umbanhowar, 2018; Ma et al., 2021; Jørgensen et al., 2021).

From this perspective, our results also highlight that warmer developmental acclimation gained secondary importance in increasing flies' heat tolerance as the stress persisted (Table 1). In addition, other works have shown warmer developmental acclimation to be dominant to increase CT_{max} of flies in ramping assays (Slotsbo et al., 2016; Kellermann et al., 2017; Schou et al., 2017). Taking these into account, it could also be that developmental and adult acclimation interact through a two-gear system: the first would uplift the potential level of expression of stress-tolerance genes, and the second would set the expression within the range allowed by the first. This would produce a dynamic response as observed: the potential increased expression of tolerance genes through warmer developmental acclimation would prove critical to survive enduring, or indeed increasing, stress exposure. By contrast, tolerance mechanisms already elevated to higher levels through warmer adult acclimation would explain the delays in the onset of heat knockdown in flies as we observed under milder, static conditions. Such a model may find support in our current understanding of the molecular-level processes related to heat stress acclimation and could be tested directly in future. Indeed, developmental acclimation has been shown to reorganize chromatin

structure and modulate the accessibility of stress responsive genes to the transcription machinery (Farkas et al., 2000; Feil and Fraga, 2012; Vihervaara et al., 2018), and thus potentially their levels and/or rates of expression. Given that different methodological approaches can mask or reveal thermal acclimation effects (Terblanche and Hoffmann, 2020), such a hypothesis remains merely speculative at this point and awaits additional scrutiny. Thermal landscape assays have more recently been proposed as a unifying methodology between static and dynamic assays (Castañeda et al., 2015; Jørgensen et al., 2019) and in this context, testing the impact of developmental/adult acclimation under similar full-factorial designs but through a thermal landscape approach would perhaps be useful to further detail these findings and our interpretations.

Finally, and quite surprisingly, we did not find that adult acclimation contributed to speeding up recovery post-heat knockdown in any marked way (Fig. 3A–C). In addition, and in contrast to our predictions, colder developmental acclimation, not warmer, drove faster heat stress recoveries (Fig. 3D–F). The heat-hardiness of flies, that is the time needed for a given fly to enter HKD at 41°C, was a poor predictor of time to recovery (Fig. 4, Figs S3 and S4, Table S1). This refutes the intuitive assumption that times to recovery from heat stress should be related to the duration of stress exposure (time needed to reach HKD). Overall, this provides indirect evidence that the mechanisms underlying heat-hardening and recovery from heat knockdown must be at least partially decoupled. Furthermore, this set of mechanisms related to recovery from heat stress seems to be leveraged specifically by colder developmental acclimation alone. Extending our interpretation by inference, increased thermal tolerance from warmer developmental acclimation may thus come at the cost of decreased recovery capabilities. Such a trade-off between heat knockdown and heat recovery may also prove to be a critical link in understanding trade-offs between basal and plastic stress resistance, an area in urgent need of further investigation (van Heerwarden and Kellermann, 2020). However, caution also needs to be exercised when inferring potential biological trade-offs, as the interactive acclimation effects have been shown to sometimes remain population-specific (Kellermann et al., 2017), and sometimes display non-linear reaction norms to temperature (Salachan et al., 2019; Sørensen et al., 2020). Thus, it will be pivotal to further document the

presence of a dynamic interaction between developmental and adult acclimation as well as a potential trade-off between increased thermal tolerance and recovery capabilities through different methodological approaches in various *D. melanogaster* strains, and across *Drosophila* species, to better understand the generality of these outcomes.

Conclusions

In summary, using a full-factorial experimental design and analysis of the effect of developmental and adult acclimation on heat knockdown resistance and knockdown recovery in *D. melanogaster*, we have shown that these two forms of acclimation occurring at different stages of development contribute in distinct ways to the dynamics of heat tolerance. Warmer adult acclimation strongly delays the onset of heat knockdown in flies exposed to constant stressful temperatures. By contrast, the effect of developmental acclimation, and the interaction between the two forms of plasticity, gradually gained importance as a function of the stress duration. Finally, we also show that, as opposed to developmental acclimation, adult acclimation had no detectable impact on the rates of recovery post heat knockdown. Thus, the mechanisms underlying the heat-hardening response (that increases the flies' initial heat tolerance), and the mechanisms underlying recovery (once knockdown has occurred and heat stress has been lifted) are likely to be at least partly distinct and could react to acclimation temperature in opposite in different ways. These results therefore provide an important framework to understand the temporal interaction of developmental and adult acclimation to promote stress tolerance in insects. They also underpin a potential ecologically and evolutionarily significant acclimation trade-off between increasing thermal tolerance and maximizing recovery rates post-stress, which could constrain the response of populations exposed to more variable and stressful microclimatic conditions.

Acknowledgement

We thank Erika Huisamen for help with fly husbandry, Henriek Bosua and Minette Karsten for their help with the experimental setup, and the anonymous referees for constructive comments on an earlier version of this paper.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: Q.W., B.L., J.S.T.; Methodology: Q.W., J.S.T.; Validation: Q.W.; Formal analysis: Q.W.; Investigation: Q.W.; Resources: B.L., J.S.T.; Writing - original draft: Q.W.; Writing - review & editing: Q.W., J.S.T.; Visualization: Q.W.; Supervision: J.S.T.; Project administration: J.S.T.; Funding acquisition: J.S.T., Q.W.

Funding

This work was supported by a Claude Leon Foundation post-doctoral fellowship (to Q.W.) and a running cost funding (to J.S.T.) from the DSI-NRF Center of Excellence for Invasion Biology (CIB).

References

- Allen, J. L., Chown, S. L., Janion-Scheepers, C. and Clusella-Trullas, S. (2016). Interactions between rates of temperature change and acclimation affect latitudinal patterns of warming tolerance. *Conserv. Physiol.* **4**, cow053. doi:10.1093/conphys/cow053
- Andersen, J. L., Manenti, T., Sørensen, J. G., MacMillan, H. A., Loeschcke, V. and Overgaard, J. (2015). How to assess *Drosophila* cold tolerance: chill coma temperature and lower lethal temperature are the best predictors of cold distribution limits. *Funct. Ecol.* **29**, 55–65. doi:10.1111/1365-2435.12310
- Angilletta, M., Jr (2009). Looking for answers to questions about heat stress: researchers are getting warmer. *Funct. Ecol.* **23**, 231–232. doi:10.1111/j.1365-2435.2009.01548.x
- Beaman, J. E., White, C. R. and Seebacher, F. (2016). Evolution of plasticity: mechanistic link between development and reversible acclimation. *Trends Ecol. Evol.* **31**, 237–249. doi:10.1016/j.tree.2016.01.004
- Braschler, B., Duffy, G. A., Nortje, E., Kritzinger-Klopper, S., du Plessis, D., Karenyi, N., Leihy, R. I. and Chown, S. L. (2020). Realised rather than fundamental thermal niches predict site occupancy: implications for climate change forecasting. *J. Anim. Ecol.* **89**, 2863–2875. doi:10.1111/1365-2656.13358
- Bush, A., Mokany, K., Catullo, R., Hoffmann, A., Kellermann, V., Sgrò, C., McEvey, S. and Ferrier, S. (2016). Incorporating evolutionary adaptation in species distribution modeling reduces projected vulnerability to climate change. *Ecol. Lett.* **19**, 1468–1478. doi:10.1111/ele.12696
- Castañeda, L. E., Rezende, E. L. and Santos, M. (2015). Heat tolerance in *Drosophila subobscura* along a latitudinal gradient: contrasting patterns between plastic and genetic responses. *Evolution* **69**, 2721–2734. doi:10.1111/evo.12757
- Colinet, H., Overgaard, J., Com, E. and Sørensen, J. G. (2013). Proteomic profiling of thermal acclimation in *Drosophila melanogaster*. *Insect Biochem. Mol. Biol.* **43**, 352–365. doi:10.1016/j.ibmb.2013.01.006
- Cooper, B. S., Williams, B. H. and Angilletta, M. J. (2008). Unifying indices of heat tolerance in ectotherms. *J. Therm. Biol.* **33**, 320–323. doi:10.1016/j.jtherbio.2008.04.001
- Crill, W. D., Huey, R. B. and Gilchrist, G. W. (1996). Within- and between-generation effects of temperature on the morphology and physiology of *Drosophila melanogaster*. *Evolution* **50**, 1205–1218. doi:10.2307/2410661
- Dahlgard, J., Loeschcke, V., Michalak, P. and Justesen, J. (1998). Induced thermotolerance and associated expression of the heat-shock protein Hsp70 in adult *Drosophila melanogaster*. *Funct. Ecol.* **12**, 786–793. doi:10.1046/j.1365-2435.1998.00246.x
- Farkas, G., Leibovitch, B. A. and Elgin, S. C. (2000). Chromatin organization and transcriptional control of gene expression in *Drosophila*. *Gene* **253**, 117–136. doi:10.1016/S0378-1119(00)00240-7
- Feil, R. and Fraga, M. F. (2012). Epigenetics and the environment: emerging patterns and implications. *Nat. Rev. Genet.* **13**, 97–109. doi:10.1038/nrg3142
- González-Tokman, D., Córdoba-Aguilar, A., Dáttilo, W., Lira-Noriega, A., Sánchez-Guillén, R. A. and Villalobos, F. (2020). Insect responses to heat: physiological mechanisms, evolution and ecological implications in a warming world. *Biol. Rev.* **95**, 802–821. doi:10.1111/brv.12588
- Jørgensen, L. B., Malte, H. and Overgaard, J. (2019). How to assess *Drosophila* heat tolerance: unifying static and dynamic tolerance assays to predict heat distribution limits. *Funct. Ecol.* **33**, 629–642. doi:10.1111/1365-2435.13279
- Jørgensen, L. B., Malte, H., Ørsted, M., Klahn, N. A. and Overgaard, J. (2021). A unifying model to estimate thermal tolerance limits in ectotherms across static, dynamic and fluctuating exposures to thermal stress. *Sci. Rep.* **11**, 1–14. doi:10.1038/s41598-021-92004-6
- Kellermann, V., Overgaard, J., Hoffmann, A. A., Flojgaard, C., Svenning, J.-C. and Loeschcke, V. (2012). Upper thermal limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. *Proc. Natl. Acad. Sci. USA* **109**, 16228–16233. doi:10.1073/pnas.1207553109
- Kellermann, V., van Heerwaarden, B. and Sgrò, C. M. (2017). How important is thermal history? Evidence for lasting effects of developmental temperature on upper thermal limits in *Drosophila melanogaster*. *Proc. R. Soc. B* **284**, 20170447. doi:10.1098/rspb.2017.0447
- Kingsolver, J. G. and Umbanhowar, J. (2018). The analysis and interpretation of critical temperatures. *J. Exp. Biol.* **221**, jeb167858. doi:10.1242/jeb.167858
- Kovacevic, A., Latombe, G. and Chown, S. L. (2019). Rate dynamics of ectotherm responses to thermal stress. *Proc. R. Soc. B* **286**, 20190174. doi:10.1098/rspb.2019.0174
- Kristensen, T. N., Kjeldal, H., Schou, M. F. and Nielsen, J. L. (2016). Proteomic data reveal a physiological basis for costs and benefits associated with thermal acclimation. *J. Exp. Biol.* **219**, 969–976. doi:10.1242/jeb.132696
- Levins, R. (1969). Thermal acclimation and heat resistance in *Drosophila* species. *Am. Nat.* **103**, 483–499. doi:10.1086/282616
- Lewis, E. B. (1960). A new standard food medium. *Drosoph. Inf. Serv.* **34**, 1–55.
- Liu, W. A., Phillips, L. M., Terblanche, J. S., Janion-Scheepers, C. and Chown, S. L. (2020). Strangers in a strange land: globally unusual thermal tolerance in *Collembola* from the Cape floristic region. *Funct. Ecol.* **34**, 1601–1612. doi:10.1111/1365-2435.13584
- Ma, C.-S., Ma, G. and Pincebourde, S. (2021). Survive a warming climate: insect responses to extreme high temperatures. *Annu. Rev. Entomol.* **66**, 163–184. doi:10.1146/annurev-ento-041520-074454
- MacMillan, H. A., Knee, J. M., Dennis, A. B., Udaka, H., Marshall, K. E., Merritt, T. J. and Sinclair, B. J. (2016). Cold acclimation wholly reorganizes the *Drosophila melanogaster* transcriptome and metabolome. *Sci. Rep.* **6**, 1–14. doi:10.1038/srep28999
- Malmendal, A., Overgaard, J., Bundy, J. G., Sørensen, J. G., Nielsen, N. C., Loeschcke, V. and Holmstrup, M. (2006). Metabolomic profiling of heat stress: hardening and recovery of homeostasis in *Drosophila*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **291**, 205–212. doi:10.1152/ajpregu.00867.2005
- Manenti, T., Cunha, T. R., Sørensen, J. G. and Loeschcke, V. (2018). How much starvation, desiccation and oxygen depletion can *Drosophila melanogaster* tolerate before its upper thermal limits are affected? *J. Insect Physiol.* **111**, 1–7. doi:10.1016/j.jinsphys.2018.09.002
- Markow, T. A. and O'Grady, P. (2005). *Drosophila: a Guide to Species Identification and Use*. Elsevier.

- Mitchell, K. A. and Hoffmann, A. A. (2010). Thermal ramping rate influences evolutionary potential and species differences for upper thermal limits in *Drosophila*. *Funct. Ecol.* **24**, 694–700. doi:10.1111/j.1365-2435.2009.01666.x
- Mitchell, K. A., Sgrò, C. M. and Hoffmann, A. A. (2011). Phenotypic plasticity in upper thermal limits is weakly related to *Drosophila* species distributions. *Funct. Ecol.* **25**, 661–670. doi:10.1111/j.1365-2435.2010.01821.x
- Mitchell, K. A., Boardman, L., Clusella-Trullas, S. and Terblanche, J. S. (2017). Effects of nutrient and water restriction on thermal tolerance: a test of mechanisms and hypotheses. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **212**, 15–23. doi:10.1016/j.cbpa.2017.06.019
- Olejnik, S. and Algina, J. (2003). Generalized eta and omega squared statistics: measures of effect size for some common research designs. *Psychol. Methods* **8**, 434. doi:10.1037/1082-989X.8.4.434
- Overgaard, J., Sørensen, J. G., Petersen, S. O., Loeschcke, V. and Holmstrup, M. (2005). Changes in membrane lipid composition following rapid cold hardening in *Drosophila melanogaster*. *J. Insect Physiol.* **51**, 1173–1182. doi:10.1016/j.jinsphys.2005.06.007
- Overgaard, J., Kristensen, T. N. and Sørensen, J. G. (2012). Validity of thermal ramping assays used to assess thermal tolerance in arthropods. *PLoS ONE* **7**, e32758. doi:10.1371/journal.pone.0032758
- Overgaard, J., Kearney, M. R. and Hoffmann, A. A. (2014). Sensitivity to thermal extremes in Australian *Drosophila* implies similar impacts of climate change on the distribution of widespread and tropical species. *Glob. Chang. Biol.* **20**, 1738–1750. doi:10.1111/gcb.12521
- Rezende, E. L., Castañeda, L. E. and Santos, M. (2014). Tolerance landscapes in thermal ecology. *Funct. Ecol.* **28**, 799–809. doi:10.1111/1365-2435.12268
- Richter, K., Haslbeck, M. and Buchner, J. (2010). The heat shock response: life on the verge of death. *Mol. Cell.* **40**, 253–266. doi:10.1016/j.molcel.2010.10.006
- Salachan, P. V., Burgaud, H. and Sørensen, J. G. (2019). Testing the thermal limits: non-linear reaction norms drive disparate thermal acclimation responses in *Drosophila melanogaster*. *J. Insect Physiol.* **118**, 103946. doi:10.1016/j.jinsphys.2019.103946
- Santos, M., Castañeda, L. E. and Rezende, E. L. (2011). Making sense of heat tolerance estimates in ectotherms: lessons from *Drosophila*. *Funct. Ecol.* **25**, 1169–1180. doi:10.1111/j.1365-2435.2011.01908.x
- Schou, M. F., Kristensen, T. N., Pedersen, A., Karlsson, B. G., Loeschcke, V. and Malmendal, A. (2017). Metabolic and functional characterization of effects of developmental temperature in *Drosophila melanogaster*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **312**, 211–222. doi:10.1152/ajpregu.00268.2016
- Sejerkilde, M., Sørensen, J. G. and Loeschcke, V. (2003). Effects of cold- and heat hardening on thermal resistance in *Drosophila melanogaster*. *J. Insect Physiol.* **49**, 719–726. doi:10.1016/S0022-1910(03)00095-7
- Sgrò, C. M., Overgaard, J., Kristensen, T. N., Mitchell, K. A., Cockerell, F. E. and Hoffmann, A. A. (2010). A comprehensive assessment of geographic variation in heat tolerance and hardening capacity in populations of *Drosophila melanogaster* from eastern Australia. *J. Evol. Biol.* **23**, 2484–2493. doi:10.1111/j.1420-9101.2010.02110.x
- Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., Dong, Y., Harley, C. D., Marshall, D. J., Helmut, B. S. et al. (2016). Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecol. Lett.* **19**, 1372–1385. doi:10.1111/ele.12686
- Slotsbo, S., Schou, M. F., Kristensen, T. N., Loeschcke, V., Sørensen, J. G. (2016). Reversibility of developmental heat and cold plasticity is asymmetric and has long-lasting consequences for adult thermal tolerance. *J. Exp. Biol.* **219**, 2726–2732. doi:10.1242/jeb.143750
- Somero, G. N. (2020). The cellular stress response and temperature: function, regulation, and evolution. *J. Exp. Zool. A Ecol. Integr. Physiol.* **333**, 379–397. doi:10.1002/jez.2344
- Sørensen, J. G., Norry, F. M., Scannapieco, A. C. and Loeschcke, V. (2005a). Altitudinal variation for stress resistance traits and thermal adaptation in adult *Drosophila buzzatii* from the New World. *J. Evol. Biol.* **18**, 829–837. doi:10.1111/j.1420-9101.2004.00876.x
- Sørensen, J. G., Nielsen, M. M., Kruhøffer, M., Justesen, J. and Loeschcke, V. (2005b). Full genome gene expression analysis of the heat stress response in *Drosophila melanogaster*. *Cell Stress Chaperones* **10**, 312. doi:10.1379/CSC-128R1.1
- Sørensen, J. G., Winther, M. L., Salachan, P. V. and MacLean, H. J. (2020). Drawing the line: Linear or non-linear reaction norms in response to adult acclimation on lower thermal limits. *J. Insect Physiol.* **124**, 104075. doi:10.1016/j.jinsphys.2020.104075
- Telonis-Scott, M., Clemson, A. S., Johnson, T. K. and Sgrò, C. M. (2014). Spatial analysis of gene regulation reveals new insights into the molecular basis of upper thermal limits. *Mol. Ecol.* **23**, 6135–6151. doi:10.1111/mec.13000
- Terblanche, J. S., Hoffmann, A. A., Mitchell, K. A., Rako, L., le Roux, P. C. and Chown, S. L. (2011). Ecologically relevant measures of tolerance to potentially lethal temperatures. *J. Exp. Biol.* **214**, 3713–3725. doi:10.1242/jeb.061283
- Terblanche, J. S. and Hoffmann, A. A. (2020). Validating measurements of acclimation for climate change adaptation. *Curr. Opin. Insect. Sci.* **41**, 7–16. doi:10.1016/j.cois.2020.04.005
- van Heerwaarden, B. and Kellermann, V. (2020). Does plasticity trade off with basal heat tolerance? *Trends Ecol. Evol.* **35**, 874–885. doi:10.1016/j.tree.2020.05.006
- van Heerwaarden, B., Kellermann, V. and Sgrò, C. M. (2016). Limited scope for plasticity to increase upper thermal limits. *Funct. Ecol.* **30**, 1947–1956. doi:10.1111/1365-2435.12687
- Vihervaara, A., Duarte, F. M. and Lis, J. T. (2018). Molecular mechanisms driving transcriptional stress responses. *Nat. Rev. Genet.* **19**, 385. doi:10.1038/s41576-018-0001-6
- Vorhees, A. S., Gray, E. M. and Bradley, T. J. (2013). Thermal resistance and performance correlate with climate in populations of a widespread mosquito. *Physiol. Biochem. Zool.* **86**, 73–81. doi:10.1086/668851