

**The ventilatory response to hypoxia and hypercapnia is absent in the neonatal
fat-tailed dunnart**

Shannon J Simpson¹, Angelina Y Fong², Kevin J Cummings³, Peter B Frappell¹

¹Adaptational and Evolutionary Respiratory Physiology Laboratory, University of Tasmania, Hobart, Australia 7005; ²Australian School of Advanced Medicine, Macquarie University, North Ryde, NSW; ³Department of Biomedical Sciences, University of Missouri-Columbia, Columbia, MO, USA

Short title: Chemoresponse in the neonatal dunnart

Corresponding author: Peter Frappell

Private Bag 3 Hobart, Tasmania, Australia 7001

peter.frappell@utas.edu.au

Symbols and abbreviations:

CO₂, carbon dioxide; °C, degrees Celsius; f, frequency of breathing; mg, milligram; min, minute; mL, millilitre; O₂, oxygen; P, days postpartum; S.E.M, standard error of the mean; TE, expiratory time; TI, inspiratory time; TI/TTOT, duty cycle; TTOT, total breath time; µl, microliter; \dot{V}_E , minute ventilation; \dot{V}_E/\dot{V}_{O_2} , convective requirement; \dot{V}_{O_2} , rate of oxygen consumption; VT, tidal volume; VT/TI, mean inspiratory flow

Abstract: (171 words)

At birth, the newborn fat-tailed dunnart relies on cutaneous gas exchange to meet metabolic demands, with continuous lung ventilation emerging several days later. We hypothesized that the delayed expression of lung ventilation (\dot{V}_E) in these animals is in part owing to a low responsiveness of the respiratory control system to blood gas perturbations. To address this hypothesis we assessed the ventilatory and metabolic response to hypoxia (10% O₂) and hypercapnia (5% CO₂) using closed-system respirometry from birth to 23 days postpartum (P). Neonatal fat-tailed dunnarts displayed no significant hypoxic or hypercapnic ventilatory responses at any age. Regardless, significant hyperventilation through a suppression of metabolic rate (\dot{V}_{O_2}) was observed at birth in response to hypercapnia and in response to hypoxia at all ages, except P12. Therefore, reliance on cutaneous gas exchange during early life may be partially attributed to reduced chemosensitivity or a lack of central integration of chemosensitive afferent information. This may be in part due to the relative immaturity of this species at birth, compared to other mammals.

Key words: chemoresponse, respiration, marsupial, neonatal development

Introduction:

In most mammals, the respiratory system is sufficiently developed at birth to function as the sole organ of gas exchange. However, in some marsupials, such as the newborn fat-tailed dunnart (body mass 13 mg), the skin is almost solely responsible for gas exchange, with discernible respiratory efforts generally absent on the day of birth (Mortola et al., 1999; Simpson et al., 2011). A central rhythm generator and sufficiently mature respiratory motoneurons and muscles are essential for coordinated ventilation to occur at birth (Feldman et al., 2003). While afferent information to the respiratory centres from lung and airway mechanoreceptors and peripheral and central chemoreceptors is not required for neuronal rhythmicity, it is important for the modulation of the depth, timing and pattern of respiration (Milsom, 1990). The integration of these components is subject to maturational changes and as such neonatal breathing is inherently unstable (Hilaire and Duron, 1999).

The respiratory response to reduced O₂ (hypoxia) or increased CO₂ (hypercapnia) is dependent on the relative size and maturity of the species at birth, and generally increases with postnatal age (Bonora et al., 1994), in part from increasing sensitivities of both central and peripheral chemoreceptors (Hanson et al., 1989; Davis et al., 2006). In contrast to the adult, the newborn typically does not sustain an increase in minute ventilation (\dot{V}_E) (hyperpnoea) in response to hypoxia, and in some cases \dot{V}_E during hypoxia falls below pre-hypoxic levels (Neubauer et al., 1990). In general, reductions in \dot{V}_E during hypoxia are associated with decreases in either the mean inspiratory flow (VT/TI), a correlate of central respiratory drive, or duty cycle (T_I/T_{TOT}). Hypometabolism also plays an important role in neonates for mitigating hypoxia (Mortola, 1993). Despite the reduced breathing in newborns subjected to

acute hypoxia, they still hyperventilate because of a larger drop in \dot{V}_{O_2} compared to \dot{V}_E . While the hypercapnic ventilatory response of most newborn species is reduced compared to adults (Carroll et al., 1993; Carroll and Fitzgerald, 1993; Davis et al., 2006), unlike hypoxia, it is sustained over time by a persistent increase in tidal volume (V_T), despite a gradual decline in the initial respiratory frequency (f) increase (Bonora et al., 1994; Cummings and Frappell, 2009). Exceptions to these patterns exist, however; with the neonatal North American opossum (another marsupial) demonstrating an unusually large ventilatory response to hypoxia and a hypometabolic response to hypercapnia (Farber, 1972) which attenuates with age (Farber et al., 1972).

Using an *in vitro* brain stem preparation, it has been demonstrated that the neonatal opossum (P5), born after a similar gestation to the fat-tailed dunnart (13 days), possesses neurons that fire rhythmically at a rate sufficient for normal breathing (Farber, 1993; Eugenin and Nicholls, 2000). In the fat-tailed dunnart, phrenic motoneurons have made contact with the diaphragm at birth, yet continuous breathing does not occur until P3 (Frappell and MacFarlane, 2006; Simpson et al., 2011). If the central pattern generator(s) of the fat-tailed dunnart fire rhythmically, as they do in the opossum, then there may be an insufficient inspiratory drive preventing the emergence of normal breathing. In the early neonatal period of the fat-tailed dunnart, reduced inspiratory drive may be due to effective cutaneous exchange and hence a lack of stimulus to the chemoreceptors, or alternatively a consequence of reduced chemoreceptor responsiveness. In this study, we therefore aimed to characterise the ventilatory and metabolic response to hypoxia and hypercapnia in the neonatal fat-tailed dunnart to determine the chemoreceptor response during the neonatal period.

Methods:

Ventilatory and metabolic responses to inspired gases (Air, hypercapnia, hypoxia) were measured in fat-tailed dunnart pouch young in different individuals at P0 (the day of birth) (N= 12, 6, 4), P5 (N= 7, 4, 4), P12 (N= 9, 6, 5) and P23 (N= 9, 4, 5) using a closed respirometry system as previously described (Frappell et al., 1989; Simpson et al., 2011). Briefly, an animal was masked and placed into a water-jacketed chamber maintained at a constant temperature of 36 °C (pouch temperature) and 100 % relative humidity; the chamber effectively separating the head and body. After an equilibration time the chamber was sealed for a known period of time depending on the mass, and hence, metabolic rate of the animal (5 – 15 minutes). Each compartment was then flushed with a known flow (21 ml.min⁻¹), and the gas passed through Nafion® tubing (dead space 0.6 ml) surrounded by a molecular sieve desiccant (crystalline metal aluminosilicate zeolite) prior to being analysed for fractional concentrations of O₂ and CO₂ by gas analysers (ADInstruments, ML205). The output of each gas analyser was recorded at 200 Hz (Chart 4.2 and PowerLab, ADInstruments, Colorado Springs). The rates of oxygen consumption (\dot{V}_{O_2}) and carbon dioxide production (\dot{V}_{CO_2}) were calculated from the time integral of the gas concentration curves multiplied by the flow and the reciprocal of the time for which the chamber was sealed (Frappell and Mortola, 2000).

Ventilation (\dot{V}_E) was measured when the chamber was sealed via a pressure transducer (Spirometer ML141, ADInstruments) connected to the head compartment, and the pressure oscillations were acquired at 200 Hz (Chart 4.2 and PowerLab, ADInstruments). The pressure oscillations associated with breathing were calibrated for volume by the injection and withdrawal of 2 µl of air and the stability of the

pressure change with each injection was also used to indicate the integrity of the seal for the chamber. When analysing \dot{V}_E , at least 50 consecutive breaths were analysed for tidal volume (VT), inspiratory time (TI), expiratory time (TE), post-inspiratory pause (TP; note that TP is fractional to TE because it represents the passive-static component of TE, achieved with closed glottis), total breath time (TTOT = TI + TE), frequency ($f = 60/\text{TTOT}$), mean inspiratory flow (VT/TI), duty cycle (TI/TTOT) and minute ventilation ($\dot{V}_E = \text{VT} \times f$). \dot{V}_E in P0 animals displayed marked instability with prolonged apnea (Simpson et al., 2011); therefore, all recorded breaths were analysed in these animals. Animals in which \dot{V}_E was not discernible were not included in the analysis. \dot{V}_{O_2} is expressed at standard temperature, pressure and humidity (STPD: 1 mL O₂ = 0.0446 mmol O₂) and volume at body temperature, pressure and humidity (BTPS).

Measurements of \dot{V}_E and \dot{V}_{O_2} were first conducted in room air (~21 % O₂, 0.03 % CO₂, balance N₂). Animals were then subjected to either hypoxia (10 % O₂, balance N₂) or hypercapnia (21 % O₂, 5 % CO₂, balance N₂) supplied by a Wosthoff gas mixing pump. In the case of hypercapnia, the animals were exposed to 5 % CO₂ for 5 minutes prior to sealing the chamber for measurement, and analysis performed within the first minute (*i.e.* at 5-6 min CO₂ exposure, allowing examination of the total response; *i.e.* that due to both peripheral and central chemoreceptor activation). When exposed to hypoxia, the chamber was sealed as soon as the gas mix had washed in and oxygen levels stabilised at 10 % O₂ so that we could analyse the early (1 minute of hypoxia) and late (6 minutes of hypoxia) response, thereby enabling the characterization of a bi-phasic response, should it occur. After measurement of the ventilatory and metabolic responses to an inspired gas, the chamber was flushed for 10 min with room air, prior to a second air measurement which was used to determine

if the animal preparation had degenerated during the experiment. In the rare event that this occurred, the entire run was rejected.

Student T-tests were used to determine whether the value obtained during a particular gas challenge was different from the value in normal room air using SPSS (version 19). The effect of age and gas condition, as well as their interaction, was assessed for each metabolic and respiratory variable using a two-way ANOVA. Significance was considered at $p < 0.05$.

Results:

Development of the breathing pattern in air

No discernible ventilatory pattern was observed in 2/3 of neonates at P0 (body weight 13 mg). Where a ventilatory pattern was detected, breathing was generally accompanied by extended periods of apnoea, such that breathing may not have been observed again for the remainder of the measurement period. By P5, the continuously expressed breathing pattern was characterised by a post-inspiratory pause, together with frequent augmented breaths, or sighs (see (Simpson et al., 2011) for representative breathing traces). There were some periods of instability as indicated by the coefficients of variation for TTOT and VT being 11% and 45% respectively. At P12, some animals had a discernible post-inspiratory pause, while others did not, leading to further increased variability in TTOT (37%) and VT (68%). Thus, P12 is presumably a time of transition from the “immature” to “adult” breathing pattern. No discernible post-inspiratory pauses were detected in P23 neonates. Spirograms demonstrating a characteristic breath, in terms of timing and amplitude, are shown for each age in Figure 1

The effects of hypercapnia on fat-tailed dunnart neonates

While there was a tendency for \dot{V}_E/\dot{V}_{O_2} to increase at all ages in response to 5 % CO₂, a significant effect was only observed at P0, owing to an unexpected but significant decrease in \dot{V}_{O_2} ($P < 0.05$; Figure 2A and C). Increasing CO₂ decreased \dot{V}_{O_2} only at P0 ($p < 0.05$), and not in older animals (Figure 2C). There was no significant hypercapnic ventilatory response at any ages; the response approached significance in animals older than P0 ($p < 0.075$) (Figure 2B). The attempted hyperpnoeic effort

observed in P5 animals exposed to 5 % CO₂ was due mainly to a significant increase in *f*, while at P23 the attempted response was dominated by a significant increase in VT (Figure 2D and E). The decrease in *f* experienced by P23 animals, however, negated any increase in \dot{V}_E . There were no significant changes in duty cycle in response to 5 % CO₂ exposure (Figure 2F). There was a significant increase in VT/TI in response to CO₂ at P12, while the effect at P23 was marginal (*p* = 0.06; Figure 2G).

The effects of hypoxia through development in the fat-tailed dunnart

Across all ages, except P12, there was a significant increase in \dot{V}_E/\dot{V}_{O_2} after 1 min of hypoxia (*p* < 0.05; Figure 3A). This effect was achieved predominantly by a decrease in \dot{V}_{O_2} (*p* < 0.05, Figure 3C). There was no increase in \dot{V}_E in response to hypoxia, regardless of the duration of exposure. Hypoxia actually suppressed \dot{V}_E in P5 and P12 animals; at P12 this suppression occurred after only 1 min (Figure 3B). The decrease in \dot{V}_E in response to hypoxia was mainly owing to a decrease in *V_T* at all ages except P23 (Fig. 3D). At P0, the fall in VT was offset by an increase in *f*, thus maintaining \dot{V}_E during hypoxia. The increase in *f* was reflected in an increase in TI/TTOT (i.e. reduced expiratory time) (Fig. 3E & 3F). The absence of a ventilatory response at any age precluded any increase in respiratory drive (VT/TI) (Fig. 3G).

In addition, analysis with two-factor ANOVA demonstrated that increasing age contributed to alterations in all respiratory and metabolic variables over the course of the neonatal period (*p* < 0.05), while the evolution of the response to the gas challenge was evident in all respiratory and metabolic outcomes except *f* and TI/TTOT during the developmental period studied. Convective requirement (\dot{V}_E/\dot{V}_{O_2}) was the only variable to demonstrate a significant interaction between age and the gas condition (*p* = 0.025).

In summary, no significant ventilatory response to hypoxia and hypercapnia was observed in the neonatal fat-tailed dunnart (Fig. 4). Regardless, a hyperventilatory response (increase in \dot{V}_E/\dot{V}_{O_2}) was observed in response to hypoxia (and hypercapnia at P0) owing to a suppression of \dot{V}_{O_2} .

Discussion:

To summarise, a very limited ventilatory response to hypoxia and hypercapnia was evident during the neonatal fat-tailed dunnart. Despite this, suppression of \dot{V}_{O_2} resulted in a hyperventilatory response (increase in \dot{V}_E/\dot{V}_{O_2}) to hypoxia at all ages, except P12, and hypercapnia at P0.

Development of the breathing pattern

The breathing pattern of the neonatal fat-tailed dunnart undergoes considerable postnatal development with the post-inspiratory pause diminishing, and V_T , TI and f increasing. Breathing was not continuously expressed in P0 neonates, and when present f was low. We have previously postulated that these movements are more likely to be akin to fetal breathing movements, preparing the musculature for breathing rather than participating in alveolar gas exchange (Simpson et al., 2011). The respiratory network of the opossum, a species born after a similar gestation to the dunnart, is capable of generating activity sufficient for normal breathing. If this is also the case in the dunnart, it may be that reduced responsiveness of respiration to changing blood gases, possibly via reduced chemoreceptor activity, contributes to their relatively reduced \dot{V}_E/\dot{V}_{O_2} compared to other newborn mammals. Regardless, the ability of the dunnart to exchange gas across the skin perhaps renders inconsequential the relatively immaturity of the respiratory control system (Simpson et al., 2011).

Hypometabolism during hypercapnia: A novel strategy

We have shown that, unlike other newborn mammals (Bonora et al., 1994; Mortola and Lanthier, 1996; Saiki and Mortola, 1996) the newborn fat-tail dunnart utilizes a

hypometabolic strategy to mitigate the effects of increasing CO₂. While the newborns of other species exhibit an attenuated ventilatory response to hypercapnia (and hypoxia) in comparison to adults, there is nearly always a discernible ventilatory response, even at birth (Mortola and Lanthier, 1996). While there was some evidence of an increase in respiratory f (P5), V_T (P23) and V_T/T_I (P12), overall, we were unable to detect any significant increase in \dot{V}_E in response to hypercapnia throughout the first 23 days of life. Unlike the majority of newborn mammals, our newborn dunnarts utilized a hypometabolic strategy in response to hypercapnia; hypometabolism is observed in most other newborn mammals during hypoxia. While a higher mass-specific metabolic rate is observed in the control (room air) P0 dunnarts when compared to the older neonates, this is not unexpected. Birth is a tumultuous event associated with drastic changes in environmental conditions. The surge of catecholamines associated with the stress of parturition generally results in newborns having a higher mass-specific \dot{V}_{O_2} (Bartlett and Areson, 1977; Mortola, 2001) with \dot{V}_{O_2} decreasing markedly within the first 24 hours after birth (Noblet and Le Dividich, 1981; Stewart et al., 1984). While previous studies have demonstrated the absence of this increased mass-specific metabolic rate in neonatal marsupials (Singer et al., 1995; MacFarlane and Frappell, 2001; Szdzuy et al., 2008), these studies were not concentrated on young in the first hours of life, where the newborn may have just recently attached to teat and not yet fully settled.

In addition to our newborn dunnarts, a hypometabolism during hypercapnia has also been described in the P5 opossum, albeit in conjunction with a hyperpnoea (Farber et al., 1972). Hypometabolism may be the preferred strategy for the opossum and dunnart because of mechanical constraints on the respiratory system related to their relative immaturity (MacFarlane et al., 2002). But it is equally plausible that

marsupials generally have reduced chemosensitivity or increased blood buffering capacity owing to their pouch development and exposure to elevated CO₂ (Frappell and Baudinette, 1995; Frappell et al., 2002). As our analysis of the hypercapnic response was limited to after 5 min exposure to CO₂ (representing the integrated response of both the peripheral and central chemoreceptors), it is possible that the ventilatory response in the dunnart peaks before 5 minutes of exposure. In neonatal rats, \dot{V}_E has been shown to decrease slightly between 3 and 5 mins though was still significantly elevated at 5 mins with respect to baseline (Cummings and Frappell, 2009). Alternatively, it may be that the reduced response we observed was due to the small size of the dunnarts. Others have described a size dependency of the \dot{V}_E response to CO₂, with larger newborns possessing a greater ventilatory response than smaller ones (Mortola and Lanthier, 1996).

The effects of hypoxia on breathing and metabolism

Hypoxia did not lead to an increase in \dot{V}_E at any age. In fact, hypoxia caused a sustained decrease in \dot{V}_E , similar to newborns of other species, including preterm humans (Alvaro et al., 1992; Mortola et al., 1992). Regardless, other than at P12, our dunnarts still hyperventilated, as the fall in \dot{V}_{O_2} was greater than the fall in \dot{V}_E . While the decrease in \dot{V}_E during hypoxia could be owing to the immaturity of the peripheral chemoreceptors, adult fat-tailed dunnarts also have a relatively weak hypoxic ventilatory response with \dot{V}_E dropping below normoxic levels (Frappell et al., 1992). Thus, the carotid body of the dunnart may be generally less responsive to hypoxia compared to other species. Alternatively, there could be an active, centrally mediated suppression of \dot{V}_E (Blanco et al., 1984; Coles and Dick, 1996).

Despite the lack of ventilatory response, the hypometabolic response observed in the fat-tailed dunnarts is sufficient to increase \dot{V}_E/\dot{V}_{O_2} for the preservation of tissue oxygenation and acid-base balance. The decreased ventilatory response to hypoxia may drive the hypometabolism via extreme hypoxemia, or the hypometabolism may be an adaptive response that negates the need for an increase in \dot{V}_E for energy conservation. Why animals did not hyperventilate at P12 is unknown, but may reflect either the high tidal volume observed in control animals (likely a result of the highly variable breathing pattern and the switch from neonatal to adult breathing in this age group) which stabilised during CO₂ exposure, or a window of development where these animals are more vulnerable to hypoxia. Such a window of vulnerability has been reported in P12 rats with an absence of increasing \dot{V}_E/\dot{V}_{O_2} , due to a decrease in tidal volume and a comparatively weak metabolic response to hypoxia; a response that was different from the rest of the first 3 postnatal weeks (Liu et al., 2006). In addition, a reduction in excitatory and increase in inhibitory neurotransmitters (Wong-Riley and Liu, 2005) as well as a significant decline in tryptophan hydroxylase and serotonin transporter immunoreactivity (Liu and Wong-Riley, 2010b, a) further suggests that normal development can contribute to a narrow window of vulnerability in rats at this age (P12), though lack of proper controls limits the interpretation of these developmental studies. Developmentally, the P12 dunnart (eye opening at P49; weening at P70) is markedly more immature when compared to the P12 rat (eye opening at P14.5; weening at P21) with lungs that are yet to commence alveolarisation (Simpson et al., 2011). However, P12 marks the period where the breathing pattern changes from the neonatal pattern (with the presence of a post inspiratory pause) to a typical adult mammalian pattern indicating that this is an important time for respiratory system development in the neonatal fat-tailed dunnart.

In summary, the reduced \dot{V}_E and \dot{V}_E/\dot{V}_{O_2} exhibited by the fat-tailed dunnart in early postnatal life may be at least in part owing to severely reduced sensitivity of the respiratory control system to hypoxia and hypercapnia. In turn, this may be due to the relative immaturity of this species, with central and peripheral chemoreceptors being underdeveloped. Despite this, these animals likely maintain tissue oxygenation and pH via a hypometabolic response. Reduced chemoresponses and low \dot{V}_E/\dot{V}_{O_2} are most likely the response to an underdeveloped lung and respiratory system and hence a heavy reliance on skin gas exchange during early postnatal life in the dunnart.

Figure 1. Spirogram demonstrating a characteristic breath for P0, 5, 12 and 23 fat-tailed dunnarts under normoxic/normocapnic conditions.

Respiratory timing variables and tidal volumes are shown for neonatal fat-tailed dunnarts at P5, 12 and 23 in black. Data for P0 (grey) include all measurable breaths.

Figure 2. The effects of hypercapnia on breathing and metabolism in the neonatal fat-tailed dunnart.

Changes in convective requirement (\dot{V}_E/\dot{V}_{O_2}) (A), mass specific minute ventilation (\dot{V}_E) (B), total mass specific rate of oxygen consumption (skin + pulmonary) (\dot{V}_{O_2}) (C), tidal volume (VT) (D), respiratory frequency (E), duty cycle (F) and respiratory drive (G) are shown after 5 minutes exposure to 5 % CO₂ in P0, 5, 12 and 23 fat-tailed dunnarts. Data are presented as means \pm 1 S.E.M. * indicates significant difference from air.

Figure 3. The effects of hypoxia on breathing and metabolism in the neonatal fat-tailed dunnart.

Changes in convective requirement (\dot{V}_E/\dot{V}_{O_2}) (A), mass specific minute ventilation (\dot{V}_E) (B), total mass specific rate of oxygen consumption (skin + pulmonary) (\dot{V}_{O_2}) (C), tidal volume (VT) (D), respiratory frequency (E), duty cycle (F) and respiratory drive (G) are shown after 1 and 6 minutes exposure to 10 % O₂ in P0, 5, 12 and 23 fat-tailed dunnarts. Data are presented as means \pm 1 S.E.M. * indicates significant difference from air.

Figure 4. Spirograms demonstrating the change in breath-timing and tidal volume in response to hypoxia and hypercapnia in developing fat-tailed dunnart neonates.

Shown are mean control values (air) at P0, 5, 12 and 23 in grey, the effects of hypercapnia (solid black) at 5 minutes and hypoxia at 1 minute (dash black) and 6 minutes (dash-dot black). Error bars represent 1 S.E.M.





Figure 1



Figure




Figure 3




Figure 4.

References

Alvaro, R., Alvarez, J., Kwiatkowski, K., Cates, D. and Rigatto, H. (1992). Small preterm infants (less than or equal to 1500 g) have only a sustained decrease in ventilation in response to hypoxia. *Pediatr Res* **32**, 403-406.

Bartlett, D. and Areson, J. (1977). Quantitative lung morphology in newborn mammals. *Respiration Physiology* **29**, 193-200.

Blanco, C. E., Hanson, M. A., Johnson, P. and Rigatto, H. (1984). Breathing pattern of kittens during hypoxia. *J Appl Physiol* **56**, 12-17.

Bonora, M., Boule, M. and Gautier, H. (1994). Ventilatory strategy in hypoxic or hypercapnic newborns. *Biol Neonate* **65**, 198-204.

Carroll, J. L. and Fitzgerald, R. S. (1993). Carotid chemoreceptor responses to hypoxia and hypercapnia in developing kittens. *Adv Exp Med Biol* **337**, 387-391.

Carroll, J. L., Bamford, O. S. and Fitzgerald, R. S. (1993). Postnatal maturation of carotid chemoreceptor responses to O₂ and CO₂ in the cat. *J Appl Physiol* **75**, 2383-2391.

Coles, S. K. and Dick, T. E. (1996). Neurons in the ventrolateral pons are required for post-hypoxic frequency decline in rats. *J Physiol* **497** (Pt 1), 79-94.

Cummings, K. J. and Frappell, P. B. (2009). Breath-to-breath hypercapnic response in neonatal rats: temperature dependency of the chemoreflexes and potential implications for breathing stability. *Am J Physiol Regul Integr Comp Physiol* **297**, R124-134.

Davis, S. E., Solhied, G., Castillo, M., Dwinell, M., Brozoski, D. and Forster, H. V. (2006). Postnatal developmental changes in CO₂ sensitivity in rats. *J Appl Physiol* **101**, 1097-1103.

Eugenin, J. and Nicholls, J. G. (2000). Control of respiration in the isolated central nervous system of the neonatal opossum, *Monodelphis domestica*. *Brain Research Bulletin* **53**, 605-613.

Farber, J. P. (1972). Development of pulmonary reflexes and pattern of breathing in the Virginia opossum. *Respiration Physiology* **14**, 278-286.

Farber, J. P. (1993). Maximum discharge rates of respiratory neurons during opossum development. *J Appl Physiol* **75**, 2040-2044.

Farber, J. P., Hultgren, H. N. and Tenney, S. M. (1972). Development of the chemical control of breathing in the Virginia opossum. *Respiration Physiology* **14**, 267-277.

Feldman, J. L., Mitchell, G. S. and Nattie, E. E. (2003). Breathing: rhythmicity, plasticity, chemosensitivity. *Annu Rev Neurosci* **26**, 239-266.

Frappell, P., Lanthier, C., Baudinette, R. V. and Mortola, J. P. (1992). Metabolism and ventilation in acute hypoxia: a comparative analysis in small mammalian species. *American Journal of Physiology* **262**, R1040-R1046.

Frappell, P. B. and Baudinette, R. V. (1995). Scaling of respiratory variables and the breathing pattern in adult marsupials. *Respiration Physiology* **100**, 83-90.

Frappell, P. B. and Mortola, J. P. (2000). Respiratory function in a newborn marsupial with skin gas exchange. *Respiration Physiology* **120**, 35-45.

Frappell, P. B. and MacFarlane, P. M. (2006). Development of the respiratory system in marsupials. *Respir Physiol Neurobiol* **154**, 252-267.

Frappell, P. B., Blevin, H. A. and Baudinette, R. V. (1989). Understanding respirometry chambers: what goes in must come out. *Journal of Theoretical Biology* **138**, 479-494.

Frappell, P. B., Baudinette, R. V., MacFarlane, P. M., Wiggins, P. R. and Shimmin, G. (2002). Ventilation and metabolism in a large semifossorial marsupial: The effect of graded hypoxia and hypercapnia. *Physiological & Biochemical Zoology* **75**, 77-82.

Hanson, M. A., Kumar, P. and Williams, B. A. (1989). The effect of chronic hypoxia upon the development of respiratory chemoreflexes in the newborn kitten. *Journal of Physiology (London)* **411**, 563-574.

Hilaire, G. and Duron, B. (1999). Maturation of the mammalian respiratory system. *Physiological Reviews* **79**, 325-360.

Liu, Q. and Wong-Riley, M. T. (2010a). Postnatal changes in the expressions of serotonin 1A, 1B, and 2A receptors in ten brain stem nuclei of the rat: implication for a sensitive period. *Neuroscience* **165**, 61-78.

Liu, Q. and Wong-Riley, M. T. (2010b). Postnatal changes in tryptophan hydroxylase and serotonin transporter immunoreactivity in multiple brainstem nuclei of the rat: implications for a sensitive period. *J Comp Neurol* **518**, 1082-1097.

Liu, Q., Lowry, T. F. and Wong-Riley, M. T. (2006). Postnatal changes in ventilation during normoxia and acute hypoxia in the rat: implication for a sensitive period. *The Journal of physiology* **577**, 957-970.

MacFarlane, P. M. and Frappell, P. B. (2001). Convection requirement is established by total metabolic rate in the newborn tammar wallaby. *Respiration Physiology* **126**, 221-231.

MacFarlane, P. M., Frappell, P. B. and Mortola, J. P. (2002). Mechanics of the respiratory system in the newborn tammar wallaby. *Journal of Experimental Biology* **205**, 533-538.

Milsom, W. K. (1990). Mechanoreceptor modulation of endogenous respiratory rhythms in vertebrates. *American Journal of Physiology* **259**, R898-R910.

- Mortola, J. P.** (1993). Hypoxic hypometabolism in mammals. *NIPS* **8**, 79-82.
- Mortola, J. P.** (2001). *Respiratory Physiology of Newborn Mammals. A Comparative Perspective*. Baltimore: John Hopkins University Press.
- Mortola, J. P. and Lanthier, C.** (1996). The ventilatory and metabolic response to hypercapnia in newborn mammalian species. *Respiration Physiology* **103**, 263-270.
- Mortola, J. P., Frappell, P. B. and Woolley, P. A.** (1999). Breathing through skin in a newborn mammal. *Nature* **397**, 660.
- Mortola, J. P., Frappell, P. B., Dotta, A., Matsuoka, T., Fox, G., Weeks, S. and Mayer, D.** (1992). Ventilatory and metabolic responses to acute hyperoxia in newborns. *American Review of Respiratory Disease* **146**, 11-15.
- Neubauer, J. A., Melton, J. E. and Edelman, N. H.** (1990). Modulation of respiration during brain hypoxia. *J Appl Physiol* **68**, 441-451.
- Noblet, J. and Le Dividich, J.** (1981). Energy metabolism of the newborn pig during the first 24 h of life. *Biology of the neonate* **40**, 175-182.
- Saiki, C. and Mortola, J. P.** (1996). Effect of CO₂ on the metabolic and ventilatory responses to ambient temperature in conscious adult and newborn rats. *Journal of Physiology* **491**, 261-269.
- Simpson, S. J., Flecknoe, S. J., Clugston, R. D., Greer, J. J., Hooper, S. B. and Frappell, P. B.** (2011). Structural and functional development of the respiratory system in a newborn marsupial with cutaneous gas exchange. *Physiological and biochemical zoology : PBZ* **84**, 634-649.
- Singer, D., Zeller, U., Hehenkamp, E., Schmidt, H. and Kuhn, H.** (1995). Suppression and activation of Kleiber's rule in the neonatal period: a comparative calorimetric investigation in preterm human and small marsupial neonates. *Eur J Physiol Suppl* **429**, R142.
- Stewart, J. H., Rose, R. J. and Barko, A. M.** (1984). Respiratory studies in foals from birth to seven days old. *Equine Vet J* **16**, 323-328.
- Szdzuy, K., Zeller, U., Renfree, M., Tzschentke, B. and Janke, O.** (2008). Postnatal lung and metabolic development in two marsupial and four eutherian species. *J Anat* **212**, 164-179.
- Wong-Riley, M. T. and Liu, Q.** (2005). Neurochemical development of brain stem nuclei involved in the control of respiration. *Respir Physiol Neurobiol* **149**, 83-98.