

1 **Behavioral correction to prevent overhydration and increase survival by larvae of the net-**
2 **spinning caddisflies in relation to water flow**

3

4 Jay A. Yoder¹, Joshua B. Benoit^{2,*}, Blake W. Nelson¹, Leighanne R. Main¹ and Jon P. Bossley³

5

6 ¹Department of Biology, Wittenberg University, Springfield, OH 45501, USA, ²Department of
7 Biological Sciences, University of Cincinnati, Cincinnati, OH 45221, USA and ³Environmental
8 Science Graduate Program, The Ohio State University, Columbus, OH 43210, USA

9

10 * Author for correspondence (joshua.benoit@uc.edu)

11

12 **ABSTRACT**

13 We report behavioral regulation of body water content in caddisfly larvae, *Hydropsyche morosa*
14 and *Cheumatopsyche pettiti*, by selecting microhabitats with different water flow rates. The
15 purpose was to examine features necessary for survival in the same apparent habitat, because
16 both co-exist in riffle areas of freshwater streams. Both species are highly sensitive to water loss
17 due to high water loss rates and depend on immersion in fresh water (hypo-osmotic) to maintain
18 water stores. In contrast to *C. pettiti*, *H. morosa* is larger, retains water more effectively, and
19 features reduced water loss rates with suppressed activation energies. When *H. morosa* was
20 confined to areas of low or no water flow, overhydration led to rapid mortality, whereas the same
21 conditions favored water balance maintenance and survival in *C. pettiti*. In attraction bioassays,
22 *H. morosa* moved and remained within areas of high water flow and *C. pettiti* preferred areas
23 with low water flow. Because water flow rates are unlikely to directly impact water gain, the
24 mechanism responsible for increased survival and water balance maintenance is likely related to
25 the impact of water flow on oxygen availability or differences in feeding ecology.

26

27 **KEY WORDS: Water exchange, Hyperosmotic, Attraction, Fresh water, Biological**
28 **indicator**

29

30 **Running title: Water balance of caddisfly larvae**

31

32 INTRODUCTION

33 Whole organism water balance is an ecologically-defining attribute and limits the ability of
34 insects to function properly in the absence of water resources (Hadley, 1994). Habitat adaptation
35 involves a complementary relationship between water balance characteristics and modifications
36 of behavior relevant to preventing drying out or overhydrating (Hadley, 1994). To maintain
37 water balance, water gain (m_S) must be equal to water loss (m_T) as defined by Wharton (1985).
38 Net water gain occurs when $m_S > m_T$, and net water loss occurs when $m_T > m_S$. The goal is to
39 maintain body water ($\Delta m = 0$), a condition that permits proper functioning and development due
40 to a lack of dehydration-induced stress. In a freshwater habitat, the m_S component is exceedingly
41 large for an aquatic insect in that they are hyperosmotic to fresh water. The water activity (a_w) of
42 the freshwater environment, $1.00a_w$, is greater than the $0.99a_w$ activity of the insect's body water
43 (0.990 - $0.997a_w$; Wharton, 1985; Sigal et al., 1991). Thus, the activity gradient of fresh water
44 results in a continuous water increase to the body by simple diffusion (Kohn, 1965; Wharton,
45 1985). To prevent overhydration, insects must either suppress cuticular permeability to prevent
46 water intake or increase the rate of water removal.

47 Net-spinning caddisfly larvae (Trichoptera: Hydropsychidae) reside at the bottoms of
48 freshwater ponds and streams on cobblestones or limbs. They have a preference for shallow riffle
49 areas with low to high current flow, in cooler, shaded areas (Wiggins, 1996; Bouchard, 2004).
50 Caddisfly larvae thrive within high quality fresh water and are utilized as bioindicators (Bonada
51 and Williams, 2002 and references therein). Appreciable amounts of water are obtained from
52 their moisture-rich food, predominately algae and detritus from the stream bottom (Snyder and
53 Hendricks, 1995) and from drinking (Sutcliffe, 1961). The emphasis for maintaining water
54 balance by caddisfly larvae focuses on increasing the water loss component, m_T , to counteract
55 the continual water influx that occurs naturally from ingestion and inward diffusion by being
56 submerged. Despite the application of caddisfly larvae to environmental studies, little work has
57 been done on the water balance physiology of caddisfly larvae. The exception is Sutcliffe's
58 (1961) work on salt balance, where it is shown that caddisfly larvae are hyperosmotic and water
59 is attracted to them.

60 In this paper, we determined water balance characteristics and conducted attraction
61 bioassays in relation to water flow for larvae of two caddisfly species that regularly co-occur in
62 freshwater streams, *Hydropsyche morosa* and *Cheumatopsyche pettiti*. Few experiments have

63 examined species comparisons based on water balance characteristics of insect larvae with the
64 exception of mosquitoes (Bradley, 1987, 2008) and midge larvae (Benoit et al. 2007; Kikawada
65 et al., 2008; Elnitsky et al., 2009). Comparative studies of water balance studies of species that
66 reside within the same habitats have also been minimal (fruit flies, Aggarwal et al., 2013;
67 Parkash et al., 2011, 2013; tsetse flies, Kleynhans and Terblanche, 2011; Terblanche et al., 2006,
68 2008; beetles, Benoit et al. 2005; mites, Benoit et al. 2008). Our hypothesis is that larvae of
69 different caddisfly species may have different water balance profiles, which likely necessitate
70 different habitat requirements to maintain water balance.

71

72 **RESULTS**

73 **Analysis of stream water and flow rates**

74 Physicochemical data of stream water were specific conductance ($524.5 \pm 12.8 \mu\text{S}/\text{cm}$),
75 dissolved oxygen ($13.6 \pm 0.59 \text{ mg}/\text{L}$ and $105.9 \pm 2.2 \%$), temperature ($22.06 \pm 0.11^\circ\text{C}$), and pH
76 (8.10 ± 0.34) ($N = \text{ten measurements/ each of three sites, mean} \pm \text{s.e.m.}$). Average stream flow
77 velocity was $4.1 \pm 1.7 \text{ km}/\text{h}$ taken near the bottom, middle, and top of the water column ($N = \text{ten}$
78 $\text{measurements/ each of three sites, mean} \pm \text{s.e.m.}$). Water flow velocities for experimental airflow
79 rates were $3 \text{ L}/\text{min} = 2.4 \pm 0.9 \text{ km}/\text{h}$; $6 \text{ L}/\text{min} = 5.7 \pm 0.7 \text{ km}/\text{h}$; and $12 \text{ L}/\text{min} = 9.1 \pm 1.6 \text{ km}/\text{h}$
80 ($N = 15 \text{ measurements, mean} \pm \text{s.e.m.}$). Oxygen content of the stream water after it was filtered
81 and autoclaved was $14.1 \pm 0.38 \text{ mg}/\text{L}$ and $108.3 \pm 2.7 \%$, indicating that water in bioassays was
82 supersaturated with oxygen ($> 100\% \text{ DO}$).

83

84 **Water balance characteristics**

85 *Hydropsyche morosa* was larger than *C. pettiti* in initial mass, dry mass, and water mass ($P <$
86 0.05 in each pairwise comparison; Table 1). In all cases, the water mass was a positive correlate
87 of the dry mass ($r^2 \geq 0.93$ for *H. morosa*; $r^2 \geq 0.91$ for *C. pettiti*; $P < 0.001$). Both species had
88 similar percentage body water content around 66% ($P > 0.05$; Table 1). Thus, *H. morosa* is
89 larger and both species have similar relative body water pools that are available for exchange
90 with their surroundings.

91 Water loss rates were more rapid for *C. pettiti* than *H. morosa* (Fig. 2, Table 1; $P < 0.05$).
92 Once *C. pettiti* lost 19.6% of their body water they were alive, but they were unable to coordinate
93 their movements and self-right (Table 1). The dehydration tolerance of *C. pettiti* was 19.6% and

94 the dehydration tolerance of *H. morosa* was 23.02% ($P > 0.05$; Table 1). Moribund *H. morosa*
95 and *C. pettiti* died if placed at 100% RH or in stream water, evidence that they had sustained an
96 irreversible level of water loss at their dehydration tolerance limit.

97 At 37°C there is a critical transition temperature (CTT) for *C. pettiti* as denoted by the
98 biphasic change on the Arrhenius plot (Fig. 3). The activation energy changes from 14.5 kJ/mol
99 in the low temperature range to 33.2 kJ/mol in the high temperature range ($P < 0.05$; Table 1).
100 Water loss shows direct Boltzmann dependence in both low and high temperature ranges ($r^2 =$
101 0.98 and 0.94, respectively; $P < 0.001$; Fig. 3). Ramp-up and ramp-down determinations yielded
102 nearly identical activation energies as given in Table 1, producing highly reproducible CTTs at
103 or near 36°C (data not shown). The 35°C CTT for *H. morosa* was not significantly different from
104 the CTT for *C. pettiti* ($P > 0.05$; Table 1). For *H. morosa*, there was a change in activation
105 energy that separated the low temperature range (9.1 kJ/mol) from the high temperature range
106 (26.4 kJ/mol) (Table 1, Fig. 3; $P < 0.05$) as a regular Boltzmann temperature function ($r^2 \geq 0.92$;
107 $P < 0.001$). Activation energies were lower for *H. morosa* than *C. pettiti* in both low and high
108 temperature ranges ($P < 0.05$; Table 1).

109

110 **Survival estimates and mass changes**

111 An experimental arena was designed using a stream water-filled specimen dish affixed with
112 tubing at one side of the dish (sector one) that created a high water flow current by bubbling air
113 (Fig. 1). 'Direct water flow' refers to the side of the dish (sector one) beneath the air tube where
114 there is a high rate of water flow (sector one). 'Indirect water flow' refers to the opposite side of
115 the dish (sector four) where the water flow is less rapid because the tube is aimed into sector one.

116 Greatest survival for *C. pettiti* occurred in indirect water flow (sector four, Fig. 1), where
117 larvae survived 8.0 days (4.5 days for 50% of larvae) compared to 5.5 days (3.5 days for 50% of
118 larvae) in direct water flow ($P < 0.05$; Fig. 4). Still water conditions also had a positive effect on
119 survival of *C. pettiti* (survival for 7.0 days, 4.0 days for 50% of larvae), with approximately a day
120 less than in indirect water flow ($P < 0.05$; Fig. 4). In contrast, direct water flow resulted in the
121 greatest survival for *H. morosa* compared to indirect water flow: 6.5 days (4.7 days for 50% of
122 larvae) versus 5.0 days (2.7 days for 50% of larvae), respectively (Fig. 4; $P < 0.05$). Survival was
123 negatively impacted in *H. morosa* by still water, which shortened survival by approximately two

124 days (survival for 3.0 days, 1.8 days for 50% of larvae) compared to survival in indirect water
125 flow ($P < 0.05$; Fig. 4).

126 *Cheumatopsyche pettiti* gained more water over time in the direct water flow than the
127 indirect water flow (Fig. 5; $P < 0.05$ in each pairwise comparison). Daily water gains in still
128 water were between the extremes of direct flow ($P < 0.05$ in each pairwise comparison) and
129 indirect flow (Fig. 5; $P < 0.05$ in each pairwise comparison). For *H. morosa*, the largest daily
130 water gains occurred for larvae in still water (Fig. 5; $P < 0.05$ in each pairwise comparison).
131 There were also large amounts of daily water gain for *H. morosa* that was held in indirect water
132 flow compared to direct water flow (Fig. 5; $P < 0.05$ in each pairwise comparison). Under still
133 and indirect water flow conditions for *H. morosa*, mass measurements were discontinued once
134 larvae died after 2-3 days (Fig. 5). Survival for *C. pettiti* did not begin to decline dramatically
135 until after the 72 hour time period of the experiment (Fig. 5).

136

137 **Behavioral responses emphasizing attraction**

138 Moving water of 3 L/min resulted in larger numbers of *C. pettiti* in sector one (Fig. 1) within a
139 day compared to untreated (no tube) or no water flow (airflow tube alone) controls (Fig. 6; $P <$
140 0.05). In all cases, a slight attraction by *C. pettiti* to sector one occurred within one hour when
141 water was flowing at any speed compared to still water controls ($P < 0.05$; Fig. 6). The one and
142 two hour responses by *H. morosa* to the low 3 L/min flow rate were similar to untreated and
143 airflow tube-only controls ($P > 0.05$), but increased substantially at 24 hours ($P < 0.05$; Fig. 6).
144 Water flow at 6 L/min and 12 L/min by *H. morosa* recruited large numbers of larvae within one
145 hour ($P < 0.05$), attracting and retaining nearly all of the larvae (Fig. 6). The heightened intensity
146 of attraction by *H. morosa* in high speed water of 6 L/min and 12 L/min remained elevated at
147 two and 24 hours (Fig. 6).

148

149 **DISCUSSION**

150 The important conclusion from this study is that larvae of *H. morosa* and *C. pettiti* display clear
151 microhabitat preferences with regard to water flow. When placed in non-preferred microhabitats,
152 larvae become overhydrated, thus indicating that inability to maintain water balance could drive
153 species-specific microhabitat preferences. Specifically, *H. morosa* has a low water loss rate and
154 prefers to reside in areas with high water flow. For *C. pettiti*, water loss rates are higher and

155 residence in areas with low water flow permits them to maintain water balance. There is also
156 thermal suitability information as implied by the critical transition temperature (CTT). The CTT
157 is within the 32-39°C of upper thermal tolerance limits for caddisfly larvae (deKozlowki and
158 Bunting, 1981; Moulton et al., 1993), thus mortality at high temperature can be attributed to a
159 sharp increase in water loss, or water gain, at or above the CTT. This water balance/behavior
160 information is relevant to use of caddisfly larvae as bioindicators of high quality water for
161 nutrient cycling, toxicological monitoring, and thermal disturbances (Bonada and Williams,
162 2002, and references therein) since turnovers in water exchange differentially affect each species.

163 From a water balance perspective, different flow rates are preferred for the two species.
164 Here we have attempted to provide a linkage between the water balance characteristics and
165 preferred flow rates, if there is one. *Cheumatopsyche pettiti* and *H. morosa* rely on continuous
166 water influx from their fresh water habitat. Their dehydration tolerance limit is unlikely to be
167 exceeded, because water is continuously entering into the body by inward diffusion, drinking,
168 and feeding on moist food. The continuous water influx is balanced by large body water losses
169 (water efflux) so that the larva functions properly at the body water content. *Cheumatopsyche*
170 *pettiti* is thus highly permeable to water and they are prevented from overhydrating by having
171 high water loss rates. *Hydropsyche morosa* with its much lower water loss rate requires that the
172 amount gained must be restricted, which suggests the presence of water-proofed cuticular barrier
173 that is unique to *H. morosa* that blocks the amount of water entry (water influx). The difference
174 in water loss rate implies that the emphasis is on water elimination for *C. pettiti* and water
175 retention for *H. morosa*. When matched with behavioral preference, implications are that the
176 water flow rate conditions of the microhabitat are opposite of the water turnover of the larva.
177 This is something that we have also found for another aquatic invertebrate, the branchiobdellids
178 (Annelida: Clitellata) that live on freshwater crayfish (Yoder et al., 2007). The species that loses
179 water the fastest (*Cambarincola fallax*) clusters in locales with least amount of water movement
180 on the crayfish, at subrostral sites at the base of the eyestalks. In contrast, a species that loses
181 water the slowest (*C. ingens*) is found preferentially in the high water flow area of the gills
182 (Yoder et al., 2007). Even though both caddisfly species have high cuticular water loss rates and
183 require direct contact with pure water, the stream flow rates significantly impact their ability to
184 maintain water balance.

185 A high water flow rate is unlikely to have an appreciable impact on the amount of water
186 that is gained by an insect, because the supply of water is not being replenished (water is the
187 supply). Hydrodynamic pressure in a high flow rate is greatest on one side of the insect (side hit
188 directly by the current) but low on the other side of the insect (not exposed to current), and this
189 would be predicted to yield a similar net effect on water gain (m_s) as still or low flow rate where
190 the hydrodynamic pressure is more equalized over the insect surface. Indeed, we found that both
191 *H. morosa* and *C. pettiti* can survive in still water, hence, they can evidently maintain proper
192 body water content within tolerable limits. This is supported by field data, where *H. morosa* and
193 *C. pettiti* have been collected in pool areas and from ponds despite the propensity by both species
194 to be more abundant in riffle areas (Bouchard, 2004). These considerations indicate that the
195 amount of water influx is constant whether *C. pettiti* is in a low or high water flow. The amount
196 of water influx for *H. morosa* is less as a tradeoff for their lower water loss rate, because they
197 cannot eliminate excess water gain as effectively as *C. pettiti*. There is enhancement of survival
198 when water gain is the least, and this occurs in regions of high flow for *H. morosa* and low flow
199 for *C. pettiti*. *Hydropsyche morosa* and *C. pettiti* function better (prolonged survival) when there
200 is at least some water flow in the area, and especially a high water flow for *H. morosa*.

201 The problem of desiccation impacting survival is a challenge encountered by many
202 aquatic insects when the stream dries. Larvae of *H. morosa* and *C. pettiti* are quite active, hence
203 lose water rapidly, and their small body size through surface area to volume properties further
204 exacerbates this dilemma. The high water loss rates and modest dehydration tolerance limit of *H.*
205 *morosa* and *C. pettiti* necessarily imply that they are hygric-suited and adapted for a moisture-
206 rich environment (Hadley, 1994). Many aquatic insects and other aquatic invertebrates prevent
207 dehydrating and remain viable by staying in constant, direct contact with a moisture-rich surface
208 (Hadley, 1994). There is little interpretative value ecologically concerning body water content
209 (i.e., not all aquatic insects have high body water contents), and for these caddisfly species it
210 approximates the mean water content (69%) of most insects (Hadley, 1994). Under drought
211 conditions, the most likely scenario is that the caddisfly larvae crawl underground (Wiggins,
212 1996; Bouchard, 2004). The differences we note in water loss rate and behavior between *H.*
213 *morosa* and *C. pettiti* should also be thought of as important resistance mechanisms that promote
214 survival of these caddisfly larvae in intermittent streams. These caddisfly larvae are driven

215 underneath the soil surface in order to satisfy an absolute moisture requirement because of their
216 high water loss rate.

217 Although our results may be a direct consequence of water balance, there are
218 physiological functions impacted by stream flow rates that likely have an impact on survival and
219 habitat preferences. The most likely possibility is that the choice experiments (Fig. 6) could be
220 explained, fully or in part, by behavioral regulation of oxygen availability. A high water flow
221 rate and increase in hydrodynamic pressure would replenish the supply of oxygen at a faster rate.
222 Higher flowing water would bring more oxygen past the caddisfly's filamentous abdominal gills
223 (there are no open spiracles in Trichoptera; Wiggins, 1996). Perhaps the larger body size of *H.*
224 *morosa* requires faster water flow rates to maintain oxygen delivery. If the oxygen requirement is
225 not met (i.e., happens in still or low water flow), then mechanisms responsible for maintaining
226 water balance begin to malfunction for energetic reasons. Overhydration and eventual death
227 would be the physiological consequence of water balance mechanisms breaking down. It also
228 seems reasonable to suggest that proper excretory function in these species requires optimal flow
229 conditions. The chloride epithelium of caddisfly larvae likely requires specific flow rates to
230 regulate chloride uptake, which is essential for maintaining water balance. There is insufficient
231 information about the excretory system of caddisflies to confirm this possibility. Additionally,
232 the negative effect on water balance in still or low flowing water could also be an artifact of
233 some primary function that recruits *H. morosa* to higher flowing waters that could relate to
234 behavioral differences between *C. pettiti* (preference and suitability for still or low water flow) in
235 feeding ecology or in predator avoidance.

236

237 **MATERIALS AND METHODS**

238 **Collection of caddisfly larvae and stream water**

239 Stream locales were selected by random block design within a 3 meter transect in riffle areas of a
240 third-order segment of Buck Creek, Clark Co., OH, USA. Larvae were collected using kick nets
241 (BioQuip Products, Rancho Dominguez, CA, USA). Larvae were identified as *Cheumatopsyche*
242 *pettiti* and *Hydropsyche morosa* and were in the final, fifth instar (keys by Merritt and Cummins,
243 1996; Wiggins, 1996). Slide-mounted vouchers are under specimen lot WUIC Nos. 1209-1229
244 (Wittenberg University, Springfield, OH, USA). Handling of larvae was done with an aspirator.
245 Dead larvae were those that did not move (legs or mouthparts) and failed to respond to stimuli

246 and crawl five body lengths when prodded when examined by 40x microscopy. Stream water
247 was collected into autoclave-sterilized (121°C, 15 psi), 1.0 L glass volumetric flasks, and filtered
248 (3M Aqua-Pure, 3M Co., St. Paul, MN, USA) before use. Physicochemical data of the stream
249 water was collected on site with a water quality sonde (YSI Environmental, Yellow Springs, OH,
250 USA) using a Flowprobe (Global Water, White Plains, NY, USA) to determine flow rate.

251

252 **Equipment, instrumentation and experimental conditions**

253 Relative humidity was maintained in glass desiccators (6.0 L; Fisher) containing anhydrous
254 CaSO₄ for 0% RH (1.5 x 10⁻²% RH; W. A. Hammond Drierite Co., Xenia, OH, USA; Toolson,
255 1978) and deionized double-distilled (DI) water for 100% RH. Relative humidity was measured
256 with a hygrometer (s.d. ± 0.5% RH; Thomas Scientific, Philadelphia, PA, USA). An
257 electrobalance (CAHN, Ventron Co., Cerritos, CA, USA; precision and accuracy were s.d. ± 0.2
258 µg and ± 6 µg at 1 mg, respectively) was used for measuring mass changes of larvae. Larvae
259 were dried to complete dryness (constant mass for three days) in a 90°C drying oven (Blue M
260 Electric Co., Chicago, IL, USA; Hadley 1994). Basic observations were conducted at 20 ± 1°C,
261 15h:9h, L:D photoperiod. Temperature for other studies varied less than ± 0.5°C.

262 Figure 1 shows the six-sector bioassay that we modified from a statistically valid, short
263 range Petri plate attraction bioassay developed by Arlian and Vyszenski-Moher (1995) and Allan
264 and Sonenshine (2002). A straight sided flat bottom specimen dish (19 cm i.d. x 7cm deep =
265 1986 L) was placed over top a paper template that had been scored into six, equal 47.2 cm²
266 sectors. The specimen dish was filled with 1 L of filtered stream water. A 1.0 cm Tygon tubing
267 (Fisher) attached to a source of compressed air and flowmeter for regulation (Rochester Gauges,
268 Dallas, TX, USA) created a bubbling water flow into sector one. No water flow of 0 L/min (still
269 water conditions) served as a control. An additional control was a specimen dish without the
270 airflow tube. Stream water, freshly autoclave-sterilized culture dishes, and unconditioned larvae
271 that had not been previously used in bioassays were utilized for each experiment.

272 All experiments were replicated ten times, ten larvae per replicate (total $N = 100$ larvae),
273 with each replicate coming from a different collection site within the stream. Data are the mean ±
274 s.e.m.

275

276 **Determination of water balance characteristics**

277 Following Wharton's (1985) standard gravimetric methods and equations, the rate of water loss
278 at 0% RH is an accurate measure of the water loss rate as it would occur while submerged in
279 water (experimental determination using ^3HOH ; Arlian and Staiger, 1979). Each larva was
280 monitored individually, without anesthesia or enclosure, and weighed in less than one minute.
281 All experiments use 4-6% pre-desiccated larvae so that mass changes reflect changes in body
282 water levels (Arlian and Eckstrand, 1975; Wharton, 1985).

283 To determine water content and water loss rate, a larva was weighed (= fresh, initial
284 mass, f), placed at 20°C and 0% RH, and re-weighed for five readings of mass. The larva was
285 then dried to constant mass (d = dry mass) in the 90°C drying oven. Dry mass was subtracted
286 from each mass measurement to convert the mass measurement into the water mass (m ; Eq. 1): m
287 = $f - d$. Percentage body water content was calculated (Eq. 2): percentage $m = 100 (f - d)/f$. Water
288 loss rate (integumental plus respiratory water loss) was calculated by fitting mass measurements
289 to the equation (Eq. 3): $m_t = m_0 e^{-kt}$; m_t is the water mass at any time t , m_0 is the initial water mass,
290 and $-k$ is the water loss. Fresh mass, dry mass, water mass and percentage body water content
291 was based on the same cohort of 100 larvae. The water loss rate was based on different cohort of
292 100 larvae.

293 To determine dehydration tolerance, a larva was weighed, held at 0% RH and 20°C, and
294 re-weighed, each time checking the larva for its ability to self-right and crawl five body lengths.
295 At the critical mass (m_c), the larva could move but could not coordinate its movements and failed
296 to self-right and crawl when placing it at 100% RH or in 15 ml of water. Percentage change in m
297 = $100 (m_c - m_0)/m_0$ (Eq. 4) was used to calculate the dehydration tolerance limit. A separate
298 cohort of 200 larvae was used for determining the dehydration tolerance limit: 100 larvae for a
299 rescue attempt by placing larvae at 100% RH and a separate group of 100 larvae for a rescue
300 attempt by placing larvae in stream water.

301 To determine activation energy (E_a) for water loss, water loss rates were determined (Eq.
302 3) with freshly killed larvae and temperature was varied. Larvae were killed by the freeze-thaw
303 method. Water loss rate was determined for the same larva experiencing a temperature increase
304 (ramp-up) or a temperature decrease (ramp-down). The Arrhenius equation (Eq. 5) was used to
305 determine the activation energy as described: $k = A e^{-E_a/(RT)}$, where k is passive water loss rate, E_a
306 is activation energy, R is universal gas constant, T is absolute temperature, and A is steric,
307 frequency factor (Gibbs, 2002). Significance of activation energy is controversial (Yoder et al.,

2005), but there is agreement on the occurrence of a critical transition temperature (CTT) when the activation energy changes (Gibbs, 2002). Water loss rate accelerates abruptly above the CTT (Gibbs, 2002). Each water loss rate determination was based on a cohort of 100 larvae in the activation energy calculation to total $N = 600$ larvae. The ramp-up and ramp-down experiment each utilized 100 larvae, tracking the water loss rate for the same cohort of 100 larvae going up, or down, the temperature scale.

314

315 **Determination of survival and attraction potential**

316 A cage for housing a larva was made with a 1.5 ml polypropylene microcentrifuge tube (Fisher) perforated with 30 holes to permit water entry into the cage. The cage was anchored with a spot of glue (Loctite low-odor; Henkel Co., Rocky Hill, CT, USA) onto a plain glass microscope slide (3" x 1" x 1mm; Fisher) to keep the cage submerged while in the bioassay arena (Fig. 1). The larva was out of test conditions for less than one minute for weighing and examination of motor activity, ability to self-right and crawl five body lengths at 40x. Percentage change in mass was calculated: (Eq. 5): $m = 100 (m_t - m_0)/m_0$, where m_t is the water mass at any time t and m_0 is the initial water mass. Treatments included placing the caged larva in non-moving water (0 flow rate) in sector one in the bioassay arena (Fig. 1), as well as in moving water with a flow rate of 3 L/min. We selected 3 L/min flow rate, because this flow rate produces a more localized flow in sector one with little to no flow in sector four compared to 6 L/min or 12 L/min that are too high. In the bioassay arena (Fig. 1), a comparison was done where the caged larva was placed in sector four ('indirect water flow'), thereby preventing the larva from being close to the source of moving water from the tubing located in sector one ('direct water flow'). Survivorship curves and mass measurements were done 20 times, five larvae at a time, each in a separate cage, to total $N = 100$.

332 Additionally, uncaged larvae were introduced, ten at time, at the center of the bioassay arena (Fig. 1). Counts of larvae in sector one (Fig. 1) were made after one hour, two hours, and 24 hours during the photophase. Treatments included exposure of larvae to 3 L/min, 6 L/min and 12 L/min water flows. There were two still water controls, one with the airflow tube in the water with 0 L/min airflow and one with no airflow tube in the water to rule out potential right-left bias. Data are the responses of 100 larvae, based on ten replicates of ten larvae.

338

339 Statistics

340 Alpha value was adjusted to a level of significance of 0.05. Arcsin transformation was done for
341 percentage data. Water balance data were compared with an analysis of covariance (ANCOVA).
342 Water loss rates and activation energies were compared with a test for the equality of slopes of
343 several regressions (Sokal and Rohlf, 1995). The number of larvae in sector one in the attraction
344 bioassays was analyzed by a means comparison, paired *t* test following an arcsin transformation
345 (Sokal and Rohlf, 1995). We did not want to have confounding effects if the survival data were
346 non-parametric, so we utilized the Kaplan-Meier survival curve with a log rank test after an
347 Abbott's correction (Sokal and Rohlf, 1995). Statistical software was SPSS 14.0 for Windows
348 (IBM, Armonk, NY, USA), Microsoft Excel (Redmond, WA, USA), and Minitab (Chicago, IL,
349 USA).

350

351 Acknowledgements

352 Not applicable.

353

354 Competing interests

355 Not applicable.

356

357 Author contributions

358 J. A.Y. conceived, designed and supervised experiments, collected and analyzed data, interpreted
359 the results, and wrote the manuscript. J. B. B., B. W. N. and L. R. Main carried out experiments,
360 collected specimens, analyzed data, and edited and proofread the manuscript. J. P. Bossley did
361 the initial taxonomic identification of the caddisfly larvae and was involved in proofreading the
362 manuscript. All authors have approved of this manuscript with confirmation in writing.

363

364 Funding

365 Funding for this project was provided, in part, by research grants to B. W. N. and L. R. M. from
366 Wittenberg University.

367

368 Supplementary material

369 Not applicable.

370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399

References

- Aggarwal, D. D., Ranga, P., Kalra, B., Parkash, R., Rashkovetsky, E. and Bantis, L. E.** (2013). Rapid effects of humidity acclimation on stress resistance in *Drosophila melanogaster*. *Comp. Biochem. Physiol. A.* **166**, 81-90.
- Allan, S. A. and Sonenshine, D. E.** (2002). Evidence of an assembly pheromone in the black-legged deer tick, *Ixodes scapularis*. *J. Chem. Ecol.* **28**, 15-27.
- Arlian, L. G. and Eckstrand, I. A.** (1975). Water balance in *Drosophila pseudoobscura*, and its ecological implications. *Ann. Entomol. Soc. Am.* **68**, 827-832.
- Arlian, L. G. and Staiger, T. E.** (1979). Water balance in the semiaquatic beetle, *Peltodytes muticus*. *J. Comp. Biochem. Physiol. A.* **62**, 1041-1047.
- Arlian, L. G. and Vyszynski-Moher, D. L.** (1995). Responses of *Sarcoptes scabiei* var. *canis* (Acari: Sarcoptidae) to lipids of mammalian skin. *J. Med. Entomol.* **32**, 34-41.
- Benoit, J. B., Yoder, J. A., Rellinger, R. J. Ark, J. T. and Keeney, G. D.** (2005) Prolonged maintenance of water balance by adult females of the American spider beetle, *Mezium affine* Boieldieu, in the absence of food and water resources. *J. Insect Physiol.* **51**, 565-573.
- Benoit, J. B., Lopez-Martinez, G., Michaud, M. R., Elnitsky, M. A., Lee, R. E., Jr. and Denlinger, D. L.** (2007). Mechanisms to reduce dehydration stress in the Antarctic midge, *Belgica antarctica*. *J. Insect Physiol.* **53**, 656-667.
- Benoit, J. B., Yoder, J. A., Lopez-Martinez, G., Elnitsky, M. A., Lee, R. E., Jr. and Denlinger, D. L.** (2008). Adaptations for the maintenance of water balance by three species of Antarctic mites. *Polar Biol.* **31**, 539-547.

- 400 **Bonada, N. and Williams, D. D.** (2002). Exploration of the utility of fluctuating asymmetry as
401 an indicator of river condition using larvae of the caddisfly *Hydropsyche morosa* (Trichoptera:
402 Hydropsychidae). *Hydrobiol.* **481**, 147-156.
- 403
- 404 **Bouchard, R. W., Jr.** (2004). Guide to Aquatic Macroinvertebrates of the Upper Midwest. St.
405 Paul: University of Minnesota.
- 406
- 407 **Bradley, T. J.** (1987) Physiology of osmoregulation in mosquitoes. *Ann. Rev. Entomol.* **32**, 439-
408 462.
- 409
- 410 **Bradley, T. J.** (2008) Saline-water insects: ecology, physiology and evolution. In *Aquatic*
411 *Insects: Challenges to Populations.* (ed. J. Lancaster and R. A. Biers), pp. 20-35. UK: CAB
412 International.
- 413
- 414 **deKozlowski, S. J. and Bunting, D. L., II.** (1981). A laboratory study on the thermal tolerance
415 of four southeastern stream insect species (Trichoptera, Ephemeroptera). *Hydrobiol.* **79**, 141-
416 145.
- 417
- 418 **Elnitsky, M. A., Benoit, J. B., Lopez-Martinez, G., Denlinger, D. L. and Lee, R. E., Jr.**
419 (2009). Osmoregulation and salinity tolerance in the Antarctic midge, *Belgica antarctica*:
420 seawater exposure confers enhanced tolerance to freezing and dehydration. *J. Exp. Biol.* **212**,
421 2864-2871.
- 422
- 423 **Gibbs, A. G.** (2002). Lipid melting and cuticular permeability: new insights into an old problem.
424 *J. Insect Physiol.* **48**, 391-400.
- 425
- 426 **Hadley, N. F.** (1994). Water Relations of Terrestrial Arthropods. New York: Academic Press.
- 427
- 428 **Kikawada, T., Saito, A., Kanamori, Y., Fujita, M., Snigórska, K., Watanabe, M. and**
429 **Okuda, T.** (2008). Dehydration-inducible changes in expression of two aquaporins in the
430 sleeping chironomid, *Polypedilum vanderplanki*. *Biochim. Biophys. Acta.* **1778**, 514-520.

- 431
- 432 **Kleynhans, E. and Terblanche, J. S.** (2011). Complex interactions between temperature and
433 relative humidity on water balance of adult tsetse (Glossinidae, Diptera): implications for climate
434 change. *Front. Physiol.* **2**:74.
- 435
- 436 **Kohn, P. G.** (1965). Tables of some physical and chemical properties of water. In *The State and*
437 *Movement of Water in Living Organisms* (ed. E. Fogg), pp. 291-318. New York: Cambridge
438 University Press.
- 439
- 440 **Merritt, R. W. and Cummins, K. W.** (1996). An Introduction to the Aquatic Insects of North
441 America. Dubuque: Kendall Hunt Publishing.
- 442
- 443 **Moulton, S. R., II, Beitinger, T. L., Stewart, K. W. and Currie, R. J.** (1993). Upper
444 temperature tolerance of four species of caddisflies (Insecta: Trichoptera). *J. Freshwater Biol.* **8**,
445 193-198.
- 446
- 447 **Parkash, R., Aggarwal, D. D., Kalra, B. and Ranga, P.** (2011). Divergence of water balance
448 mechanisms in two melanic *Drosophila* species from the western Himalayas. *Comp. Biochem.*
449 *Physiol. A.* **158**, 531-541.
- 450
- 451 **Parkash, R., Aggarwal, D. D., Singh, D., Lambhod, C. and Ranga, P.** (2013). Divergence of
452 water balance mechanisms in two sibling species (*Drosophila simulans* and *D. melanogaster*):
453 effects of growth temperatures. *J. Comp. Physiol. B.* **183**, 359-378.
- 454
- 455 **Sigal, M. D., Machin, J. and Needham, G. R.** (1991). Hyperosmotic oral fluid secretion during
456 active water vapour absorption and during desiccation-induced storage-excretion by the unfed
457 female tick *Amblyomma americanum*. *J. Exp. Biol.* **157**, 585-591.
- 458
- 459 **Snyder, C. D. and Hendricks, A. C.** (1995). Effect of seasonally changing feeding habits on
460 whole-animal mercury concentrations in *Hydropsyche morosa* (Trichoptera: Hydropsychidae).
461 *Hydrobiol.* **299**, 115-123.

- 462
- 463 **Sokal, R. R. and Rohlf, F. J.** (1995). *Biometry: The Principles and Practice of Statistics in*
464 *Biological Research*. San Francisco: W. H. Freeman.
- 465
- 466 **Sutcliffe, D. W.** (1961). Studies on salt and water balance in caddis larvae (Trichoptera): I.
467 Osmotic and ionic regulation of body fluids in *Limnephilus affinis* Curtis. *J. Exp. Biol.* **38**, 501-
468 519.
- 469
- 470 **Terblanche, J. S., Clusella-Trullas, S., Deere, J. A. and Chown, S. L.** (2008). Thermal
471 tolerance in a south-east African population of the tsetse fly *Glossina pallidipes* (Diptera:
472 Glossinidae): implications for forecasting climate change impacts. *J. Insect Physiol.* **54**, 114-127.
- 473
- 474 **Terblanche, J. S., Klok, C. J., Krafur, E. S. and Chown, S. L.** (2006). Phenotypic plasticity
475 and geographic variation in thermal tolerance and water loss of the tsetse *Glossina pallidipes*
476 (Diptera: Glossinidae): implications for distribution modelling. *Am. J. Trop. Med. Hyg.* **74**, 786-
477 794.
- 478
- 479 **Toolson, E. C.** (1978). Diffusion of water through the arthropod cuticle: thermodynamic
480 consideration of the transition phenomenon. *J. Therm. Biol.* **3**, 69-73.
- 481
- 482 **Wharton, G. W.** (1985). Water balance of insects. In *Comprehensive Insect Physiology,*
483 *Biochemistry and Pharmacology*, Vol. 4. (ed. G. A. Kerkut and L. I. Gilbert), pp. 565-603.
484 Oxford: Pergamon Press.
- 485
- 486 **Wiggins, G. B.** (1996). *Larvae of the North American Caddisfly genera (Trichoptera)*. Toronto:
487 University of Toronto Press.
- 488
- 489 **Yoder, J. A., Benoit, J. B., Rellinger, E. J. and Ark, J. T.** (2005). Letter to the Editors: Critical
490 transition temperature and activation energy with implications for arthropod cuticular
491 permeability. *J. Insect Physiol.* **51**, 1063-1065.
- 492

493 **Yoder, J. A., Tank, J. L., Brown, B. L. and Hobbs, H. H. III.** (2007). Water exchange
 494 pertaining to host attachment sites and stream preference in crayfish-associated branchiobdellids,
 495 *Cambarincola fallax* and *Cambarincola ingens* (Annelida: Clitellata). *Hydrobiologia* **592**, 523-
 496 533.

497

498 **TABLES**

499

500 **Table 1. Water balance characteristics of larvae of the caddisfly, *Cheumatopsyche pettiti***
 501 **and *Hydropsyche morosa*.** Values are the mean \pm s.e.m. ($N = 100$, ten replicates of ten larvae
 502 each or at each of the six test temperatures in the activation energy calculation).

503

| 504 Characteristic | <i>C. pettiti</i> | <i>H. morosa</i> | cf. Fig. |
|---|-------------------|------------------|----------|
| 506 Water content | | | |
| 507 Initial mass, f (mg) | 11.03 \pm 0.46 | 41.74 \pm 0.32 | |
| 508 Dry mass, d (mg) | 3.84 \pm 0.09 | 13.62 \pm 0.11 | |
| 509 Water mass, m (mg) | 7.19 \pm 0.36 | 28.12 \pm 0.44 | |
| 510 Water content, m/d (mg) | 1.87 | 2.06 | |
| 511 Water content (%) | 65.19 \pm 0.52 | 67.37 \pm 0.61 | |
| 512 Water loss | | | |
| 513 20°C (%/min) | 0.45 \pm 0.02 | 0.27 \pm 0.05 | 2 |
| 514 Dehydration tolerance | | | |
| 515 Water mass when moribund, m_c (mg) | 5.78 \pm 0.19 | 21.65 \pm 0.15 | |
| 516 Dehydration tolerance limit (%) | 19.61 \pm 0.49 | 23.02 \pm 0.62 | |
| 517 Critical transition temperature (°C) | 37.1 \pm 1.2 | 34.7 \pm 1.8 | 3 |
| 518 Activation energy, E_a (kJ/mol) < CTT | 14.5 \pm 0.5 | 9.1 \pm 0.7 | 3 |
| 519 Activation energy, E_a (kJ/mol) > CTT | 33.2 \pm 0.9 | 26.4 \pm 0.6 | 3 |

520

521

522 **FIGURE LEGENDS**

523

524

525 **Fig. 1. Experimental set up for exposing different water flow rates to larvae,**
526 ***Cheumatopsyche pettiti* and *Hydropsyche morosa*.** The airflow tube was fixed to an airflow
527 detector/regulator attached to a laboratory source of filtered compressed air. Photo credit: B.
528 Nelson and L. Main.

529

530 **Fig. 2. Proportion of water mass lost by larvae, *Cheumatopsyche pettiti* and *Hydropsyche***
531 ***morosa*, at 20°C, 0% RH.** Slope of the regression line is the water loss rate: m_t , water mass at
532 any time t , and m_0 , initial water mass. Error bars lie within confines of graph symbols (\pm s.e.m \leq
533 0.011). Each point is the mean of 100 larvae.

534

535 **Fig. 3. Water loss-temperature relationship for freshly killed larvae, *Cheumatopsyche pettiti***
536 **and *Hydropsyche morosa*.** Slope of the regression is $-E_a/R$ for calculating the activation energy:
537 K , absolute temperature; E_a , activation energy; R , gas constant. Error bars are within symbols
538 used on the graph (\pm s.e.m. \leq 0.004). Each point is the mean of 100 larvae.

539

540 **Fig. 4. Survivorship curves for starved, caged larvae of *Cheumatopsyche pettiti* and**
541 ***Hydropsyche morosa* exposed to different water flows using the set up as in Figure 1 (20°C,**
542 **15h:9h, L:D), with a flow rate of 3 L/min going into sector one.** Still, no water flow (0 L/min;
543 caged larva placed in sector one, but airflow turned off); direct, caged larva placed in sector one,
544 exposure to high water flow from airflow tube; indirect, caged larva placed in sector four with
545 airflow occurring in sector one. Each point is the mean of 100 larvae (\pm s.e.m. \leq 5.8).

546

547 **Fig. 5. Changes in body water mass under exposure to water flow (20°C, 15h:9h, L:D) for**
548 **larvae, *Cheumatopsyche pettiti* and *Hydropsyche morosa*, set up in the bioassay described by**
549 **Figure 1.** Flow rate was set at 3 L/min going into sector one. Larvae were caged during exposure
550 in sector one (direct), sector four (indirect), or sector one (still; no water flow). Larvae were
551 taken out of cages for mass determinations. Each point is the mean of 100 larvae (\pm s.e.m. \leq 2.1).

552

553 **Fig. 6. Attraction to sector one in response to water flows by larvae, *Cheumatopsyche pettiti***
554 **and *Hydropsyche morosa* (experimental set up is shown in Fig. 1; 20°C, 15h:9h, L:D).**

555 Untreated, no airflow tube was inserted into the water; 0 L/min, airflow tube inserted in water,
556 but airflow was turned off. Larvae were introduced at the center of the arena. Each test exposure
557 is the mean of 100 larvae (\pm s.e.m. \leq 4.3).











