

Microbiome symbionts and diet diversity incur costs on the immune system of insect larvae

Indrikis A. Krams^{1,2,3}, Sanita Kecko⁴, Priit Jõers⁵, Giedrius Trakimas^{4,6}, Didzis Elferts⁷, Ronalds Krams⁴, Severi Luoto^{8,9}, Markus J. Rantala¹⁰, Inna Inashkina¹¹, Dita Gudrā¹¹, Dāvids Fridmanis¹¹, Jorge Contreras-Garduño¹², Lelde Grantiņa-Ieviņa¹³, Tatjana Krama^{4,14}

¹ Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

² Department of Zoology and Animal Ecology, Faculty of Biology, University of Latvia, Rīga, Latvia

³ University of Tennessee, Department of Psychology, Knoxville, USA

⁴ Department of Biotechnology, Institute of Life Sciences and Technology, Daugavpils University, Daugavpils, Latvia

⁵ Institute of Molecular and Cell Biology, University of Tartu, Tartu, Estonia

⁶ Institute of Biosciences, Vilnius University, Vilnius, Lithuania

⁷ Department of Botany and Ecology, Faculty of Biology, University of Latvia, Rīga, Latvia

⁸ English, Drama and Writing Studies, University of Auckland, Auckland, New Zealand

⁹ School of Psychology, University of Auckland, Auckland, New Zealand

¹⁰ Department of Biology & Turku Brain and Mind Centre, University of Turku, Turku, Finland

¹¹ Latvian Biomedical Research and Study Centre, Riga, Latvia

¹² Escuela Nacional de Estudios Superiores Unidad Morelia, Universidad Nacional Autónoma de México, Morelia, Mexico

¹³ Institute of Food Safety, Animal Health and Environment BIOR, Rīga, Latvia

¹⁴ Department of Plant Protection, Institute of Agricultural and Environmental Sciences, Estonian University of Life Science, Tartu, Estonia

*Corresponding author indrikis.krams@ut.ee

Summary statement

This study suggests that having high numbers of midgut symbionts may bring costs to a host organism, which is a crucial finding for understanding the evolution of host-symbiont interactions.

ABSTRACT

Communities of symbiotic microorganisms that colonize the gastrointestinal tract play an important role in food digestion and protection against opportunistic microbes. Diet diversity increases the number of symbionts in the intestines, a benefit that is considered to impose no cost for the host organism. However, less is known about the possible immunological investments that hosts have to make in order to control the infections caused by symbiont populations that increase due to diet diversity. By using taxonomical composition analysis of the *16S rRNA* V3 region, we show that *Enterococci* are the dominating group of bacteria in the midgut of the larvae of the greater wax moth (*Galleria mellonella*). We found that the number of colony-forming units (CFU) of *Enterococci* and expressions of certain immunity-related antimicrobial peptide (AMP) genes such as *Gallerimycin*, *Gloverin*, *6-tox*, *Cecropin-D* and *Galiomicin* increased in response to a more diverse diet, which in turn decreased the encapsulation response of the larvae. Treatment with antibiotics significantly lowered the expression of all AMP genes. Diet and antibiotic treatment interaction did not affect the expression of *Gloverin* and *Galiomicin* AMP genes, but significantly influenced the expression of *Gallerimycin*, *6-tox* and *Cecropin-D*. Taken together, our results suggest that diet diversity influences microbiome diversity and AMP gene expression, ultimately affecting an organism's capacity to mount an immune response. Elevated basal levels of immunity-related genes (*Gloverin* and *Galiomicin*) might act as a prophylactic against opportunistic infections and as a mechanism that controls the gut symbionts. This would indicate that a diverse diet imposes higher immunity costs on organisms.

KEY WORDS: Antimicrobial peptides, Bacterial endosymbionts, Encapsulation response, Diet diversity, *Galleria mellonella*, Immunity

INTRODUCTION

Pathogens are a constant threat to organisms (Schmid-Hempel, 2011), and so they need a surveillance system that recognizes and attacks the intruders. These may consist of viral, bacterial, fungal and parasitic infections or even the body's own non-cooperative cells such as cancer. The innate immunity system is important because it starts acting immediately after a pathogen breaches the host's defense system (Janeway et al., 2005). The responses of the innate immunity system are more general than the more pathogen-specific responses of the adaptive immune system. The production of antimicrobial peptides (AMPs) is an early component of an innate immune response towards bacterial and fungal infections. AMPs are a group of antibiotics that impose a lethal effect against invading organisms by interfering with their basal biochemical functions (Zasloff, 2002; Brogden, 2005). Various AMPs modulate pathogen load and prevent the occurrence of an infection (Kaneko et al., 2007).

Immune function is expensive in terms of energy expenditure (e.g., Lochmiller and Deerenberg, 2000; Ardia et al., 2012), immunopathology (e.g., Råberg et al., 1998) and depletion of somatic resources (Adamo et al., 2008; Krams et al., 2012). This suggests that there are trade-offs in the allocation of resources between immune function and other components of fitness (e.g., Sheldon and Verhulst, 1996; Gonzalez-Santoyo and Cordoba-Aguilar, 2012). Access to food and the nutritional quality of food are of particular importance in trade-offs between life history traits and nutrients. For example, the nutritional state of the host and the nutritional quality of its diet may have a profound effect on life history trade-offs and the ability to resist an infection (Alonso-Alvarez and Tella, 2001; Moret and Schmid-Hempel, 2000; Ponton et al., 2013; Povey et al., 2013). This suggests that food limitation and a low quality diet are likely to compromise the host's immune response (Smits et al., 2017). However, different levels of food limitation may produce different immune response patterns (e.g., Krams et al., 2014). For example, brief food deprivation may lead to an increase in the expression of AMP genes in the fruit fly *Drosophila melanogaster*, even in the absence of pathogens (Becker et al., 2010). This response is the opposite of what would be predicted if the immune system gradually declined as resources dwindle. Moreover, recent research shows that intracellular immune signaling pathways are interconnected with nutrient signaling pathways (e.g. in invertebrates, Becker et al., 2010; Adamo et al., 2016; in vertebrates, Odegaard and Chawla, 2013), providing an evolutionary explanation for the complex effects of food deprivation on immunity.

Epithelial cells of the gastrointestinal tract are covered with a large population of normally nonpathogenic bacteria. A fundamental part of digestion, these nonpathogenic bacteria are known as ‘the microbiota’. Associations between the host and its microbiota are often mutualistic, which facilitates the establishment of persistent colonization of host organs and provides a variety of benefits to the hosts (Kau et al., 2011; Huang and Douglas, 2015; Masson et al., 2016). The microbiota competes with opportunistic microbes for nutrients and for attachment sites on epithelial surfaces by producing various antimicrobial substances. When the microbiota is affected by antibiotic treatment, pathogens may replace mutualistic microorganisms and cause diseases such as gastroenteritis, metabolic imbalance, inflammatory bowel disease and colorectal cancer (see Garrett et al., 2010). Moreover, the host’s failure to regulate the survival and numbers of mutualistic microorganisms can allow normally nonpathogenic bacteria to grow excessively and also cause disease. A balance between bacterial growth and efficiency of nutrient processing in the gut and intestines is maintained by immunity mechanisms, such as generation of reactive oxygen species (De Block and Stoks, 2008; Jones et al., 2012) and AMPs (Login et al., 2011; Johnston and Rolff, 2015; Makarova et al., 2016). Overall, the composition of gut microbiomes is structured by (the host’s) diet (Muegge et al., 2011). Some studies show that diet diversity correlates positively with microbiome diversity, leading the microbiome to be more adaptable to perturbations (reviewed in Heiman and Greenway, 2016, but see Bolnick et al., 2014). Previous studies on Lepidoptera (Broderick et al., 2004; Xiang et al., 2006; Pinto-Tomás et al., 2011; Gayatri Priya et al., 2012; Mason and Raffa, 2014) confirm the positive relationship between diet diversity and larval midgut community composition. The prediction arises that food limitations potentially decrease microbiome diversity. This may relieve the organism’s investment in immune mechanisms of microbiota control in general and expression of AMP genes in particular, a factor overlooked in eco-immunological research.

It has been demonstrated that dysbiosis caused by the oral delivery of antibiotics eliminates the midgut microbiome of the larvae of the greater wax moth (*Galleria mellonella*) and decreases the transcriptional activation of AMPs (Mukherjee et al., 2013). In this study, we expected that AMP genes in the midgut of greater wax moth larvae would be less expressed when grown on simple food of low nutritional value than when grown on macronutrient/energy-rich food. We also fed larvae with an antibiotic cocktail to test whether AMP genes’ expression during dysbiosis is similar to the transcriptional activation of AMPs of larvae raised on nutrient-poor food.

In *G. mellonella*, *Enterococci mundtii* (syn. *Streptococcus faecalis* Andrewes and Horder) is considered to be the dominating bacterium found in the midgut of the host (Jarosz, 1979; Johnston and Rolff, 2015). The dominance of just one microorganism can be explained by the antibacterial properties of their larval food, which consists of bee-wax and honey, and the ability of *E. mundtii* to survive under these conditions. This microbe is a heritable nutrient-providing symbiont that is transmitted vertically – from mother to offspring (Chen et al., 2016) – in *G. mellonella* (Bucher, 1963; Johnston and Rolff, 2015). We analyzed the V3 region of 16S rDNA to determine the taxonomical composition of microorganisms in the midgut samples of the larvae of *G. mellonella*. Using conventional bacterial culturing, we investigated the numbers of *Enterococci* in the midgut microbiota of the larvae grown either on diverse or simple food, and either force-fed with antibiotics or receiving no antibiotic treatment (control group). Food-related stress has been shown to affect the arms of the immune system via an antagonistic cross-regulation between different components of the immune system (Krams et al., 2012). To find any possible costs of AMP gene upregulation in *G. mellonella*, we tested whether higher levels of AMP gene expressions suppress encapsulation ability (Kangassalo et al. 2016; Krams et al. 2016; Kecko et al. 2017), an important defence mechanism against parasitic protozoans and metazoans, fungi, mites and parasitoids such as wasp eggs or larvae (Gillespie et al., 1997; Lavine and Strand, 2002; Krams et al., 2017).

MATERIALS AND METHODS

Insects and food quality

We studied a captive population of *G. mellonella* consisting of individuals collected from natural populations in Estonia in summer 2014. Moths were reared in 2.4 litre plastic boxes at $28 \pm 1^\circ\text{C}$ in the dark in Sanyo MIR-253 incubators. To study the effects of diet diversity on the expression of AMP genes of *G. mellonella* larvae, we assigned them to groups differing in the macronutritional diversity and energetic value of food. Each larva was placed individually into a plastic container (50 ml) with a lid and wire-mesh to allow ventilation and to prevent it from escaping. The larvae were kept on contrasting diets between hatching and post-hatch day 25 (Fig. 1).

The ‘diverse diet’ group received only food of high nutritional value and diversity, provided *ad libitum* (Fig. 1). The ‘simple diet’ group received only food of low nutritional value and diversity, provided *ad libitum*. The diverse diet consisted of a mix of equal proportions of honey, glycerol, bee-wax, dried milk, wheat flour, dry yeast, distilled water and two servings of corn meal. Foods in both diets were autoclaved at 121°C for 15 min. Under natural conditions, increased diet diversity typically means that a higher diversity of opportunistic microorganisms is ingested by a host. This was not the case in our study, however, since the larvae received bacteria-free food. The amount of energy associated with this food was estimated as c. 16.90 kJ/g by a combustion calorimeter C 2000 basic (IKA® - Werke GmbH & Co. KG, Germany). The ‘simple diet’ consisted of natural bee-wax with a 5% admixture of corn meal. Bee-wax is a natural polymer produced by bees, and it is considered to have an extremely low nutritional value. However, we observed the ability of some wild progenitors of our study population to reproduce solely on bee-wax. Thus, the larvae of the simple food group received a slightly better food than pure wax.

On post-hatch day 26, one half of the larvae in both food groups received peroral antibiotics to eliminate their microbiota. These individuals were treated with antibiotics and antifungal drugs, which was done twice within an 8-hour interval. We force-fed each larva with 10 µl of antibiotic cocktail consisting of water-soluble forms of ampicillin (Sigma-Aldrich, U.S.A.) (code: A5354, concentration 50 mg/ml; Gram-positive bacteria including *Streptococcus pneumonia*, *S. pyogenes*, *Staphylococcus aureus*, *Enterococci*, and Gram-negative bacteria including *Enterobacteriaceae*), erythromycin (Sigma-Aldrich, U.S.A.) (code: 45703, concentration 1 mg/ml; Gram-positive bacteria: *Streptococci*, *Staphylococci* and Gram-negative bacteria and some fungi), gentamicin (Sigma-Aldrich, U.S.A.) (code: G-1397 code, concentration 50 mg/ml; Gram-negative bacteria including *Pseudomonas*, *Escherichia coli*, *Enterobacter aerogenes*, *Proteus* spp., *Serratia marcescens*, *Klebsiella pneumonia* and Gram-positive *Staphylococcus*), kanamycin (Sigma-Aldrich) (code: K0254, concentration 50 mg/ml; Gram-negative bacteria including *E. coli*, *Proteus* spp., *S. marcescens*, *K. pneumonia*), and ketoconazole (Sigma-Aldrich, U.S.A.) solution in dimethyl sulfoxide (code: K0600000, antifungal agent given in concentration of 1 µg/larva). The antibacterial cocktail has previously been documented to significantly decrease the expression of AMP genes and stress-related genes in *G. mellonella* (Mukherjee et al., 2013).

We extracted RNA from the larvae 24 h after they received the first dose of antibiotics. We also extracted RNA from the larvae raised on diverse and simple foods that did not receive any antibiotics – these individuals were force-fed 10 µl of distilled water. In total, we had four experimental groups and one reference group of the greater wax moth. The experimental groups were as follows: (1) a control group raised on a diverse diet, (2) a control group raised on a simple diet, (3) an antibiotic treatment group raised on a diverse diet, (4) an antibiotic treatment group raised on a simple diet. The larvae of the reference group were raised on a simple diet. These larvae were force-fed 10 µl of distilled water every 3 h and received the antibiotics cocktail three times within a 12-hour interval to eliminate their microbiome. RNA was extracted from these larvae 24 h after they received the first dose of antibiotics.

RNA extraction and quantitative real-time PCR of immunity-related gene expression

Collected larvae were chilled on ice for 15 min, surface sterilized with 70% ethanol and then dissected to collect their midguts. We pooled midguts from ten individual larvae for each group. RNA was obtained from five replicates of each of the four groups (200 larvae in total). The midguts were homogenized in 1 ml of Trizol reagent (Sigma-Aldrich, U.S.A.), and RNA was extracted according to the manufacturer's recommendations. RNA integrity was confirmed by ethidium bromide gel staining, and quantities were determined spectrophotometrically.

Levels of steady-state transcripts were determined from cDNA samples by real-time quantitative RT-PCR using $\Delta\Delta C_t$ protocol with the 7500 Real-Time PCR System (Applied Biosystems, U.S.A.) and SYBR Green PCR mix (Qiagen, The Netherlands). First-strand cDNA was synthesized using 5 µg of RNA, random hexamers and RevertAidTM M-MuLV Reverse Transcriptase (Fermentas, Lithuania) according to the manufacturer's protocols, and subsequently diluted with nuclease-free water (Fermentas, Lithuania) 10 to 100 times. Amplification mixtures (25 µl) contained 2µl template cDNA, 2xSYBR Green Master Mix buffer (12,5 µl) (Life Technologies, U.S.A.) and 2µM forward and reverse primer.

Five target genes were investigated, coding for AMPs: *Gloverin* (strong activity against gram-positive bacteria and weak activity against Gram-negative bacteria) (forward primer: AGATGCACGGTCCTACAG, reverse primer: GATCGTAGGTGCCTTGTG), *Gallerimycin* (strong effect against filamentous fungi) (forward primer:

GAAGTCTACAGAATCACACGA, reverse primer: ATCGAAGACATTGACATCCA) (Schuhmann et al., 2003), *6-tox* (an atypical defensin-derived immune-related peptide expressed in midgut against invading bacteria) (forward primer: GACGAACTGCGAAGAATTATC, reverse primer: TGTCTGTCTTGAGTTGCATATTG) (Lee et al., 2010), *Galiomycin* (strong antifungal effect and limited effect against bacteria) (forward primer: GTGCGACGAATTACACCTC, reverse primer: TACTCGCACCAACAATTGAC) and *Cecropin-D* (strong activity against Gram-negative bacteria and fungi, weak activity against Gram-positive bacteria) (forward primer: CTGCGCCATGTTCTTCA, reverse primer: TCGCATCTCTGATCCTCTG) (Melo et al., 2013).

Cycling conditions were 95°C for 5 min, followed by 42 cycles of: 95°C 5 sec, annealing 10 sec, 72°C 20 sec. An initial touchdown of 1°C per cycle from 65°C for the first 5 cycles resulted in optimal amplification for all loci. High Resolution Melting analysis performed at the end of each run allowed each PCR to be checked for the presence of the expected product. We are confident that the efficiency was between 98.0% and 100% for all primer pairs. Every cDNA sample was normalized against two housekeeping genes: *18S rRNA* (AF286298; forward primer: CACATCCAAGGAAGGCAG, reverse primer: AGTGTACTCATTCCGATTACGA) and translation elongation factor 1-alpha (*EF1a*; AF423811; forward primer: AACCTCCTTACAGTGAATCC, reverse primer: ATGTTATCTCCGTGCCAG) (Vogel et al., 2011) using geNorm software (Vandesompele et al., 2002). The calculated gene expression stability coefficient *M* was applied to the assay results of the genes of interest. The primers were obtained from Metabion International AG (Planegg, Germany).

***16S V3 rRNA* gene amplification and Ion ProtonTM semiconductor sequencing**

The larvae used for gene sequencing received the same food as the ‘diverse diet’ group, matching the conditions of *ad libitum* food availability in natural bee hives. DNA was extracted using Fast DNA Spin Kit for Soil (MP Biomedicals, LLC, U.S.A.) according to the manufacturer’s guidelines. The concentration of extracted DNA was measured using Qubit® 2.0 Fluorometer High Sensitivity Assay (Life Technologies, USA). The amount, average size and quality of the DNA were assessed using electrophoresis in 1.2% agarose gels. DNA was obtained from six laboratory populations and we used the midguts of six individual larvae per each sample (36 larvae in total).

rRNA V3 region was amplified separately by reverse (Probio_Uni_R 5'-ATTACCGCGGCTGCT-3') and forward (Probio_Uni_F 5'-CCTACGGGRSGCAGCAG-3') primers (Milani et al., 2013). Both primers were tagged with 10-11 bp unique barcode labels along with the adapter sequence (5'-CCATCTCATCCCTGCGTGTCTCCGAC-3'). PCR amplification was performed by GeneAmp® PCR System 9700 (Thermo Fisher Scientific, U.S.A.) with the PCR conditions used as follows: 98°C for 30 sec, 35 cycles of 98°C for 10 sec, 67°C for 15 sec, 72°C for 15 sec with the final extension at 72°C for 7 minutes. *16S rRNA* PCR products were then quantified, pooled and purified using NucleoMag® NGS Clean-Up and Size Select kit (Macherey-Nagel, Germany). The quality and acquired amount of *16S rRNA* V3 amplicons was assessed by Agilent High Sensitivity DNA kit on Agilent 2100 BioAnalyzer (Agilent Technologies, U.S.A.).

Prior to clonal amplification, each library was diluted to 8 pM and pooled. Sample emulsion PCR, emulsion breaking and enrichment were performed using the Ion PI™ Hi-Q™ OT2 Kit (Life Technologies, U.S.A.), following the manufacturer's instructions. The complete sample was loaded onto a PI™ chip v3 and sequenced on the Ion Proton™ Semiconductor Sequencer for 520 cycles employing the Ion PI™ Hi-Q™ Sequencing 200 Kit. Bidirectional sequencing was performed (i.e. sequence reads started from forward and reverse PCR primers), but reads were not paired. Each run was expected to produce approximately 240,000 reads. After the sequencing run was completed, the individual sequence reads were filtered by the Proton software to remove low quality sequences. Sequences matching the Proton 3' adaptor were automatically trimmed. All Proton quality-approved, trimmed and filtered data were exported as bam files.

Sequencing data analysis was carried out using QIIME v.1.8.0. and UPARSE v.7.0.1001 pipeline to quality-filter and cluster 16S rRNA amplicon sequences (Pylro et al., 2014). Quality control retained sequences with the mean sequence quality score >20. Operational Taxonomic Units (OTUs) were built at 97% sequence identity with uclust (Edgar, 2010). Taxonomic assignment to the lowest possible rank was performed with RDP (Wang et al., 2007), using the Greengenes (DeSantis et al., 2006) (<http://greengenes.secondgenome.com>) reference dataset (gg_otus-13_8 release). Alpha diversity measure – Shannon diversity index was calculated within the QIIME environment.

Conventional bacterial culturing

Midguts from six individual larvae were dissected and pooled for each of 20 replicates in four groups (480 larvae in total) and used for microbiological analysis. We homogenized 0.1 ml of the midgut samples with 1 ml of a sterile 0.1% peptone water solution using a laboratory paddle sampler for 3 min. Serial dilutions of the midgut homogenate were prepared with sterile peptone water and plated in duplicates on the Bile Esculin Azide Agar, a selective *Enterococcus* agar (Sigma-Aldrich, U.S.A.). *Enterococci* were further determined by the BBL Crystal Identification Systems Gram-positive ID KIT (Becton, Dickinson and Company, U.S.A.) (e.g., Von Baum et al., 1998).

Immune assays

The strength of encapsulation response was measured in the control larvae raised on (1) the ‘diverse diet’ (n = 28) and (2) the ‘simple diet’ (n = 27). We also used *G. mellonella* larvae of the antibiotic treatment group raised on (3) the ‘diverse diet’ (n = 29) and (4) the ‘simple diet’ (n = 26). However, these were different larvae than those used for RNA extraction and conventional bacterial culturing.

The immune treatment *G. mellonella* larvae on post-hatch day 25 consisted of an immune system challenge induced with a disinfected nylon monofilament implant (2 mm length, 0.18 mm diameter, knotted at one end) which was inserted between the 3rd and 4th sternite of the cuticle of the larvae (Krams et al., 2014; Kecko et al. 2017) for 10 h at 28 ± 0.5 °C. The implant was removed after this treatment. The strength of the encapsulation response was measured as the darkness of the nylon filament insert after it had been dried. Insect immune systems respond to this challenge by attempting to encapsulate the implant in a coating of cellular materials and chemical deposits as if the insert were a foreign body / parasite (e.g., Rantala *et al.*, 2000). The stronger the immune response to the insert, the darker the encapsulation and the higher its grayscale value (Yourth et al., 2001; Krams et al., 2011, 2017; Daukste et al. 2012) due to phenoloxidase enzyme production activated by the immune response, resulting in melanization of the capsule (Ratcliffe *et al.*, 1985). In the greater wax moth, the encapsulation response has been confirmed to be associated with parasitoid (Kryukova *et al.*, 2011), fungal (Dubovskiy *et al.*, 2011, 2013) and bacterial infections (Grizanova *et al.*, 2014). Lightness was assessed from photographs of the inserts taken from three directions under constant light conditions using a Zeiss Lumar V12 Stereo (Carl Zeiss, Jena, Germany) microscope. Digital images were analyzed using the Image J

software (<http://rsbweb.nih.gov/ij/>; Abramoff *et al.*, 2004). Before the insertion of the monofilament implant, we calibrated the reflectance of the implant to zero level.

Data analysis

As the Brown-Forsythe test showed a violation of the homogeneity of variances of gene expressions and encapsulation rates, we used generalized linear models (GLM) with gamma error structure and an inverse link function to compare gene expressions and encapsulation rates. Diet, treatment and their interaction were used as independent variables. If an interaction showed a significant effect, the Games-Howell post hoc test was used to account for unequal variances. The analysis was performed using software R 3.3.2 (R Core Team 2016) libraries' onewaytests (Dag *et al.*, 2016) and userfriendlyscience (Peters, 2016). We compared the *Enterococci* CFU in diverse and simple diet groups (without antibiotic treatment) using t-test. Data are expressed as the mean \pm SD where applicable.

RESULTS

The GLM analysis showed that expressions of all the AMP genes were significantly higher in the 'diverse diet' group than in the 'simple diet' group (*Gallerimycin*: GLM, LR Chisq₁ = 235.709, $P < 0.0001$; *6-tox*: GLM, LR Chisq₁ = 40.384, $P < 0.0001$; *Galiomicin*: GLM, LR Chisq₁ = 22.491, $P < 0.0001$; *Cecropin-D*: GLM, LR Chisq₁ = 171.380, $P < 0.0001$; *Gloverin*: GLM, LR Chisq₁ = 23.858, $P < 0.0001$). Treatment with antibiotics was associated with significantly lower expressions of all AMP genes compared with groups without antibiotics treatment (*6-tox*: GLM, LR Chisq₁ = 77.948, $P < 0.0001$; *Cecropin-D*: GLM, LR Chisq₁ = 201.231, $P < 0.0001$; *Gallerimycin*: GLM, LR Chisq₁ = 288.633, $P < 0.0001$; *Galiomicin*: GLM, LR Chisq₁ = 61.111, $P < 0.0001$; *Gloverin*: GLM, LR Chisq₁ = 29.028, $P < 0.0001$). Diet and treatment interaction showed significant effect on expression of *Gallerimycin* (GLM, LR Chisq₁ = 7.322, $P = 0.0068$), *6-tox* (GLM, LR Chisq₁ = 5.068, $P = 0.0244$) and *Cecropin-D* (GLM, LR Chisq₁ = 6.983, $P = 0.0082$) gene expression while *Gloverin* (GLM, LR Chisq₁ = 3.260, $P = 0.071$) and *Galiomicin* (GLM, LR Chisq₁ = 0.438, $P = 0.5079$) gene expression did not differ significantly between interaction groups, suggesting that diet alone has the potential to increase the expression of these two genes.

Post hoc tests revealed that the AMP genes were expressed at significantly higher levels in the ‘diverse diet’ group (mean₁ ± SD) than in the ‘simple diet’ group (mean₂ ± SD) when the larvae of *G. mellonella* were not treated with antibiotics (*Gallerymicin*: mean₁ ± SD = 16.96 ± 3.88 vs mean₂ ± SD = 1.15 ± 0.2, *P* = 0.003; *6-tox*: mean₁ ± SD = 8.08 ± 1.12 vs mean₂ ± SD = 0.82 ± 0.05, *P* < 0.001; *Cecropin-D*: mean₁ ± SD = 11.46 ± 2.27 vs mean₂ ± SD = 0.30 ± 0.07, *P* = 0.001; Fig. 1). However, we did not find any significant differences between the ‘diverse diet’ group under antibiotics treatment and ‘simple diet’ group under antibiotics treatment in expressions of the AMP genes (all *P* > 0.05, Fig. 1A-C). In general, the AMP genes were similarly expressed between the ‘simple diet without antibiotics treatment’ group and the ‘diverse diet’ and ‘simple diet’ groups with antibiotics treatment (*P* > 0.05). The exception was *6-tox* which was expressed significantly more in the ‘simple diet without antibiotics’ group than in the ‘diverse diet with antibiotics’ group (*P* = 0.006, Fig. 1). Expressions of *Cecropin-D*, *6-tox* and *Gallerymicin* genes in the ‘diverse diet without antibiotics’ group were significantly higher than those in the diverse and simple diet groups with antibiotics (all *P* < 0.005, Fig. 1).

Sequencing the midgut samples of *G. mellonella* by Ion Proton™ semiconductor sequencer resulted in 248,073 sequences per one sample, but the number of sequences retained after the quality control was 177,742. Taxonomical composition analysis of the *16S rRNA* V3 region revealed that the most prevalent genus in the microbial community associated with the midgut of *G. mellonella* larvae was *Enterococcus* (c. 80% of the sequences), while the relative abundance of the family *Enterococcaceae* accounted for 87% of microbiome (Supplementary material).

The highest number of CFU of *Enterococci* was found in the ‘diverse diet without antibiotics’ group ($7.6 \times 10^6 \pm 14.70 \times 10^6$ CFU/ml; mean ± SD), while the number of *Enterococci* CFU was significantly lower in the ‘simple diet’ group ($0.8 \times 10^3 \pm 1.5 \times 10^3$ CFU/ml; mean ± SD) (t-test: $t(38) = 2.31$, *P* = 0.027). We did not find any bacteria in the midgut of the individuals in the antibiotics treatment groups grown on diverse and simple diets (Fig. 2).

The encapsulation response was affected by diet (GLM, LR Chisq₁ = 371.42, *P* < 0.0001), treatment (GLM, LR Chisq₁ = 24.78, *P* < 0.0001) and interaction between diet and treatment (GLM, LR Chisq₁ = 7.32, *P* = 0.007). The Games-Howell post hoc test showed that there are significant differences (all *P*s < 0.01, Fig. 3) between the ‘simple diet without antibiotics treatment’ group (mean ± SD = 29.04 ± 2.47), the ‘simple diet with antibiotics

treatment' group (mean \pm SD = 35.88 \pm 4.26), the 'diverse diet without antibiotics' group (mean \pm SD = 19.50 \pm 3.17) and the 'diverse diet with antibiotics' group (mean \pm SD = 19.76 \pm 3.16). The only exception occurred between the 'diverse diet without antibiotics' and the 'diverse diet with antibiotics treatment' groups which did not differ significantly ($P > 0.05$, Fig. 3).

DISCUSSION

Alpha diversity measurements of the relative abundance of bacteria have shown that *Enterococci* are the most common group of microorganisms in the midgut microbiome of *G. mellonella* (Jarosz, 1979; Johnston and Rolff, 2015). Our results confirm those findings and support other findings on how the elimination of nutrients reduces investments in the immune system (Alonso-Alvarez and Tella, 2001; Krams et al., 2014), while also reducing symbiont numbers and microbiota diversity (David et al., 2014; Carmody et al., 2015; Sonnenburg et al., 2016). This highlights the importance of food diversity and quality in shaping intestinal microbiota, which may have profound effects on organismal growth trajectories and the evolution of reproductive trade-offs (Lazzaro and Rolff, 2011).

The results of this study show that when force-fed with antibiotics at larval stage, *G. mellonella* developed basal expression levels of immunity-related AMP genes. We suggest that these levels reflect 'surveillance' activity of the immune system in the midgut in the absence of symbionts. Interestingly, the basal expression of *Cecropin-D*, *Gallerimycin* and *6-tox* of larvae force-fed with antibiotics did not differ from expressions of those genes under conditions when the larvae received a simple diet without antibiotics. This shows that investment in the production of AMPs remains the same between conditions of microbial dysbiosis and the diet consisting of simple/low-energy food despite the difference in symbiont numbers in the midgut. This result suggests the leading role of diet played in the expression of AMPs. In contrast, the AMP genes *6-tox*, *Cecropin-D*, *Galiomicin*, *Gallerimycin* and *Gloverin* were all significantly upregulated in the group of larvae grown on a diverse diet – those harboring the highest number of *Enterococci* symbionts. Hence, the elevated gene expressions correlated with the increased diet diversity and the number of *Enterococci* symbionts as revealed by conventional bacterial culturing. This positive association between the increased number of symbionts and elevated immune-related gene expression shows that diet quality/diversity may result in a higher number of symbionts

involved in the digestion of nutrients, and this presumably requires more control over symbionts by means of AMP proteins.

The gastrointestinal microbiome is considered to be more diverse in the intestines of healthy organisms, while a loss in species diversity is a common finding in several disease states (for a review, see Heiman and Greenway, 2016). Symbionts may, however, become harmful if they grow and reproduce uncontrollably. Symbionts may therefore consume more nutrients than would be normally expected to maintain symbiotic relationships, causing bacterial overgrowth in intestines/midgut if not properly controlled by the immune system of the host (Tamboli et al., 2004; Erdogan and Rao, 2015; Fujimori, 2015; Moos et al., 2016). For example, insects are known to use protective bacterial symbionts to provide additional defense against their predators and parasitoids (Oliver et al., 2010; Polin et al., 2014). However, these symbioses have been shown to induce costs. For example, while bacterial symbionts provide the pea aphid (*Acyrtosiphon pisum*) strong protection against parasitoids, this symbiosis becomes costly in the presence of predatory ladybirds because aphids infected by protective symbionts express less defensive anti-predator responses (Polin et al., 2014). Luo et al. (2017) have recently investigated the ecological effects of a non-essential secondary endosymbiont *Regiella insecticola* on different clones of the English grain aphid (*Sitobion avenae*) based on the life table data. They showed that *R. insecticola* can decelerate the normal development of the hosts. Similarly, our results suggest that high numbers of symbiotic *Enterococci* bacteria (Johnston and Rolff, 2015) are involved in elevation of costly expressions of immune-related genes (for example, *Gloverin*), an outcome that may be further influenced by the presence of other microorganisms undetected / not evaluated in our study. The ‘surveillance’ of symbionts and the activation and production of AMPs in cases when the symbionts become too abundant are important functions. These functions might incur costs, since the larvae would need to divert resources and energy away from growth and other organismal needs in order to carry out those functions. A fine-grained analysis of these potential trade-offs requires a more detailed experimental approach in future studies.

Our findings are indirectly aligned with previous studies concerning the role of host immunity in shaping microbiota. The gut microbiome often represents a complex and highly dynamic biological system (e.g., Lanan et al., 2016). Symbiont communities within the gut lumen may be unstable due to the constant influx of food particles (Blum et al., 2013) and invasions of other non-symbiotic food-associated bacteria (Jones et al., 2013; Cariveau et al., 2014). Thus, the number of bacteria and the diversity of insect microbiome can be explained by bacterial competition in the midgut (Greenberg and Kandlowden, 1972). Another

hypothesis suggests that the immune system of the host plays an important role in the formation and maintenance of bacterial assemblages in the midgut of insects (Russell and Dunn, 1996; Caccia et al., 2016). More recent work shows that host and symbiont communities cooperatively interact to maintain the midgut microbiota in a symbiotic balance (Johnston and Rolff, 2015). Our results suggest that all of the above ideas are worth testing by manipulating the levels of food resources available to the host, which we found to be an important agent affecting symbiont numbers and the strength of AMP gene expression.

Importantly, our results suggest that food diversity has the potential to influence the immune system, as shown by the elevated expression of *Gloverin* (responsible for suppressing Gram-positive bacteria such as *Enterococci*) and *Galiomicin* AMP genes. This may be adaptive under natural conditions because the larvae of *G. mellonella* often invade only hives with low honey bee populations or hives where bees have already died from a disease (Barjac and Thomson, 1970). However, the hives are not kept as clean as those maintained by healthy bee colonies. This suggests that the food of *G. mellonella* is often contaminated by various bacteria and fungi. Thus, a diverse diet in nature might indicate a higher probability of acquiring opportunistic infections, while an elevated expression of *Gloverin* and *Galiomicin* AMP genes may be considered to be a prophylactic response to pathogens (Barnes and Siva-Jothy, 2000).

It has recently been shown that a high-energy/diverse diet decreases the strength of encapsulation response, which indicates that food quality is the primary determinant of encapsulation response in the larvae of *G. mellonella* (Krams et al., 2014). Since the larvae grow the fastest when raised on a high-quality diet, it was suggested that longer larval development on low-quality diet requires enhanced encapsulation reactions (Krams et al., 2014). The present study, however, reveals one more mechanism that may link the diverse / high quality diet with weaker encapsulation responses in *G. mellonella*. We show that the elevated expression of all AMP genes is associated with the diverse diet and that this result was independent of the antibiotics treatment. Interestingly, the encapsulation response was stronger in the 'simple diet with antibiotics treatment' group, which indicates that the encapsulation response is dependent also on the presence of bacteria, but only when grown on the 'simple diet'. This shows that the 'diverse diet' itself provides a signal of possible bacterial contamination which decreases encapsulation rate and elevates the AMP protection because the latter is more important in fighting opportunistic infections. However, what needs to be investigated in the future is whether the relationship between the production of AMP peptides and the strength of encapsulation response is a result of stress-induced

suppression of humoral immune response manifesting through an antagonistic cross-regulation between different components of the immune system (Sapolsky et al., 2000; Dhabhar, 2009) or whether the encapsulation response and AMP gene expressions are a part of the same protective mechanism. For example, Nagai et al. (2000) found that Tachyplestin, a major chitin-binding AMPs of the horseshoe crab (*Tachypleus tridentatus*), induce the synthesis of phenoloxidase which is a crucial part of the host defense system including the synthesis of melanin and facilitation of encapsulation response. Phenoloxidase, a copper-containing enzyme, is known to be responsible for initiating the biosynthesis of melanin which is used during encapsulation response (e.g., Hoffmann et al. 1999).

In conclusion, while symbiotic interactions are supposed to be mutually beneficial to the host and its symbionts, this study suggests that high numbers of midgut symbionts may induce ecological costs to their insect hosts. This conclusion is based on the finding that a diverse diet (and, consequently, a higher number of microbiome symbionts) activates the part of the immune system of *G. mellonella* larvae which is responsible for the production of AMPs. More specifically, we found that the CFU of *Enterococci* and expressions of *Gallerimycin*, *Gloverin*, *6-tox*, *Cecropin-D* and *Galiomicin* increased in response to a more diverse diet. However, quantifying and revealing the exact mechanisms of these costs as a part of ecological trade-offs remain to be done in future research. This task will require studies to disentangle the possible effects caused by a diverse diet (Huang and Douglas, 2015). This can be accomplished by eliciting the prophylactic activation of the immune system and analyzing the relationship between diet diversity and the number of symbionts, which seemingly needs to be controlled by the host. Finally, food availability is an important growth determinant not only for the host but also for its symbionts. Since the availability of food resources has the potential to affect symbiont numbers and microbiota species composition, these effects need to be taken into account while designing and performing future experiments. This type of work is crucial in order to increase our understanding of the ecological effects that food resources have on immunity/growth/reproduction trade-offs under the immense complexity of food webs and their dynamic interactions with the microbiota.

Acknowledgements

We thank Dr. Inese Kivleniece and Dr. Jolanta Vrublevska for maintaining a laboratory stock of greater wax moth.

Competing interests

The authors declare no competing or financial interests.

Author contributions

I.A.K., S.K., G.T., M.J.R., I.I., D.G., D.F. and L.G-I. designed the study; I.A.K., S.K., R.K., G.T., T.K. and J.D performed research; I.A.K., S.K., P.J., I.I., D.G., D.F., D.E., L.M. and J.C-G. analyzed and interpreted data; I.A.K., S.K., D.E., S.L., and J.C-G. wrote the manuscript.

Funding

The study was funded by a grant from the Latvian Council of Science (# 290/2012) to SK, TK, IAK. The Estonian Research Council supported TK, IAK (PUT1223) and PJ (PUT573).

References

- Adamo, S. A., Roberts, J. L., Easy, R. H. and Ross, N. W.** (2008). Competition between immune function and lipid transport for the protein apolipoprotein III leads to stress-induced immunosuppression in crickets. *J. Exp. Biol.* **211**, 531–538.
- Adamo, S. A., Davies, G., Easy, R., Kovalko, I. and Turnbull, K. F.** (2016). Reconfiguration of the immune system network during food limitation in the caterpillar *Manduca sexta*. *J. Exp. Biol.* **219**, 706–718.
- Alonso-Alvarez, C., and Tella, J. L.** (2001). Effects of experimental food restriction and body-mass changes on the avian T-cell-mediated immune response. *Can. J. Zool.* **79**, 101–105.
- Ardia, D. R., Gantz, J.E., Schneider, B.C. and Strebel, S.** (2012). Costs of immunity in insects: an induced immune response increases metabolic rate and decreases antimicrobial activity. *Funct. Ecol.* **26**, 732–739.
- Barjac, H. and Thomson, J. V.** (1970). A new serotype of *Bacillus thuringiensis*: *Bacillus thuringiensis* var. *thompsoni* (serotype 11). *J. Invertebr. Pathol.* **15**, 141–144.
- Barnes, A.I. and Siva-Jothy, M.T.** (2000). Density-dependent prophylaxis in the mealworm beetle *Tenebrio molitor* L. (Coleoptera: Tenebrionidae): cuticular melanization is an indicator of investment in immunity. *Proceedings of the Royal Society London B* **267**, 177–182.
- Becker, T., Loch, G., Beyer, M., Zinke, I., Aschenbrenner, A. C., Carrera, P., Inhester, T., Schultze, J. L. & Hoch, M.** (2010). FOXO-dependent regulation of innate immune homeostasis. *Nature* **463**, 369–373.
- Blum, J. E., Fischer, C. N., Miles, J. and Handelsman, J.** (2013). Frequent replenishment sustains the beneficial microbiome of *Drosophila melanogaster*. *mBio* **4**, e00860–13.
- Bolnick, D. I., Snowberg, L. K., Hirsch, P. E., Lauber, C. L., Knight, R., Caporaso, J. G. and Svanbäck, R.** (2014). Individuals' diet diversity influences gut microbial diversity in two freshwater fish (threespine stickleback and Eurasian perch). *Ecol. Lett.* **17**, 979–987.
- Broderick, N. A., Raffa, K.F., Goodman, R. M. and Handelsman, J.** (2004). Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. *Applied and Environmental Microbiology* **70**, 293–300.
- Brogden, K. A.** (2005). Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nature Reviews Microbiology* **3**, 238–250.

- Bucher, G. E.** (1963). Survival of populations of *Streptococcus faecalis* Andrewes and Horder in the gut of *Galleria mellonella* (Linnaeus) during metamorphosis, and transmission of the bacteria to the filial generation of the host. *Journal of Insect Pathology* **5**, 336–343.
- Caccia, S., Di Lelio, I., La Storia, A., Marinelli, A., Varricchio, P., Franzetti, E., Banyuls, N., Tettamanti, G., Casartelli, M., Giordana, B., Ferré, J., Gigliotti, S., Ercolini, D. and Pennacchio, F.** (2016). Midgut microbiota and host immunocompetence underlie *Bacillus thuringiensis* killing mechanism. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 9486–9491.
- Cariveau, D. P., Powell, J. E., Koch, H., Winfree, R. and Moran, N. A.** (2014). Variation in gut microbial communities and its association with pathogen infection in wild bumble bees (*Bombus*). *ISMEJ* **8**, 2369–2379.
- Carmody, R. N., Gerber, G. K., Luevano, J. M., Gatti, D. M., Somes, L, Svenson, K. L. and Turnbaugh, P. J.** (2015). Diet dominates host genotype in shaping the murine gut microbiota. *Cell Host and Microbe* **17**, 72–84.
- Chen, B., The, B-S., Sun, C., Hu, S., Lu, X., Boland, W. and Shao, Y.** (2016). Biodiversity and activity of the gut microbiota across the life history of the insect herbivore *Spodoptera littoralis*. *Scientific Reports* **6**, 29505. doi: 10.1038/srep29505
- Dag, O., Dolgun, A. and Konar, N. M.** (2016) One way tests: One-Way Independent Groups Design. R package version 1.2. <https://CRAN.R-project.org/package=onewaytests>
- David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, E., Ling, A. V., Devlin, A. S., Varma, Y., Fischbach, M. A., Biddinger, S. B., Dutton, R. J. and Turnbaugh, P. J.** (2014). Diet rapidly and reproducibly alters the humans gut microbiome. *Nature* **505**, 559–563.
- Daukste, J., Kivleniece, I., Krama, T., Rantala, M. J. and Krams, I.** (2012). Senescence in immune priming and attractiveness in a beetle. *Journal of Evolutionary Biology* **25**, 1298–1304.
- De Block, M. and Stoks, R.** (2008). Short-term larval food stress and associated compensatory growth reduce adult immune function in a damselfly. *Ecological Entomology* **33**, 796–801.
- DeSantis, T. Z., Hugenholtz, P. and Larsen, N.** (2006). Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology* **72**, 5069–5072.

- Dubovskiy, I.M., Grizanova, E.V., Ershova, N.S., Rantala, M.J. and Glupov, V.V.** (2011). The effects of dietary nickel on the detoxification enzymes, innate immunity and resistance to the fungus *Beauveria bassiana* in the larvae of the greater wax moth *Galleria mellonella*. *Chemosphere* **85**, 92–96.
- Dubovskiy, I.M., Whitten, M.M.A., Kryukov, V.Y., Yaroslavtseva, O.N., Grizanova, E.V., Greig, C., Mukherjee, K., Vilcinskis, A., Mitkovets, P.V., Glupov, V.V. and Butt, T.M.** (2013). More than a colour change: insect melanism, disease resistance and fecundity. *Proceedings of the Royal Society London B* **280**, 20130584.
- Edgar, R. C.** (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461.
- Erdogan, A. and Rao, S. S.** (2015). Small intestinal fungal overgrowth. *Current Gastroenterology Reports* **17**, 16.
- Fujimori, S.** (2015). What are the effects of proton pump inhibitors on the small intestine? *World Journal of Gastroenterology* **21**, 6817–6819.
- Garrett, W. S., Gordon, J. I. and Glimcher, L. H.** (2010). Homeostasis and inflammation in the intestine. *Cell* **140**, 859–870.
- Gayatri Priya, N., Ojha, A., Kajla, M. K., Raj, A. and Rajagopal, R.** (2012). Host plant induced variation in gut bacteria of *Helicoverpa armigera*. *PLoS ONE* **7**, e30768.
- Gillespie, J. P., Kanost, M. R. and Trenczek, T.** (1997). Biological mediators of insect immunity. *Annual Review of Entomology* **42**, 611–643
- Gonzalez-Santoyo, I. and Cordoba-Aguilar, A.** (2012). Phenoloxidase: a key component of the insect immune system. *Entomology Experimentalis et Applicata* **142**, 1–16.
- Greenberg, B. and Klowden, M.** (1972). Enteric bacterial interactions in insects. *American Journal of Clinical Nutrition* **25**, 1459–1466.
- Grizanova, E.V., Dubovskiy, I.M., Whitten, M.M.A. and Glupov, V.V.** (2014). Contributions of cellular and humoral immunity of *Galleria mellonella* larvae in defence against oral infection by *Bacillus thuringiensis*. *Journal of Invertebrate Pathology* **119**, 40–46.
- Heiman, M. L. and Greenway, F. L.** (2016). A healthy gastrointestinal microbiome is dependent on dietary diversity. *Molecular Metabolism* **5**, 317–320.
- Hoffmann, J.A., Kafatos, F.C., Janeway, C.A. and Ezekowitz, R.A.** (1999). Phylogenetic perspectives in innate immunity. *Science* **284**, 1313–1318.
- Huang, J-H. and Douglas, A. E.** (2015). Consumption of dietary sugar by gut bacteria determines *Drosophila* lipid content. *Biology Letters* **11**, 20150469.

- Janeway, C., Travers, P., Walport, M. and Shlomchik, M.** (2005). *Immunobiology*. Garland Science, New York.
- Jarosz, J.** (1979). Gut flora of *Galleria mellonella* suppressing ingested bacteria. *Journal of Invertebrate Pathology* **34**, 192–198.
- Johnston, P. R. and Rolff, J.** (2015). Host and symbiont jointly control gut microbiota during complete metamorphosis. *PLoS Pathogen* **11(11)**, e1005246. doi:10.1371/journal.ppat.1005246.
- Jones, R. M., Mercante, J. W. and Neish, A.S.** (2012). Reactive oxygen production induced by the gut microbiota: pharmacotherapeutic implications. *Current Medical Chemistry* **19**, 1519–1529.
- Jones, R. T., Vetter, S. M., Montenieiri, J., Holmes, J., Bernhardt, S. A. and Gage, K.L.** (2013). *Yersinia pestis* infection and laboratory conditions alter flea-associated bacterial communities. *ISMEJ* **7**, 224–228.
- Kaneko, Y., Furukawa, S., Tanaka, H. and Yamakawa, M.** (2007). Expression of antimicrobial peptide genes encoding enbocin and gloverin isoforms in the silkworm, *Bombyx mori*. *Bioscience, Biotechnology and Biochemistry* **71**, 2233–2241.
- Kangassalo, K., Kosonen, K., Pölkki, M., Sorvari, J., Krams, I., Rantala, M.J.** (2016). Immune challenge has a negative effect on cuticular darkness in the mealworm beetle, *Tenebrio molitor*. *Annales Zoologici Fennici* **53**, 255–262.
- Kau, A. L., Ahem, P. P., Griffin, N. W., Goodman, A. L. and Gordon, J. L.** (2011). Human nutrition, the gut microbiome and the immune system. *Nature* **474**, 327–336.
- Kecko, S., Mihailova, A., Kangassalo, K., Elferts, D., Krama, T., Krams, R., Luoto, S., Rantala, M.J. and Krams, I.A.** (2017). Sex-specific compensatory growth in the larvae of the greater wax moth *Galleria mellonella*. *Journal of Evolutionary Biology*, DOI: 10.1111/jeb.13150.
- Krams, I., Daukšte, J., Kivleniece, I., Krama, T., Rantala, M.J., Ramey, G. and Šauša, L.** (2011). Female choice reveals terminal investment in male mealworm beetles, *Tenebrio molitor*, after a repeated activation of immune system. *Journal of Insect Science* **11**, 56.
- Krams, I., Vrublevska, J., Cirule, D., Kivleniece, I., Krama, T., Rantala, M.J., Sild, E. and Hõrak, P.** (2012). Heterophil/lymphocyte ratios predict the magnitude of humoral immune response to a novel antigen in great tits (*Parus major*). *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative Physiology* **161**, 422–428.

- Krams, I., Kecko, S., Kangassalo, K., Moore, K.F., Jankevics, E., Inashkina, I., Krama, T., Lietuvietis, V., Meija, L. and Rantala, M.J.** (2014). Effects of food quality on trade-offs among growth, immunity and survival in the greater wax moth (*Galleria mellonella*). *Insect Science* **22**, 431–439.
- Krams, I., Burghardt, G.M, Krams, R, Trakimas, G., Kaasik, A., Luoto, S., Rantala, M.J. and Krama T.** (2016). A dark cuticle allows higher investment in immunity, longevity and fecundity in a beetle upon a simulated parasite attack. *Oecologia* **182**, 99–109.
- Krams, I.A., Krama, T., Trakimas, G., Kaasik, A., Rantala, M.J. and Škute, A.** (2017). Reproduction is costly in an infected aquatic insect. *Ethology, Ecology & Evolution* **29**, 74–84.
- Kryukova, N.A., Dubovskiy, I.M., Chertkova, E. A., Vorontsova, Ya.L., Slepneva, I.A. and Glupov, V.V.** (2011). The effect of *Habrobracon hebetor* venom on the activity of the prophenoloxidase system and the generation of reactive oxygen species and encapsulation in the haemolymph of *Galleria mellonella* larvae. *Journal of Insect Physiology* **57**, 796–800.
- Lanan, M. C., Rodrigues, P. A. P., Agellon, A., Jansma, P. and Wheeler, D. E.** (2016). A bacterial filter protects and structures the gut microbiome of an insect. *ISMEJ* **10**, 1866–1876.
- Lavine, M.D. and Strand, M.R.** (2002). Insect hemocytes and their role in immunity. *Insect Biochemistry and Molecular Biology* **32**, 1295–1309.
- Lazzaro, B. P. and Rolff, J.** (2011). Danger, microbes, and homeostasis. *Science* **332**, 43–44.
- Lee, J. H., Park, S., Chae, K-S. and Lee, I. H.** (2010). *Galleria mellonella* 6-Tox gene, putative immune related molecule in Lepidoptera. *International Journal of Industrial Entomology* **21**, 127–132.
- Lochmiller, R. L. and Deerenberg, C.** (2000). Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos* **88**, 87–98.
- Login, F. H., Balmand, S., Vallier, A., Vincent-Monegat, C., Vigneron, A., Weiss-Gayet, M., Rochat, D. and Heddi, A.** (2011). Antimicrobial peptides keep insect endosymbionts under control. *Science* **334**, 362–365 (doi:10.1126/science.1209728).
- Luo, C., Luo, K., Meng, L., Wan, B., Zhao, H. & Hu, Z.** (2017). Ecological impact of a secondary bacterial symbiont on the clones of *Sitobion avenae* (Fabricius) (Hemiptera: Aphididae). *Scientific Reports* **7**, 40754.

- Makarova, O., Rodriguez-Rojas, A., Eravci, M., Weise, C., Dobson, A., Johnston, P. and Rolff, J.** (2016). Antimicrobial defence and persistent infection in insects revisited. *Philosophical Transactions of the Royal Society London B* **371**, 20150296. <http://dx.doi.org/10.1098/rstb.2015.0296>.
- Mason, C. J. and Raffa, K. F.** (2014). Acquisition and structuring of midgut bacterial communities in gypsy moth (Lepidoptera: Erebidae) larvae. *Environmental Entomology* **43**, 595–604.
- Masson, F., Zaidman-Remy, A. and Heddi, A.** (2016). Antimicrobial peptides and cell processes tracking endosymbiont dynamics. *Philosophical Transactions of the Royal Society London B* **371**, 20150298. (doi:10.1098/rstb.2015.0298).
- Melo, N.Rd., Abdrahman, A., Greig, C., Mukherjee, K., Thornton, C., Ratcliffe, N. A., Vilcinskas, A. and Butt, T. M.** (2013). Myriocin significantly increases the mortality of a non-mammalian model host during *Candida* pathogenesis. *PLoS ONE*, **8**, e78905.
- Milani, C., Hevia, A., Feroni, E., Duranti, S., Turrone, F., Lugli, G. A., Sanchez, B., Martín, R., Gueimonde, M., van Sinderen, D., Margolles, A. and Ventura, M.** (2013). Assessing the Fecal Microbiota: An optimized ion torrent 16S rRNA gene-based analysis protocol. *PLoS ONE* **8**, e68739.
- Moos, W. H., Faller, D. V., Harpp, D. N., Kanara, I., Pernokas, J., Powers, W. R. and Steliou, K.** (2016). Microbiota and Neurological Disorders: A Gut Feeling. *Biores Open Access* **5**: 137–145.
- Moret, Y. and Schmid-Hempel, P.** (2000). Survival for immunity: the price of immune system activation for bumblebee workers. *Science* **290**, 1166–1168.
- Muegge, B. D., Kuczynski, J. and Knights, D.** (2011). Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science* **332**, 970–974.
- Mukherjee, K., Raju, R., Fischer, R. and Vilcinskas, A.** (2013). *Galleria mellonella* as a model host to study gut microbe homeostasis and brain infection by the human pathogen *Listeria monocytogenes*. *Advances in Biochemical Engineering / Biotechnology* **135**, 27–39.
- Nagai, T., Osaki, T. and Kawabata, S.** (2001). Functional conversion of hemocyanin to phenoloxidase by horseshoe crab antimicrobial peptides. *Journal of Biological Chemistry* **276**, 27166–27170.
- Odegaard, J. I. and Chawla, A.** (2013). The immune system as a sensor of the metabolic state. *Immunity* **38**, 644–654.

- Oliver, K. M., Degnan, P. H., Burke, G. R. and Moran, N. A.** (2010). Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. *Annual Review of Entomology* **55**, 247–266.
- Peters, G.** (2016). User friendly science: Quantitative analysis made accessible_. R package version 0.5-2, URL: <http://userfriendlyscience.com>.
- Pinto-Tomás, A. A. A., Sittenfeld, A., Uribe-Iorio, L., Mora, M., Janzen, D. H., Goodman, R. M., Holly, M. & Simon, H. M.** (2011). Comparison of midgut bacterial diversity in tropical caterpillars (Lepidoptera: Saturniidae) fed on different diets. *Environ. Entomol.* **40**, 1111–1122.
- Polin, S., Simon, J-C. and Outreman, Y.** (2014). An ecological cost associated with protective symbionts of aphids. *Ecology and Evolution* **4**, 836–840.
- Ponton, F., Wilson, K., Holmes, A. J., Cotter, S. C., Raubenheimer, D. and Simpson, S. J.** (2013). Integrating nutrition and immunology: a new frontier. *Journal of Insect Physiology* **59**, 130–137.
- Povey, S., Cotter, S. C., Simpson, S. J. and Wilson, K.** (2013). Dynamics of macronutrient self-medication and illness-induced anorexia in virally infected insects. *Journal of Animal Ecology* **83**, 245–255.
- Pyro, V. S., Roesch, L. F. W., Morais, D. K., Clark, I. M., Hirsch, P. R., Tótola, M. R.** (2014). Data analysis for 16S microbial profiling from different benchtop sequencing platforms. *Journal of Microbiological Methods* **107**, 30–37.
- R Core Team.** **R:** A language and environment for statistical computing. (2016). R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Råberg, L., Grahn, M., Hasselquist, D. and Svensson, E.** (1998). On the adaptive significance of stress-induced immunosuppression. *Proceedings of the Royal Society London B* **265**, 1637–1641.
- Rantala, M.J., Koskimaki, J., Taskinen, J., Tynkkynen, K. and Suhonen, J.** (2000). Immunocompetence, developmental stability and wingspot size in the damselfly *Calopteryx splendens* L. *Proceedings of the Royal Society London B* **267**, 2453–2457.
- Ratcliffe, N.A., Rowley, A.F., Fitzgerald, S.W. and Rhodes, C.P.** (1985). Invertebrate immunity: basic concepts and recent advances. *International Review of Cytology* **97**, 183–350.
- Russell, V. and Dunn, P. E.** (1996). Antibacterial proteins in the midgut of *Manduca sexta* during metamorphosis. *Journal of Insect Physiology* **42**, 65–71.

- Schmid-Hempel, P.** (2011). *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford University Press.
- Schuhmann, B., Seitz, V., Vilcinskas, A. and Podsiadlowski, L.** (2003). Cloning and expression of gallerimycin, an antifungal peptide expressed in immune response of greater wax moth larvae, *Galleria mellonella*. *Archives of Insect Biochemistry and Physiology* **53**, 125-133.
- Sheldon, B. C. and Verhulst, S.** (1996). Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends in Ecology and Evolution* **11**, 317–321.
- Smits, S. A., Leach, J., Sonnenburg, E. D., Gonzalez, C. G., Lichtman, J. S., Reid, G., Knight, R., Manjurano, A., Changalucha, J., Elias, J. L. and Dominguez-Bello, M. G.** (2017). Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania. *Science* **357**, 802–806.
- Sonnenburg, E. D., Smits, S. A., Tikhonov, M., Higginbottom, S. K., Wingreen, N. S. and Sonnenburg, J. L.** (2016). Diet-induced extinctions in the gut microbiota compound over generations. *Nature* **529**, 212–215.
- Tamboli, C. P., Neut, C., Desreumaux, P. and Colombel, J.F.** (2004). Dysbiosis in inflammatory bowel disease. *Gut* **53**, 1–4.
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. and Speleman, F.** (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* **3**:research0034.1.
- Vogel, H., Altincicek, B., Glockner, G. and Vilcinskas, A.** (2011). A comprehensive transcriptome and immune-gene repertoire of the lepidopteran model host *Galleria mellonella*. *BMC Genomics* **12**, 308.
- Von Baum, H., Klemme, F. R., Geiss, H. K. and Sonntag, H-G.** (1998) Comparative evaluation of a commercial system for identification of gram-positive cocci. *European Journal of Clinical Microbiology and Infectious Diseases* **17**, 849–852.
- Wang, Q., Garrity, G. M., Tiedje, J. M. and Cole, J. R.** (2007). Naïve bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* **73**, 5261–5267.
- Xiang, H., Wei, G., Jia, S., Huang, J., Miao, X., Zhou, Z., Zhao, L. and Huang, L.** (2006). Microbial communities in the larval midgut of laboratory and field populations of cotton boll-worm (*Helicoverpa armigera*). *Can. J. Microbiol.* **52**, 1085–1092.

- Yourth, C.P., Forbes, M.R. and Smith, B.P.** (2001). On understanding variation in immune expression of the damselflies *Lestes* spp. *Canadian Journal of Zoology* **79**, 815–821.
- Zasloff, M.** (2002). Antimicrobial peptides of multicellular organisms. *Nature* **415**, 389–395.

Figures




Fig. 1. The transcription levels of five AMP genes: *6-tox* (A), *Cecropin-D* (B), *Gallerimycin* (C), *Galiomicin* (D) and *Gloverin* (E) in the midgut of the greater wax moth larvae grown on a diverse and a simple diet with and without antibiotic treatment. The transcription levels of the AMP genes were determined by a quantitative real-time RT-PCR analysis and are shown relative to the expression levels of the reference group in which the microbiome was eliminated. Results were normalized against the expression of the housekeeping *18S rRNA* and *EF1* genes and represent means of six independent determinations and standard deviations. *** indicates significant main effects of diet and antibiotic treatment ($P < 0.0001$). X indicates significant interaction between diet and treatment ($P < 0.05$). Lower-case letters 'a', 'b' and 'c' denote significant differences by post

hoc tests at $P < 0.05$. For example, the 'bc' bar significantly differs from the 'a' bar, while the 'bc' bar does not differ from the 'b' and 'c' bars.




Fig. 2. The *Enterococci* counts of colony forming units (CFU)/ml in the midgut samples of *G. mellonella* in four experimental treatments. Thick lines represent medians, while the boxes show 25-75 percentiles.




Fig. 3. Encapsulation rate (average \pm SD) of the greater wax moth larvae grown on the ‘diverse diet’ and the ‘simple diet’ with and without antibiotic treatment. X indicates significant interaction between diet diversity and treatment ($P < 0.01$). Lower-case letters ‘a’, ‘b’ and ‘c’ denote significant differences by post hoc tests at $P < 0.01$.