

Avian thermoregulation in the heat: phylogenetic variation among avian orders in evaporative cooling capacity and heat tolerance

Ben Smit^{1,*}

Maxine C. Whitfield²

William A. Talbot³

Alexander R. Gerson⁴

Andrew E. McKechnie^{2,5}

Blair O. Wolf³

¹Department of Zoology and Entomology, Rhodes University, P.O. Box 94 Grahamstown 6040, South Africa

²DST-NRF Centre of Excellence at the Percy FitzPatrick Institute, Department of Zoology and Entomology, University of Pretoria, Private Bag X20, Hatfield 0028, South Africa

³UNM Biology Department, University of New Mexico, MSC03-2020, Albuquerque, NM 87131-0001, USA

⁴Department of Biology, University of Massachusetts, Amherst, MA 01003, USA

⁵South African Research Chair in Conservation Physiology, National Zoological Gardens of South Africa, P.O. Box 754, Pretoria 0001, South Africa

*Author for correspondence (b.smit@ru.ac.za)

Keywords: evaporative heat loss, heat dissipation, upper critical limits of thermoneutrality, panting, gular flutter

Summary statement

We show that avian evaporative cooling efficiency and heat tolerance display substantial taxonomic variation that are, unexpectedly, not systematically related to the use of panting versus gular flutter processes.

Abstract

Little is known about the phylogenetic variation of avian evaporative cooling efficiency and heat tolerance in hot environments. We quantified thermoregulatory responses to high air temperature (T_a) in ~100-g representatives of three orders: African cuckoo (*Cuculus gularis*, Cuculiformes), lilac-breasted roller (*Coracias caudatus*, Coraciiformes), and Burchell's starling (*Lamprotornis australis*, Passeriformes). All three species initiated respiratory mechanisms to increase evaporative heat dissipation when body temperature (T_b) approached 41.5°C in response to increasing T_a , with gular flutter observed in cuckoos and panting in rollers and starlings. Resting metabolic rate (RMR) and evaporative water loss (EWL) increased by quantitatively similar magnitudes in all three species, although maximum rates of EWL were proportionately lower in starlings. Evaporative cooling efficiency [defined as the ratio of evaporative heat loss (EHL) to metabolic heat production (MHP)] generally remained below 2.0 in cuckoos and starlings, but reached a maximum of ~3.5 in rollers. The high value for rollers reveals a very efficient evaporative cooling mechanism, and is similar to EHL/MHP maxima for similarly sized columbids which very effectively dissipate heat via cutaneous evaporation. This unexpected phylogenetic variation among the orders tested in the physiological mechanisms of heat dissipation is an important step toward determining the evolution of heat tolerance traits in desert birds.

Introduction

Evaporative heat loss (EHL) is the only mechanism whereby birds can maintain body temperature (T_b) below lethal limits in hot environments where environmental temperature exceeds T_b . Rapid increases in evaporative water loss (EWL) are a ubiquitous avian response to such conditions (Bartholomew and Cade, 1963; Dawson and Bartholomew, 1968; Smith et al., 2015; Whitfield et al., 2015). Water requirements for thermoregulation have thus shaped the ecology and evolution of birds living in hot environments, and provide the basis for important trade-offs between dehydration and hyperthermia avoidance in arid-zone species (Smit et al., 2013; Tieleman and Williams, 2000; Tieleman and Williams, 2002; Tieleman et al., 2003).

Avian heat tolerance and evaporative cooling efficiency (quantified as the maximum ratio of heat dissipated evaporatively, EHL, to that generated metabolically, MHP) appears to vary substantially among orders (Lasiewski and Seymour, 1972; Smith et al., 2015), within orders (McKechnie et al., 2016a; McKechnie et al., 2017), and even within species (Noakes et al., 2016). For example, recent studies have shown that the efficiency of evaporative cooling is generally very high ($EHL/MHP > 2.0$) in members of the orders Columbiformes and Caprimulgiformes (McKechnie et al., 2016a; O'Connor et al., 2017; Smith et al., 2015; Talbot et al., 2017), but less so in Passeriformes, Pteroclidiformes, Galliformes and Strigiformes, where efficiency rarely exceeds 2.0 (Bartholomew et al., 1968; McKechnie et al., 2016b; McKechnie et al., 2017; Smith et al., 2015; Smith et al., 2017; Whitfield et al., 2015). Narrow phylogenetic sampling means we still have an incomplete understanding of the diversity of avian EHL mechanisms and their functional significance for heat tolerance.

Body mass represents one of the most prominent sources of variation in heat tolerance and the efficiency of EHL (McKechnie and Wolf, 2010), but even species of similar size can still show substantial variation. For example, Lasiewski and Seymore (1972) showed that four similarly-sized species from different orders (*Ploceus cucullatus*, Passeriformes; *Excalfactoria chinensis*, Galliformes; *Scardafella inca*, Columbiformes; *Phalaenoptilus nuttallii*, Caprimulgiformes) show substantial variation in terms of the magnitude of elevations in T_b , metabolism, EWL, and EHL/MHP during heat exposure. Recent studies have suggested that some of the mechanisms underlying this variation may include differential reliance on

respiratory versus cutaneous pathways of evaporative cooling (McKechnie et al., 2016a), and morphological variation (e.g. bill size) (Danner et al., 2016; Tattersall et al., 2009; van de Ven et al., 2016). Species relying on cutaneous evaporative pathways, such as Columbiformes, are thought to show negligible metabolic heat production associated with evaporative cooling demands (McKechnie et al., 2016a). In contrast, species relying on respiratory evaporative cooling (especially panting) may incur greater metabolic costs in their efforts to dissipate heat (Bartholomew et al., 1968; Lasiewski and Seymour, 1972). Gular flutter (gaping while pulsating the hyoid bone to ventilate the buccal cavity) has been argued to require less energy than panting (gaping while using rapid movements of the thorax and abdominal cavities) (Bartholomew et al., 1962; Dawson, 1958; Lasiewski and Bartholomew, 1966); the gular flutter process may thus enhance efficiency of EHL. The use of gular flutter *versus* panting further seems to vary among taxa (Calder and Schmidt-Nielsen, 1967), but the phylogenetic distribution of these evaporative cooling processes remains poorly understood.

Variation in the rate and efficiency of evaporative cooling may reflect physiological costs, such as risk of dehydration and hyperthermia (McKechnie and Wolf, 2010), but may also reflect behavioural [e.g. reduced foraging efficiency correlated with heat dissipation efforts (du Plessis et al., 2012)] and ecological costs [e.g. reliance of surface water sources (Smit et al., 2016)]. Thus far, evaporative cooling efficiency and heat tolerance are best-studied in the Passeriformes and Columbiformes, and data from more under-represented orders are needed to further our understanding of the evolutionary drivers and mechanisms involved in thermoregulation in the heat.

Here we present data on thermoregulation under very hot conditions in three 100-g species from three orders: Cuculiformes, Coraciiformes and Passeriformes. The Cuculiformes and Coraciiformes remain largely unstudied in terms of thermoregulation at high environmental temperatures; both represent diverse taxa that include many species occupying hot, arid regions. To the best of our knowledge, the only thermoregulatory data from a cuculiform under hot conditions is for the roadrunner, *Geococcyx californianus* (Calder and Schmidt-Nielsen, 1967). In the latter study, high chamber humidity may have reduced the efficiency of EHL at air temperatures (T_a) below T_b (Gerson et al., 2014). Recent advances in carbon dioxide and water analysers have allowed us to maintain low humidity levels during testing by using very high flow rates, thus avoiding the complications associated with high humidity that plagued early studies (Gerson et al., 2014; Lasiewski et al., 1966; Smith et al., 2015; Whitfield et al., 2015). We further describe the mechanism of respiratory EHL (panting or gular flutter) used by these three species, and predict that, compared to panting, the use of

gular flutter is strongly associated with improved evaporative cooling efficiency and heat tolerance. In addition, our inclusion of a large passerine (>100 g) allows for more rigorous analyses of the scaling of traits related to heat tolerance and evaporative cooling in this speciose taxon (McKechnie et al., 2017).

Methods

Study species and study sites

We measured resting metabolic rate (RMR), evaporative water loss (EWL), and T_b in species representing three avian orders in the southern Kalahari Desert in the Northern Cape province of South Africa, an arid area with a mean annual rainfall of ~200 mm and summer daily maximum T_a ranging from ~20-43°C (Whitfield et al., 2015). We followed the methodology of Whitfield et al. (2015) to quantify thermoregulatory responses to high T_a in a field laboratory from January to April 2012 at Wildsgenot Game Ranch (27°04'S, 21°23'E), and from January to March 2013 at Leeupan Game Ranch (26°58'S, 21°50'E).

We trapped six African cuckoos (*Cuculus gularis* Stephens; order Cuculiformes, hereafter cuckoos) with a mean \pm s.d. body mass (M_b) of 109.6 ± 5.6 g (range 99.0 - 116.0 g) at Leeupan Game Ranch during early 2013 (austral summer) using mist nets. We trapped 10 lilac-breasted rollers (*Coracias caudatus* Linnaeus; Order Coraciiformes, hereafter rollers) with a M_b of 95.4 ± 8.5 g (78.2 - 110.35 g); two individuals were trapped in February 2012 at Wildsgenot Game Ranch, and the remaining individuals were trapped at Leeupan Game Ranch between January 2013 and March 2013. We trapped seven Burchell's starlings (*Lamprotornis australis* Smith; order Passeriformes, hereafter starlings) with a M_b of 109.1 ± 9.3 g (85.4 - 116.9 g) using mist nets and/or flap traps baited with tenebrionid beetle larvae on Wildsgenot Game Ranch in February 2012, and Leeupan Game Ranch from December 2013 to February 2013. Measurements were carried out on the same day of capture. If birds were not tested within the first three hours after being trapped, they were housed in cages constructed of shade-cloth, and were provided with tenebrionid beetle larvae and water ad libitum, until physiological measurements commenced.

Birds were held in respirometry chambers for 2-3 hr, a period that typically limited M_b loss to <5% of initial M_b (mean M_b loss during measurements was $4.0 \pm 1.8\%$ of initial values) and time in captivity did not exceed 24 hr, after which birds were released at the site of capture following Whitfield et al. (2015). All experimental procedures were approved by the Animal Ethics Committee of the University of Pretoria (protocol EC071-11) and the Institutional Animal Care and Use Committee of the University of New Mexico (12-1005370-MCC). A

permit to trap the birds was issued by the Northern Cape Department of Environmental Affairs (ODB 008/2013).

Gas exchange and temperature measurements

We used the same experimental set-up as described by Whitfield et al. (2015) to obtain gas exchange, T_a and T_b measurements. We measured T_b using calibrated temperature-sensitive passive integrated transponder (PIT) tags (Biomark, Boise, ID, USA) which were injected intraperitoneally into the abdominal cavity of each bird shortly after capture. We monitored T_b throughout gas exchange measurements, using a reader and transceiver system (model FS2001, Biomark, Boise, ID, USA). We obtained carbon dioxide production (\dot{V}_{CO_2}) and EWL measurements over the T_a range of 25-56°C (depending on the species), also using the same experimental setup as described by Whitfield et al. (2015). All three species were placed individually in 9-L plastic chambers, and stood on a platform of plastic mesh 10 cm above a 1-cm layer of mineral oil to trap excreta. We used flow rates between 9 and 30 L min⁻¹ for cuckoos; 9 and 70 L min⁻¹ for rollers, and 10 and 55 L min⁻¹ for starlings, depending on the experimental T_a , in order to keep chamber humidity below 5 ppt. As was the case in previous studies using the same methodology (Smith et al. 2015, Whitfield et al. 2015, McKechnie et al. 2016a, McKechnie et al. 2016b), birds remained calmer at very high T_a when we increased flow rate (and hence decreased humidity).

Experimental protocol

Following the protocol described by Whitfield et al. (2015), we exposed birds to progressively higher T_a , with increments of approximately 5°C between 25 and 40°C, and 2°C increments between 40°C and the maximum T_a (ranging from 48 to 56°C depending on the species). Tests were conducted during the day, as this was the active phase of all three species. Birds spent between 10 and 30 min at each T_a value. We continually monitored birds during measurements using a live video feed and an infrared light source (Whitfield et al., 2015). We recorded behavioural responses of birds within the respirometry chambers every two minutes; an activity state score of the individual (e.g. calm, turning around or jumping) was recorded, as well as respiratory EHL mechanisms, including panting (defined as gaping) and gular flutter (defined as an obvious pulsation of hyoid bone in the gular area while gaping). For each individual, we recorded the air temperature and body temperature at which heat dissipation behaviour was initiated.

We followed Whitfield et al. (2015) in terminating test runs when a bird a) exhibited prolonged escape behaviour such as agitated jumping, pecking and/or wing flapping), b) showed signs of a loss of coordination or balance accompanied by $T_b > 44^\circ\text{C}$, or c) exhibited a decrease in EWL and RMR accompanied by an uncontrolled increase in T_b . In the last instance, a bird was considered to have reached its upper limit of heat tolerance, and the T_a associated with the onset of these signs of heat stress was considered the thermal endpoint for that individual. When a bird reached any the conditions described above, we removed the individual from the chamber by hand, gently rubbed a cotton pad soaked in ethanol onto its body, and held it in front of an air-conditioner producing chilled air in order to facilitate rapid heat loss (Whitfield et al., 2015).

Data analyses

Data were analysed following Whitfield et al. (2015). We calculated physiological estimates by extracting, EWL and T_b as the lowest 5 min mean at each T_a using Expedata (Sable Systems, Las Vegas NV, USA). We present whole-animal values, although we also calculated the slope of mass-specific EWL vs. T_a to compare our values with the allometric equation presented by McKechnie and Wolf (2010). We followed McKechnie et al. (2017) by converting rates of EWL to evaporative heat loss (W) assuming a latent heat of vaporisation of water of 2.406 J mg^{-1} at 40°C (Tracy et al., 2010). Since none of our study species have crops we assumed all birds were post-absorptive at the time of measurements, but we were unable to confirm this. Measurements typically took place more than 1 hr after capture, and food (mealworm larvae) was only offered to individuals not tested within 3 hrs of capture. We therefore assumed a respiratory exchange ratio of 0.85, representative of a mix of carbohydrate and lipid metabolism in post-absorptive birds (Walsberg and Wolf 1995), and converted rates of \dot{V}_{CO_2} to metabolic rate (Watt, W) using $24.4 \text{ J mL}^{-1} \text{ CO}_2$ (Withers 1992).

We used segmented linear regression models fitted in the *R* package *segmented* (Muggeo, 2009) to estimate inflection T_a in the physiological data. Although our use of these models violates assumptions of independent samples, we used segmented models purely to aid us in identifying inflection T_a values in physiological variables. We subsequently used linear mixed effects models that included individual identity as a random factor in the *R* package *nlme* (Pinheiro et al., 2009), to obtain estimates of physiological variables as a function of T_a , using subsets of the data above respective inflection T_a . We contrasted the slopes of physiological parameters obtained from these mixed effects models among species, by including data

previously collected from laughing doves (*Stigmatopelia senegalensis*, 89.4 ± 13.0 g, hereafter doves) studied at the same time and site as our species (McKechnie et al., 2016a), and freckled nightjars (*Caprimulgus tristigma*, 64.7 ± 6.3 g, hereafter nightjars) studied from a desert site in Namaqualand, South Africa (O'Connor et al., 2017). Thermoregulatory data in the doves and nightjars were collected in the same manner as described here, although we recalculated physiological estimates of \dot{V}_{CO_2} , EWL and T_b in the dove, as the lowest 5 min mean at each T_a , to facilitate comparison with our study species and the nightjar. Our small sample size of species prevented us from conducting more rigorous statistical analyses that test phylogenetic inertia and, if necessary, account for phylogenetic relatedness. We calculated the upper and lower 95% confidence limits (CL) of each coefficient, and considered species to differ quantitatively in the parameters when there was no overlap in the 95% CL.

Results

Body temperature

The T_b measured at T_{as} below the respective T_{uc} of each species varied by 0.7°C , from a mean of $40.0 \pm 0.6^\circ\text{C}$ in cuckoos to $40.7 \pm 0.8^\circ\text{C}$ in rollers (Table 1). The inflection of T_a above which T_b increased was $< 37^\circ\text{C}$ in all three species (Table 1), with T_b increasing linearly and significantly above the inflection T_{as} (cuckoo, $t_{1,26}=14.275$, $p<0.001$; rollers, $t_{1,56}= 12.10$, $p<0.001$; starlings, $t_{1,28}=7.71$, $p<0.001$) (Fig. 1). The maximum T_a reached was 50°C for both cuckoos and starlings, and 56°C for rollers. Any further increases in T_a resulted in these individuals becoming severely agitated. Average maximum T_b ranged between 42.8° and 44.0°C in the three species at the highest T_{as} tested. In two calm cuckoos, T_b exceeded 44°C at $T_a = 46^\circ$ and 48°C , respectively, whereupon they lost their ability to right themselves and the experiment was terminated immediately. With the exception of $T_b > 44^\circ\text{C}$ in one roller individual (with no ill-effects), T_b remained $< 44^\circ\text{C}$ in calm rollers and starlings at the maximum T_{as} . At the highest T_a shared among the three species (46°C), rollers maintained a lower T_b than cuckoos and starlings (Table 1). The mean T_a at which birds initiated respiratory EHL mechanisms varied from 35.9°C in starlings to $\sim 41^\circ\text{C}$ in cuckoos and rollers, typically as T_b approached 41.5°C in all three species (Table 1). Our observations of the cuckoos revealed that gaping was always accompanied by gular flutter, which was visibly evident from the pulsating throat and moving hyoid apparatus. Rollers and starlings panted while gaping (the gape is wide in rollers) and we did not observe gular flutter (hyoid movement) in either species.

Resting metabolic rate

One cuckoo showed abnormally high RMR values (ranging from 1.03 W at $T_a = 30^\circ\text{C}$, to 5.20 W at $T_a = 50^\circ\text{C}$, i.e. $> 6 \times$ s.d. greater than the mean for the remaining cuckoos), and we excluded data from this individual from our RMR analyses and descriptive statistics. Minimum RMR values measured at $T_a < 35^\circ\text{C}$, below the respective T_{uc} s of each species, varied from 0.64 ± 0.20 W and 0.69 ± 0.13 W in cuckoos and rollers, respectively, to 1.19 ± 0.31 W in starlings (Table 1; mass-specific values are provided in supplementary information, Table S1). In rollers and starlings the T_{uc} (i.e., T_a inflection) ranged from 31°C in starlings to 47.5°C in rollers. No clear inflection point could be identified for cuckoos, and RMR increased across the entire range of T_a tested (Fig. 2). In all three species, RMR increased linearly and significantly at $T_a > T_{uc}$ (cuckoo, $t_{1,23}=2.67$, $p<0.05$; rollers, $t_{1,25} = 2.59$, $p<0.05$; starlings, $t_{1,29}=4.13$, $p<0.001$) (Fig. 2). Maximum RMR observed at maximum T_a was higher in the starlings, with an average of 1.91 W, and lowest in the rollers at ~ 1.10 W. Maximum RMR was 0.96 W and 2.77 W in two cuckoo individuals, respectively (Table 1). The average magnitudes of RMR elevations above estimated thermoneutral levels (max. RMR / min. RMR) were generally below 1.6, with the exception of a single cuckoo that showed a 3-fold increase in RMR (Table 1). At the highest test T_a shared among the three species, RMR was quantitatively similar in cuckoos and rollers, but ~ 2 -fold higher in the starlings (Table 1).

Evaporative water loss

Minimum EWL values measured at T_{as} below the respective T_{uc} of each species ranged from 0.48 ± 0.14 g h⁻¹ in cuckoos to 1.00 ± 0.67 g h⁻¹ in starlings (Table 1; mass-specific values, TableS1). The inflection T_a varied by more than 10 °C among the three species; inflection T_a was 33.3°C in starlings and greater than 42°C in cuckoos and rollers (Table 1). Whereas cuckoos and starlings showed stable EWL rates below their respective inflection T_{as} , rollers showed a shallow but significant increase in EWL to 1.65 ± 0.35 g h⁻¹ at $T_a = 44.7^\circ\text{C}$ ($t_{1,22}=3.22$, $p<0.01$; Fig. 3). Above the inflection T_a EWL increased linearly and significantly in all three species (cuckoo, $t_{1,20}=13.22$, $p<0.001$; rollers $t_{1,33}=11.64$, $p<0.001$; starlings, $t_{1,28}=11.133$, $p<0.001$; Fig. 3). Maximum elevations in EWL were around 5-fold (>5.5 -fold in cuckoos and rollers) above thermoneutral values in all three species (Fig. 3, Table 1). Whereas rollers showed maximum EHL/MHP ratios exceeding 3 at the maximum T_{as} tested, cuckoos and starlings generally showed values below 2 (Fig. 4). At the highest shared T_a cuckoos and rollers showed similar EWL rates and evaporative efficiencies (Table 1). In contrast, starlings showed EWL of ~ 3.40 g h⁻¹, and a slightly lower efficiency compared to the former species (Table 1).

Variation in thermoregulation at high air temperatures

The T_b of cuckoos and doves increased more rapidly compared to rollers and nightjars (95% CL did not overlap, Fig. 5, Table 2). Starlings showed an intermediate slope of T_b , with 95% CL overlapping with most other species. The slope of RMR at T_a above thermoneutrality was similar among species, with the exception of nightjars which showed almost no increase in RMR above 40°C and no overlap in 95% CL with any of the other species (Fig. 5, Table 2). The slope of EWL above the inflection T_a did not differ among cuckoos, starlings and doves, but was significantly steeper in rollers and significantly shallower in nightjars compared to all other species (Fig. 5, Table 2). The slopes of EHL/MHP were similar in doves and starlings, and significantly lower than in all the other species (Fig. 5; Table 2). Nightjars showed a steeper EHL/MHP slope compared to rollers and cuckoos, yet overlapped marginally with the 95% CL of these species.

Discussion

All three species studied investigated here showed elevations in T_b , RMR and EWL at $T_a > 40^\circ\text{C}$, qualitatively consistent with typical avian patterns. Despite their similarity in M_b , the three study species showed substantial variation in patterns of T_b , RMR and EWL at high T_a . The comparatively shallow elevations in RMR in rollers were associated with the highest evaporative efficiency, and this species tolerated the highest T_a . Rollers showed very small elevations in T_b , associated with high evaporative efficiency. These patterns in rollers are unexpected since they appear to use panting as the primary EHL pathway. In contrast, at lower T_a both starlings and cuckoos showed sharp increases in T_b and RMR as well as lower evaporative efficiencies. Yet, the T_a at which these responses were initiated, patterns of T_b regulation and EWL elevations were strikingly different among these two species. Additionally, our five-order comparison reveals that all species, with the exception of caprimulgids (O'Connor et al., 2017), clearly showed elevated metabolic costs of heat dissipation (as defined by the slope of RMR above the inflection point) (Fig. 5). These findings illustrate how little we know about the phylogenetic variation in avian evaporative cooling pathways, and specifically the functional significance, selective pressures and evolution of these seemingly diverse pathways for heat tolerance.

Body temperature and heat tolerance limits

The mean T_b of cuckoos at T_a below 35°C ($39.8 \pm 0.30^\circ\text{C}$) was 1.5°C lower than the mean active phase T_b values for Cuculiformes (41.3°C), but the mean T_b s of starlings and rollers were within 0.5°C of the mean values reported for Passeriformes and Coraciiformes, respectively (Prinzinger et al., 1991). In all three species, the inflection T_a for T_b was $< 40^\circ\text{C}$ (Fig. 1). These values are lower than the inflection T_a s reported for columbids, quail, and smaller passerines (generally $T_a \sim 40^\circ\text{C}$) (McKechnie et al., 2016a; Smith et al., 2015; Whitfield et al., 2015). Two species of Caprimulgiform also show a T_b inflection at $T_a < 40^\circ\text{C}$ (O'Connor et al., 2017).

In calm birds, maximum T_b was $>44.5^\circ\text{C}$ in a cuckoo at $T_a = 49.6^\circ\text{C}$, 44.1°C in a starling at $T_a = 49.2^\circ\text{C}$ and 44.6°C in a roller at $T_a = 53^\circ\text{C}$ (Fig. 1). The two cuckoos lost coordination at $T_b > 44^\circ\text{C}$, suggesting this taxon may have lower critical thermal limits than Passeriformes and Columbiforms, where several species have been shown to tolerate $T_b > 45^\circ\text{C}$ (Dawson, 1954; McKechnie et al., 2016a; Smith et al., 2015; Smith et al., 2017; Whitfield et al., 2015). Several smaller passerines also reached thermal limits as T_a approached 50°C (McKechnie et al., 2017; Smith et al., 2017; Whitfield et al., 2015). However, $T_a = 50^\circ\text{C}$ is probably an underestimation of the heat tolerance limits of cuckoos and starlings, as we did not reach T_a at which calm individuals showed a plateau in EWL accompanied by an increasing T_b (Whitfield et al., 2015). In contrast to cuckoos and starlings, rollers defended T_b over a much larger T_a - T_b gradient, tolerating $T_a \approx 56^\circ\text{C}$, with a mean $T_b \approx 43^\circ\text{C}$ (Fig. 1). The T_b patterns of the coraciiform in our study at T_a approaching 60°C was therefore quantitatively similar to the thermoregulatory patterns observed in Columbiformes (McKechnie et al., 2016a; Smith et al., 2015).

Resting metabolic rate

Individual variation in RMR tended to be greater in starlings and rollers compared to cuckoos. Some of this variation may be explained by the larger M_b ranges of starlings and rollers compared with cuckoos, but our data do not allow for a rigorous analyses of the scaling effects. Nevertheless, within species, individuals showed similar increasing trends in RMR elevations in slope estimates. The upper inflection T_a in RMR versus T_a , i.e., the T_{uc} indicating increased energetic costs of respiratory EHL, generally occurs at $T_a < 40^\circ\text{C}$ (Calder and Schmidt-Nielsen, 1967; Dawson and Bennett, 1973; McNab and Bonaccorso, 1995; Tieleman and Williams, 2002; Whitfield et al., 2015). In all three species studied here, T_{uc} was not clearly related to

increased efforts of respiratory EHL, unlike the case in Burchell's sandgrouse (McKechnie et al., 2016b). Our data suggest that the initiation of gular flutter or panting was related to a threshold T_b of $\sim 41.5^\circ\text{C}$, rather than a threshold T_a (Table 1). The cuckoos in our study did not show a clear upper inflection T_a , contrasting with the marked T_{uc} associated with the onset of gular flutter in Burchell's sandgrouse (*Pterocles burchelli*) (McKechnie et al., 2016b). At the T_a value at which we observed cuckoos initiating gular flutter ($\sim 42^\circ\text{C}$, Table 1), RMR had already increased by more than 50% above minimum resting levels. Similarly, in the starlings, the inflection T_a for RMR was well below the T_a at which this species initiated gaping behaviour, and lower than values in smaller passerines (Whitfield et al. 2015). The patterns in these two species are perhaps explained by a Q_{10} effect associated with T_b elevations, but our small sample sizes precluded us from drawing any firm conclusions; it is noteworthy that Weathers (1981) showed that, in several passerines, elevations in T_b do not affect RMR, but this idea has not been rigorously tested. In contrast to starlings and cuckoos, rollers maintained a low and stable RMR up to $T_a = 47^\circ\text{C}$ (Fig. 2), much higher than the T_a associated with gaping. This suggests that rollers kept metabolic heat loads low during the initial stages of panting. Our estimate of the upper T_{uc} in rollers is thus substantially higher than typically observed in passerines, which also make use of panting (McKechnie et al., 2017; Weathers, 1981; Whitfield et al., 2015).

Our data reveal considerable overlap in the slope of RMR increases above the T_{uc} in all three species studied here when compared to a similarly-sized dove (*S. senegalensis*) (Fig. 5, Table 2). The authors of at least one previous study noted that the metabolic costs in species that pant (e.g. village weavers, *Ploceus cucullatus*) are similar to those in species which gular flutter (e.g. Chinese painted quails, *Excalfactoria chinensis*), and additionally cutaneous EHL [e.g. Inca doves, *Scardafella inca* (Lasiewski and Seymour, 1972)]. However, one striking difference among the species we compare here is that doves and rollers reached maximum RMR at higher T_a and maintained lower T_b compared to cuckoos and starlings. Fractional elevations in RMR at $T_a = 50^\circ\text{C}$ compared to thermoneutral values were as high as 60% in starlings, 25% in cuckoos, and $\sim 10\%$ in rollers and doves. These increases fall broadly between values previously reported for passerines [30-90%; (McKechnie et al., 2017; Whitfield et al., 2015) and columbids ($\sim 7\%$ (McKechnie et al., 2016a) over a similar T_a range (30°C to 50°C). The lack of metabolic elevations at T_a s above those where gular flutter was initiated in the nightjars ($T_a \sim 40^\circ\text{C}$) contrasts strongly with all the above-mentioned species (Fig. 5). Metabolic rates at high temperature thus seem not related to the two modes (gular flutter versus panting) these phylogenetically diverse species employ.

Evaporative heat loss

Patterns of EWL followed the typical avian pattern, where EWL is minimal at T_a around 25–35°C and increases approximately linearly at T_a s approaching or exceeding 40°C. The variation in inflection T_a and slope of EWL above these inflections is difficult to explain, and suggests the capacity to dissipate heat evaporatively may vary greatly among taxa of similar size.

Maximum capacity for evaporative cooling, expressed as EHL/MHP, was qualitatively related to the maximum T_a reached by our three study species. On average, cuckoos and starlings evaporated less than 200% of metabolic heat production at the maximum T_a reached (Fig. 4) – similar to patterns observed in passerines (McKechnie et al., 2017; Smith et al., 2017; Whitfield et al., 2015). The comparatively high RMR of starlings likely constrained their evaporative cooling efficiency as they had to evaporate water almost twice as rapidly compared to the other species to dissipate 100% of their metabolic heat. This may have broad-scale implications for heat loss in the order Passeriformes, especially in light of a recent study demonstrating high basal metabolism in this order (Londoño et al., 2015). Cuckoos showed a slope of EHL/MHP versus T_a only marginally shallower than rollers, yet despite their low RMR at $T_a < 35^\circ\text{C}$ did not achieve a high evaporative efficiency (Fig. 4); even though a very calm cuckoo dissipated 200% of its metabolic heat load at $T_a = 50^\circ\text{C}$, the T_b of this individual was already $\sim 43^\circ\text{C}$.

Rollers showed a mean maximum EHL/MHP of 3.6 (> 4 in one individual) (Fig. 4), quantitatively similar to values reported in Columbiformes (2.3 - 4.7) (McKechnie et al., 2016a; Smith et al., 2015). In fact, at $T_a > 50^\circ\text{C}$ rollers evaporated a greater fraction of metabolic heat compared to medium-sized columbids (140-g *S. capicola*, and 90-g *S. senegalensis*) occupying the same Kalahari Desert habitat (McKechnie et al., 2016a). This result is surprising as panting is often argued to be a metabolically costly process compared to gular flutter (Bartholomew et al., 1962; Whitfield et al., 2015). We suspect the pronounced heat tolerance in rollers may be functionally linked to the long periods they spend perched in exposed sites while hunting (Herremans, 2005; Smit et al., 2016). We do not know whether rollers rely on cutaneous evaporation at high T_a , although we propose that evaporative cooling may be enhanced by the large gape of rollers increasing the buccal surface area for EWL (personal observation); a combination of large gape [$\sim 15\%$ of skin surface area, (Cowles and Dawson, 1951)] and high evaporative efficiency also is common to the caprimulgids, but it would be interesting to investigate the contribution of CEWL to total EWL in rollers. Our understanding of phylogenetic variation in the partitioning of EWL into cutaneous and respiratory pathways

remains very poorly studied, and beside for a few studies focussing on Columbiformes and Passeriformes (McKechnie and Wolf, 2004; Ro and Williams, 2010; Tieleman and Williams, 2002; Wolf and Walsberg, 1996), there are limited empirical data on how increased reliance on CEWL at high T_a reduces the energetic demands for respiratory heat dissipation.

Preliminary comparisons of phylogenetic variation in thermoregulation under hot conditions

Although our study was not designed to directly quantify gular flutter versus panting in our study species, our results, combined with those of (Lasiewski and Seymour, 1972), reveal that the mechanisms of respiratory heat dissipation are not clearly related to evaporative cooling capacity and tolerance of high T_a . Even Columbiformes, thought to rely greatly on cutaneous EHL seem to suffer reductions in evaporative efficiency when they start incorporating gular flutter or panting mechanisms [e.g., *S. capicola*, and *S. senegalensis*, (McKechnie et al., 2016a)]. Calder and Schmidt-Nielsen (Calder and Schmidt-Nielsen, 1967) suggested the process of gular flutter may differ among taxa and determine the metabolic costs of ventilating the buccal cavity. These costs may include processes where the frequency and amplitude of breathing cycles are in synchrony with gular flutter rates [e.g. Strigiformes, Columbiformes and Cuculiformes (Bartholomew et al., 1968)], or asynchronous where rapid gular flutter is accompanied by slow breathing rates [e.g. some Pelicaniformes and some Caprimulgiformes (Bartholomew et al., 1968; Lasiewski and Bartholomew, 1966; Lasiewski and Seymour, 1972)]. Bartholomew et al. (1968) further suggested that species maintaining gular flutter rates asynchronous to breathing rates, such as the poorwill (*Phalaenoptilus nuttalli*), achieve very high evaporative cooling efficiencies compared to species where breathing is completely synchronised with gular fluttering (probably no more costly than achieved through panting alone). To the best of our knowledge, the role these different modes of gular flutter play in the efficiency of EHL has not been further investigated.

The phylogenetic variation in modes of avian respiratory EHL pathways is interesting, and there is no clear explanation for the dichotomy between gular flutter and/or panting among avian taxa. The prevalence of gular flutter in phylogenetically older taxa, such as Paleognathae and most basal Neognathae, suggests it is a plesiomorphic trait. This may indicate that the function of the hyoid bone, which ventilates the gular area, was modified in some clades of the Neognathae (e.g. Coraciiformes and Passeriformes). Interestingly, Psitticiformes [parrots, closely related to Passeriformes, (Hackett et al., 2008) are the only avian taxon in which lingual flutter (movement of tongue) is known to supplement panting to ventilate the buccal area

(Bucher, 1981; Bucher, 1985). However, to the best of our knowledge, no studies have examined whether lingual flutter differs in efficiency from panting alone.

In conclusion, substantial phylogenetic variation exists in avian heat tolerance and evaporative cooling capacity, an observation reiterated by our comparisons among representatives of five orders. This variation may have far-reaching implications for the ecology and evolution of birds that routinely have to maintain T_b below environmental temperature. The consequences of this variation for water balance and risk of over-heating during very hot weather are strikingly illustrated by the observation that, among the species considered here at $T_a \sim 50^\circ\text{C}$, rates of EWL expressed in term of body mass loss vary ~ 2 -fold; from an equivalent of $2.3\% M_b \text{ h}^{-1}$ in nightjars to $4.6\% M_b \text{ h}^{-1}$ in starlings. Moreover, at the same T_a the T_b of starlings is $\sim 3^\circ\text{C}$ higher than that of nightjars. Quantifying phylogenetic diversity in avian thermal physiology under hot conditions is critical for testing hypotheses regarding physiological adaptation, and predicting vulnerability to the higher T_a and more frequent and intense heat waves associated with anthropogenic climate change (Albright et al., 2017). For instance, the 140-fold differences in slopes of RMR and EWL with increasing T_a , (RMR $0.0003 \text{ mW } ^\circ\text{C}^{-1}$ in nightjars to $0.0417 \text{ mW } ^\circ\text{C}^{-1}$ in starlings) and 3.6-fold (EWL $0.093 \text{ g h}^{-1} ^\circ\text{C}^{-1}$ in nightjars to $0.337 \text{ g h}^{-1} ^\circ\text{C}^{-1}$ in rollers) suggests that, even within similarly-sized species, the consequences of the predicted 4°C increase in extreme T_a maxima for the 21st Century (IPCC, 2007; IPCC, 2011) will vary substantially. The current phylogenetic sampling of data on avian thermoregulatory responses to heat is poor, and a thorough review of the functional roles of these mechanisms as determinants of evaporative cooling efficiency will require sampling more avian taxa from currently unstudied orders.

Acknowledgements

The Scholtz and de Bruin families (South Africa) allowed us to conduct this research on their properties. We also thank Michelle Thompson, Matthew Noakes, Ryan O'Connor, and Mateo Garcia for assistance in the field and laboratory.

Competing interests

The authors declare no competing financial interests.

Author contributions

B.O.W., A.E.M. and B.S. designed the study. M.C.W., B.S., W.A.T. and B.O.W. collected data, B.S. and A.R.G. analysed the data. B.S. wrote the manuscript.

Funding

This material is based on work supported by the National Science Foundation under IOS-1122228 to B. O. Wolf. Any opinions, findings and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

Abbreviations

CEWL cutaneous evaporative water loss

EHL evaporative heat loss

EWL evaporative water loss

M_b body mass

MHP metabolic heat production

RMR Resting metabolic rate

CL confidence limit

T_a air temperature

T_b body temperature

\dot{V}_{CO_2} carbon dioxide production

\dot{V}_{O_2} oxygen consumption

References

- Albright, T. P., Mutiibwa, D., Gerson, A. R., Smith, E. K., Talbot, W. A., O'Neill, J. J., McKechnie, A. E. and Wolf, B. O.** (2017). Mapping evaporative water loss in desert passerines reveals an expanding threat of lethal dehydration. *Proc. Natl. Acad. Sci.* **114**, 2283–2288.
- Bartholomew, G. A. and Cade, T. J.** (1963). The water economy of land birds. *Auk* **80**, 504–539.
- Bartholomew, G. A., Hudson, J. W. and Howell, T. R.** (1962). Body temperature, oxygen consumption, evaporative water loss, and heart rate in the poor-will. *Condor* **64**, 117–125.
- Bartholomew, G. A., Lasiewski, R. C. and Crawford, E. C.** (1968). Patterns of panting and gular flutter in cormorants, pelicans, owls, and doves. *Condor* **70**, 31–34.
- Bucher, T. L.** (1981). Oxygen consumption, ventilation and respiratory heat loss in a parrot, *Bolborhynchus lineola*, in relation to ambient temperature. *J. Comp. Physiol.* **142**, 479–488.
- Bucher, T. L.** (1985). Ventilation and oxygen consumption in *Amazona viridigenalis*: a reappraisal of “resting” respiratory parameters in birds. *J. Comp. Physiol.* **155**, 269–276.
- Calder, W. A. and Schmidt-Nielsen, K.** (1967). Temperature regulation and evaporation in the pigeon and the roadrunner. *Am. J. Physiol.* **213**, 883–889.
- Cowles, R. B. and Dawson, W. R.** (1951). A cooling mechanism of the Texas nighthawk. *Condor* **53**, 19–22.
- Danner, R. M., Gulson-Castillo, E. R., James, H. F., Dzielski, S. A., Frank, D. C., Sibbald, E. T. and Winkler, D. W.** (2016). Habitat-specific divergence of air conditioning structures in bird bills. *Auk* **134**, 65–75.
- Dawson, W. R.** (1954). Temperature regulation and water requirements of the brown and Abert towhees, *Pipilo fuscus* and *Pipilo aberti*. In *University of California Publications in Zoology* (ed. Bartholomew, G. A., Crescitelli, F., Bullock, T. H., Furgason, W. H., and Schechtman, A. M.), pp. 81–123. Berkeley: University of California Press.
- Dawson, W. R.** (1958). Relation of Oxygen Consumption and Evaporative Water Loss to Temperature in the Cardinal. *Physiol. Zool.* **31**, 37–48.
- Dawson, W. R. and Bartholomew, G. A.** (1968). Temperature regulation and water economy of desert birds. In *Desert biology* (ed. Brown, G. W.), p. New York: Academic

Press.

- Dawson, W. R. and Bennett, A. F.** (1973). Roles of metabolic level and temperature regulation in the adjustment of western plumed pigeons (*Lophophaps ferruginea*) to desert conditions. *Comp. Biochem. Physiol.* **44A**, 249–266.
- du Plessis, K. L., Martin, R. O., Hockey, P. A. R., Cunningham, S. J. and Ridley, A. R.** (2012). The costs of keeping cool in a warming world: implications of high temperatures for foraging, thermoregulation and body condition of an arid-zone bird. *Glob. Chang. Biol.* **18**, 3063–3070.
- Gerson, A. R., Smith, E. K., Smit, B., McKechnie, A. E. and Wolf, B. O.** (2014). The impact of humidity on evaporative cooling in small desert birds exposed to high air temperatures. *Physiol. Biochem. Zool.* **87**, 782–795.
- Hackett, S. J., Kimball, R. T., Reddy, S., Bowie, R. C. K., Braun, E. L., Braun, M. J., Chojnowski, J. L., Cox, W. A., Han, K.-L., Harshman, J., et al.** (2008). A phylogenomic study of birds reveals their evolutionary history. *Science* (80-.). **320**, 1763–1768.
- Herremans, M. L. J.** (2005). Lilac-breasted Roller. In *Roberts Birds of Southern Africa VIIIth Edition* (ed. Hockey, P. A. R., Dean, W. R. J., and Ryan, P. G.), p. Cape Town: The Trustees of the John Voelcker Bird Book Fund.
- IPCC** (2007). *Climate change 2007: Fourth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press.
- IPCC** (2011). *Intergovernmental Panel on Climate Change special report on managing the risks of extreme events and disasters to advance climate change adaptation*. Cambridge: Cambridge University Press.
- Lasiewski, R. C. and Bartholomew, G. A.** (1966). Evaporative cooling in the poor-will and the tawny frogmouth. *Condor* **68**, 253–262.
- Lasiewski, R. C. and Seymour, R. S.** (1972). Thermoregulatory responses to heat stress in four species of birds weighing approximately 40 grams. *Physiol. Zool.* **45**, 106–118.
- Lasiewski, R. C., Acosta, A. L. and Bernstein, M. H.** (1966). Evaporative water loss in birds - I. Characteristics of the open flow method of determination, and their relation to estimates of thermoregulatory ability. *Comp. Biochem. Physiol.* **19**, 445–457.
- Londoño, G. A., Chappell, M. A., del Rosario Castañeda, M., Jankowski, J. E. and Robinson, S. K.** (2015). Basal metabolism in tropical birds: latitude, altitude, and the “pace of life”. *Funct. Ecol.* **29**, 338–346.
- McKechnie, A. E. and Wolf, B. O.** (2004). Partitioning of evaporative water loss in white-

- winged doves: plasticity in response to short-term thermal acclimation. *J. Exp. Biol.* **207**, 203–210.
- McKechnie, A. E. and Wolf, B. O.** (2010). Climate change increases the likelihood of catastrophic avian mortality events during extreme heat waves. *Biol. Lett.* **6**, 253–256.
- McKechnie, A. E., Whitfield, M. C., Smit, B., Gerson, A. R., Smith, E. K., Talbot, W. A., McWhorter, T. J. and Wolf, B. O.** (2016a). Avian thermoregulation in the heat: efficient evaporative cooling allows for extreme heat tolerance in four southern Hemisphere columbids. *J. Exp. Biol.* **219**, 2145–2155.
- McKechnie, A. E., Smit, B., Whitfield, M. C., Noakes, M. J., Talbot, W. A., Garcia, M., Gerson, A. R. and Wolf, B. O.** (2016b). Avian thermoregulation in the heat: evaporative cooling capacity in an archetypal desert specialist, Burchells sandgrouse (*Pterocles burchelli*). *J. Exp. Biol.* **219**, 2137.
- McKechnie, A. E., Gerson, A. R., McWhorter, T. J., Smith, E. K., Talbot, W. A. and Wolf, B. O.** (2017). Avian thermoregulation in the heat: evaporative cooling in five Australian passerines reveals within-order biogeographic variation in heat tolerance. *J. Exp. Biol.* **220**, 2436–2444.
- McNab, B. K. and Bonaccorso, F. J.** (1995). The energetics of Australasian swifts, frogmouths and nightjars. *Physiol. Zool.* **68**, 245–261.
- Muggeo, V. M. R.** (2009). Segmented: an R package to fit regression models with broken-line relationships.
- Noakes, M. J., Wolf, B. O. and McKechnie, A. E.** (2016). Seasonal and geographical variation in heat tolerance and evaporative cooling capacity in a passerine bird. *J. Exp. Biol.* **219**, 859–869.
- O'Connor, R. S., Wolf, B. O., Brigham, R. M. and McKechnie, A. E.** (2017). Avian thermoregulation in the heat: efficient evaporative cooling in two southern African nightjars. *J. Comp. Physiol. B* **187**, 477–491.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and RDevelopmentCoreTeam** (2009). nlme: Linear and nonlinear mixed effects models. R package version 3.1-94.
- Prinzinger, R., Preßmar, A. and Schleucher, E.** (1991). Body temperature in birds. *Comp. Biochem. Physiol.* **99A**, 499–506.
- Ro, J. and Williams, J. B.** (2010). Respiratory and cutaneous water loss of temperate-zone passerine birds. *Comp. Biochem. Physiol. A* **156**, 237–246.
- Smit, B., Harding, C. T., Hockey, P. A. R. and McKechnie, A. E.** (2013). Adaptive thermoregulation during summer in two populations of an arid-zone passerine. *Ecology*

94, 1142–1154.

- Smit, B., Zietsman, G., Martin, R. O., Cunningham, S. J., McKechnie, A. E. and Hockey, P. A. R.** (2016). Behavioural responses to heat in desert birds: implications for predicting vulnerability to climate warming. *Clim. Chang. Responses* **3**, 9.
- Smith, E. K., O'Neill, J., Gerson, A. R. and Wolf, B. O.** (2015). Avian thermoregulation in the heat: resting metabolism, evaporative cooling and heat tolerance in Sonoran Desert doves and quail. *J. Exp. Biol.* **218**, 3636–3646.
- Smith, E. K., O'Neill, J. J., Gerson, A. R., McKechnie, A. E. and Wolf, B. O.** (2017). Avian thermoregulation in the heat: resting metabolism, evaporative cooling and heat tolerance in Sonoran Desert songbirds. *J. Exp. Biol.* **220**, 3290–3300.
- Talbot, W. A., McWhorter, T. J., Gerson, A. R., McKechnie, A. E. and Wolf, B. O.** (2017). Avian thermoregulation in the heat: evaporative cooling capacity of arid-zone Caprimulgiformes from two continents. *J. Exp. Biol.* **220**, 3488 LP-3498.
- Tattersall, G. J., Andrade, D. V and Abe, A. S.** (2009). Heat exchange from the toucan bill reveals a controllable vascular thermal radiator. *Science* (80-.). **325**, 468–470.
- Tieleman, B. I. and Williams, J. B.** (2000). The adjustment of avian metabolic rates and water fluxes to desert environments. *Physiol. Biochem. Zool.* **73**, 461–479.
- Tieleman, B. I. and Williams, J. B.** (2002). Cutaneous and respiratory water loss in larks from arid and mesic environments. *Physiol. Biochem. Zool.* **75**, 590–599.
- Tieleman, B. I., Williams, J. B. and Bloomer, P.** (2003). Adaptation of metabolic rate and evaporative water loss along an aridity gradient. *Proc. R. Soc. London* **270**, 207–214.
- Tracy, C. R., Welch, W. R., Pinshow, B. and Porter, W. P.** (2010). *Properties of air: a manual for use in biophysical ecology, 4th edn.* Madison, WI.
- van de Ven, T. M. F. N., Martin, R. O., Vink, T. J. F., McKechnie, A. E. and Cunningham, S. J.** (2016). Regulation of heat exchange across the hornbill beak: functional similarities with toucans? *PLoS One* **11**, e0154768.
- Weathers, W. W.** (1981). Physiological thermoregulation in heat-stressed birds: consequences of body size. *Physiol. Zool.* **54**, 345–361.
- Whitfield, M. C., Smit, B., McKechnie, A. E. and Wolf, B. O.** (2015). Avian thermoregulation in the heat: scaling of heat tolerance and evaporative cooling capacity in three southern African arid-zone passerines. *J. Exp. Biol.* **218**, 1705–1714.
- Wolf, B. O. and Walsberg, G. E.** (1996). Respiratory and cutaneous evaporative water loss at high environmental temperatures in a small bird. *J. Exp. Biol.* **199**, 451–457.

Tables:

Table 1: Physiological variables (mean \pm s.d., n) related to thermoregulation at the air temperature (T_a) range we tested African cuckoo (*Cuculus gularis*; cuckoo), lilac-breasted roller (*Coracias caudatus*; roller), and Burchell's starling (*Lamprotornis australis*; starling). Minimum values represent lowest value for each individual at $T_a < 35^\circ\text{C}$, or the upper critical limit of thermoneutrality (T_{uc}) for each species. Maximum values were obtained at the highest T_a tested; when $n < 3$, we also show average values at the second highest T_a tested. We further indicate mean values for the highest T_a ($n \geq 3$), i.e. $T_a = 46^\circ\text{C}$, shared by the three species.

Physiological variable	cuckoo	roller	starling
Body mass (g)	110.00 \pm 0.02 (5)	94.50 \pm 8.50	109.10 \pm 9.30 (5)
Body temperature (T_b)			
Minimum T_b ($^\circ\text{C}$)	40.04 \pm 0.57 (6)	40.69 \pm 0.82 (11)	40.65 \pm 0.75 (5)
Inflection T_a ($^\circ\text{C}$)	32.8	36.7	34.2
T_b vs T_a slope	0.2790	0.1613	0.1843
Maximum T_b ($^\circ\text{C}$)	44.0 (2)	43.58 \pm 0.84 (5)	42.81 \pm 1.03 (3)
	43.7 (2)	43.40 \pm 0.68 (2)	43.18 (2)
Maximum T_a ($^\circ\text{C}$)	48 (2)	54 (5)	48 (3)
	50 (2)	56 (2)	50 (2)
T_b at onset of panting/gular flutter ($^\circ\text{C}$)	41.56 \pm 0.57 (5)	41.31 \pm 0.53 (9)	41.46 \pm 0.23 (5)
T_a at onset of panting/gular flutter ($^\circ\text{C}$)	40.89 \pm 0.79 (5)	41.18 \pm 1.94 (9)	35.92 \pm 4.45 (5)
Resting metabolic rate (RMR)			
Minimum RMR (W)	0.643 \pm 0.198 (5)	0.69 \pm 0.13 (8)	1.189 \pm 0.311 (5)
T_{uc} ($^\circ\text{C}$)	na	47.5	31
RMR slope (mW $^\circ\text{C}^{-1}$)	0.0224	0.0396	0.0417
Maximum RMR (W)	2.773 (1)	1.00 \pm 0.24 (4)	1.538 \pm 0.215 (3)
	0.955 (1)	1.07 \pm 0.17 (2)	1.913 (2)
Max. RMR / Min. RMR	3.06 (1)	1.45 (4)	1.29 (3)
	1.24 (1)	1.55 (2)	1.61 (2)
Evaporative water loss (EWL)			
Minimum EWL (g h^{-1})	0.481 \pm 0.144 (6)	0.85 \pm 0.38 (8)	1.00 \pm 0.67
Inflection T_a ($^\circ\text{C}$)	42.7	44.7	33.3
EWL slope (g h^{-1} $^\circ\text{C}^{-1}$)	0.2383	0.3370	0.2306
Maximum EWL (g h^{-1})	2.72 (2)	4.81 \pm (4)	3.33 \pm 0.51 (3)
	3.09 (2)	5.41 \pm (2)	4.97 (2)
Max. EWL / Min. EWL	5.66 (2)	5.66 (4)	3.33 (3)
	6.57 (2)	6.36 (2)	4.97 (2)
Minimum EHL/MHP	0.382 \pm 0.157 (5)	0.72 \pm 0.43 (8)	0.29 \pm 0.29
Maximum EHL/MHP	0.80 (1)	3.35 \pm 0.75 (4)	1.70 \pm 0.280 (3)
	2.13 (1)	3.48 \pm 0.85 (2)	1.90 (2)
EHL/MHP vs T_a slope	0.1590	0.1709	0.0704

Shared T_a = 46°C

<i>T_b (°C)</i>	43.00 ± 0.95 (5)	41.96 ± 1.30 (8)	42.61 ± 0.36 (4)
<i>RMR (W)</i>	0.857 ± 0.111 (4)	0.830 ± 0.333 (8)	1.700 ± 0.378 (4)
<i>EWL (g hr⁻¹)</i>	1.88 ± 0.28 (5)	1.86 ± 0.38 (8)	3.40 ± 1.15 (4)
<i>EHL/MHP (W)</i>	1.49 ± 0.24 (4)	1.57 ± 0.32 (8)	1.31 ± 0.20 (4)

Table 2: Slopes and upper and lower 95% confidence intervals (CI) for African cuckoos (cuckoo; *Cuculus gularis*, this study), Lilac-breasted rollers (roller; *Coracias caudatus*, this study), Burchell's starlings (starling; *Lamprotornis australis*, this study), laughing doves [dove; *Spilopelia senegalensis* (McKechnie et al., 2016a)], and Freckled nightjars [nightjar; *Caprimulgis tristigma* (O'Connor et al., 2017)] derived from linear mixed effects models. Non-overlapping 95% CI are indicated by different symbols.

Species	RMR (mW °C ⁻¹)	EWL (g h ⁻¹ °C ⁻¹)	EHL/MHP (mW/°C ⁻¹)	T _b (°C °C ⁻¹)
Cuckoo	0.0224 [0.0059, 0.0389] ^a	0.2383 [0.2030, 0.2736] ^a	0.1590 [0.1041, 0.2139] ^a	0.2790 [0.2408, 0.3172] ^{a,b}
Roller	0.0396 [0.0096, 0.0696] ^a	0.3370 [0.2802, 0.3938] ^b	0.1709 [0.1342, 0.2076] ^a	0.1613 [0.1353, 0.1874] ^{c,d}
Starling	0.0417 [0.0220, 0.0613] ^a	0.2306 [0.1900, 0.2712] ^a	0.0704 [0.0610, 0.0798] ^b	0.1843 [0.1375, 0.2311] ^{b,c,d}
Dove	0.0382 [0.0190, 0.0574] ^a	0.1999 [0.1583, 0.2415] ^a	0.0803 [0.0687, 0.0919] ^b	0.2367 [0.1920, 0.2814] ^{a,b}
Nightjar	0.0003 [-0.0015, 0.0021] ^b	0.093 [0.0812, 0.1048] ^c	0.224 [0.2044, 0.2436] ^a	0.151 [0.1334, 0.1687] ^d

Figures




Figure 1: **Body temperature (T_b) in African cuckoos (*Cuculus gularis*; n=6), lilac-breasted rollers (*Coracias caudatus*; n=10), and Burchell's starlings (*Lamprotornis australis*; n=7) as a result of air temperature (T_a).** The trendlines represent relationships between T_b and T_a above an upper inflection T_a (see methods). The slopes and intercepts were calculated using linear mixed-effects models. Significant relationships are represented by asterisks: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.




Figure 2: **Resting metabolic rate (RMR) in African cuckoos (*Cuculus gularis*; n=5), lilac-breasted rollers (*Coracias caudatus*; n=10), and Burchell's starlings (*Lamprotornis australis*; n=7) as a result of air temperature (T_a).** The trendlines represent relationships between T_b and T_a above an upper inflection T_a (see methods). The slopes and intercepts were calculated using linear mixed-effects models. Significant relationships are represented by asterisks: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.




Figure 3: **Evaporative water loss (EWL) in African cuckoos (*Cuculus gularis*; n=6), lilac-breasted rollers (*Coracias caudatus*; n=10), and Burchell's starlings (*Lamprotornis australis*; n=7) as a result of air temperature (T_a).** The trendlines represent relationships between T_b and T_a above an upper inflection T_a (see methods). The slopes and intercepts were calculated using linear mixed-effects models. For lilac-breasted rollers two significant relationships are shown ($25\text{ }^\circ\text{C} < T_a < 44.7\text{ }^\circ\text{C}$; and $T_a > 44.7\text{ }^\circ\text{C}$). Significant relationships are represented by asterisks: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.




Figure 4: **Ratio of evaporative heat loss (EHL) to metabolic heat production (MHP) in African cuckoos (*Cuculus gularis*; n=6), lilac-breasted rollers (*Coracias caudatus*; n=10), and Burchell's starlings (*Lamprotornis australis*; n=7) as a result of air temperature (T_a).** The trendlines represent relationships between T_b and T_a above an upper inflection T_a (see methods). The slopes and intercepts were calculated using linear mixed-effects models. Significant relationships are represented by asterisks: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.




Figure 5: Physiological responses as a result of temperature in five desert species making use of varying modes of respiratory heat dissipation. These include African cuckoos (cuckoo; *Cuculus gularis*, this study) and laughing doves [dove; *Spilopelia senegalensis* (McKechnie et al. 2016a)] make use of gular flutter probably synchronised with panting. Lilac-breasted rollers (roller; *Coracias caudatus*, this study) and Burchell's starlings (starling; *Lamprotornis australis*, this study) make use of panting only. Freckled nightjars [nightjar; *Caprimulgis tristigma* (O'Connor et al., 2017)] make use of rapid gular, but probably asynchronously to breathing rates. The slopes and 95% confidence levels of the relationship for each species are displayed in Table 2).

Table S1: Mass-specific values for resting metabolic rate (RMR) and evaporative water loss (EWL) related to thermoregulation at the air temperature (T_a) range we tested African cuckoo (*Cuculus gularis*; cuckoo), lilac-breasted roller (*Coracias caudatus*; roller), and Burchell's starling (*Lamprotornis australis*; starling). Minimum values represent lowest value for each individual at $T_a < 35^\circ\text{C}$, or the upper critical limit of thermoneutrality (T_{uc}) for each species. Maximum values were obtained at the highest T_a tested; when $n < 3$, we also show average values at the second highest T_a tested. We also indicate mean values for the highest T_a ($n \geq 3$), i.e. $T_a = 46^\circ\text{C}$, shared by the three species.

Physiological variable	Cuckoo	Roller	starling
<i>Resting metabolic rate (RMR)</i>			
Minimum RMR (mW g^{-1})	5.85 ± 1.79 (5)	8.91 ± 1.11 (6)	10.76 ± 1.98 (5)
Maximum RMR (mW g^{-1})	25.37 (1) 8.62 (1)	10.55 ± 3.61 (3) 11.32 (2)	14.15 ± 3.45 (3) 16.39 (2)
<i>Evaporative water loss (EWL)</i>			
Minimum EWL ($\text{mg g}^{-1} \text{h}^{-1}$)	4.45 ± 1.45 (6)	8.28 ± 2.82 (6)	8.92 ± 5.67 (5)
EWL slope ($\text{mg g}^{-1} \text{h}^{-1} \text{ }^\circ\text{C}^{-1}$)	2.10	4.03	2.07
Maximum EWL ($\text{mg g}^{-1} \text{h}^{-1}$)	30.09 (1) 27.14 (1)	52.91 ± 8.95 (3) 57.36 (30.19 ± 3.36 (3) 42.41 (2)