

Allosteric mechanisms underlying the adaptive increase in hemoglobin-oxygen affinity of the bar-headed goose

Agnieszka Jendroszek^{1,3}, Hans Malte¹, Cathrine B. Overgaard¹, Kristian Beedholm¹, Chandrasekhar Natarajan², Roy E. Weber¹, Jay F. Storz², Angela Fago^{1*}

¹Department of Bioscience - Zoophysiology, Aarhus University, Denmark

²School of Biological Sciences, University of Nebraska, United States

³Present address: Department of Molecular Biology and Genetics – DANDRITE, Aarhus University, Denmark

***Corresponding author** (angela.fago@bios.au.dk)

Keywords: high-altitude, blood, oxygen transport, allostery, adaptation, hypoxia

Summary statement

We unravel the functional mechanism responsible for the adaptive increase in hemoglobin-oxygen affinity and its allosteric regulation in bar-headed goose, a hypoxia-tolerant species renowned for its high-altitude migratory flights.

Abstract

The high blood-O₂ affinity of the bar-headed goose (*Anser indicus*) is an integral component of the biochemical and physiological adaptations that allow this hypoxia-tolerant species to undertake migratory flights over the Himalayas. The high blood-O₂ affinity of this species was originally attributed to a single amino acid substitution of the major hemoglobin (Hb) isoform, HbA, which was thought to destabilize the low-affinity T-state, thereby shifting the T-R allosteric equilibrium towards the high-affinity R-state. Surprisingly, this mechanistic hypothesis has never been addressed using native proteins purified from blood. Here, we report a detailed analysis of O₂ equilibria and kinetics of native major HbA and minor HbD isoforms from bar-headed goose and greylag goose (*Anser anser*), a strictly lowland species, to identify and characterize the mechanistic basis for the adaptive change in Hb function. We find that HbA and HbD of bar-headed goose have consistently higher O₂ affinities than those of the greylag goose. The corresponding Hb isoforms of the two species are equally responsive to physiological allosteric cofactors and have similar Bohr effects. Thermodynamic analyses of O₂ equilibrium curves according to the two-state MWC model revealed higher R-state O₂ affinities in the bar-headed goose Hbs, associated with lower O₂ dissociation rates, compared to the greylag goose. Conversely, the T-state was not destabilized and the T-R allosteric equilibrium was unaltered in bar-headed goose Hbs. The physiological implication of these results is that increased R-state affinity allows for enhanced O₂ saturation in the lungs during hypoxia, but without impairing O₂ delivery to tissues.

Introduction

Using flight paths that crest the Himalayas, bar-headed geese (*Anser indicus*) make a roundtrip annual migration between summer breeding grounds in Central Asia and wintering grounds in the Indian subcontinent (Bishop et al., 2015; Hawkes et al., 2013; Hawkes et al., 2011). Given the high energetic costs of flapping flight (Ward et al., 2002), the ability of bar-headed geese to fly at elevations of over 9000 m has earned them the respect of comparative physiologists and mountaineers alike (Hochachka and Somero, 2002; Scott, 2011; Scott et al., 2015). The physiological capacity of bar-headed geese for high-altitude flight derives from modifications of numerous convective and diffusive steps in the O₂-transport pathway (Scott, 2011; Scott et al., 2015). One of the best documented physiological adaptations of this hypoxia-tolerant species involves an increase in hemoglobin (Hb)-O₂ affinity (Black and Tenney, 1980; Jessen et al., 1991; Meir and Milsom, 2013; Natarajan et al., 2018; Petschow et al., 1977; Weber et al., 1993). The elevated Hb-O₂ affinity plays a key role in hypoxia adaptation by enhancing pulmonary O₂ loading, which helps to maintain a high arterial O₂ saturation in spite of the reduced O₂ partial pressure (P_{O₂}) of inspired air (Scott and Milsom, 2006). The fact that bar-headed geese have evolved a derived increase in Hb-O₂ affinity relative to lowland congeners, such as the greylag goose (*Anser anser*), is consistent with a general trend, as avian taxa native to extremely high elevations tend to have higher Hb-O₂ affinities than their lowland relatives (Galen et al., 2015; Natarajan et al., 2016; Natarajan et al., 2015; Projecto-Garcia et al., 2013; Storz, 2016; Zhu et al., 2018).

Previous studies of bar-headed goose Hb have focused exclusively on the major HbA isoform ($\alpha^A_2\beta^A_2$), which typically accounts for 70-80% of total Hb in the adult red blood cells of waterfowl species, with the minor HbD isoform ($\alpha^D_2\beta^A_2$) accounting for the remaining fraction (Grispo et al., 2012; Natarajan et al., 2015; Opazo et al., 2015). Protein-engineering studies have demonstrated that the increased O₂-affinity of bar-headed goose HbA is largely attributable to a rare substitution at the intradimer $\alpha_1\beta_1$ interface of the tetrameric HbA, $\alpha^A119\text{Pro}\rightarrow\text{Ala}$ (Jessen et al., 1991; Weber et al., 1993), but two other α -chain substitutions, $\alpha^A18\text{Gly}\rightarrow\text{Ser}$ and $\alpha^A63\text{Ala}\rightarrow\text{Val}$, also make sizable contributions (Natarajan et al., 2018). Perutz noted that the $\alpha^A119\text{Pro}\rightarrow\text{Ala}$ substitution eliminates a van der Waals interaction between $\alpha^A119\text{Pro}$ and $\beta^A55\text{Leu}$, residues located on opposing subunits of the same $\alpha_1\beta_1$ interface, and suggested that the loss of this atomic contact destabilizes the low-affinity T-state conformation of the Hb tetramer (Perutz, 1983). According to the two-state Monod-Wyman-Changeaux (MWC) allosteric model (Monod et al., 1965), destabilization of the low-affinity T-state (the prevailing conformation of deoxy Hb) has the effect of shifting the T-R allosteric equilibrium between the two quaternary conformations in favor of the high-affinity R-state (the prevailing conformation of oxy Hb). Thus, according to Perutz's hypothesis (Perutz, 1983), the increased O₂-affinity of bar-headed goose HbA is caused by a single, large-effect substitution that reduces the MWC allosteric constant L (the ratio of the two conformational states [T]/[R] of the deoxy Hb) and increases the O₂ association equilibrium constant for T-state Hb, K_T . According to this mechanism, the O₂ association equilibrium constant for R-state Hb, K_R , should remain unaltered.

Although this postulated mechanism for high-altitude adaptation has never been tested in native Hbs, early experiments involving recombinant human Hb with the engineered $\alpha^A119\text{Pro}\rightarrow\text{Ala}$ mutation appeared to support Perutz's hypothesis (Jessen et al., 1991; Weber et al., 1993). Nonetheless, there are reasons to suspect that our understanding of the mechanistic basis of this classic example of biochemical adaptation is still far from complete. First, given the function-altering effects of the other α^A -chain substitutions,

$\alpha^A18\text{Gly}\rightarrow\text{Ser}$ and $\alpha^A63\text{Ala}\rightarrow\text{Val}$ (Natarajan et al., 2018), information about the effects of $\alpha^A119\text{Pro}\rightarrow\text{Ala}$ alone cannot be expected to provide a complete picture. Second, mutagenesis experiments have revealed that identical Hb mutations often have different functional effects on different genetic backgrounds (Kumar et al., 2017; Natarajan et al., 2013; Storz, 2018; Tufts et al., 2015). Thus, the measured effects of engineered mutations in recombinant human Hb (Jessen et al., 1991; Weber et al., 1993) may not perfectly recapitulate the effects of the same mutational changes in the Hbs of other species, especially in taxa as divergent as birds and mammals (human Hb and bar-headed goose HbA differ at 89 of 287 sites in each $\alpha\beta$ half-molecule). In light of these considerations, we performed a set of thermodynamic (equilibrium) and kinetic experiments to identify and characterize the allosteric mechanism responsible for the evolved increase in the O_2 -affinity of bar-headed goose HbA and its physiological regulation by anions and pH. Specifically, we measured O_2 equilibrium curves and effects of allosteric cofactors in purified, native iso-Hbs of bar-headed goose and greylag goose and analyzed these curves to estimate allosteric parameters (i.e. L , K_T and K_R) describing Hb function according to the two-state MWC model (Monod et al., 1965). To provide a complete assessment of the factors underlying evolved differences in blood O_2 -affinity, we examined the functional properties of the major HbA isoform as well as the minor HbD isoform of both species.

Materials and methods

Blood samples and Hb purification

Blood was sampled from a single, adult specimen of bar-headed goose and greylag goose housed in outdoors pens (Galten, Aarhus). Animals were gently restrained and after exposing the brachial vein at the wing elbow joint, blood was collected by using 5-ml heparinized syringes, with the needle pointing towards the tip of the wing. The whole procedure took less than 2 min. All animal procedures were approved by the Danish Law for Animal Experimentation (permit 2018-15-0201-01507).

We used an aliquot of blood from the same animal for genotyping the adult-expressed globin genes (GenBank accession numbers: MH375701-MH375710). We isolated total RNA using the RNeasy kit (Qiagen). We amplified full-length cDNAs for the major adult-expressed globin genes using a OneStep RT-PCR kit (Qiagen), and we then cloned RT-PCR products using the TOPO TA Cloning Kit (Life Technologies), and we sequenced at least five clones per gene (α^A , α^D , β^A) to recover both alleles of each globin gene (Natarajan et al., 2016; Natarajan et al., 2015). Red blood cells (RBCs) were separated from plasma by centrifugation (3000 g, 15 min), washed in 0.9 % NaCl, and lysed on ice (30 min) by adding a 4-fold volume of ice-cold 10 mM Hepes, pH 7.4, 0.5 mM EDTA. The resulting hemolysate was centrifuged (12000 g, 20 min) and the supernatant loaded on a PD-10 5-ml desalting column (GE Healthcare Life Sciences) equilibrated with 10 mM Hepes, pH 7.4, 0.5 mM EDTA. Prior to loading, samples were added NaCl (to 0.2 M final concentration), to facilitate removal of endogenous organic phosphates (stripping). Avian Hb isoforms HbA and HbD were then separated by anion exchange chromatography on a HiTrap Q HP 5-ml column connected to a Äkta Pure Chromatography System (GE Healthcare Life Sciences) equilibrated with 10 mM Tris-HCl pH 8.6, 0.5 mM EDTA and eluted with a linear gradient of 0-250 mM NaCl, at a flow rate of 1 ml min^{-1} . Eluate absorbance was monitored at 415 and 280 nm. This protocol also removed residual endogenous phosphates (Bonaventura et al., 1999; Rollema and Bauer, 1979). The purity of HbA and HbD isoforms was verified by isoelectric-focusing on precast polyacrylamide gels (pH range 3-9) using the

PhastSystem (GE Healthcare Life Sciences). Purified HbA and HbD were then desalted by dialysis (Slide-A-Lyzer, Thermo Scientific) against 10 mM Hepes, pH 7.6, 0.5 mM EDTA, concentrated by ultrafiltration (Amicon Ultra Centrifugal Devices, Millipore) to a final heme concentration > 1 mM and stored in aliquots at -80 °C. Heme concentration (oxy derivative) was determined by absorption spectra (400-700 nm), and lack of heme oxidation was verified by the relative intensities of absorbance peaks at 575 and 541 nm of the oxy form (Van Assendelft and Zijlstra, 1975).

O₂ equilibrium curves: determination of P_{50} and cooperativity

Purified HbA and HbD (0.3 mM heme) were dissolved in 0.1 M Hepes, pH 7.4, in the absence (stripped) and presence of the allosteric effectors KCl (0.1 M) and inositol hexaphosphate (IHP, 0.15 mM) added separately or in combination. IHP is a chemical analog of the avian RBC organic phosphate inositol pentaphosphate (Johnson and Tate, 1969; Rollema and Bauer, 1979). O₂ equilibria were measured in duplicate at 37 °C in 5- μ l samples using a thin-layer modified diffusion chamber described in detail elsewhere (Damsgaard et al., 2013; Janecka et al., 2015; Tufts et al., 2015; Weber, 1981; Weber, 1992; Weber et al., 1993). Absorbance at 436 nm was recorded to derive the fractional O₂ saturation after stepwise equilibration with humidified gas mixtures (ultrapure N₂ and O₂ or air) of known O₂ tension (PO_2 , torr). Absolute absorption data were samples at a rate of 1 Hz, and PO_2 values were entered into the data series using in-house made software (Beedholm, 2018). Using the same program, saturation values were determined on the fly from relative absorbance and the nominally set PO_2 values. An assumption was made that any drift in recorded absorption values was linear over the recording interval. For each Hb isoform and buffer condition, P_{50} and n_{50} (O₂ tension and Hill cooperativity coefficient at half-saturation, respectively) were derived (mean \pm SE) for by non-linear fitting of 4-6 saturation data points (O₂ saturation range \sim 0.2-0.8) using the SigmaPlot 12 (Systat.com) software. Data were fitted ($r^2 > 0.99$) according to the sigmoidal Hill's equation $Y = PO_2^{n_{50}} / (P_{50}^{n_{50}} + PO_2^{n_{50}})$, where Y is the fractional saturation. The allosteric binding of IHP to each Hb (0.3 mM heme) was analysed from plots of P_{50} vs. IHP concentration (range 0-0.75 mM). For HbD, the IHP equilibrium dissociation constant (corresponding to the IHP concentration eliciting half of the maximal P_{50}) was derived from non-linear fitting to a sigmoidal Hill's equation ($r^2 > 0.97$ and $r^2 > 0.99$ for the bar-headed goose and greylag goose HbD, respectively). The Bohr effect was measured under identical buffer conditions as described above, in the absence and presence of IHP, and in the pH range 6.7-7.8 for HbA and 6.7-7.4 for HbD. In these pH ranges, the Bohr factor ($\Phi = \Delta \log P_{50} / \Delta \text{pH}$) was constant under all conditions ($r^2 > 0.99$).

O₂ equilibrium curves: determination of MWC allosteric parameters

To perform a detailed analysis of the allosteric regulation of O₂ binding in HbA and HbD in terms of the two-state MWC model (Monod et al., 1965) it is necessary to cover a broad range of fractional saturations, including values approaching zero ($< 5\%$) and full ($> 95\%$) O₂ saturation. To this end, we performed 5-9 separate experiments as described above in the absence (stripped) and presence of saturating IHP concentration (0.75 mM), each experiment covering partially overlapping saturations. Data from all experiments were collectively analyzed by non-linear curve fitting using the MWC allosteric model (Monod et al., 1965). This model assumes the presence of an equilibrium between a low-affinity T-state and a high-affinity R-state, with K_T and K_R (torr⁻¹) as O₂ association equilibrium constants, respectively, and with the equilibrium constant L describing the [T]/[R] ratio in the absence of O₂ (or any other heme ligand).

Following a previous study (Weber et al., 1993), the allosteric parameters K_T , K_R and L were thus derived under each buffer condition by fitting ($r^2 > 0.99$) the saturation function (Eq. 1) to the data:

$$Y = \frac{L K_T P (1 + K_T P)^3 + K_R P (1 + K_R P)^3}{L (1 + K_T P)^4 + (1 + K_R P)^4} \quad (\text{Eq. 1})$$

where P is the PO_2 . Saturation data were corrected by extrapolation for possible incomplete saturation with pure O_2 or desaturation with pure N_2 , as described previously (Fago et al., 1997).

Stopped-flow kinetic measurements of O_2 dissociation rates

O_2 dissociation rates (k_{off} , s^{-1}) of HbA and HbD were measured by stopped-flow using an OLIS RSM 1000 UV/Vis rapid-scanning spectrophotometer (OLIS, Bogart, GA, USA) equipped with an OLIS data collection software, as described (Helbo and Fago, 2012). In these experiments, one syringe of the stopped-flow contained 10 μM oxy Hb in 200 mM Hepes buffer pH 7.4, in the absence (stripped) and presence of 10 μM IHP alone and with 6.6 mM KCl (using the same heme/KCl ratio as in equilibrium experiments). The other syringe contained 40 mM sodium dithionite freshly prepared in deoxygenated buffer. The content of the two syringes was rapidly mixed at a 1:1 ratio at 37 °C by stopped-flow and the O_2 dissociation reaction (where dithionite acts as an O_2 scavenger) was followed by monitoring absorbance at 431 nm (peak of deoxy Hb). The reaction was completed within 0.15 s. Traces were best fitted ($r^2 > 0.99$) by a monoexponential fitting using the GraphPad Prism 6.01, yielding the O_2 dissociation rate (k_{off} , s^{-1}) under each buffer condition. Rates are presented as mean \pm SE from 8-9 and 4-5 separate experiments for HbA and HbD, respectively.

Gel filtration experiments

Apparent molecular weights of oxy and deoxy HbA and HbD were determined by gel-filtration chromatography using a Superdex 75 10/300 GL column connected to a Äkta Pure Chromatography System (GE Healthcare Life Sciences) and equilibrated with 50 mM Tris-HCl, pH 7.4, 0.15 M NaCl, 0.5 mM EDTA (elution buffer). Hb was loaded at a concentration of 0.3 mM heme. Flow rate was 0.5 ml min^{-1} and absorbance of the eluate was monitored at 280 and 414 nm. In experiments with deoxy Hb, elution buffer was bubbled with N_2 for at least 2 h before chromatographic runs and contained 0.3 mg ml^{-1} sodium dithionite, to remove possible residual oxygen (Fago and Weber, 1995; Storz et al., 2015; Weber et al., 2013). The effect of IHP on the molecular weight of the Hb was assessed by adding IHP (0.75 mM) to the elution buffer. Samples (0.3 mM heme) of adult human Hb (Sigma Aldrich), *Scapharca inaequivalvis* HbI (kindly provided by Prof. E. Chiancone) and horse myoglobin (Sigma Aldrich) were loaded to estimate elution times corresponding to tetrameric, dimeric and monomeric globin structures, respectively. As tetrameric human Hb tends to dissociate into dimers in gel filtration experiments, where it displays a lower apparent molecular weight (Ackers, 1964; Chiancone, 1968), these chromatographic experiments provide estimates of the type of quaternary assembly but not strictly of molecular weights (Fago et al., 2018; Kumar et al., 2017).

Results

Cloning and sequencing of the adult-expressed globin genes from the bar-headed goose and greylag goose specimens confirmed previously documented amino acid substitutions that distinguish the HbA and HbD isoforms of the two species (Hiebl, 1987; McCracken et al., 2010; Oberthür, 1982) (Fig. 1). An alignment of orthologous sequences from 12 additional waterfowl species in the subfamily Anserinae revealed that all amino acid replacements in the α^A - and α^D -globin genes were specific to the bar-headed goose lineage. By contrast, the β^A -globin replacements occurred in the common ancestor of greylag goose and all other lowland *Anser* species after divergence from bar-headed goose (Fig. 1) (Natarajan et al., 2018). In addition to the six fixed differences in the three adult-expressed globin genes of bar-headed and greylag goose (four affecting HbA and two affecting HbD), the bar-headed goose specimen was heterozygous at two sites α^A12 (Gly/Ala) and β^A125 (Asp/Glu) (Fig. 1). The allelic polymorphism at α^A12 has been documented previously (McCracken et al., 2010). Available sequence data from bar-headed geese (Hiebl, 1987; McCracken et al., 2010; Oberthür, 1982) indicate that α^A12 Ala and β^A125 Glu represent low-frequency amino acid variants. Since the HbD isoforms of bar-headed goose and greylag goose have not been examined previously, it is noteworthy that α^D -globin orthologs of the two species are distinguished by replacements at α^D3 (Thr→Ser) and α^D48 (Leu→Val) (Fig. 1). A sequence alignment of α^A and α^D from both species is shown in supplementary Fig. S1.

We purified native HbA and HbD isoforms from bar-headed goose and greylag goose hemolysates by anion exchange chromatography (Fig. S2). The HbA:HbD ratios estimated from the chromatographic profiles were ~86:14 in the bar-headed goose and ~90:10 in the greylag goose, consistent with data from other waterfowl taxa (Grispo et al., 2012; Natarajan et al., 2015; Opazo et al., 2015). O₂ equilibrium curves of purified bar-headed goose HbA and HbD were left-shifted relatively to the respective Hbs of greylag goose (Fig. 2A) and showed consistently lower P_{50} values under all effector conditions examined (Table S1). Specifically, in the presence of IHP and KCl, the P_{50} values of bar-headed goose HbA and HbD were 30 % and 31%, respectively, similarly lower than the corresponding greylag goose isoHbs. O₂ binding was cooperative in all cases ($n_{50} = 1.51-3.67$) (Table S1), reflecting normal allosteric T-R shift upon oxygenation, with no evidence of super-cooperativity (Hill coefficients > 4) described for some avian Hbs (Riggs, 1998). Under physiologically relevant conditions resembling those in the RBCs, i.e. in the presence of IHP and KCl, bar-headed goose HbA had a much lower P_{50} (34.71 torr) than that (49.37 torr) of the greylag goose (Table S1). These P_{50} values are highly consistent with those reported for blood (Petschow et al., 1977). Stripped (anion-free) bar-headed goose HbA had also a slightly lower P_{50} than greylag goose HbA (3.72 and 4.24 torr, respectively; Table S1). Together, these data indicate that the difference in intrinsic O₂ affinities (i.e. in the absence of allosteric anionic cofactors) between the HbA isoforms of the two species is greatly amplified in the presence of the physiological cofactor IHP, which had strong effects on HbA and HbD oxygenation (Fig. 2A,B) (Table S1). The minor HbD had a consistently higher O₂ affinity than HbA in each of the two species (Table S1).

In both species the Bohr effect of the two Hb isoforms was highly similar and was slightly enhanced by IHP (Fig. 2C,D; Table S1), as expected from anion-induced increase in Bohr proton binding. The Bohr factors were in good agreement with those reported earlier for native HbA of the two species (Rollema and Bauer, 1979) and for the recombinant human Hb wild type and α^A119 Pro→Ala mutant (Weber et al., 1993).

For both Hb isoforms, the anion IHP was a more potent allosteric effector than chloride, as it induced a larger right-shift in the O₂ equilibrium curve (Fig. 2A,B). Moreover, O₂ equilibrium curves were less right-shifted with KCl and IHP than with IHP alone (Fig. 2), indicating that these anions compete for binding sites on the Hb molecule. For the HbA and HbD isoforms of both species, we quantified the allosteric effect of IHP by measuring O₂ equilibrium curves in the presence of increasing IHP concentrations (Fig. 3). These experiments revealed that IHP binding to the HbA of both species was essentially stoichiometric, as full saturation was achieved at a 1:1 molar ratio of IHP to Hb tetramer (Fig. 3A). This finding indicates that at equimolar ratio of protein to effector there is essentially no free IHP, reflecting a very high affinity for this effector (i.e. very low equilibrium dissociation constant < 10⁻⁸ M), fully consistent with previous studies (Rollema and Bauer, 1979). The HbD isoforms of both species also bound IHP with high affinity (albeit lower than for HbA), with estimated values (means ± SE) of equilibrium dissociation constants of 5.0 ± 0.1 × 10⁻⁵ M (bar-headed goose HbD) and 5.2 ± 0.5 × 10⁻⁵ M (greylag goose HbD) (Fig. 3B).

The O₂ dissociation rate of oxy Hb (k_{off} , s⁻¹) was in all cases higher in the presence of IHP (Table 1). HbA from bar-headed goose showed lower O₂ dissociation rates than greylag goose HbA under all conditions (Table 1), indicating that the amino acid substitutions that distinguish this isoHb in the two species alter the dissociation pathway of heme-bound O₂. Bar-headed goose HbA and HbD dissociated O₂ at nearly identical rates under the same conditions, whereas in the greylag goose HbA dissociated O₂ slightly faster than HbD (Table 1). The fact that kinetic traces were best fitted by monoexponential functions further indicated that α and β hemes dissociated O₂ at similar rates.

Having established that both HbA and HbD had higher O₂ affinity in the bar-headed goose than in the greylag goose, especially in the presence of IHP (Table S1, Figs. 2,3), we then examined the origin of this functional difference in terms of the two-state MWC allosteric model (Monod et al., 1965). To this end, we measured O₂ equilibrium curves in the absence and presence of saturating IHP to cover a wide span of O₂ saturations between zero and full saturation, and we fitted the data according to the MWC model (Eq. 1) (Fig. 4). The derived allosteric MWC parameters (Table S2) included the O₂ association equilibrium constants of T- and R-state (K_T and K_R , respectively, torr⁻¹) and the allosteric constant L , the [T]/[R] ratio in the absence of ligand. When comparing HbA of bar-headed goose and greylag goose, K_R was consistently higher in the bar-headed goose (Fig. 4E) by approximately 12% and 26% in the absence and presence of IHP, respectively (Table S2). In addition, K_T of the bar-headed goose HbA was consistently lower (Fig. 4C), whereas the allosteric constant L was the same (Fig. 4G). The minor isoform HbD of the bar-headed goose had higher K_T and K_R than HbD of the greylag goose (Fig. 4D,F), but highly similar L (Fig. 4H). For both HbA and HbD of each species, the IHP strongly decreased both the T-state O₂ association equilibrium constant, K_T (Fig. 4C,D) and the R-state O₂ association constant, K_R (Fig. 4E,F), while also increasing L (Fig. 4G,H). These findings indicate that IHP binds preferentially to the T state, thereby decreasing its O₂ affinity and shifting the T-R equilibrium of the deoxy Hb towards the T-state (increasing L), but that it also affects the R-state. Comparison of MWC parameters for HbD with those for HbA (Table S2, Fig. 4) shows lower K_R (Fig. 4E,F) but higher K_T (Fig. 4C,D) and lower L (Fig. 4G,H) in the latter isoform. Higher K_T and lower L explain the higher overall O₂ affinity of HbD compared to HbA in terms of P_{50} (Table S1).

To determine whether IHP may affect O₂ binding properties via changes in the structural assembly of the Hb, we examined the apparent molecular weight of oxy and deoxy HbA and HbD of both species by gel-filtration in the absence and presence of IHP. Although HbA and HbD were mainly present in the tetrameric

form under all conditions (Fig. 5), an extended 'tail' observed in the elution profile in the absence of IHP indicated the presence of a small fraction of dissociation products (caused by protein dilution during column migration) that almost disappeared in deoxygenated conditions and in the presence of IHP. For comparison, human Hb maintained the same elution profile regardless of IHP (Fig. 5). Taken together, these results indicate that HbA and HbD from bar-headed and greylag goose may form less tight subunit interactions than human Hb and that IHP binding stabilizes the tetrameric assembly considerably.

Discussion

A key finding of this study is that the evolved increase in O_2 affinity (decrease in P_{50}) of bar-headed goose HbA compared to greylag goose HbA is attributable to an increased O_2 association equilibrium of the R-state (higher K_R) (Fig. 4E) and not from a less stable T-state, as earlier proposed (Perutz, 1983). The HbA of the two species have in fact identical L values under identical conditions (Fig. 4G), which reflects a similar stability of the T state relative to the R state. The higher R-state O_2 association constant of the bar-headed goose HbA derives, at least in part, from a slower O_2 dissociation of the fully saturated protein compared to the greylag goose HbA (Table 1). This property may reflect the slightly different subunit orientation found in the crystal structure of the bar-headed goose oxy HbA compared to human Hb, although the α and β heme pockets appeared very similar (Zhang et al., 1996). The higher R-state O_2 association constant of the bar-headed goose HbA has important physiological implications. Since Hb is in the R-state in the lungs, where the O_2 tension is high, the higher K_R of bar-headed goose HbA would enhance O_2 loading in the upper part of the blood O_2 equilibrium curve, well after the T-R transition has taken place at intermediate saturations. Furthermore, the T-state HbA has a lower O_2 association equilibrium constant (lower K_T) in the bar-headed goose than in the greylag goose (Fig. 4C), a property that would favor O_2 unloading at the low O_2 tensions prevailing in working muscles. However, this decrease in K_T is not large enough to offset the concomitant increase in K_R , and the resulting overall O_2 affinity (in terms of P_{50}) is therefore higher in the bar-headed goose HbA compared to the greylag goose HbA (Fig. 2A). Taken together, our results show how opposite functional effects on each of the two quaternary states of the major HbA isoform may act together to improve O_2 transport (increase the difference between loading and unloading) in the bar-headed goose at high altitudes. In contrast to our functional analysis, introduction of the $\alpha^A119\text{Pro}\rightarrow\text{Ala}$ mutation into recombinant human Hb had the effect of increasing the T-state O_2 association constant and decreasing L , while leaving the R-state O_2 association constant unaffected (Weber et al., 1993) (Table S2). These changes, although still consistent with a decrease in P_{50} , would have the unwanted *in vivo* consequence of disfavoring unloading of O_2 to tissues, while leaving O_2 loading at the lungs almost unaffected.

Another factor incrementing arterial blood O_2 saturation in the bar-headed goose is the respiratory alkalosis initially experienced at low inspired PO_2 (Black and Tenney, 1980; Lague et al., 2016), which would increase overall O_2 affinity via the Bohr effect (Table S1). However, given that bar-headed and greylag goose hemoglobins display highly similar Bohr factors, the increment in blood O_2 saturation in the two goose species would be similar for a given increment in blood pH. Moreover, studies on acid-base regulation in bar-headed and greylag goose reveal the onset of a metabolic acidosis at low inspired PO_2 (Scott et al., 2007; Lague et al., 2016).

Our analysis in terms of the MWC model shows that IHP causes marked reductions in the O₂ association equilibrium constant of the T (Fig. 4C,D) and R states (Fig. 4E,F) and a strong shift of the T-R state equilibrium towards the T state, as reflected by increased L (Fig. 4G,H). These results indicate that IHP binding to HbA and HbD induces conformational changes that affect heme-O₂ binding within each of the two quaternary states. These functional effects of IHP can be ascribed in part to enhanced heme-O₂ dissociation rates (Table 1). Binding of IHP to the HbA of both species was particularly strong and followed stoichiometric binding (Fig. 3A), supporting earlier reports of extremely high affinities for IPP (Rollema and Bauer, 1979). By virtue of the strong allosteric effect of IPP on HbA oxygenation (Rollema and Bauer, 1979), altitude-related change in IPP levels in the RBCs will effectively cause marked left- or right-shifts in the blood O₂ equilibrium curves. It should be borne in mind that the bar-headed goose is exposed to high altitude only during seasonal migrations, while spending long periods of the year at low to moderate altitudes (Bishop et al., 2015; Hawkes et al., 2013; Hawkes et al., 2011; Scott et al., 2015). Changes in RBC levels of organic phosphates result in immediate and reversible adjustments in blood O₂ transport properties to altitude variations, as found in species, where the levels of organic phosphates change during hypoxia (Fago, 2017; Weber, 2007; Weber and Fago, 2004). Whether IPP levels of RBCs change significantly upon altitude exposure in the bar-headed goose remains to be determined. Although some studies indicate that IPP levels in bird RBCs remain constant (Jaeger and McGrath, 1974; Lutz, 1980), others have reported slightly increased arterial O₂ content in bar-headed geese reared at high altitudes (Lague et al., 2016) that suggest decreased IPP content in RBCs.

The structure of the IPP binding site in the bar-headed goose provides a basis for these functional effects. The IPP binding site corresponds to a constellation of positively charged amino acid residues located in the cleft between the β subunits (Arnone and Perutz, 1974; Grispo et al., 2012; Riccio, 2001; Tamburrini et al., 2000). These positive residues are neutralized when bound to negatively charged IPP or IHP. Our data on stripped HbA and HbD (Fig. 4) indicate that, in the absence of IHP, extensive electrostatic repulsions in this cluster of charged residues effectively destabilize the T-state and shift the T-R equilibrium towards the R-state, as reflected by a low L (Fig. 4G,H; Table S2). This may explain why the low-salt crystal structure of bar-headed goose deoxy HbA is essentially in the R state (Liang et al., 2001). When IHP neutralizes this positively charged patch, the T-R equilibrium shifts towards the low-affinity T-state (as shown by the large increase in L) (Fig. G,H), and the O₂ affinity consequently decreases (Fig. 2A,B). The view that an excess of positively charged residues located at specific protein clusters may effectively promote Hb T-R switches is not new and has been invoked earlier to explain other allosteric effects in vertebrate Hbs, such as chloride effects (Bonaventura et al., 1994), Root effect (Mylvaganam et al., 1996), reverse Bohr effect and high phosphate sensitivities (Fago et al., 1995). In addition, excess of positive charges at the IPP binding site cluster may explain the higher tendency to dissociate into dimers in geese Hbs compared to human Hb in gel filtration experiments, a tendency that is reversed by IHP (Fig. 5). The less defined patch of positively charged residues at the IPP-binding site of HbD compared to HbA (Grispo et al., 2012) is moreover in agreement with the lower binding affinity of IHP to HbD (Fig. 3B).

The minor isoform HbD has an overall higher O₂ affinity (i.e. lower P_{50}) than HbA (Table S1), consistent with reports from previous studies of isoHb differentiation in birds (Grispo et al., 2012; Natarajan et al., 2016; Natarajan et al., 2015; Opazo et al., 2015; Zhu et al., 2018). However, the MWC analysis revealed that HbD has a lower K_R than HbA (Table S2, Fig. 4), and is therefore less suited to enhance O₂ loading at the lungs at high altitude. Moreover, HbD is equally sensitive to pH changes as HbA (Fig. 2C,D; Table S1) but it is less

sensitive than HbA to IHP (Fig. 3), and by inference less responsive to potential hypoxia-induced changes in IPP levels in RBC. Thus, although HbA and HbD of bar-headed goose both evolved similar increases in O₂-affinity relative to the corresponding isoHbs of greylag goose, the allosteric properties of HbA appear better suited to high-altitude O₂ transport. Since the two isoHbs share the same β^A -chains, the observed differences in O₂-affinity and effector sensitivity must be attributable to the net effect of 61 amino acid substitutions that distinguish the α^A - and α^D -globin paralogs (Fig. S1).

Conclusions

Our experimental data confirm that the major isoform HbA of bar-headed goose has a higher O₂-affinity than that of greylag goose, consistent with previous studies (Black and Tenney, 1980; Jessen et al., 1991; Petschow et al., 1977). For the first time we document a quantitatively similar species difference in O₂-affinity of the minor HbD isoform. Most importantly, the experimental data shed light on the functional mechanism responsible for the evolved changes in bar-headed goose HbA and HbD.

Of the amino acid substitutions that distinguish HbA and HbD of bar-headed goose from the corresponding Hb isoforms of greylag goose, phylogenetic evidence clearly indicates that all α^A and α^D substitutions occurred in the bar-headed goose lineage, whereas all β^A substitutions occurred in common ancestor of all lowland *Anser* species (including *Anser anser*) after diverging from the line of descent leading to bar-headed goose (Natarajan et al., 2018) (Fig. 1). In the case of the major HbA isoform, site-directed mutagenesis experiments involving reconstructed ancestral Hbs demonstrated that the evolved increase in HbA O₂ affinity of bar-headed goose compared to greylag goose (Fig. 2; Table S1) is mainly attributable to the net effect of the three α^A -substitutions that are specific to bar-headed goose at α^A sites 18, 63, and 119 (Fig. 1) (Natarajan et al., 2018). Since HbA and HbD share the same β^A -chain subunit, it is equally clear that the evolved increase in HbD O₂-affinity (Fig. 2; Table S1) is attributable to the independent or joint effects of the two amino acid replacements at α^D sites 3 and 48 in the α^D -globin of bar-headed goose (Fig. 1). Thus, substitutions in the α -type chains of both isoHbs contribute to the increased overall blood-O₂ affinity of the high-flying bar-headed goose relative to the lowland greylag goose.

Our combination of equilibrium and kinetic measurements provide novel insights into the functional mechanism responsible for the evolved increase in Hb-O₂ affinity in the bar-headed goose. Contrary to Perutz's hypothesis (Perutz, 1983), the increased O₂-affinity of bar-headed goose HbA is not attributable to an evolved change in the equilibrium constant between the T and R conformations of the protein, L , as HbA of bar-headed goose and greylag goose are identical in this regard (Fig. 4G). Instead, in the presence of physiological anion concentrations, the increased HbA O₂-affinity of bar-headed goose is primarily attributable to an increase in K_R , the O₂ association constant for Hb in the R-state (Fig. 4E), thereby favoring loading at the lungs. This increase in K_R overrides a concomitant decrease in K_T , the O₂ association constant for Hb in the T-state (Fig. 4C), which favors O₂ unloading to severely hypoxic tissues. The increased O₂-affinity of bar-headed goose HbD is primarily attributable to combined increases in K_R and K_T , but not in L (Fig. 4D-H). The particularly strong allosteric effect of organic phosphates on the O₂ affinity of HbA suggests a potential adaptive role for blood O₂ transport during rapid changes in altitudes.

The elevated Hb-O₂ affinity of bar-headed goose is an iconic, textbook example of biochemical adaptation (Hochachka and Somero, 2002; Scott et al., 2015). The findings presented here provide us with a more complete understanding of the mechanistic basis of the evolved change in protein function.

Acknowledgments

We thank Elin E. Petersen (Aarhus) for skilled technical assistance.

Competing interests

The authors declare no competing interests.

Funding

This work was funded by the National Institutes of Health / National Heart, Lung and Blood Institute (HL087216 to JFS), the National Science Foundation (MCB- 1517636 and RII Track-2 FEC-1736249 to JFS), and the Danish Council for Independent Research, Natural Sciences (4181-00094 to AF).

References

- Ackers, G. K.** (1964). Molecular exclusion and restricted diffusion processes in molecular-sieve chromatography. *Biochemistry* **3**, 723-730.
- Arnone, A. and Perutz, M. F.** (1974). Structure of inositol hexaphosphate–human deoxyhaemoglobin complex. *Nature* **249**, 34-36.
- Beedholm, K.** (2018) "Spektrosampler", Aarhus University, Denmark, <http://www.marinebioacoustics.com/sharing/spektrosampler.zip>
- Bishop, C. M., Spivey, R. J., Hawkes, L. A., Batbayar, N., Chua, B., Frappell, P. B., Milsom, W. K., Natsagdorj, T., Newman, S. H., Scott, G. R. et al.** (2015). The roller coaster flight strategy of bar-headed geese conserves energy during Himalayan migrations. *Science* **347**, 250-254.
- Black, C. P. and Tenney, S. M.** (1980). Oxygen transport during progressive hypoxia in high-altitude and sea-level waterfowl. *Respir. Physiol.* **39**, 217-239.
- Bonaventura, C., Arumugam, M., Cashion, R., Bonaventura, J. and Moo-Penn, W. F.** (1994). Chloride masks effects of opposing positive charges in Hb A and Hb Hinsdale (β 139 Asn \rightarrow Lys) that can modulate cooperativity as well as oxygen affinity. *J. Mol. Biol.* **239**, 561-568.
- Bonaventura, C., Ferruzzi, G., Tesh, S. and Stevens, R. D.** (1999). Effects of S-Nitrosation on oxygen binding by normal and sickle cell hemoglobin. *J. Biol. Chem.* **274**, 24742-24748.
- Chiancone, E.** (1968). Dissociation of hemoglobin into subunits. II. Human oxyhemoglobin: gel filtration studies. *J. Biol. Chem.* **243**, 1212-1219.
- Damsgaard, C., Storz, J. F., Hoffmann, F. G. and Fago, A.** (2013). Hemoglobin isoform differentiation and allosteric regulation of oxygen-binding in the turtle *Trachemys scripta*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **305**, R961-R967.
- Fago, A.** (2017). Functional roles of globin proteins in hypoxia-tolerant ectothermic vertebrates. *J. Appl. Physiol.* **123**, 926-934.
- Fago, A., Bendixen, E., Malte, H. and Weber, R. E.** (1997). The anodic hemoglobin of *Anguilla anguilla*. Molecular basis for allosteric effects in a Root-effect hemoglobin. *J. Biol. Chem.* **272**, 15628-15635.
- Fago, A., Carratore, V., Prisco, G. d., Feuerlein, R. J., Sottrup-Jensen, L. and Weber, R. E.** (1995). The cathodic hemoglobin of *Anguilla anguilla*. Amino acid sequence and oxygen equilibria of a reverse Bohr effect hemoglobin with high oxygen affinity and high phosphate sensitivity. *J. Biol. Chem.* **270**, 18897-18902.
- Fago, A., Rohlfig, K., Petersen, E. E., Jendroszek, A. and Burmester, T.** (2018). Functional diversification of sea lamprey globins in evolution and development. *Biochim. Biophys. Acta (BBA) - Proteins and Proteomics* **1866**, 283-291.
- Fago, A. and Weber, R. E.** (1995). The hemoglobin system of the hagfish *Myxine glutinosa*: aggregation state and functional properties. *Biochim. Biophys. Acta* **1249**, 109-115.
- Galen, S. C., Natarajan, C., Moriyama, H., Weber, R. E., Fago, A., Benham, P. M., Chavez, A. N., Cheviron, Z. A., Storz, J. F. and Witt, C. C.** (2015). Contribution of a mutational hot spot to hemoglobin adaptation in high-altitude Andean house wrens. *Proc. Natl. Acad. Sci. USA* **112**, 13958-13963.
- Grispo, M. T., Natarajan, C., Projecto-Garcia, J., Moriyama, H., Weber, R. E. and Storz, J. F.** (2012). Gene duplication and the evolution of hemoglobin isoform differentiation in birds. *J. Biol. Chem.* **287**, 37647-37658.
- Hawkes, L. A., Balachandran, S., Batbayar, N., Butler, P. J., Chua, B., Douglas, D. C., Frappell, P. B., Hou, Y., Milsom, W. K., Newman, S. H. et al.** (2013). The paradox of extreme high-altitude migration in bar-headed geese *Anser indicus*. *Proc. Royal Soc. B: Biological Sciences* **280**, 20122114.
- Hawkes, L. A., Balachandran, S., Batbayar, N., Butler, P. J., Frappell, P. B., Milsom, W. K., Tseveenmyadag, N., Newman, S. H., Scott, G. R., Sathiyaselvam, P. et al.** (2011). The trans-Himalayan flights of bar-headed geese (*Anser indicus*). *Proc. Natl. Acad. Sci. USA* **108**, 9516-9519.
- Helbo, S. and Fago, A.** (2012). Functional properties of myoglobins from five whale species with different diving capacities. *J. Exp. Biol.* **215**, 3403-3410.

- Hiebl, I. B., G.; Schneeganss, D.** (1987). The primary structures of the major and minor hemoglobin-components of adult Andean goose (*Chloephaga melanoptera*, Anatidae): the mutation Leu →Ser in position 55 of the beta-chains. *Biol. Chem. Hoppe Seyler* **368**, 1559-1569.
- Hochachka, P. W. and Somero, G. N.** (2002). Biochemical Adaptation. Mechanism and Process in Physiological Evolution. Oxford: Oxford University Press.
- Jaeger, J. J. and McGrath, J. J.** (1974). Hematologic and biochemical effects of simulated high altitude on the Japanese quail. *J. Appl. Physiol.* **37**, 357-361.
- Janecka, J. E., Nielsen, S. S., Andersen, S. D., Hoffmann, F. G., Weber, R. E., Anderson, T., Storz, J. F. and Fago, A.** (2015). Genetically based low oxygen affinities of felid hemoglobins: lack of biochemical adaptation to high-altitude hypoxia in the snow leopard. *J. Exp. Biol.* **218**, 2402-2409.
- Jessen, T. H., Weber, R. E., Fermi, G., Tame, J. and Braunitzer, G.** (1991). Adaptation of bird hemoglobins to high altitudes: demonstration of molecular mechanism by protein engineering. *Proc. Natl. Acad. Sci. USA* **88**, 6519-6522.
- Johnson, L. F. and Tate, M. E.** (1969). Structure of "phytic acids". *Can. J. Chem.* **47**, 63-73.
- Kumar, A., Natarajan, C., Moriyama, H., Witt, C. C., Weber, R. E., Fago, A. and Storz, J. F.** (2017). Stability-mediated epistasis restricts accessible mutational pathways in the functional evolution of avian hemoglobin. *Mol. Biol. Evol.* **34**, 1240-1251.
- Lague, S. L., Chua, B., Farrel, A. P., Wang, Y. and Milsom W. K.** (2016). Altitude matters: differences in cardiovascular and respiratory responses to hypoxia in bar-headed geese reared at high and low altitudes. *J. Exp. Biol.* **219**, 1974-1984.
- Liang, Y., Hua, Z., Liang, X., Xu, Q. and Lu, G.** (2001). The crystal structure of bar-headed goose hemoglobin in deoxy form: the allosteric mechanism of a hemoglobin species with high oxygen affinity. *J. Mol. Biol.* **313**, 123-137.
- Lutz, P. L.** (1980). On the oxygen affinity of bird blood. *Amer. Zool.* **20**, 11.
- McCracken, K. G., Barger, C. P. and Sorenson, M. D.** (2010). Phylogenetic and structural analysis of the HbA (α^A/β^A) and HbD (α^D/β^A) hemoglobin genes in two high-altitude waterfowl from the Himalayas and the Andes: Bar-headed goose (*Anser indicus*) and Andean goose (*Chloephaga melanoptera*). *Mol. Phylogen. Evol.* **56**, 649-658.
- Meir, J. U. and Milsom, W. K.** (2013). High thermal sensitivity of blood enhances oxygen delivery in the high-flying bar-headed goose. *J. Exp. Biol.* **216**, 2172-2175.
- Monod, J., Wyman, J. and Changeux, J. P.** (1965). On the nature of allosteric transitions: a plausible model. *J. Mol. Biol.* **12**, 88-118.
- Mylvaganam, S. E., Bonaventura, C., Bonaventura, J. and Getzoff, E. D.** (1996). Structural basis for the Root effect in haemoglobin. *Nature Struct. Biol.* **3**, 275-283.
- Natarajan, C., Hoffmann, F. G., Weber, R. E., Fago, A., Witt, C. C. and Storz, J. F.** (2016). Predictable convergence in hemoglobin function has unpredictable molecular underpinnings. *Science* **354**, 336-339.
- Natarajan, C., Inoguchi, N., Weber, R. E., Fago, A., Moriyama, H. and Storz, J. F.** (2013). Epistasis among adaptive mutations in deer mouse hemoglobin. *Science* **340**, 1324-1327.
- Natarajan, C., Jendroszek, A., Kumar, A., Weber, R. E., Tame, J. R. H., Fago, A. and Storz, J. F.** (2018). Molecular basis of hemoglobin adaptation in the high-flying bar-headed goose. *PLOS Genet.* **14**, e1007331.
- Natarajan, C., Projecto-Garcia, J., Moriyama, H., Weber, R. E., Munoz-Fuentes, V., Green, A. J., Kopuchian, C., Tubaro, P. L., Alza, L., Bulgarella, M. et al.** (2015). Convergent evolution of hemoglobin function in high-altitude Andean waterfowl involves limited parallelism at the molecular sequence level. *PLoS Genet.* **11**, e1005681.
- Oberthür, W. B., G.; Würdinger, I.** (1982). Hemoglobins, XLVII. Hemoglobins of the bar-headed goose (*Anser indicus*): primary structure and physiology of respiration, systematic and evolution. *Hoppe Seylers Z Physiol. Chem.* **363**, 581-590.

- Opazo, J. C., Hoffmann, F. G., Natarajan, C., Witt, C. C., Berenbrink, M. and Storz, J. F.** (2015). Gene turnover in the avian globin gene families and evolutionary changes in hemoglobin isoform expression. *Mol. Biol. Evol.* **32**, 871-887.
- Perutz, M. F.** (1983). Species adaptation in a protein molecule. *Mol. Biol. Evol.* **1**, 1-28.
- Petschow, D., Wurdinger, I., Baumann, R., Duhm, J., Braunitzer, G. and Bauer, C.** (1977). Causes of high blood O₂ affinity of animals living at high altitude. *J. Appl. Physiol.* **42**, 139-143.
- Projecto-Garcia, J., Natarajan, C., Moriyama, H., Weber, R. E., Fago, A., Cheviron, Z. A., Dudley, R., McGuire, J. A., Witt, C. C. and Storz, J. F.** (2013). Repeated elevational transitions in hemoglobin function during the evolution of Andean hummingbirds. *Proc. Natl. Acad. Sci. USA* **110**, 20669-20674.
- Riccio, A. T., M.; Giardina, B.; di Prisco, G.** (2001). Molecular dynamics analysis of a second phosphate site in the hemoglobins of the seabird, South Polar skua. Is there a site-site migratory mechanism along the central cavity? *Biophys. J.* **81**, 1938-1946.
- Riggs, A. F.** (1998). Self-association, cooperativity and supercooperativity of oxygen binding by hemoglobins. *J. Exp. Biol.* **201**, 1073-1084.
- Rollema, H. S. and Bauer, C.** (1979). The interaction of inositol pentaphosphate with the hemoglobins of highland and lowland geese. *J. Biol. Chem.* **254**, 12038-12043.
- Scott, G. R. and Milsom, W. K.** (2007). Control of breathing and adaptation to high altitude in the bar-headed goose. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **293**, R379-R391.
- Scott, G. R.** (2011). Elevated performance: the unique physiology of birds that fly at high altitudes. *J. Exp. Biol.* **214**, 2455-2462.
- Scott, G. R., Hawkes, L. A., Frappell, P. B., Butler, P. J., Bishop, C. M. and Milsom, W. K.** (2015). How bar-headed geese fly over the himalayas. *Physiology* **30**, 107-115.
- Scott, G. R. and Milsom, W. K.** (2006). Flying high: a theoretical analysis of the factors limiting exercise performance in birds at altitude. *Respir. Physiol. Neurobiol.* **154**, 284-301.
- Storz, J. F.** (2016). Hemoglobin-oxygen affinity in high-altitude vertebrates: is there evidence for an adaptive trend? *J. Exp. Biol.* **219**, 3190.
- Storz, J. F.** (2018). Compensatory mutations and epistasis for protein function. *Current Opinion in Structural Biology* **50**, 18-25.
- Storz, J. F., Natarajan, C., Moriyama, H., Hoffmann, F. G., Wang, T., Fago, A., Malte, H., Overgaard, J. and Weber, R. E.** (2015). Oxygenation properties and isoform diversity of snake hemoglobins. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **309**, R1178-R1191.
- Tamburrini, M., Riccio, A., Romano, M., Giardina, B. and di Prisco, G.** (2000). Structural and functional analysis of the two haemoglobins of the Antarctic seabird *Catharacta maccormicki*. *Eur. J. Biochem.* **267**, 6089-6098.
- Tufts, D. M., Natarajan, C., Revsbech, I. G., Projecto-Garcia, J., Hoffmann, F. G., Weber, R. E., Fago, A., Moriyama, H. and Storz, J. F.** (2015). Epistasis constrains mutational pathways of hemoglobin adaptation in high-altitude pikas. *Mol. Biol. Evol.* **32**, 287-298.
- Van Assendelft, O. W. and Zijlstra, W. G.** (1975). Extinction coefficients for use in equations for spectrophotometric analysis of hemoglobin mixtures. *Anal. Biochem.* **69**, 43-48.
- Ward, S., Bishop, C. M., Woakes, A. J. and Butler, P. J.** (2002). Heart rate and the rate of oxygen consumption of flying and walking barnacle geese (*Branta leucopsis*) and bar-headed geese (*Anser indicus*). *J. Exp. Biol.* **205**, 3347-3356.
- Weber, R. E.** (1981). Cationic control of O₂ affinity in lugworm erythrocrucorin. *Nature* **292**, 386-387.
- Weber, R. E.** (1992). Use of ionic and zwitterionic (Tris/BisTris and HEPES) buffers in studies on hemoglobin function. *J. Appl. Physiol.* **72**, 1611-1615.
- Weber, R. E.** (2007). High-altitude adaptations in vertebrate hemoglobins. *Resp. Physiol. Neurobiol.* **158**, 132-142.

Weber, R. E. and Fago, A. (2004). Functional adaptation and its molecular basis in vertebrate hemoglobins, neuroglobins and cytoglobins. *Respir. Physiol. Neurobiol.* **144**, 141-159.

Weber, R. E., Fago, A., Malte, H., Storz, J. F. and Gorr, T. A. (2013). Lack of conventional oxygen-linked proton and anion binding sites does not impair allosteric regulation of oxygen binding in dwarf caiman hemoglobin. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **305**, R300-R312.

Weber, R. E., Jessen, T. H., Malte, H. and Tame, J. (1993). Mutant hemoglobins (alpha 119-Ala and beta 55-Ser): functions related to high-altitude respiration in geese. *J. Appl. Physiol.* **75**, 2646-2655.

Zhang, J., Hua, Z., Tame, J. R. H., Lu, G., Zhang, R. and Gu, X. (1996). The crystal structure of a high oxygen affinity species of haemoglobin (bar-headed goose haemoglobin in the oxy form). *J. Mol. Biol.* **255**, 484-493.

Zhu, X., Guan, Y., Signore, A. V., Natarajan, C., DuBay, S. G., Cheng, Y., Han, N., Song, G., Qu, Y., Moriyama, H. et al. (2018). Divergent and parallel routes of biochemical adaptation in high-altitude passerine birds from the Qinghai-Tibet Plateau. *Proc. Natl. Acad. Sci. USA* **115**, 1865.

Figures




Figure 1. Amino acid replacements at eight sites that distinguish the HbA and HbD isoforms of the high-flying bar-headed goose (*Anser indicus*) and the strictly lowland greylag goose (*Anser anser*). Amino acid residues at the same sites are shown for 12 other lowland waterfowl species in the subfamily Anserinae and the most recent common ancestor (MRCA) of the genus *Anser*, see (Natarajan et al., 2018). Given the known phylogenetic relationships of these 12 waterfowl species, the multiple alignment clearly indicates that all amino acid replacements in α^A - and α^D -globin were specific to the bar-headed goose lineage, whereas the β^A -globin replacements occurred in the common ancestor of greylag goose and all other lowland *Anser* species after divergence from bar-headed goose. In addition to the six fixed differences in the three adult-expressed globin genes (four affecting HbA and two affecting HbD), bar-headed geese harbor low-frequency allelic polymorphisms at two sites α^A12 (Gly/Ala) and β^A125 (Asp/Glu) (Hiebl, 1987; McCracken et al., 2010; Oberthür, 1982).




Figure 2. Representative O₂ equilibrium curves (A,B) and Bohr effect (C,D) of purified HbA (A,C) and HbD (B,D) from bar-headed goose (BHG, closed symbols) and greylag goose (GLG, open symbols). Data were obtained at a heme concentration of 0.3 mM, 37 °C, in 0.1 M HEPES buffer, pH 7.4, in the absence (stripped) and presence of the allosteric effectors KCl (0.1 M) and inositol hexaphosphate (IHP, 0.15 mM) added separately or in combination, as indicated. Lines indicate fitting of the data according to the Hill's sigmoidal equation (A,B) to derive O₂ affinity (P_{50} , torr) and cooperativity (n_{50}) parameters, and to linear regression (C,D) to derive the Bohr factor (slope of the plot). Parameters from these experiments are listed in supplementary Table S1.




Figure 3. Effect of IHP on O₂ affinity (P_{50} , torr) of HbA (A) and HbD (B) of bar-headed goose (BHG, closed symbols) and greylag goose (GLG, open symbols). Data for HbA (A) followed a 1:1 stoichiometric binding of IHP to HbA (described by straight lines intersecting at 1:1 stoichiometry), whereby the affinity (equilibrium dissociation constant $<10^{-8}$ M) cannot be estimated. Data for HbD (B) were best fitted according to sigmoidal equations (lines), yielding the Hb-IHP binding affinity as the IHP concentration at half-saturation. In both panels, the 1:1 tetrameric Hb to IHP ratio is indicated by a vertical dotted line. Data were obtained at a heme concentration of 0.3 mM, 37 °C, in 0.1 M Hepes buffer, pH 7.4.











Figure 4. O₂ equilibrium curves of purified HbA (A) and HbD (B) from bar-headed goose (BHG, closed symbols) and greylag goose (GLG, open symbols) covering a wide range of saturations (approaching zero and 100%) to derive allosteric parameters describing Hb function according the MWC model.

Measurements from 5-9 separate experiments were carried out at 0.3 mM heme concentration, 37 °C, in 0.1 M Hepes buffer, pH 7.4, in the absence (stripped) and presence of inositol hexaphosphate (IHP, 0.15 mM). The curves show fitting of all measured data under a given condition according to the saturation function (Eq. 1) describing the MWC model (parameters of this analysis are reported in Table S3). Bar graphs indicate derived parameters (mean ± SE) $\log K_T$ (C,D) and $\log K_R$ (E,F), the O₂ association equilibrium constants of T- and R-states, respectively, and L (G,H), the allosteric equilibrium constant describing the [T]/[R] ratio in the absence of O₂. Closed bars, bar-headed goose (BHG); Open bars, greylag goose (GLG).




Figure 5. Gel filtration elution profiles of HbA and HbD from bar-headed goose (BHG) and greylag goose (GLG) and of human adult HbA in the absence (continuous line) and presence of IHP (discontinuous line), under oxygenated (red) and deoxygenated (blue) conditions. Dotted vertical lines indicate the elution peaks of tetrameric Hb (T), dimeric Hb (D) and monomeric Mb (M) globins.

Table 1. O₂ dissociation rates (k_{off} , s⁻¹) of HbA and HbD from bar-headed goose and greylag goose. Rates were measured by stopped-flow in the absence (stripped) and presence of IHP and KCl, separately and in combination. Conditions were (after mixing) 5 μM heme, 0.2 M Hepes buffer, pH 7.4, IHP = 5 μM, KCl = 3.3 mM, 37 °C. Values are reported as mean ± SE. Number of replicates for each condition is in parentheses.

Species	Conditions	HbA	HbD
		k_{off} (s ⁻¹)	k_{off} (s ⁻¹)
Bar-headed goose	stripped	86.5 ± 1.6 (9)	83.3 ± 1.4 (4)
	IHP	143.3 ± 3.1 (9)	143.9 ± 3.7 (5)
	IHP+KCl	104.4 ± 4.1 (9)	101.1 ± 3.3 (4)
Greylag goose	stripped	102.6 ± 3.8 (9)	86.6 ± 1.0 (4)
	IHP	172.0 ± 12.4 (8)	134.7 ± 8.1 (5)
	IHP+KCl	114.5 ± 3.0 (8)	101.8 ± 2.9 (5)

Supplementary material

Figure S1. Alignment of α -type globin sequences from bar-headed goose and greylag goose. The α^A - and α^D -globins comprise the α -type subunits of the major HbA and the minor HbD isoforms, respectively.



Figure S2. Purification of HbA and HbD by anion exchange chromatography of bar-headed goose (A, left panel) and graylag goose (B, left panel) hemolysates, and corresponding analysis of peaks by isoelectric focusing (right panels). In the chromatograms (absorbance, continuous lines), peak 1 passes unbound through the column and corresponds to the major HbA, peak 3 corresponds to the minor HbD, while peak 2 contains HbA, presumably due to protein washout at the onset of the salt gradient (dashed line).




Table S1. O₂ affinity (P_{50} , torr), Hill cooperativity coefficient (n_{50}) and Bohr effect (expressed as Bohr factor $\Phi = \Delta \log P_{50} / \Delta \log \text{pH}$) of HbA and HbD from bar-headed goose and graylag goose. Experiments were performed in the absence (stripped) and presence of 0.1 M KCl and 0.15 mM IHP, added separately and in combination. Experiments were done at a heme concentration of 0.3 mM, 37 °C, in 0.1 M HEPES buffer, pH 7.4. Data are means of duplicate experiments (variation was < 10%), except for the Bohr effect, where Φ is derived from linear regression (linear plots of Fig. 2E,F). n.d., not determined.

Species	Conditions	HbA				HbD			
		P_{50} (torr)	$\log P_{50}$	n_{50}	Φ	P_{50} (torr)	$\log P_{50}$	n_{50}	Φ
Bar-headed goose	stripped	3.72	0.57	2.35	-0.36	2.79	0.45	2.15	-0.35
	KCl	4.38	0.64	2.58	n.d.	3.20	0.51	2.13	n.d.
	IHP	46.22	1.66	3.21	-0.55	29.84	1.47	2.94	-0.45
	IHP+KCl	34.71	1.54	3.67	n.d.	24.29	1.39	2.95	n.d.
Greylag goose	stripped	4.24	0.63	2.29	-0.35	4.60	0.66	1.93	-0.39
	KCl	6.05	0.78	3.00	n.d.	6.47	0.81	1.51	n.d.
	IHP	60.22	1.78	2.94	-0.45	45.28	1.66	3.18	-0.43
	IHP+KCl	49.37	1.69	2.78	n.d.	35.24	1.55	3.50	n.d.

Table S2. Thermodynamic MWC allosteric parameters for O₂ binding of HbA and HbD from bar-headed goose and graylag goose in the absence (stripped) and presence of 0.15 mM IHP, fitted within 95% confidence limits. Experiments (shown in Fig. 4) were performed at 37 °C (heme concentration 0.3 mM, 0.1 M HEPES buffer, pH 7.4). MWC parameters of recombinant adult human Hb wild type (wt) and $\alpha 119\text{Pro} \rightarrow \text{Ala}$ mutant obtained under stripped conditions at 25 °C (heme concentration 0.146 mM, 0.1 M HEPES, pH ~7.25) (Weber et al., 1993) are shown for comparison. P_{50} , O₂ tension at half-saturation; P_m , median O₂ tension; n_{50} and n_{\max} , half-saturation and maximal Hill's cooperativity coefficients; K_T and K_R (torr⁻¹), O₂ association equilibrium constant of T- and R-states, respectively; $c = K_T/K_R$; ΔG (kcal mol⁻¹), free energy of cooperativity ($\Delta G = -RT \ln K_T/K_R$, where R is the gas constant and T is the absolute temperature). Fitting correlation coefficient r^2 is indicated.

Parameter	Bar-headed goose				Greylag goose				Human Hb at 25°C (Weber et al. 1993)	
	HbA		HbD		HbA		HbD		Hb wt	Hb $\alpha 119\text{Pro} \rightarrow \text{Ala}$
	stripped	IHP	stripped	IHP	stripped	IHP	stripped	IHP	stripped	stripped
P_{50} (torr)	3.37	44.68	2.74	29.39	4.10	57.30	4.05	47.21	2.00	1.12
P_m (torr)	3.64	43.82	2.98	29.27	4.31	54.46	4.39	47.04	1.84	1.05
n_{50}	2.67	3.44	1.97	3.05	2.57	3.09	2.17	3.01		
n_{\max}	2.74	3.45	1.93	3.05	2.60	3.13	2.13	3.01		
K_T (torr ⁻¹)	0.012	0.002	0.040	0.003	0.022	0.003	0.023	0.002	0.170	0.448
K_R (torr ⁻¹)	1.20	0.76	0.86	0.38	1.07	0.61	0.68	0.22	6.22	6.22
$\log L$	2.57	6.08	1.62	4.18	2.65	6.10	1.90	4.08	4.24	3.28
c	0.010	0.002	0.046	0.009	0.021	0.004	0.033	0.010		
ΔG (kcal mol ⁻¹)	2.69	3.77	1.65	2.89	2.33	3.30	1.90	2.83	2.13	1.56
r^2	0.998	0.998	0.998	0.999	0.998	0.998	0.998	0.998		

References

Weber, R. E., Jessen, T. H., Malte, H. and Tame, J. (1993). Mutant hemoglobins (alpha 119-Ala and beta 55-Ser): functions related to high-altitude respiration in geese. *J. Appl. Physiol.* **75**, 2646-2655.