

Prolonged survival out of water is linked to a slow pace of life in a selfing amphibious fish

Turko, Andy J.^{1,2*}, Doherty, Justine E.¹, Yin-Liao, Irene¹, Levesque, Kelly¹, Kruth, Perryn¹,
Holden, Joseph M.³, Earley, Ryan L.³, and Wright, Patricia A.¹

¹Department of Integrative Biology, University of Guelph, Guelph, ON, Canada, N1G 2W1

²Great Lakes Institute for Environmental Research, University of Windsor, Windsor, ON,
Canada, N9B 3P4 (present address)

²Department of Biological Sciences, University of Alabama, Tuscaloosa, AL, USA, 35487

*Corresponding author: ajturko@gmail.com

Key words: metabolism, life-history, trade-offs, *Kryptolebias marmoratus*, mangrove rivulus

Summary Statement

Intrinsically low metabolic rates increased survival and fitness of an amphibious fish when access to water was limited, and there was no apparent cost when water was abundant.

Abstract

Metabolic rate and life history traits vary widely both among and within species reflecting trade-offs in energy allocation, but the proximate and ultimate causes of variation are not well understood. We tested the hypothesis that these trade-offs are mediated by environmental heterogeneity, using isogenic strains of the amphibious fish *Kryptolebias marmoratus* that vary in the amount of time each can survive out of water. Consistent with pace of life theory, the strain that survived air exposure the longest generally exhibited a “slow” phenotype including the lowest metabolic rate, largest scope for metabolic depression, slowest consumption of energy stores, and least investment in reproduction under standard conditions. Growth rates were fastest in the otherwise “slow” strain, however. We then tested for fitness trade-offs between “fast” and “slow” strains using microcosms where fish were held with either constant water availability or under fluctuating conditions where water was absent for half of the experiment. Under both conditions the “slow” strain grew larger and was in better condition, and under fluctuating conditions the “slow” strain produced more embryos. However, the “fast” strain had larger adult population sizes under both conditions, indicating that fecundity is not the sole determinant of population size in this species. We conclude that genetically based differences in pace of life of amphibious fish determine survival duration out of water. Relatively “slow” fish tended to perform better under conditions of limited water availability, but there was no detectable cost under control conditions. Thus, pace of life differences may reflect a conditionally neutral instead of antagonistic trade-off.

1. Introduction

Life history and metabolic rate are often conceptually linked to form pace of life syndromes, in which organisms span a continuum of “fast” (rapid growth, early age at maturity, high metabolism) to “slow” (slow growth, delayed maturity, low metabolism) lifestyles (Ricklefs and Wikelski, 2002; Réale et al., 2010; Arnqvist et al., 2017; Auer et al., 2018). Pace of life varies substantially both among and within species, but the mechanistic causes and ultimate consequences of this variation are not well understood (e.g. Stearns, 1992; Burton et al., 2011; Careau and Garland, 2012). Variation between fast and slow lifestyles is thought to reflect trade-offs in energy allocation, for example to growth versus reproduction. Energy allocation depends first on energy acquisition and resource availability, thus spatial or temporal environmental heterogeneity in resource availability is thought to be a key factor that produces and maintains variation in pace of life (Mueller and Diamond, 2001; Reid et al., 2011). However, the direction and mechanistic relationships between environmental conditions, life history, and metabolic rate remain unclear (Koons et al., 2008; Burton et al., 2011; Auer et al., 2018).

Organisms that inhabit extreme and variable environments are useful systems for understanding the mechanisms that generate and maintain variation in pace of life (Passow et al., 2017). One of the most abrupt and dramatic changes in the physical environment experienced by any animal occurs when amphibious fishes move between water and land (Dejours, 1988; Graham, 1997; Wright and Turko, 2016). Among the many challenges faced by fishes out of water, capturing and consuming prey is particularly problematic, as the low density of air makes suction feeding difficult (Heiss et al., 2018). Thus, most amphibious fishes must rely on internal energy stores when out of water, although there are exceptions (Heiss et al., 2018). Therefore, the ability to deeply depress metabolism is thought to be a key factor enabling prolonged survival out of water in amphibious fishes such as the African

lungfish *Protopterus aethiopicus*, which reduce O₂ consumption by about 80% while aestivating on land (Guppy and Withers, 1999). Many species do not aestivate, however, and the mechanisms that underlie variation in emersion tolerance are not well understood. In non-aestivating species, a relatively low overall metabolic rate would conserve resources (e.g. Killen et al., 2011) and could extend survival out of water. Thus, environmental variation in water availability may be a key factor that causes and maintains intra-specific variation in metabolism and life history traits of amphibious fishes, but this hypothesis has not been tested.

Our objective was to understand the proximate and ultimate reasons why some amphibious fishes survive out of water longer than others. First, we tested the hypothesis that the amount of time amphibious fishes can spend out of water is limited by their pace of life. This hypothesis predicts that emersion-tolerant fish will have relatively slow metabolic and growth rates, reduced consumption of energy stores, and low levels of activity and reproductive output. We then tested whether the costs and benefits of different paces of life ultimately depend on environmental conditions – i.e. there is an antagonistic trade-off between emersion tolerance and aquatic performance. Specifically, we investigated the prediction that a relatively “fast” lifestyle would be favoured when food and water were constantly available, while a “slow” pace of life would be favoured in a fluctuating environment frequently lacking water.

We tested these hypotheses using the amphibious mangrove rivulus *Kryptolebias marmoratus*, one of only two known self-fertilizing hermaphroditic vertebrates (Harrington, 1961; Avise and Tatarenkov, 2015) - the other is the sister species *K. hermaphroditus* (Costa, 2011). This “selfing” reproductive system allows the production of large numbers of isogenic, effectively clonal individuals and ultimately enables repeated experiments on the same genotype (Tatarenkov et al., 2010; Turko et al., 2011; Earley et al., 2012). *Kryptolebias*

marmoratus can survive more than 66 d out of water in leaf litter or packed nose-to-tail within rotting mangrove logs (Taylor et al., 2008). There is no evidence that *K. marmoratus* aestivates when on land (Ong et al., 2007; Blanchard et al., 2018), but they are largely inactive (Turko et al., 2014; Turko et al., 2017) and do not eat (Pronko et al., 2013; Wells et al., 2015). We first measured energy use and oxygen uptake in isogenic strains acclimated to terrestrial conditions for 21 d. Then, fish from “fast” and “slow” strains (relatively high or low metabolic rate, estimated by rates of oxygen uptake) were reared together for 12 months in microcosms in which water was either always present, or was absent for random periods totaling 6 months, and the number, condition, and reproductive output of fish was measured.

2. Materials and Methods

2.1 Animals

For all physiological experiments, *Kryptolebias marmoratus* (Poey) hermaphrodites were raised individually in the Hagen Aqualab, University of Guelph, in 120 mL plastic holding cups. Fish from the isogenic strains 50.91 (“Belize”, from Twin Cayes, Papa Gabriel, Belize), SLC8E (“Florida”, from St. Lucie County, Florida, USA), and HON11 (“Honduras”, from Bay Islands, Utila, Honduras) were used in these experiments (Tatarenkov et al., 2010). Fish were kept at 25°C, 15‰ salinity, with 12 h: 12 h light: dark cycle and were fed *Artemia* nauplii 3 times per week. Fish were fed to satiation for three consecutive days immediately prior to all experiments but were not fed for the duration of the experiments. Fish in microcosms were fed live *Artemia* nauplii three times per week but food was not added to fluctuating microcosms when water was not present. All experiments were approved by the University of Guelph animal care committee. Emersion tolerance, metabolic rate, and energy use experiments were conducted in 2015-2016, microcosm experiments were conducted in 2016-2017, and genetic identification of these fish occurred in 2017-2018.

2.2 Emersion tolerance

To measure emersion tolerance, size-matched fish ($n = 20/\text{strain}$) were terrestrially acclimated on moist filter paper (Ong et al., 2007), and survival was monitored at least once per day (Wells et al., 2015). For ethical reasons, the experiment was terminated when 20% of each strain remained as statistically significant differences among strains were clear at this point; remaining fish were euthanized with tricaine methanesulfonate (MS222; 500 mg L⁻¹).

2.3 Metabolic rate

To test the hypothesis that emersion tolerance depends on metabolic rate, O₂ consumption ($n = 8\text{-}10/\text{strain}$) was compared in water (control) and during air exposure using intermittent flow respirometry (Loligo Systems WITROX 4, Viborg, Denmark; 2.5 mL chambers carefully cleaned with ethanol prior to each trial) as described elsewhere (Livingston et al., 2018; Sutton et al., 2018) with the following modifications. Aquatic O₂ consumption of each individual was measured in triplicate (3 h chamber acclimation with flow-through normoxic water, 12-15 min recordings, 10 min flushing periods; measurements occurred between 11:00-13:00 h), then water was drained from the chambers and aerial O₂ consumption was measured once per day for 7 d at the same time (12:00-16:00 h) to minimize the effect of diurnal metabolic rhythms (Rodela and Wright, 2006). Humidified air (100% relative humidity) was introduced into the chambers between measurements. Fish were weighed before and after the 7d experiment and the average mass was used for statistical analyses and to standardize O₂ uptake (mass of Belize fish = $0.071 \pm 0.004\text{g}$, Florida fish = $0.072 \pm 0.003\text{g}$, Honduras fish = $0.076 \pm 0.003\text{g}$; no difference among groups $p = 0.64$). To measure O₂ uptake after three weeks in air, an initial O₂ consumption rate in water was first determined for a separate group of fish, then fish were air-exposed for 21 d and O₂ uptake was measured in air ($n = 10/\text{strain}$). For these fish, mass was measured immediately after each measurement

of O₂ uptake (mass of Belize fish before = 0.070 ± 0.005 g, after = 0.56 ± 0.003 g, Florida fish before = 0.078 ± 0.006 g, after = 0.060 ± 0.006 g, Honduras fish before = 0.099 ± 0.005 g, after = 0.089 ± 0.006 g; among strains $p = 0.001$, before vs after $p < 0.001$). Activity of the fish was not observed during these measurements. Microbial respiration, measured after each trial, was negligible.

2.4 Energy reserves and consumption

To test whether the amount of energy reserves and/or energy use was related to emersion tolerance, we measured activity, overall body condition, and body composition. To measure activity, fish ($n = 8/\text{strain}$) were photographed every 5 s for 1 h (between 12:00 – 13:00 h) at six time points (1 h, and 1, 3, 7, 14, 21 d out of water). Activity on land consists of discrete jumps, so activity was quantified as the proportion of photos in which fish had changed location between consecutive frames (Turko et al., 2014). Fulton's K , a general index of body condition (Froese, 2006) was calculated for control fish in water ($n = 6/\text{strain}$), and in a separate group of fish ($n = 12/\text{strain}$) that were terrestrially acclimated for 21 d. To measure body composition, independent groups of fish were required because the small size of *K. marmoratus* precluded measuring multiple energy reserves in the same sample. Fish held in water (control) or air (21 d) were euthanized (MS222), blotted dry, weighed, and snap frozen in liquid nitrogen (for glycogen and protein determination) or dried (48 h at 50°C) for lipid analysis and measurement of water content. Glycogen content of whole fish ($n = 8-9/\text{strain}$) was measured enzymatically (Bergmeyer et al., 1974). Whole body lipid stores ($n = 6-12/\text{strain}$) were measured by chloroform extraction (Junior and Peixoto, 2013). Crude protein ($n = 6-10/\text{strain}$) was measured using the Kjeldahl method (AOAC, 1995). Tecator Kjeltex digestion and distillation units (Foss, Eden Prairie, MN, USA) were used for protein analysis and the percentage of total nitrogen was determined based on a dry matter basis ($\%N \times 6.25$)

(Bureau et al., 2000). Body water content ($n = 6-12/\text{strain}$) was calculated by subtracting wet from dry body mass and dividing by the wet mass.

To calculate energy use during emersion, lipid, glycogen, and protein utilization were each calculated by subtracting the average mass-specific energy stores (mg g^{-1}) in terrestrially-acclimated fish (E_{terr}) from those of the control fish (E_{con}), accounting for changes in body water content (W) and the average change in body mass (ΔBM), according to the formula:

$$\text{Energy consumed} = (E_{\text{con}})(1 - W_{\text{con}}) - (E_{\text{terr}})(1 - W_{\text{terr}})(\Delta\text{BM})$$

All data used for these calculations is provided in Table S1. Total energy use (i.e. E_{con} and E_{terr} in the equation above) was calculated by adding the energy contained in the consumed glycogen (17 kJ g^{-1}), lipid (37 kJ g^{-1}), and protein (17 kJ g^{-1}). Overall standard deviations were calculated using standard methods of error propagation, and effective degrees of freedom were calculated using the Welch-Satterthwaite equation (JCGM, 2008). Using these values, we compared energy use among the isogenic lineages with one-way ANOVA and *post hoc* Holm–Sidak tests. Data were \ln transformed when necessary to meet assumptions of normality and equal variance.

2.5 Life history traits

Routinely collected records from our *K. marmoratus* colony were used to compare overall embryo production, clutch size, and age at first reproduction among strains. To assess cumulative reproductive output, we only used data from fish that hatched within one year of those used for experiments, released at least one embryo, were never used for any experiments, and survived for over 18 months (Belize $n = 42$, Florida $n = 27$, Honduras $n = 20$). Age at first reproduction was determined for a larger subset of fish that simply released an embryo prior to use in any experiments (Belize $n = 90$, Florida $n = 72$, Honduras $n = 90$).

2.6 Microcosms

We used a 12-month microcosm experiment to test the hypothesis that there is an environmentally mediated trade-off between emersion tolerance and metabolic rate. Belize and Honduras fish ($n = 3/\text{strain}/\text{microcosm}$) were placed into each microcosm at the start of the experiment. Fish were size-matched to minimize performance differences in aggressive/competitive interactions between the strains (Earley and Hsu, 2008), which resulted in Belize fish being slightly older (434 ± 9.4 versus 325 ± 17 days old) but of similar mass (Figure S1) at the beginning of the experiment. All fish were sexually mature and had released at least one embryo in the laboratory colony before being placed in a microcosm to standardize reproductive status. We maintained constant water levels in control microcosms ($n = 10$), while fluctuating microcosms ($n = 10$) were drained and refilled (every 1-3 weeks, randomly assigned) such that water was absent for half of the experiment. Microcosms were constructed from 9 L plastic boxes ($38 \times 24 \times 14$ cm) filled half-way with 15‰ brackish water. The bottom was covered with soft filtration media to provide a moist substrate in the fluctuating treatment when water was absent, and 20 pieces of grey plastic pipe (3 cm length, 1.5 cm diameter) and three green acrylic yarn mops were added to provide shelter. Emersion periods never exceeded 3 weeks because our emersion tolerance data showed that 100% of fish survived this duration, and the goal of this experiment was to test whether sub-lethal effects could mediate a trade-off between “fast” and “slow” fish. Water levels were altered (fluctuating) or water was refreshed (control) via permanently-installed plastic tubing under the filter media to minimize disturbance.

After the 12-month experiment, all adult fish were euthanized (MS222), photographed (for length measurements), weighed, and a piece of caudal fin tissue was fixed in DNA preservative (0.25 M ethylenediaminetetraacetic acid, 20% dimethyl sulfoxide, NaCl saturated, pH = 7.5; Seutin et al., 1991) for determination of genetic identity. Gonads and

liver were dissected and weighed to assess condition. Gills were removed to test whether gill surface area was related to pace of life differences. The number and length of gill filaments was measured in whole mounts of the left-side arches. Sex was assessed based on external morphology (Scarsella et al., 2018) and appearance of ovarian tissue in the gonads. All embryos were collected and fixed in DNA preservative (see above) for genetic identification. No larvae or juvenile fish were found.

The genetic strain (i.e. Belize or Honduras) of each adult and embryo was determined using previously described protocols and microsatellite markers (Mackiewicz et al., 2006; Tatarenkov et al., 2010). Genomic DNA was extracted and purified using a commercially available kit according to manufacturer's instructions (GeneJET DNA Purification Kit, Fisher Scientific). Microsatellite "R18" from Mackiewicz et al. (2006) was used to differentiate between strains, as this is one of the three most divergent loci between Belize and Honduran fish and amplified more consistently than the other most divergent loci (R3 and R34). PCR products were run on an acrylamide gel (5%, 3000 V for 2 h at 55°C), and were manually scored for strain identity. We were able to identify 100% of the adult fish, however, some embryos (20 of 81 from control, 209 of 979 from fluctuating microcosms) could not be confidently assigned to either strain, probably due to low DNA content. These unidentified embryos were excluded from statistical analysis.

2.7 Statistical Analysis

Survival out of water was compared among strains using a Kaplan-Meier survival analysis. Rates of oxygen consumption were compared among strains and over time using both ANOVA (comparison of mass-corrected rates) and a linear model that included body mass as a covariate (R package lme; Pinheiro et al., 2014). Energy stores and energy consumption was compared among strains using one- or two-way ANOVA as appropriate,

followed by Holm–Sidak *post hoc* tests. Life history data (embryo production, age at first reproduction) was not normally distributed and was therefore analyzed using a Kruskal–Wallis ANOVA on ranks. Growth rates of Honduras versus Belize fish were compared using ANCOVA (dependent variables: length, mass; covariate: age). Body size, condition, and organ sizes of microcosm fish were compared using two-way ANOVA. Embryo production between strains and treatments was compared using ANCOVA to account for different numbers of adults in each microcosm at the end of the experiment. Critical $\alpha = 0.05$ for all tests, throughout the text values are given as means \pm SEM.

3. Results

Survival out of water was significantly longer in the Honduras strain relative to the Belize and Florida strains (Kaplan–Meier log-rank statistic = 11.198, $p = 0.004$; Figure 1). The Honduras strain had consistently lower (by 30–50%) rates of O_2 consumption in both water and air relative to Belize and Florida strains over 7 d both when mass-corrected rates were compared directly (ANOVA, $F_{2,24} = 9.96$, $p < 0.001$), and when body mass was included as a covariate in the statistical model (lme, $\chi^2 = 23.02$, $p < 0.001$; Figure 2A). Overall, O_2 consumption increased in all strains after 2 and 3 d in air compared to other timepoints (lme, $\chi^2 = 23.21$, $p = 0.003$, interaction $p = 0.20$; Figure 2A). In the 21d terrestrial acclimation experiment, mass-specific O_2 consumption of Honduras fish was again lower (by ~40%) than that of Belize and Florida fish. This difference was significant when mass-corrected values were compared with ANOVA ($F_{2,27} = 4.81$, $p = 0.016$), and approached statistical significance when mass was included as a covariate in the statistical model (lme, $\chi^2 = 4.34$, $p = 0.11$; Figure 2B). This discrepancy between the different analyses is likely because the Honduras fish used in this experiment were slightly larger (~30%) than the other strains (see

methods). After 21 d in air, rates of O₂ consumption decreased by 44% (lme, $\chi^2 = 19.85$, $p < 0.0001$, interaction $p = 0.45$; Figure 2B).

There was no difference in initial body condition among the three strains, but terrestrial acclimation for 21 d resulted in significantly lower condition factor of Belize and Florida, but not Honduras, fish (two-way ANOVA $F_{2,47} = 3.35$, interaction $p = 0.024$; Figure 3).

Similarly, in paired measurements of different individuals, Honduras fish lost significantly less body mass than other strains after 21 d out of water (ANOVA $F_{2,27} = 6.92$, $p = 0.004$, Figure S2). All three strains in water had similar levels of glycogen, lipid, and protein stores at the beginning of the experiment (two-way ANOVA, all $p > 0.05$; Figure S3). After 21 d out of water there was no difference among strains in lipid use (ANOVA, $F_{2,83} = 0.92$, $p = 0.40$; Figure 4A), but Florida fish used more glycogen than the other strains (ANOVA, $F_{2,191} = 6.72$, $p = 0.002$; Figure 4B). Florida fish also consumed the most protein; Honduras fish consumed the least (ANOVA, $F_{2,779} = 47.60$, $p < 0.001$; Figure 4C). Overall, Honduras fish used significantly less energy over 21 d on land compared to Belize and Florida fish (ANOVA, $F_{2,65} = 5.21$, $p = 0.008$; Figure 4D). Water content was significantly elevated in all strains after 21 days on land (two-way ANOVA, $F_{1,53} = 75.31$, $p < 0.001$, interaction $p = 0.60$; Figure S2D). Activity generally decreased over 21 d out of water, but differed among the strains only at the 1 h time point (two-way ANOVA $F_{2,16} = 5.67$, interaction $p = 0.001$; Figure S4).

In long-term studies of our laboratory colony, Belize fish produced more embryos (Kruskal-Wallis ANOVA on ranks $H_2 = 10.61$, $p = 0.005$; Figure 5A) and had larger mean clutches (Kruskal-Wallis ANOVA on ranks $H_2 = 20.12$, $p < 0.001$; Figure 5B) than Florida or Honduras fish over the first 18 months of life, while age at first reproduction tended to be earlier (Kruskal-Wallis ANOVA on ranks $H_2 = 6.91$, $p = 0.032$; Figure 5C).

Belize and Florida fish generally showed similar trends, so we focused on a comparison of Belize and Honduras fish for our followup microcosm experiments. At the beginning of these experiments, Honduras fish were heavier (ANCOVA $F_{1,287} = 163.75$, $p < 0.0001$; Figure S1A) and longer (ANCOVA $F_{1,287} = 83.60$, $p < 0.0001$; Figure S1B) than Belize fish at a given age, indicating a faster growth rate.

At the end of the microcosm experiment, there was no significant difference in the total number of fish between control and fluctuating microcosms ($F_{1,39} = 0.10$, $p = 0.76$), but there were significantly more Belize than Honduras fish overall ($F_{1,39} = 7.92$, $p = 0.008$, interaction $p = 0.76$; Figure 6A). Honduras fish were longer (two-way ANOVA, $F_{1,110} = 49.95$, $p < 0.001$; Figure 6B), heavier (two-way ANOVA, $F_{1,110} = 80.40$, $p < 0.001$; Figure 6C), and in better condition (Fulton's K, two-way ANOVA, $F_{1,110} = 36.24$, $p < 0.001$; Figures 6D, S5A). Gonado-somatic index was also significantly higher in Honduras fish (two-way ANOVA, $F_{1,110} = 93.76$, $p < 0.001$; Figures 6E, S5B), but there was no difference in hepato-somatic index (two-way ANOVA, $F_{1,110} = 1.4$, $p = 0.23$; Figures 6F, S5C). Fish from control microcosms were longer (two-way ANOVA, $F_{1,110} = 168.04$, $p < 0.001$), heavier (two-way ANOVA, $F_{1,110} = 214.96$, $p < 0.001$), in better condition (two-way ANOVA, $F_{1,110} = 42.02$, $p < 0.001$) and had larger gonads (two-way ANOVA, $F_{1,110} = 7.77$, $p = 0.006$) than fish from fluctuating microcosms, but there was no difference in hepato-somatic index ($F_{1,110} = 3.70$, $p = 0.057$; Figures 6, S5). After controlling for body length, total gill filament length was not different between strains (ANCOVA, $F_{1,82} = 1.22$, $p = 0.27$; Figure S6) or microcosm conditions (ANCOVA, $F_{1,82} = 0.85$, $p = 0.36$; Figure S6). Almost 11-fold more embryos were recovered from fluctuating (89.8 ± 7.6) versus control (8.1 ± 1.4) microcosms (Mann-Whitney, $U = 0$, $p < 0.001$). After accounting for the number of adult fish in each microcosm, embryo quantity was dependent on a significant strain-by-treatment interaction (ANCOVA, $F_{1,32} = 4.36$, $p = 0.045$). Honduras fish produced significantly more embryos than Belize fish

under fluctuating conditions (Tukey, $p = 0.021$; Figure 7), but there was no difference between strains under control conditions ($p = 0.98$). Two embryos collected from a single fluctuating microcosm were heterozygous at the microsatellite locus we used for identification. No adult males were found in the microcosm that contained the heterozygous embryos. Seven adult males were present at the end of the experiment, each in a different microcosm (five control, two fluctuating).

4. Discussion

Using a self-fertilizing amphibious fish, we were able to directly relate survival out of water to genetically-based metabolic and life history phenotypes. In support of the proximate hypothesis that the overall pace of life determines the length of time non-aestivating *K. marmoratus* can survive out of water, we found that emersion tolerance was negatively associated with metabolic rate and the rate of fuel use, but not the size of initial energy reserves. The emersion-tolerant Honduras strain of *K. marmoratus* also produced fewer offspring under normal laboratory conditions, due to an increased age of first reproduction and smaller average clutch size, consistent with a slow pace of life. We used long-term microcosms to test the ultimate hypothesis that a sublethal trade-off between emersion tolerance and aquatic performance mediates the relative advantages of different paces of life. Contrary to the prediction made by this hypothesis, “slow” Honduras fish did not outcompete relatively “fast” Belize fish (in terms of adult population size) under conditions of low water availability. However, Honduras fish produced more embryos than Belize fish in the water-limited condition, in support of the trade-off hypothesis. Honduras fish also tended to be larger and in better condition regardless of water availability, suggesting an overall advantage to low metabolic rate under our experimental conditions.

4.1 Metabolism and emersion tolerance

The rate of O₂ consumption in Honduras fish was consistently ~40% lower than the other strains in both water and air, suggesting that this strain has an inherently slow metabolism that is genetically based. This difference was highly statistically significant in our 7d terrestrial acclimation experiment ($p < 0.001$), but was marginally non-significant ($p = 0.11$) in our 21d experiment despite being of similar magnitude (~40%). Furthermore, the scope of metabolic depression after 21 d in air in Honduras fish was larger (58% reduction) than either the Belize (-31%) or Florida (-44%) strains, which likely allowed Honduras fish to conserve protein stores. Previous studies have suggested that *K. marmoratus* maintain or increase metabolic rate for several days after moving from water to land, but those experiments did not examine fish that were out of water for longer than 7 days (Ong et al., 2007; Blanchard et al., 2018). While probably helpful for survival during prolonged emersion, the scope of metabolic depression we found in *K. marmoratus* is smaller than has been measured in classically aestivating amphibious fishes such as *P. aethiopicus*, *Synbranchus marmoratus*, and *Lepidogalaxias salamandroides* that reduce O₂ consumption by 65-80% during months-long aestivation in mud (Guppy and Withers, 1999). Interestingly, the control rate of O₂ consumption in Honduras fish was almost identical to the depressed rate in the other two strains. One possibility is that the constitutively low metabolic rate of the Honduras strain evolved via the genetic assimilation of metabolic plasticity, which would be expected if these fish were found in habitats that regularly dried (Pigliucci et al., 2006; Lande, 2009). More research is required to understand the mechanism by which Honduras fish achieve low metabolic rate (e.g. increased efficiency or reduced expenditure), especially given the high growth rate of this strain. Overall, however, constitutive expression of a low baseline metabolic rate, combined with the ability to further reduce metabolism during extended

periods without water, probably allows the Honduras strain to prolong survival out of water by conserving energy reserves.

There was a small but consistent increase in O₂ consumption after 2-3 d out of water across all three strains we investigated, even though fish remained largely motionless. Consistent with this finding, previous work found increased rates of CO₂ excretion in *K. marmoratus* over 5 d of terrestrial acclimation (Ong et al., 2007). This transient increase in metabolic rate may reflect the energetic cost of mounting phenotypically flexible responses during air exposure, such as gill remodelling (Ong et al., 2007), cutaneous angiogenesis (Cooper et al., 2012; Blanchard et al., 2018) and enlargement of cutaneous ionocytes (LeBlanc et al., 2010).

The pattern of energy use varied among strains in a manner consistent with the differences in whole animal O₂ consumption. In independent experiments, Honduras fish lost the least amount of body mass relative to the other two strains and showed no change in condition factor after 21 d out of water. Protein catabolism was also lowest in Honduras fish. Generally, teleosts use protein and some lipids as fuel sources when food is not restricted, but glycogen and lipids are more important during starvation (Jobling, 1994; Moyes and West, 1995). However, African lungfish defend glycogen stores during aestivation, perhaps to facilitate rapid recovery when routine metabolism must be restored (Frick et al., 2008). After 21 days of emersion, all three strains of mangrove rivulus we tested had consumed a large fraction of their glycogen (~80% of initial) and lipid (~61%) stores, but only some protein (17-29%, depending on strain). This pattern resembles that of a typical starving teleost, rather than an aestivating lungfish. Presumably, the Florida and Belize fish, with higher metabolic rates, began to metabolise protein earlier in the emersion period than Honduras fish, resulting in greater consumption at the 21 d timepoint. In Honduras fish, larger protein reserves after 21 d of air exposure, in addition to fueling continued emersion, may also help preserve

locomotor ability when these fishes return to water, similar to inactive hibernating mammals that protect protein stores to minimize impairment of skeletal muscle function when they emerge from winter dens (e.g. Hindle et al., 2015).

There were no differences in initial energy stores among the Belize, Florida, and Honduras strains. One possible explanation is that there may be costs to carrying large energy stores, such as increased attractiveness to predators (Jensen et al., 2012) or decreased locomotory performance (Gibb et al., 2013). Alternatively, fish in the wild may detect environmental cues that indicate the onset of the dry season and respond by increasing internal energy stores (Griffiths and Kirkwood, 1995; Schultz and Conover, 1997). In our laboratory setting, however, such anticipatory feeding would not have been possible as the fish were given no signals that emersion was imminent.

4.2 Emersion tolerance trade-offs

Pace of life theory makes conflicting predictions about the direction of correlations between metabolic rate, growth, and reproductive investment (Burton et al., 2011). According to the acquisition model, a fast metabolism allows more resources to be acquired, leading to faster growth and increased reproductive investment (Mathot and Dingemanse, 2015). Conversely, the allocation model suggests that limited resources are split between various physiological processes via trade-offs; negative relationships between each of metabolic rate, growth, and reproduction are thus expected (Burton et al., 2011). We found that Honduras fish, with the lowest metabolic rate, produced fewer embryos but grew faster than Belize fish under standard laboratory rearing conditions, suggesting a trade-off consistent with the allocation model. Furthermore, Honduras fish were much larger than Belize fish in both microcosm conditions. Although faster growth is typically correlated with high metabolic rates in animals (Allen et al., 2016), the negative relationship between growth and metabolic

rate we found in *K. marmoratus* has also been found in some other fishes (Alvarez and Nicieza, 2005; Norin and Malte, 2011).

Our results support, in part, the hypothesis that differences in metabolic rate between mangrove rivulus strains are ultimately caused and maintained by a trade-off between emersion tolerance and aquatic performance. This hypothesis predicts that the “slow” Honduras fish would have higher fitness than Belize fish in fluctuating conditions when water was periodically unavailable, and this was indeed the case by several metrics. Honduras fish were larger, in better condition, and produced more embryos than Belize fish in the fluctuating condition. However, Honduras fish were also larger in the control microcosms where there was no difference between strains in embryo production, in contrast to the prediction that Honduras fish would have lower fitness in constant aquatic conditions. Our results are consistent with many other studies of local adaptation, which often find that phenotypes that are advantageous under some conditions have neutral consequences in other environments, especially in highly heterogeneous environments (Bono et al., 2017).

It is not clear whether “fast” and “slow” phenotypes have effectively equal fitness under fully aquatic conditions, or whether there are situations that would favour the “fast” Belize phenotype. One limitation to our microcosm experiment was that fish were unable to disperse. The natural habitat of mangrove rivulus and other rivuline killifishes typically consists of a mosaic of small, intermittent pools (Turko and Wright, 2015; Furness et al., 2018; Sutton et al., 2018). A high metabolic rate is often correlated with high boldness and activity (e.g. Killen et al., 2011; Gangloff et al., 2017), and perhaps Belize fish are more likely to leave water and disperse to unoccupied habitats. We have previously found that metabolic rate of *K. marmoratus* was positively associated with frequency of emersion behaviour (Turko et al., 2018). Embryo production by Belize fish was highest of any strain in our laboratory colony, typical of a dispersal phenotype. Furthermore, when embryo

production in the microcosm experiment is standardized to gonad mass, rather than the number of adults, the Belize strain produced almost twice as many embryos per gram of gonad compared to Honduras fish. However, we found that Belize fish had the lowest activity of any strain after forced air exposure, opposite to the prediction made by the dispersal phenotype hypothesis. We also found nearly 9-fold more embryos in fluctuating vs. control conditions, despite the larger overall size of control fish. We think it is unlikely that this disparity reflects differences in reproductive output, but instead is the result of cannibalism under aquatic conditions (Wells et al., 2015). Our experiment ended when fluctuating microcosms had been without water for one week, so embryos deposited terrestrially during this period could not be consumed by the suction-feeding mangrove rivulus (Heiss et al., 2018) unlike embryos under control conditions. Presumably, most of the embryos produced under fluctuating conditions were cannibalized each time water returned, explaining the very low levels of recruitment we observed. Thus, Belize fish may live a traditionally “fast” pace of life that favours reproduction over growth, but the inability to escape competition/embryo cannibalism from the larger Honduras fish in our microcosms nullified the advantages of this life history strategy. High metabolic rates are often correlated with aggression (Réale et al., 2010); thus the fast Belize phenotype may be superior competitors in situations of relative food scarcity. Our conditions of *ad libitum* food provisioning would have failed to detect such a benefit to increased metabolic rate.

The androdioecious mating system of mangrove rivulus is thought to provide these fish a mechanism to benefit from both self-fertilization and outcrossing, depending on the context (Ellison et al., 2011; Avise and Tatarenkov, 2015). This “best of both worlds” hypothesis predicts that animals in relatively suitable habitats should self-fertilize to maximize the genetic inheritance by the offspring, while those in less suitable habitats should opt for outcrossing to increase the genetic variation of progeny so that some offspring will be better

suited to the environmental conditions. In our experiments, this could be reflected as an increased number of male Belize fish and higher rate of outcrossing in fluctuating microcosms, while in control conditions more Honduras males and outcrossing would be predicted (due to environmental mismatching). We found no evidence consistent with these predictions. Only 7 of 114 adult fish were male, and 2 of 979 identified embryos were the result of outcrossing between strains. One possibility is that despite being without water for 6 of 12 months, the fluctuating microcosms were not sufficiently stressful for the benefits of outcrossing to outweigh the cost of reduced genetic inheritance. Alternatively, the mating system of adult mangrove rivulus may not be responsive to environmental conditions and could instead rely on developmental plasticity or stochastic epigenetic effects that act on early life stages.

We found two embryos that were heterozygous at the microsatellite locus we studied, indicating that these resulted from outcrossing. The current view is that males are required for outcrossing (Avisé and Tatarenkov, 2015; Furness et al., 2015), but no males were found in the microcosm that contained both of these outcrossed embryos. One possibility is that a male(s) was present when the embryos were fertilized, but subsequently died and decomposed before the microcosms were sampled. Alternatively, these embryos resulted from outcrossing between a hermaphroditic individual of each strain. If so, these two embryos would represent a hermaphrodite-driven outcrossing rate of ~0.2% given that we genotyped 831 embryos, consistent with the lack of outcrossing reported by Furness et al. (2015) in a sample of 173 embryos.

4.3 Conclusions and perspective

Our findings show that pace of life was associated with emersion tolerance in *K. marmoratus* out of water, which allowed the “slow” Honduras strain to survive out of water

for an average of 12 days (~21%) longer than the relatively “fast” Belize strain. Furthermore, Honduras fish had a lower metabolic rate under control aquatic conditions, indicating genetic divergence between strains that could possibly be the result of genetic assimilation of metabolic plasticity (Pigliucci et al., 2006; Lande, 2009). These results are consistent with other work showing generally “slow” lifestyles in extremophile fishes (Passow et al., 2017). We did not detect an obvious cost to emersion tolerance in our microcosm experiment, consistent with other studies of local adaptation that have often found conditional neutrality of variable traits instead of antagonistic trade-offs (Bono et al., 2017). We also did not push environmental extremes to the limit, as we were interested in investigating only sub-lethal effects, and life history trade-offs are often only revealed under extreme environmental conditions (Lemaître et al., 2015).

Recently, we discovered genetically divergent wild populations of *K. marmoratus* occupying abiotically distinct habitats (high versus low water availability), and the phenotypes of these populations matched those predicted by pace of life theory (Turko et al., 2018). The population that inhabits an ephemeral pond (no water during the dry season) tended to be larger, in better body condition, and had metabolic rates that were ~30% lower than fish from a nearby site with higher water availability. Together with our laboratory data, these findings support the idea that environmental heterogeneity is an important factor that drives differences in metabolic rate and pace of life.

Acknowledgements

We thank Doug Fudge, Todd Gillis, and Graham Scott for helpful discussions regarding experimental design, and Matt Cornish, Mike Davies, and Bianca Cisternino for animal care.

The authors declare that they have no conflicts of interest.

Author contributions

AJT, JED, RLE, and PAW conceived the study. AJT, JED, IY-L, KL, PK, JH, and RLE carried out the experiments, AJT and JED analyzed the data. AJT wrote the draft manuscript, all authors contributed to the manuscript and gave approval for publication.

Funding

Funding was provided by a Natural Sciences and Engineering Research Council of Canada (NSERC) Canada Graduate Scholarship to AJT and an NSERC Discovery grant to PAW.

Data availability

Data will be provided as a supplementary file.

References

- Allen, D., Rosenfeld, J. and Richards, J.** (2016). Physiological basis of metabolic trade-offs between growth and performance among different strains of rainbow trout. *Canadian Journal of Fisheries and Aquatic Science*, **73**, 1493-1506.
- Alvarez, D. and Niecieza, A. G.** (2005). Is metabolic rate a reliable predictor of growth and survival of brown trout (*Salmo trutta*) in the wild? *Canadian Journal of Fisheries and Aquatic Science*, **62**, 643-649.
- AOAC** (1995) Official Methods of Analysis: Official Method for Protein. Method No. 920.87. Association of Official Analytical Chemists, Washington DC.
- Arnqvist, G., Stojković, B., Rönn, J. L. and Immonen, E.** (2017). The pace-of-life: A sex-specific link between metabolic rate and life history in bean beetles. *Functional Ecology*, **31**, 2299-2309.
- Auer, S. L., Dick, C. A., Metcalfe, N. B., and Reznick, D. N.** (2018). Metabolic rate evolves rapidly and in parallel with the pace of life history. *Nature Communications*, **9**, 14.
- Avise, J. C. and Tatarenkov, A.** (2015). Population genetics and evolution of the mangrove rivulus *Kryptolebias marmoratus*, the world's only self-fertilizing hermaphroditic vertebrate. *Journal of Fish Biology*, **87**, 519-538.
- Bergmeyer, H. U., Berndt, E., Schmidt, F., and Stork, H.** (1974). D-Glucose: determination with hexokinase and glucose-6-phosphate dehydrogenase. In *Methods of enzymatic analysis*, 2nd Ed (ed HU Bergmeyer) pp. 1196-1201. New York, NY: Academic Press, Inc.
- Blanchard, T. S., Whitehead, A., Dong, Y. W., and Wright, P. A.** (2019). Phenotypic flexibility in respiratory traits is associated with improved aerial respiration in an amphibious fish out of water. *Journal of Experimental Biology*, **222**, jeb186486.

- Bono, L. M., Smith Jr., L. B., Pfennig, D. W., and Burch, C. L.** (2017). The emergence of performance trade-offs during local adaptation: insights from experimental evolution. *Molecular Ecology*, **26**, 1720-1733.
- Bureau, D. P., Harris, A. M., Bevan, D. J., Simmons, L. A., Azevedo, P. A., and Cho, C. Y.** (2000). Feather meals and meat and bone meals from different origins as protein sources in rainbow trout (*Oncorhynchus mykiss*) diets. *Aquaculture*, **181**, 281-291.
- Burton, T., Killen, S. S., Armstrong, J. D., and Metcalfe, N. B.** (2011). What causes intraspecific variation in resting metabolic rate and what are its ecological consequences? *Proceedings of the Royal Society B*, **278**, 3465-3473.
- Careau, V., and Garland, T.** (2012). Performance, personality, and energetics: correlation, causation, and mechanism. *Physiological and Biochemical Zoology*, **85**, 543-571.
- Cooper, C. A., Litwiller, S. L., Murrant, C. L., and Wright, P. A.** (2012). Cutaneous vasoregulation during short- and long-term aerial acclimation in the amphibious mangrove rivulus, *Kryptolebias marmoratus*. *Comparative Biochemistry and Physiology B*, **161**, 268-274.
- Costa, W. J. E. M.** (2011). Identity of *Rivulus ocellatus* and a new name for a hermaphroditic species of *Kryptolebias* from south-eastern Brazil (Cyprinodontiformes: Rivulidae). *Ichthyological Explorations of Freshwaters*, **22**, 185-192.
- Dejours, P.** (1976). Water versus air as the respiratory media. In *Respiration of Amphibious Vertebrates* (ed. G. M. Hughes), pp. 1-15. London: Academic Press.
- Earley, R. L., and Hsu, Y.** (2008). Reciprocity between endocrine state and contest behavior in the killifish, *Kryptolebias marmoratus*. *Hormones and Behaviour*, **53**, 442-451.
- Earley, R. L., Hanninen, A. F., Fuller, A., Garcia, M. J., & Lee, E. A.** (2012). Phenotypic plasticity and integration in the mangrove rivulus (*Kryptolebias marmoratus*): A prospectus. *Integrative and Comparative Biology*, **52**, 814-827.

- Ellison, A., Cable, J., and Consuegra, S.** (2011). Best of both worlds? Association between outcrossing and parasite loads in a selfing fish. *Evolution*, **65**, 3021-3026.
- Frick, N. T., Bystriansky, J. S., Ip, Y. K., Chew, S. F., and Ballantyne, J. S.** (2008). Lipid, ketone body and oxidative metabolism in the African lungfish, *Protopterus dolloi* following 60 days of fasting and aestivation. *Comparative Biochemistry and Physiology A*, **151**, 93-101.
- Froese, R.** (2006). Cube law, condition factor and weight–length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, **22**, 241-253.
- Furness, A. I., Tatarenkov, A., and Avise, J. C.** (2015). A genetic test for whether pairs of hermaphrodites can cross-fertilize in a selfing killifish. *Journal of Heredity*, **106**, 749-752.
- Furness, A. I., Reznick, D. N., Tatarenkov, A., and Avise, J. C.** (2018) The evolution of diapause in *Rivulus (Laimosemion)*. *Zoological Journal of the Linnean Society*, **184**, 773-790.
- Gibb, A. C., Ashley-Ross, M. A., & Hsieh, S. T.** (2013). Thrash, flip, or jump: The behavioral and functional continuum of terrestrial locomotion in teleost fishes. *Integrative and Comparative Biology*, **53**, 295-306.
- Graham, J. B.** (1997). Air-Breathing Fishes: Evolution, Diversity, and Adaptation. San Diego: Academic Press.
- Griffiths, D., & Kirkwood, R. C.** (1995). Seasonal variation in growth, mortality, and fat stores of roach and perch in Lough Neagh, Northern Ireland. *Journal of Fish Biology*, **47**, 537–554.
- Guppy, M. and Withers, P.** (1999). Metabolic depression in animals: physiological perspectives and biochemical generalizations. *Biological Reviews*, **74**, 1-40.
- Harrington, R. W.** (1961). Oviparous hermaphroditic fish with internal self-fertilization. *Science*, **134**, 1749-1750.

- Heiss, E., Aerts, P., and Van Wassenbergh, S.** (2018). Aquatic–terrestrial transitions of feeding systems in vertebrates: a mechanical perspective. *Journal of Experimental Biology*, **221**, jeb154427.
- Hindle, A. G., Otis, J. P., Epperson, L. E., Hornberger, T. A., Goodman, C. A., Carey, H. V., & Martin, S. L.** (2015). Prioritization of skeletal muscle growth for emergence from hibernation. *Journal of Experimental Biology*, **218**, 276-284.
- Jensen, K., Mayntz, D., Toft, S., Clissold, F. J., Hunt, J., Raubenheimer, D., & Simpson, S. J.** (2012). Optimal foraging for specific nutrients in predatory beetles. *Proceedings of the Royal Society B*, **279**, 2212-2218.
- Jobling, M.** (1994). Fish bioenergetics. London: Chapman and Hall.
- Joint Committee for Guides in Metrology (JCGM)** (2008). Evaluation of measurement data – Guide to the expression of uncertainty in measurement. Retrieved from: www.bipm.org/utls/common/documents/jcgm/JCGM_100_2008_E.pdf
- Junior, R. S. L. and Peixoto, P. E. C.** (2013). Males of the dragonfly *Diastatops obscura* fight according to predictions from game theory models. *Animal Behaviour*, **85**, 663-669.
- Killen, S. S., Marras, S., and McKenzie, D. J.** (2011). Fuel, fasting, fear: routine metabolic rate and food deprivation exert synergistic effects on risk-taking in individual juvenile European sea bass. *Journal of Animal Ecology*, **80**, 1024-1033.
- Koons, D. N., Metcalf, C. J., & Tuljapurkar, S.** (2008). Evolution of delayed reproduction in uncertain environments: a life-history perspective. *American Naturalist*, **172**, 797-805.
- Lande, R.** (2009). Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *Journal of Evolutionary Biology*, **22**, 1435-1446.
- LeBlanc, D. M., Wood, C. M., Fudge, D. S., & Wright, P. A.** (2010). A fish out of water: gill and skin remodeling promotes osmo- and ionoregulation in the mangrove killifish *Kryptolebias marmoratus*. *Physiological and Biochemical Zoology*, **83**, 932-949.

- Lemaître, J.- F., Berger, V., Bonenfant, C., Douhard, M., Gamelon, M., Plard, F., and Gaillard, J.- M.** (2015). Early-late life trade-offs and the evolution of ageing in the wild. *Proceedings of the Royal Society B*, **282**, 20150209.
- Livingston, M. D., Bhargav, V. V., Turko, A. J., Wilson, J. M., & Wright, P. A.** (2018). Widespread use of emersion and cutaneous ammonia excretion in Aplocheiloid killifishes. *Proceedings of the Royal Society B*, 20181496.
- Mackiewicz, M., Tatarenkov, A., Perry, A., Martin, J. R., Elder, J. F., Bechler, D. L., and Avise, J. C.** (2006). Microsatellite documentation of outcrossing between inbred laboratory strains of the self-fertilizing mangrove killifish (*Kryptolebias marmoratus*). *Journal of Heredity*, **97**, 508-513.
- Mathot, K. J., and Dingemanse, N. J.** (2015). Energetics and behavior: Unrequited needs and new directions. *Trends in Ecology and Evolution*, **30**, 199-206.
- Moyes, C. D. and West, T. G.** (1995). Exercise metabolism of fish. In: *Metabolic Biochemistry. Biochemistry and Molecular Biology of Fish*, vol. 4 (eds PW Hochachka, TP Mommsen) Amsterdam, NL: Elsevier Science.
- Mueller, P. and Diamond, J.** (2001). Metabolic rate and environmental productivity: well-provisioned animals evolved to run and idle fast. *Proceedings of the National Academy of Sciences*, **98**, 2550-2554.
- Norin, T. and Malte, H.** (2011). Repeatability of standard metabolic rate, active metabolic rate and aerobic scope in young brown trout during a period of moderate food availability. *Journal of Experimental Biology*, **214**, 1668-1675.
- Ong, K. J., Stevens, E. D., and Wright, P. A.** (2007). Gill morphology of the mangrove killifish (*Kryptolebias marmoratus*) is plastic and changes in response to terrestrial air exposure. *Journal of Experimental Biology*, **210**, 1109-1115.

- Passow, C. N., Arias-Rodriguez, L., and Tobler, M.** (2017). Convergent evolution of reduced energy demands in extremophile fish. *PLoS ONE*, **12**, e0186935.
- Pigliucci, M., Murren, C. J., & Schlichting, C. D.** (2006). Phenotypic plasticity and evolution by genetic assimilation. *Journal of Experimental Biology*, **209**, 2362-2367.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team** (2014). nlme: linear and nonlinear mixed effects models. R package version 3.1-117. Available at <http://CRAN.R-project.org/package=nlme>.
- Pronko, A. J., Perlman, B. M., and Ashley-Ross, M. A.** (2013). Launches, squiggles and pounces, oh my! The water-land transition in mangrove rivulus (*Kryptolebias marmoratus*). *Journal of Experimental Biology*, **216**, 3988-3995.
- Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., and Montiglio, P. O.** (2010). Personality and the emergence of the pace-of-life syndrome concept at the population level. *Philosophical Transactions of the Royal Society B*, **365**, 4051-4063.
- Reid, D., Armstrong, J. D., and Metcalfe, N. B.** (2011). Estimated standard metabolic rate interacts with territory quality and density to determine the growth rates of juvenile Atlantic salmon. *Functional Ecology*, **25**, 1360-1367.
- Ricklefs, R. E. and Wikelski, M.** (2002). The physiology/life-history nexus. *Trends in Ecology and Evolution*, **17**, 462-468.
- Rodela, T. M. and Wright, P. A.** (2006). Metabolic and neuroendocrine effects on diurnal urea excretion in the mangrove killifish *Rivulus marmoratus*. *Journal of Experimental Biology*, **209**, 2704-2712.
- Scarsella, G. E., Gresham, J. D., and Earley, R. L.** (2018). Relationships between external sexually dimorphic characteristics and internal gonadal morphology in a sex changing fish. *Journal of Zoology*, **305**, 133-140.

- Schultz, E. T. and Conover, D. O.** (1997). Latitudinal differences in somatic energy storage: adaptive responses to seasonality in an estuarine fish (Atherinidae: *Menidia menidia*). *Oecologia*, **109**, 516-529.
- Seutin, G., White, B. N., and Boag, P. T.** (1991). Preservation of avian blood and tissue samples for DNA analyses. *Canadian Journal of Zoology*, **69**, 82-90.
- Stearns, S. C.** (1992). *The Evolution of Life Histories*. London: Oxford University Press.
- Sutton, A. O., Turko, A. J., McLaughlin, R. L., and Wright, P. A.** (2018). Behavioral and physiological responses of an amphibious, euryhaline mangrove fish to acute salinity exposure. *Copeia*, **106**, 305-311.
- Tatarenkov, A., Ring, B. C., Elder, J. F., Bechler, D. L., and Avise, J. C.** (2010). Genetic composition of laboratory stocks of the self-fertilizing fish *Kryptolebias marmoratus*: a valuable resource for experimental research. *PloS One*, **5**, e12863.
- Taylor, D. S., Turner, B. J., Davis, W. P., and Chapman, B. B.** (2008). A novel terrestrial fish habitat inside emergent logs. *American Naturalist*, **171**, 263-266.
- Turko, A. J., and Wright, P. A.** (2015). Evolution, ecology and physiology of amphibious killifishes (Cyprinodontiformes). *Journal of Fish Biology*, **87**, 815-835.
- Turko, A. J., Earley, R. L., and Wright, P. A.** (2011). Behaviour drives morphology: voluntary emersion patterns shape gill structure in genetically identical mangrove rivulus. *Animal Behaviour*, **82**, 39-47.
- Turko, A. J., Robertson, C. E., Bianchini, K., Freeman, M., and Wright, P. A.** (2014). The amphibious fish *Kryptolebias marmoratus* uses different strategies to maintain oxygen delivery during aquatic hypoxia and air exposure. *Journal of Experimental Biology*, **217**, 3988-3995.

- Turko, A. J., Kültz, D., Fudge, D., Croll, R. P., Smith, F. M., Stoyek, M. R., & Wright, P. A.** (2017). Skeletal stiffening in an amphibious fish out of water is a response to increased body weight. *Journal of Experimental Biology*, **220**, 3621-3631.
- Turko, A. J., Tatarenkov, A., Currie, S., Earley, R. L., Platek, A., Taylor, D. S., and Wright, P. A.** (2018). Emersion behaviour underlies variation in aquatic respiratory function in the amphibious fish *Kryptolebias marmoratus*. *Journal of Experimental Biology*, **221**, jeb.168039.
- Wells, M. W., Turko, A. J., and Wright, P. A.** (2015). Fish embryos on land: terrestrial embryo deposition enhances development in the amphibious fish *Kryptolebias marmoratus*. *Journal of Experimental Biology*, **218**, 3249-3256.
- Wright, P. A. and Turko A. J.** (2016). Amphibious fishes: Evolution and phenotypic plasticity *Journal of Experimental Biology*, **219**, 2245-2259.

Figures




Figure 1. Survival of different strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* out of water. Different letters represent significant differences between strains ($p < 0.05$).




Figure 2. Rates of O₂ consumption in different strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* in water and in air. O₂ consumption was measured (A) daily over 7 d of terrestrial acclimation, and (B) after 21 d out of water. Asterisks denote a significant overall difference between the Honduran strain versus the other two strains ($p < 0.05$), and daggers indicate significant overall differences in metabolic rate compared to the value in water ($p < 0.05$). Data are presented as means \pm s.e.m.




Figure 3. Fulton's condition factor in different strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* in water under normal conditions (control) and after 21 d out of water (terrestrial). For each boxplot, the bold horizontal line in the middle of each box represents the median, the top and bottom of the box represent the quartiles (i.e. 25th and 75th percentiles), whiskers show the highest and lowest values within 1.5x the interquartile range, and points show values beyond the range of the whiskers. Asterisks represent significant differences within a strain ($p < 0.05$).




Figure 4. Fuel use after 21 d of terrestrial acclimation in 3 strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus*. (A) lipid use, (B) glycogen use, (C) protein use, and (D) total energy consumed. Data are presented relative to wet mass. Different letters represent significant differences ($p < 0.05$) between strains. Data are presented as means \pm s.e.m.




Figure 5. Reproductive measures of 3 strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* over the first 18 months of life. (A) Total embryo production, (B) mean clutch size, and (C) age at first reproduction. For each boxplot, the bold horizontal line in the middle of each box represents the median, the top and bottom of the box represent the quartiles (i.e. 25th and 75th percentiles), whiskers show the highest and lowest values within 1.5x the interquartile range, and points show values beyond the range of the whiskers. Different letters represent significant differences ($p < 0.05$) between strains.




Figure 6. Number, body size and condition of *Kryptolebias marmoratus* after 12 months in fully aquatic (control) or periodically drained (fluctuating) microcosms. (A) Number of adult fish collected, (B) standard length, (C) wet mass, (D) Fulton's condition factor, (E) gonado-somatic index (GSI), and (F) hepato-somatic index (HSI). The dashed line in (A) indicates the number of fish of each strain initially placed in each microcosm. For each boxplot, the bold horizontal line in the middle of each box represents the median, the top and bottom of the box represent the quartiles (i.e. 25th and 75th percentiles), whiskers show the highest and lowest values within 1.5x the interquartile range, and points show values beyond the range of the whiskers. Different letters represent a significant overall difference between treatment conditions, and asterisks denote a significant overall difference between strains ($p < 0.05$).




Figure 7. Number of embryos recovered from microcosms after 12 months. More embryos were released by the Belize versus Honduras strain under fluctuating conditions ($p < 0.05$).

Table S1. Body composition data used to calculate energy use in three distinct genetic lineages of *Kryptolebias marmoratus* after 21 d of terrestrial acclimation.

lineage	variable	treatment	mean	SD	n
50.91 (Belize)	glycogen (mg g ⁻¹ dry mass)	control	61.687	21.35	9
		terrestrial	19.445	5.64	6
	lipid (mg g ⁻¹ dry mass)	control	109.711	25.84	6
		terrestrial	49.481	28.55	11
	protein (mg g ⁻¹ dry mass)	control	598.168	19.98	10
		terrestrial	602.463	27.05	9
	body water (proportion)	control	0.762	0.009	6
		terrestrial	0.785	0.013	12
	Δ body mass (final initial ⁻¹)	n/a	0.819	0.09	10
SLC8E (Florida)	glycogen (mg g ⁻¹ dry mass)	control	74.838	36.45	8
		terrestrial	17.315	11.25	9
	lipid (mg g ⁻¹ dry mass)	control	129.411	18.36	6
		terrestrial	89.090	26.84	12
	protein (mg g ⁻¹ dry mass)	control	598.132	18.68	6
		terrestrial	626.655	26.45	7
	body water (proportion)	control	0.751	0.011	6
		terrestrial	0.779	0.010	12
	Δ body mass (final initial ⁻¹)	n/a	0.765	0.12	10
HON11 (Honduras)	glycogen (mg g ⁻¹ dry mass)	control	60.320	21.30	9
		terrestrial	14.627	2.10	9
	lipid (mg g ⁻¹ dry mass)	control	100.960	17.49	6
		terrestrial	46.069	14.04	12
	protein (mg g ⁻¹ dry mass)	control	605.562	19.01	10
		terrestrial	607.166	21.29	10
	body water (proportion)	control	0.771	0.005	6
		terrestrial	0.792	0.004	12
	Δ body mass (final initial ⁻¹)	n/a	0.908	0.05	10




Figure S1. Size at age for two strains of *Kryptolebias marmoratus*. (A) Body mass and (B) standard length. Honduras fish were significantly heavier ($p < 0.0001$) and longer ($p < 0.0001$) than Belize fish at a given age.




Figure S2. Change in body mass of 3 strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* after 21 d terrestrial acclimation. The bold horizontal line in the middle of each box represents the median, the top and bottom of the box represent the quartiles (i.e. 25th and 75th percentiles), whiskers show the highest and lowest values within 1.5x the interquartile range, and points show values beyond the range of the whiskers. Different letters represent significant differences ($p < 0.05$) among strains.




Figure S3. Energy reserves and water content in 3 strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* under normal housing conditions (control) and after 21 d terrestrial acclimation. (A) lipids, (B) glycogen, (C) crude protein, and (D) water content. All values are relative to wet mass. For each boxplot, the bold horizontal line in the middle of each box represents the median, the top and bottom of the box represent the quartiles (i.e. 25th and 75th percentiles), whiskers show the highest and lowest values within 1.5x the interquartile range, and points show values beyond the range of the whiskers. Asterisks denote significant differences ($p < 0.05$) after terrestrial acclimation, and different letters indicate significant overall differences among strains ($p < 0.05$).




Figure S4. Time spent moving (%) in different strains (Belize, Florida, Honduras) of *Kryptolebias marmoratus* over 21 d of terrestrial acclimation. Different letters indicate significant differences among strains at the 1 h time point ($p < 0.05$). Within the Honduras strain, activity at 1 h is significantly higher than at the 7 d and 14 d timepoints ($p < 0.05$). Within the Florida strain, activity at 1 h is significantly higher than all other timepoints ($p < 0.05$). There were no differences in the Belize strain over time ($p > 0.05$). Data are presented as means \pm s.e.m.




Figure S5. Body and organ mass of Belize and Honduras strains of *Kryptolebias marmoratus* after 12 months in fully aquatic (control) or periodically drained (fluctuating) microcosms. (A) Body mass, (B) gonad mass, and (C) liver mass as a function of body length. Body mass was significantly greater in Honduras than Belize fish ($p < 0.05$), and under control versus fluctuating conditions ($p < 0.05$, interaction $p > 0.05$). Gonad mass was higher in Honduras than Belize fish ($p < 0.05$), and not affected by treatment ($p > 0.05$). Liver mass was not affected by strain or treatment (both $p > 0.05$).




Figure S6. Total length of gill filaments in Belize and Honduras strains of *Kryptolebias marmoratus* after 12 months in fully aquatic (control) or periodically drained (fluctuating) microcosms. There were no significant effects of strain or environment ($p > 0.05$).